

Ehrenfest breakdown of the mean-field dynamics of Bose gases

Xizhi Han (韩希之)¹ and Biao Wu (吴飙)^{1,2,3,*}

¹*International Center for Quantum Materials, School of Physics, Peking University, Beijing 100871, China*

²*Collaborative Innovation Center of Quantum Matter, Beijing 100871, China*

³*Wilczek Quantum Center, College of Science, Zhejiang University of Technology, Hangzhou 310014, China*

(Received 20 June 2015; published 12 February 2016)

The unstable mean-field dynamics of a Bose gas is shown to break down at time $\tau_h = (c_1/\gamma) \ln N$, where γ is the Lyapunov exponent of the mean-field theory, N is the number of bosons, and c_1 is a system-dependent constant. The breakdown time τ_h is essentially the Ehrenfest time that characterizes the breakdown of the correspondence between classical and quantum dynamics. This breakdown can be well described by a quantum fidelity defined for one-particle reduced density matrices. Our results are obtained with the formalism in particle-number phase space and are illustrated with a triple-well model. The logarithmic quantum-classical correspondence time may be verified experimentally with Bose-Einstein condensates.

DOI: [10.1103/PhysRevA.93.023621](https://doi.org/10.1103/PhysRevA.93.023621)

I. INTRODUCTION

The nonlinear Gross-Pitaevskii equation (GPE), as a mean-field theory, has been the dominant tool in describing the dynamics of Bose-Einstein condensates (BECs) in ultracold atomic gases [1,2]. However, we face a quandary when the mean-field dynamics of a BEC becomes dynamically unstable or chaotic [3–9]: On the one hand, one may regard this instability as an unphysical artifact resulting from the mean-field approximation, since the exact dynamics of a BEC is governed by the many-body Schrödinger equation, which is linear and thus does not allow chaos; on the other hand, the dynamical instability was observed in experiments [10–15] and it has been proved with mathematical rigor that the GPE describes correctly not only the ground state but also the dynamics of a BEC in the large- N limit (N is the number of bosons) [16,17].

Our aim in this work is to resolve this fundamental dilemma. Our study shows that the mean-field theory (the GPE) is only valid up to time

$$\tau_h = \frac{c_1}{\gamma} \ln N + o(\ln N), \quad (1)$$

where $\gamma > 0$ is the Lyapunov exponent of the mean-field dynamics and c_1 is a constant that depends only on systems. With this time scale, the dilemma is resolved: On the one hand, in the large- N limit ($N \rightarrow \infty$), τ_h goes to infinity and thus the GPE is always valid just as proved rigorously in Ref. [17]; on the other hand, the time τ_h increases with N only logarithmically and it is not a long time for a typical BEC experiment. For example, for the system studied in Ref. [3], the Lyapunov time $\tau_\gamma = 1/\gamma \sim 1$ ms. As the number of atoms in a BEC prepared in a typical experiment is around 10^4 , we have $\tau_h \sim 10$ ms. As a result, the dynamical instability or the breakdown of the mean-field dynamics can be easily observed in a typical experiment as reported in Ref. [12].

This time scale τ_h is essentially the Ehrenfest time, which is the time that the correspondence between classical and quantum dynamics breaks down [18,19]. The usual Ehrenfest

time $\tau_{\text{Eh}} = (c_1/\gamma) \ln(A/\hbar)$, where γ is the Lyapunov exponent of the classical motion and A is a typical action [19]. The similarity is due to the fact that the GPE can be regarded as a classical equation in the large- N limit [20]. Therefore, our result paves the way to experimental investigation of a fundamental relation in the quantum-classical correspondence, the logarithmic behavior of the Ehrenfest time, as N can be varied in experiments.

We cast the quantum dynamics onto the particle-number phase space (PNPS), which is a rearrangement of Fock states. In this phase space, for a nearly coherent state and in the large- N limit, the quantum many-body dynamics is equivalent to an ensemble of mean-field dynamics. When the mean-field motion is regular, mean-field trajectories will stay together and the Bose gas remains coherent. If the mean-field motion is unstable or chaotic, mean-field trajectories will soon separate from each other exponentially, leading to decoherence of the Bose gas and breakdown of the mean-field theory. Thus, there are two distinct types of quantum dynamics, whose difference can be characterized by the quantum fidelity for one-particle reduced density matrices (RDMs).

We investigate the Ehrenfest breakdown numerically in the system of a BEC in a triple-well potential [21–25], which may be the simplest BEC model that embraces chaotic mean-field dynamics. With this model, we verify numerically the Ehrenfest time and show that our quantum fidelity can well capture the characteristics of two different types of quantum dynamics.

The mean-field instability or breakdown has been discussed in the literature [3,5–7,26–31]. However, a general and explicit relation between mean-field chaos, the number of particles, and breakdown time is still lacking. Further, in PNPS not only can such a breakdown be understood intuitively and quantitatively, but the significance of a local phase structure is also apparent, the distortion of which leads to decoherence.

II. PARTICLE-NUMBER PHASE SPACE

In Ref. [20] it is shown that many quantum systems become classical in the large- N limit. A dilute Bose gas belongs to this class of quantum systems: Its dynamics becomes classical and it is well described by the mean-field GPE in the large- N limit. In this section we introduce PNPS, where

*wubiao@pku.edu.cn

this quantum-classical correspondence in the large- N limit becomes transparent.

A. Definition

Any quantum state $|\Psi\rangle$ of a system of N identical bosons with M single-particle states can be regarded as a wave function $\varphi(\mathbf{x})$ over an $(M-1)$ -dimensional lattice space, which we call particle-number phase space, via

$$\varphi(\mathbf{x}) \equiv \langle 0 | \prod_{i=1}^M \frac{\hat{a}_i^{Nx_i}}{\sqrt{(Nx_i)!}} |\Psi\rangle, \quad (2)$$

where the x_i are entries of the M -dimensional vector \mathbf{x} , $Nx_i \in \{0, \dots, N\}$ for $1 \leq i \leq M$, and $\sum_{i=1}^M x_i = 1$. The \hat{a}_i^\dagger and \hat{a}_i are the creation and annihilation operators for the i th single-particle state, with $[\hat{a}_i, \hat{a}_j^\dagger] = \delta_{ij}$ and $\hat{n}_i \equiv \hat{a}_i^\dagger \hat{a}_i$. The continuous limit of PNPS is a hyperplane in $[0, 1]^M$ (defined by the constraint $\sum_{i=1}^M x_i = 1$), where we can define (for i from 1 to M)

$$\langle x_i \rangle \equiv \int d\mathbf{x} x_i |\varphi(\mathbf{x})|^2, \quad (3)$$

$$\langle (\Delta x_i)^2 \rangle \equiv \int d\mathbf{x} (x_i - \langle x_i \rangle)^2 |\varphi(\mathbf{x})|^2 \quad (4)$$

to characterize the average position and spread of the distribution $|\varphi(\mathbf{x})|^2$ over PNPS, given $|\Psi\rangle$ normalized. Of course for any finite N the integral should be interpreted as a summation over all \mathbf{x} in PNPS.

As an example of our particular interest, we examine an $SU(M)$ coherent state $|\Psi\rangle_c$ in PNPS:

$$|\Psi\rangle_c \equiv \frac{1}{\sqrt{N!}} \left(\sum_{i=1}^M \psi_i a_i^\dagger \right)^N |0\rangle, \quad (5)$$

where $\sum_i |\psi_i|^2 = 1$. In such a case we say that $|\psi\rangle$ (an M -dimensional vector with ψ_i as its entries) is the mean-field state of the $SU(M)$ coherent state $|\Psi\rangle_c$. It is straightforward to show for this coherent state $|\Psi\rangle_c$ that

$$\langle x_i \rangle = |\psi_i|^2, \quad \langle (\Delta x_i)^2 \rangle = |\psi_i|^2 (1 - |\psi_i|^2) / N, \quad (6)$$

which indicates that the coherent state $|\Psi\rangle_c$ corresponds to a localized distribution $|\varphi(\mathbf{x})|^2$ in PNPS that peaks around $(|\psi_1|^2, |\psi_2|^2, \dots, |\psi_M|^2)$ with a vanishing spread at large N .

The wave function $\varphi(\mathbf{x})$ in PNPS has a phase structure. For any \mathbf{x} and \mathbf{y} in PNPS,

$$\arg \varphi(\mathbf{x}) - \arg \varphi(\mathbf{y}) = N \sum_{i=1}^M (x_i - y_i) \arg \psi_i \pmod{2\pi}, \quad (7)$$

which shows a wave vector \mathbf{k} : $k_i = N \arg \psi_i \propto N$. This phase structure is important as it will give us an estimate of the time τ_h in our later discussion. It is worth noting that when $N \rightarrow \infty$, there is no limit of the wave function $\varphi(\mathbf{x})$ because its wave vector \mathbf{k} diverges.

Overall, we find that the coherent state corresponds to a single-peaked wave packet with plane-wave phase structure in PNPS. In the following, we discuss quantum dynamics in

PNPS and its relation to the mean-field dynamics. Note that similar formalisms were also used in other contexts [32,33].

B. Dynamics

Consider the following Hamiltonian of a Bose gas

$$\hat{\mathcal{H}} = \sum_{i,j=1}^M \left\{ H_{ij}^0 \hat{a}_i^\dagger \hat{a}_j + \frac{U_{ij}}{N} \hat{a}_i^\dagger \hat{a}_j^\dagger \hat{a}_j \hat{a}_i \right\}, \quad (8)$$

where $H_{ij}^0 = H_{ji}^{0*}$, $U_{ij} = U_{ij}^*$ and $U_{ij} = U_{ji}$. Corresponding to the Schrödinger equation $i\partial_t |\Psi\rangle = \hat{\mathcal{H}} |\Psi\rangle$, there is an equation of motion (EOM) for $\varphi(\mathbf{x}; t)$ in PNPS [Eq. (A1) in the Appendix]. We are especially interested in the dynamics of a nearly coherent state $\varphi(\mathbf{x}; t)$, which is prescribed as the following two conditions.

(i) The distribution $|\varphi(\mathbf{x}; t)|^2$ is localized such that $1/N \ll \sqrt{\langle (\Delta x_i)^2 \rangle} \ll 1$ for all $i = 1, 2, \dots, M$.

(ii) A local wave vector $\mathbf{k}(\mathbf{x}; t) \equiv \nabla_{\mathbf{x}} \arg \varphi(\mathbf{x}; t)$ exists in PNPS and varies insignificantly over a scale of $1/N$, i.e., $|\partial_{x_i} k_j| \ll N$ for all $i, j = 1, 2, \dots, M$.

With these two conditions and keeping only finite terms in the large- N limit, an approximate [to $O(1)$] EOM for $\varphi(\mathbf{x}; t)$ in PNPS can be derived [see Eq. (A2)]. Mathematically, there are δ -function solutions to this EOM [Eq. (A2)]:

$$\varphi(\mathbf{x}; t) = \exp[i\alpha(t)] \prod_{i=1}^M \delta(x_i - x_i^0(t)) \exp[ik_i^0(t)x_i]. \quad (9)$$

In these δ -function solutions, $x_i^0(t)$ and $k_i^0(t)$ satisfy the equation

$$i\partial_t \rho = [\hat{\mathcal{H}}_{\text{MF}}, \rho], \quad (10)$$

where $\mathcal{H}_{\text{MF},ij}(t) \equiv H_{ij}^0 + 2U_{ij}\rho_{ij}(t)$ and

$$\rho_{ij}(t) \equiv \sqrt{x_i^0(t)x_j^0(t)} e^{i[k_i^0(t) - k_j^0(t)]/N}. \quad (11)$$

This is just the mean-field EOM for the one-particle RDM.

Conditions (i) and (ii) reflect our expectations of nearly coherent states [see Eqs. (6) and (7)]. The existence of δ -function solutions corresponds to the established result that for any time t_0 , when $N \rightarrow \infty$, coherent states at $t = 0$ stay coherent when $t = t_0$ [17].

The results above can be interpreted as follows: At large N , for any initial state satisfying the two conditions, its time evolution may be regarded as the superposition of mean-field dynamics of δ functions, since any function in PNPS can be decomposed into a superposition of a cloud of δ functions. This is similar to the quantum dynamics of a single-particle wave packet in real space: It can be regarded as a cloud of classical particles and each of them follows Newton's EOM.

As the quantum-classical correspondence between a quantum wave packet and a classical particle will break down at the Ehrenfest time, the correspondence between one state in PNPS and its mean-field description, one δ -function solution [see Eq. (9)], will also fail when the mean-field trajectories of the δ functions in the cloud diverge. The breakdown time τ_h can be estimated using a conventional strategy in quantum

chaos as in Ref. [19]. Essentially, before the breakdown the wave packet of nearly coherent states in PNPS expands in the form of $\exp \gamma t$, where γ is the Lyapunov exponent of the mean-field dynamics. According to Eq. (6), for $t < \tau_h$,

$$\Delta(t) \equiv \sqrt{\frac{1}{M} \sum_{i=1}^M \langle (\Delta x_i)^2 \rangle (t)} \propto \frac{e^{\gamma t}}{\sqrt{N}}. \quad (12)$$

In addition, there is a consistent mean-field description only if local wave vectors across the wave packet are almost equal, that is,

$$\kappa t N^{-\lambda} \Delta(t) \ll 1, \quad (13)$$

where $\kappa N^{-\lambda}$ is the average rate of growth of curvature $\partial_{x_i} k_j$ and the N dependence is written explicitly. Substituting (12) into (13), we have

$$\gamma t + \ln t + \ln \kappa \ll (\lambda + \frac{1}{2}) \ln N. \quad (14)$$

The Ehrenfest time τ_h in Eq. (1) is obtained with $c_1 = \lambda + \frac{1}{2}$, which is independent of N or γ . Numerical verification of this relation will be presented later.

Note that it is well known that the quantum-classical correspondence may last far beyond the Ehrenfest time (see, e.g., Ref. [34]). Similarly, it is possible that the mean-field theory remains valid even after our first estimate $t = \tau_h$; this interesting and special topic will be left for future study.

III. EXAMPLE OF THE TRIPLE-WELL MODEL

We now illustrate our results with an example. Consider a BEC in a ring-shaped triple-well potential [25]. Under tight-binding approximation, the second-quantized Hamiltonian is [as a specific case of Eq. (8)]

$$\hat{\mathcal{H}} = -\frac{1}{2} \sum_{1 \leq i, j \leq 3}^{i \neq j} \hat{a}_i^\dagger \hat{a}_j + \frac{c}{2N} \sum_{i=1}^3 \hat{a}_i^\dagger \hat{a}_i^\dagger \hat{a}_i \hat{a}_i, \quad (15)$$

where c is the on-site interaction strength. For this system $M = 3$. Its corresponding nonlinear mean-field EOM is

$$i \frac{d}{dt} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} = \begin{pmatrix} c|\psi_1|^2 & -1/2 & -1/2 \\ -1/2 & c|\psi_2|^2 & -1/2 \\ -1/2 & -1/2 & c|\psi_3|^2 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix}. \quad (16)$$

Shown in Fig. 1 is a Poincaré section of the above mean-field dynamics, where two kinds of motion are evident: The central regular region is surrounded by a chaotic sea. The conjugate variables used in plotting Fig. 1 are (J_1, θ_1) and (J_2, θ_2) , which are defined as $J_1 = |\psi_1|^2 - |\psi_3|^2$, $J_2 = |\psi_3|^2$, $\theta_1 = \arg \psi_2 - \arg \psi_1$, and $\theta_2 = 2 \arg \psi_2 - \arg \psi_1 - \arg \psi_3$.

The quantum dynamics of this model can also be computed rather easily. The evolution of $|\varphi(x)|^2$ in PNPS is plotted in Fig. 2, where two types of quantum dynamics are clearly observed. In Figs. 2(a) and 2(b) an initial coherent state, which is a Gaussian-like wave packet in PNPS, shows no significant expansion or distortion during dynamical evolution. In Figs. 2(c) and 2(d) the situation is drastically different: A similar-looking initial coherent state expands and becomes dramatically distorted after a certain time. The difference is




FIG. 1. Poincaré section of the classical (mean-field) triple-well Hamiltonian with conjugate variables (J_1, θ_1) and (J_2, θ_2) at $\theta_2 = 0$, $\theta_2 < 0$, $c = 1.25$, and $E \approx 0.708$. The + represents a state in the central regular region and the * represents a state in the chaotic sea.

caused by the fact that the initial state in Fig. 2(a) corresponds to a mean-field state in the regular region in Fig. 1 while the one in Fig. 2(c) corresponds to a mean-field state in the chaotic region.

It is obvious that the mean-field theory cannot describe the dramatic quantum dynamics shown in Figs. 2(c) and 2(d). Such a failure or breakdown of the mean-field theory due to rapid decoherence has long been noticed in the literature [28–31]. In Ref. [30] a remedy was tried unsuccessfully to bridge the gap between the mean-field theory and the exact quantum theory. In




FIG. 2. Plot of $|\varphi(x_1, x_2, x_3; t)|^2$ for the quantum triple-well model with $N = 40$. The two axes are $x_1 \in [0, 1]$ and $x_2 \in [0, 1]$ ($x_3 = 1 - x_1 - x_2$). Red regions are of larger $|\varphi|^2$. (a) Initial state corresponding to the mean-field state denoted by + in Fig. 1, (b) the + state after evolving dynamically for $t = 14.5$, (c) initial state corresponding to the * state in Fig. 1, and (d) the * state at $t = 14.5$.

this work we have shown that there exists a general time scale τ_h in terms of the Lyapunov exponent and the number of bosons beyond which the mean-field theory fails. In the following we introduce a quantum fidelity to distinguish the two types of quantum dynamics shown in Fig. 2 without using mean-field formalism and confirm the time scale τ_h numerically.

A. Quantum fidelity

To quantify the loss of coherence in the quantum evolution as shown in Fig. 2(d), we introduce the following quantum fidelity \mathcal{F}_q for the one-particle RDM $\hat{\rho}$ and $\hat{\chi}$:

$$\mathcal{F}_q(\hat{\rho}, \hat{\chi}) \equiv \frac{1}{N^2} \text{tr} \hat{\rho}^\dagger \hat{\chi}. \quad (17)$$

For a quantum state $|\Psi(t)\rangle$, its one-particle RDM can be explicitly written as

$$\sum_{i,j} |i\rangle \langle \Psi(t) | \hat{a}_i^\dagger \hat{a}_j | \Psi(t) \rangle \langle j|. \quad (18)$$

There are three reasons to use this quantum fidelity.

(a) Experimentally, we are often interested in the one-particle RDM.

(b) It allows us to define coherence \mathcal{C} :

$$\mathcal{C}(\hat{\rho}) \equiv \mathcal{F}_q(\hat{\rho}, \hat{\rho}), \quad (19)$$

where $\hat{\rho}$ is the one-particle RDM for $|\Psi\rangle$. The coherence \mathcal{C} can quantify how coherent the state $|\Psi\rangle$ is, as $\mathcal{C}(\hat{\rho}) = 1$ if and only if $|\Psi\rangle$ is a coherent state as in Eq. (5).

(c) It returns to the mean-field fidelity for coherent states, i.e., $\mathcal{F}_q(\hat{\rho}, \hat{\chi}) = \mathcal{F}_{\text{MF}}(\psi, \phi) \equiv |\langle \phi | \psi \rangle|^2$ if $\hat{\rho}$ and $\hat{\chi}$ are one-particle RDMs for coherent states $|\Psi\rangle_c$ and $|\Phi\rangle_c$, and ψ and ϕ are mean-field states of $|\Psi\rangle_c$ and $|\Phi\rangle_c$ [see the discussion under Eq. (5)]. Therefore, before the Ehrenfest breakdown \mathcal{F}_q essentially captures mean-field characteristics, especially the Lyapunov exponent, which distinguishes regular and chaotic mean-field trajectories.

B. Numerical results

The numerical simulation aims at verifying our theoretical understanding as discussed: For a coherent initial state, at the beginning the mean-field dynamics agrees with the quantum evolution, producing even the same growth of discrepancy between states; however, long-time exponential growth is not allowed by quantum mechanics, so there exists an Ehrenfest time τ_h beyond which the mean-field and quantum correspondence fails. Such a failure is due to the decoherence of quantum states; the breakdown time τ_h is given in Eq. (1).

We choose a coherent initial state $|\Psi(t=0)\rangle_c$ with the one-particle RDM $\hat{\rho}(t=0)$, whose corresponding mean-field state is $|\psi(t=0)\rangle$. Then we slightly perturb the mean-field state into $|\tilde{\psi}(t=0)\rangle$ and generate the corresponding coherent state $|\tilde{\Psi}(t=0)\rangle_c$ and RDM $\hat{\rho}(t=0)$. Next we observe the evolution of quantum fidelity between these two states, which allows us to calculate the Lyapunov exponent. Of course, $|\psi(t)\rangle$ and $|\tilde{\psi}(t)\rangle$ evolve according to the mean-field




FIG. 3. (a) Quantum and mean-field fidelities. Solid lines are $1 - \mathcal{F}_{\text{MF}}(\psi(t), \tilde{\psi}(t))$, dashed lines are $1 - \mathcal{F}_q(\hat{\rho}(t), \hat{\rho}(t)) / \sqrt{\mathcal{F}_q(\hat{\rho}(t), \hat{\rho}(t)) \mathcal{F}_q(\hat{\rho}(t), \hat{\rho}(t))}$ for $N = 80$, and dotted lines are for $N = 40$. The \mathcal{F}_q is normalized to better show the correspondence. (b) Coherence $\mathcal{C}(\hat{\rho}(t))$. Curves show the decay of coherence of quantum many-body states in (a). In the simulation, $c = 1.25$, $E \approx 0.708$, $\theta_1 = \pi$, $\theta_2 = 0$, and $\|\psi - \tilde{\psi}\|_{t=0} \approx 10^{-4}$. The lower set of lines in (a) and the corresponding upper set of lines in (b) are for the integrable case $J_1 = 0$; the upper set in (a) and the corresponding lower set in (b) are for the chaotic case $J_1 = 0.5$.

equations (16), $|\Psi(t)\rangle$ and $|\tilde{\Psi}(t)\rangle$ evolve according to the quantum Hamiltonian in Eq. (8), and $\hat{\rho}(t)$ and $\hat{\rho}(t)$ are obtained from $|\Psi(t)\rangle$ and $|\tilde{\Psi}(t)\rangle$, respectively. The $1 - \mathcal{F}_q(\hat{\rho}(t), \hat{\rho}(t))$ and $1 - \mathcal{F}_{\text{MF}}(\psi(t), \tilde{\psi}(t))$ are shown in Fig. 3(a), where we see that the mean-field fidelity \mathcal{F}_{MF} coincides with \mathcal{F}_q for small t , as expected.

However, we also observe in Fig. 3(a) that there is an Ehrenfest time τ_h when \mathcal{F}_q and \mathcal{F}_{MF} start to visibly disagree. Cases for different N and γ are plotted in Fig. 3(a), where we can see that as N increases or γ decreases, τ_h gets longer. This qualitatively agrees with the scaling of the Ehrenfest time. In Fig. 3(b) it is observed that although τ_h is different for different N and γ , τ_h is approximately the time when the coherence $\mathcal{C}(\hat{\rho}(t))$ drops below 98%. This confirms our understanding that the failure of correspondence between the mean-field and quantum descriptions is the result of decoherence of quantum states.




FIG. 4. (a) Time evolution of coherence $C(\hat{\rho}(t))$ for the integrable $J_1 = 0$ and the chaotic $J_1 = 0.6$ trajectories. Here $c = 1.25$, $E \approx 0.708$, $\theta_1 = \pi$, $\theta_2 = 0$, and the time step in the simulation is 10^{-3} . The C of integrable $J_1 = 0$ (the solid line in the top) remains high while the C of chaotic $J_1 = 0.6$ (other solid lines) drops quickly ($N = 20, 30, \dots, 80$ from left to right). The dashed line is $C = 98\%$. (b) Linear fitting of $\ln \Delta(t)/\Delta(0) + \ln t + \ln \kappa = c_1 \ln N$ for the chaotic initial state in (a) with $c = 1, 1.25$, and 2 and N from 20 to 80 . Data points are calculated when C drops to 98% . The slope $c_1 \approx 0.6$ is found to be independent of c or γ .

Based on such an understanding, we can quantitatively define the Ehrenfest time in this example as the time when the coherence $C(\hat{\rho}(t))$ drops below 98% . Examples of decay of $C(\hat{\rho}(t))$ are illustrated in Fig. 4(a), where the Ehrenfest time τ_h is measured when $C(\hat{\rho}(t))$ drops below the dashed line. By varying N and c , we verify the relation (14) [with \ll replaced by $=$], which leads to Eq. (1), in Fig. 4(b). A linear fitting between $\ln \Delta(t)/\Delta(0) + \ln t$ and $\ln N$ is found with a constant slope $\lambda + \frac{1}{2} \approx 0.6$ [see Eq. (14)], suggesting $c_1 \approx 0.6$ in Eq. (1). Note that here γt is replaced by $\ln \Delta(t)/\Delta(0)$ for numerical convenience.

IV. CONCLUSION

To summarize, we have answered an intriguing question: When does the mean-field approximation of a dilute Bose gas remain valid as the system evolves? Our answer is that for unstable trajectories the mean-field dynamics breaks down at

the Ehrenfest time $\tau_h = (c_1/\gamma) \ln N$. The study was facilitated by introducing particle-number phase space, where one can see easily that the correspondence between many-body quantum dynamics and mean-field dynamics is similar to the usual quantum-classical correspondence.

As N can be varied in BEC experiments, it is now possible to experimentally measure the logarithmic behavior of the Ehrenfest time. One can compare physical observables in the experiment with their theoretical mean-field values and measure the Ehrenfest time when their discrepancy exceeds a threshold. Bose-Einstein condensates with unstable or chaotic mean-field descriptions are suitable for such experiments; for example, spinor BECs [35,36] may be a good candidate system.

ACKNOWLEDGMENTS

This work was supported by the National Basic Research Program of China (Grants No. 2013CB921903 and No. 2012CB921300) and the National Natural Science Foundation of China (Grants No. 11274024, No. 11334001, and No. 11429402).

APPENDIX: QUANTUM EOM IN PNPS AND ITS MEAN-FIELD APPROXIMATION

For the Hamiltonian in Eq. (8), the Schrödinger equation in PNPS reads ($\hbar = 1$)

$$\begin{aligned}
 i \partial_t \varphi(\mathbf{x}; t) = & \sum_i H_{ii}^0 N x_i \varphi(\mathbf{x}; t) + \sum_i U_{ii} N x_i (x_i - \epsilon) \varphi(\mathbf{x}; t) \\
 & + \sum_{i \neq j} H_{ij}^0 N \sqrt{x_i (x_j + \epsilon)} \varphi(\mathbf{x} + \epsilon \mathbf{e}^{ij}; t) \\
 & + \sum_{i \neq j} U_{ij} N x_i x_j \varphi(\mathbf{x}; t), \quad (\text{A1})
 \end{aligned}$$

where $\epsilon \equiv 1/N$ and \mathbf{e}^{ij} is an M -dimensional vector $e_k^{ij} \equiv -\delta_{ik} + \delta_{jk}$, with $k = 1, 2, \dots, M$. We are especially interested in the dynamics of a nearly coherent state. With conditions (i) and (ii) in Sec. II B and $N \rightarrow \infty$, Eq. (A1) becomes

$$\begin{aligned}
 i \partial_t \varphi = & \sum_i H_{ii}^0 N x_i \varphi + \sum_i U_{ii} N x_i (x_i - \epsilon) \varphi \\
 & + \sum_{i \neq j} H_{ij}^0 N \sqrt{x_i x_j} \varphi \exp[i(k_j - k_i)\epsilon] \\
 & + \sum_{i \neq j} H_{ij}^0 \frac{1}{2} \sqrt{\frac{x_i}{x_j}} \varphi \exp[i(k_j - k_i)\epsilon] \\
 & + \sum_{i \neq j} H_{ij}^0 \sqrt{x_i x_j} [(\partial_j - \partial_i) |\varphi|] \frac{\varphi}{|\varphi|} \exp[i(k_j - k_i)\epsilon] \\
 & + \sum_{i \neq j} U_{ij} N x_i x_j \varphi + o(1), \quad (\text{A2})
 \end{aligned}$$

where $\partial_i \equiv \frac{\partial}{\partial x_i}$ and $k_i(\mathbf{x}; t)$ is the local wave vector of wave function φ at $(\mathbf{x}; t)$, as discussed in condition (ii). The argument of all k_i and φ is $(\mathbf{x}; t)$, which is omitted.

Now we assume a δ -function solution as in Eq. (9). By setting the coefficients before δ , $\partial_t \delta$, and the derivatives of coefficients before δ (which is necessary to reflect the plane-wave phase structure) on both sides equal, keeping finite terms in the large- N limit, we obtain

$$\partial_t x_i^0 = 2 \operatorname{Im} \sum_j H_{ij}^0 \sqrt{x_i^0 x_j^0} \exp [i(k_j^0 - k_i^0)\epsilon], \quad (\text{A3})$$

$$\partial_t k_i^0 \epsilon = -\operatorname{Re} \sum_j H_{ij}^0 \sqrt{\frac{x_j^0}{x_i^0}} \exp [i(k_j^0 - k_i^0)\epsilon] - 2 \sum_j U_{ij} x_j^0, \quad (\text{A4})$$

where the argument t of all x_i^0 and k_i^0 is omitted for brevity. Lengthy but straightforward calculations will verify that Eqs. (A3) and (A4) are equivalent to Eq. (10), which is same as the mean-field EOM for the one-particle RDM $\hat{\rho}$.

-
- [1] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
- [2] V. I. Yukalov, *Laser Phys. Lett.* **1**, 435 (2004).
- [3] B. Wu and Q. Niu, *Phys. Rev. A* **64**, 061603 (2001).
- [4] A. Smerzi, A. Trombettoni, P. G. Kevrekidis, and A. R. Bishop, *Phys. Rev. Lett.* **89**, 170402 (2002).
- [5] J. Liu, W. Wang, C. Zhang, Q. Niu, and B. Li, *Phys. Lett. A* **353**, 216 (2006).
- [6] G. Manfredi and P.-A. Hervieux, *Phys. Rev. Lett.* **100**, 050405 (2008).
- [7] J. Reslen, C. E. Creffield, and T. S. Monteiro, *Phys. Rev. A* **77**, 043621 (2008).
- [8] Q. Thommen, J. C. Garreau, and V. Zehnlé, *Phys. Rev. Lett.* **91**, 210405 (2003).
- [9] P. Buonsante, R. Franzosi, and V. Penna, *Phys. Rev. Lett.* **90**, 050404 (2003).
- [10] S. Burger, F. S. Cataliotti, C. Fort, F. Minardi, M. Inguscio, M. L. Chiofalo, and M. P. Tosi, *Phys. Rev. Lett.* **86**, 4447 (2001).
- [11] B. Wu and Q. Niu, *Phys. Rev. Lett.* **89**, 088901 (2002).
- [12] L. Fallani, L. De Sarlo, J. E. Lye, M. Modugno, R. Saers, C. Fort, and M. Inguscio, *Phys. Rev. Lett.* **93**, 140406 (2004).
- [13] N. Gemelke, E. Sarajlic, Y. Bidel, S. Hong, and S. Chu, *Phys. Rev. Lett.* **95**, 170404 (2005).
- [14] W. Zhang, D. L. Zhou, M.-S. Chang, M. S. Chapman, and L. You, *Phys. Rev. Lett.* **95**, 180403 (2005).
- [15] I. Bloch, *Nat. Phys.* **1**, 23 (2005).
- [16] E. H. Lieb, R. Seiringer, and J. Yngvason, *Phys. Rev. A* **61**, 043602 (2000).
- [17] L. Erdős, B. Schlein, and H.-T. Yau, *Phys. Rev. Lett.* **98**, 040404 (2007).
- [18] G. Berman and G. Zaslavsky, *Physica A* **91**, 450 (1978).
- [19] P. G. Silvestrov and C. W. J. Beenakker, *Phys. Rev. E* **65**, 035208 (2002).
- [20] L. G. Yaffe, *Rev. Mod. Phys.* **54**, 407 (1982).
- [21] K. Nemoto, C. A. Holmes, G. J. Milburn, and W. J. Munro, *Phys. Rev. A* **63**, 013604 (2000).
- [22] R. Franzosi and V. Penna, *Phys. Rev. A* **65**, 013601 (2001).
- [23] R. Franzosi and V. Penna, *Phys. Rev. E* **67**, 046227 (2003).
- [24] B. Liu, L.-B. Fu, S.-P. Yang, and J. Liu, *Phys. Rev. A* **75**, 033601 (2007).
- [25] Q. Guo, X. Chen, and B. Wu, *Opt. Express* **22**, 19219 (2014).
- [26] X. Luo, Q. Xie, and B. Wu, *Phys. Rev. A* **76**, 051802 (2007).
- [27] S. Habib, K. Shizume, and W. H. Zurek, *Phys. Rev. Lett.* **80**, 4361 (1998).
- [28] A. Vardi and J. R. Anglin, *Phys. Rev. Lett.* **86**, 568 (2001).
- [29] B. Gertjerenken, S. Arlinghaus, N. Teichmann, and C. Weiss, *Phys. Rev. A* **82**, 023620 (2010).
- [30] C. Weiss and N. Teichmann, *Phys. Rev. Lett.* **100**, 140408 (2008).
- [31] I. Březinová, A. U. J. Lode, A. I. Streltsov, O. E. Alon, L. S. Cederbaum, and J. Burgdörfer, *Phys. Rev. A* **86**, 013630 (2012).
- [32] P. Buonsante, V. Penna, and A. Vezzani, *Phys. Rev. A* **84**, 061601 (2011).
- [33] P. Buonsante, R. Burioni, E. Vescovi, and A. Vezzani, *Phys. Rev. A* **85**, 043625 (2012).
- [34] Z. P. Karkuszewski, J. Zakrzewski, and W. H. Zurek, *Phys. Rev. A* **65**, 042113 (2002).
- [35] M.-S. Chang, Q. Qin, W. Zhang, L. You, and M. S. Chapman, *Nat. Phys.* **1**, 111 (2005).
- [36] W. Zhang, D. L. Zhou, M.-S. Chang, M. S. Chapman, and L. You, *Phys. Rev. A* **72**, 013602 (2005).