

Quantum walks and non-Abelian discrete gauge theory

Pablo Arnault,^{1,*} Giuseppe Di Molfetta,^{2,†} Marc Brachet,³ and Fabrice Debbausch¹

¹*LERMA, Observatoire de Paris, PSL Research University, Centre National de la Recherche Scientifique, Sorbonne Universités, UPMC Université Paris 6, UMR 8112, F-75014 Paris, France*

²*Departamento de Física Teórica and IFIC, Universidad de Valencia-CSIC, 46100-Burjassot, Valencia, Spain*

³*Centre National de la Recherche Scientifique, Laboratoire de Physique Statistique, École Normale Supérieure, 75231 Paris Cedex 05, France*

(Received 6 May 2016; published 19 July 2016)

A family of discrete-time quantum walks (DTQWs) on the line with an exact discrete $U(N)$ gauge invariance is introduced. It is shown that the continuous limit of these DTQWs, when it exists, coincides with the dynamics of a Dirac fermion coupled to usual $U(N)$ gauge fields in two-dimensional spacetime. A discrete generalization of the usual $U(N)$ curvature is also constructed. An alternate interpretation of these results in terms of superimposed $U(1)$ Maxwell fields and $SU(N)$ gauge fields is discussed in the Appendix. Numerical simulations are also presented, which explore the convergence of the DTQWs towards their continuous limit and which also compare the DTQWs with classical (i.e., nonquantum) motions in classical $SU(2)$ fields. The results presented in this paper constitute a first step towards quantum simulations of generic Yang-Mills gauge theories through DTQWs.

DOI: [10.1103/PhysRevA.94.012335](https://doi.org/10.1103/PhysRevA.94.012335)

I. INTRODUCTION

Discrete-time quantum walks (DTQWs) are unitary quantum automata and can be viewed as formal generalizations of classical random walks. They were first considered in a systematic way by Meyer [1], following the seminal work of Feynman and Hibbs [2] and Aharonov *et al.* [3]. DTQWs have been realized experimentally with a wide range of physical objects and setups [4–10], and are studied in a large variety of contexts, ranging from fundamental quantum physics [10,11] to quantum algorithmics [12,13], solid-state physics [14–17], and biophysics [18,19].

It has been shown recently that the continuous limit of several DTQWs coincides with the dynamics of Dirac fermions coupled to electromagnetic [20–22] and relativistic gravitational fields [23–26]. Though these fields are naturally gauge fields, they are not generic Yang-Mills gauge fields. Indeed, electromagnetism is based on the Abelian gauge group $U(1)$, while relativistic gravitational fields are not Yang-Mills gauge fields, since they are represented by a metric, and not by a connection. The aim of this paper is to exhibit and study DTQWs the continuous limit of which coincides with the dynamics of a fermion coupled to Yang-Mills $U(N)$ gauge fields.

To make things definite and as simple as possible, we focus on one-dimensional (1D) DTQWs. The minimal 1D DTQWs have a two-dimensional coin space. Their wave functions thus have two components, one propagating towards the left and one towards the right. To take into account the internal degrees of freedom associated to $U(N)$ gauge invariance, we consider 1D DTQWs with coin space of dimension $2N$, i.e., $2N$ -component wave functions. Half of the wave-function components propagates towards the left, and the other half towards the right. The so-called mixing operator advancing the walk in time is represented by a $2N \times 2N$ time- and space-dependent unitary matrix.

We introduce 1D DTQWs with $2N$ components which admit an exact discrete $U(N)$ gauge invariance and build for the DTQWs a discrete equivalent \mathcal{F} of the usual Yang-Mills curvature F . We then prove that the limit of these DTQWs, when it exists, coincides with the dynamics of Dirac fermions coupled to $U(N)$ gauge fields and that the discrete curvature \mathcal{F} tends towards F in the continuous limit.

These formal computations are complemented by numerical simulations. These address the convergence of the DTQWs towards their continuous limit and the correspondence with classical (i.e., nonquantum) trajectories in Yang-Mills fields [27]. The paper concludes by a brief summary and a discussion of the main results. Finally, the Appendix elaborates on the fact that a $U(N)$ gauge field can be viewed as the superposition of a $U(1)$ Maxwell field and an $SU(N)$ gauge field, and reinterprets our results in that alternate context.

II. THE DTQWS AND THEIR GAUGE INVARIANCE

A. The DTQWs

We consider DTQWs defined over discrete time and infinite discrete one-dimensional space. Instants are labeled by $j \in \mathbb{N}$ and space points are labeled by $p \in \mathbb{Z}$. The coin space of the DTQWs has dimension $2N$. Given a certain orthonormal basis in this space, the wave functions Ψ of the walks are represented by $2N$ components and we group these components into two N -component sets ψ^- and ψ^+ , which represent those parts of Ψ which propagate, respectively, to the left and to the right. The evolution equations read

$$\begin{bmatrix} \psi_{j+1,p}^- \\ \psi_{j+1,p}^+ \end{bmatrix} = \mathbf{B}(\theta, P_{j,p}, Q_{j,p}) \begin{bmatrix} \psi_{j,p+1}^- \\ \psi_{j,p-1}^+ \end{bmatrix}, \quad (1)$$

with

$$\begin{aligned} \mathbf{B}(\theta, P, Q) &= (\mathbf{C}(\theta) \otimes \mathbf{1}_N) \times \begin{bmatrix} P & 0 \\ 0 & Q \end{bmatrix} \\ &\equiv \begin{bmatrix} (\cos \theta) P & (i \sin \theta) Q \\ (i \sin \theta) P & (\cos \theta) Q \end{bmatrix}, \end{aligned} \quad (2)$$

*pablo.arnault@upmc.fr

†giuseppe.di.molfetta@uv.es

where \otimes is the so-called Kronecker (or tensorial) product for matrices, and P and Q are elements of $U(N)$. These walks are unitary, i.e., $\Pi_j = \sum_p |\Psi_{j,p}|^2$ is independent of j .

In the continuous limit, the parameter θ will code for the mass of the fermion and the matrices P and Q will code for the potential of the $U(N)$ gauge field to which the fermion is coupled.

B. Discrete $U(N)$ gauge invariance and discrete curvature

The DTQWs defined by Eq. (1) admit a discrete local $U(N)$ gauge invariance. Indeed, consider the local gauge transformation $\Psi_{j,p} = (\mathbf{1}_2 \otimes G_{j,p}^{-1}) \Psi'_{j,p}$, where $G_{j,p}$ is some matrix of $U(N)$. Equation (1) is kept invariant under this transformation, that is,

$$\begin{bmatrix} \Psi'_{j+1,p} \\ \Psi'_{j+1,p+1} \end{bmatrix} = \mathbf{B}(\theta, P'_{j,p}, Q'_{j,p}) \begin{bmatrix} \Psi'_{j,p+1} \\ \Psi'_{j,p-1} \end{bmatrix}, \quad (3)$$

provided that we set

$$\begin{aligned} P'_{j,p} &= G_{j+1,p} P_{j,p} G_{j,p+1}^{-1}, \\ Q'_{j,p} &= G_{j+1,p} Q_{j,p} G_{j,p-1}^{-1}. \end{aligned} \quad (4)$$

The above gauge invariance suggests that $R = (P, Q)$ is the discrete equivalent of the usual continuous $U(N)$ gauge potentials. This will be confirmed in Sec. III, where the continuous limit of Eq. (1) will be derived. We now wish to build out of R an object \mathcal{F} defined on the spacetime lattice, which generalizes for DTQWs the usual curvature (field-strength) tensor [28] F of standard gauge fields. This will be done by searching for an object whose transformation law under a change of gauge resembles the transformation law of F . Let

$$\begin{aligned} U_{j,p}(R) &= Q_{j,p}^\dagger P_{j,p}, \\ V_{j,p}(R) &= Q_{j,p} P_{j-1,p-1}, \end{aligned} \quad (5)$$

whose transformation under a change of gauge reads

$$\begin{aligned} U'_{j,p}(R') &= G_{j,p-1} U_{j,p}(R) G_{j,p+1}^\dagger, \\ V'_{j,p}(R') &= G_{j+1,p} V_{j,p}(R) G_{j-1,p}^\dagger, \end{aligned} \quad (6)$$

involving shifts of $G_{j,p}$ only in the spatial (temporal) dimension for $U_{j,p}(R)$ [$V_{j,p}(R)$], while these shifts were mixed in the transformation laws of Eq. (4). From these equations, we can write transformation laws involving the $2 \times 2 = 4$ discrete-spacetime neighbors of $G_{j,p}$:

$$\begin{aligned} U'_{j+1,p}(R') &= G_{j+1,p-1} U_{j+1,p}(R) G_{j+1,p+1}^\dagger, \\ U'_{j-1,p}(R') &= G_{j-1,p-1} U_{j-1,p}(R) G_{j-1,p+1}^\dagger, \\ V'_{j,p-1}(R') &= G_{j+1,p-1} V_{j,p-1}(R) G_{j-1,p-1}^\dagger, \\ V'_{j,p+1}(R') &= G_{j+1,p+1} V_{j,p+1}(R) G_{j-1,p+1}^\dagger, \end{aligned} \quad (7)$$

from which we can build

$$\mathcal{F}_{j,p}(R) = U_{j-1,p}^\dagger(R) V_{j,p-1}^\dagger(R) U_{j+1,p}(R) V_{j,p+1}(R), \quad (8)$$

whose transformation law reads

$$\mathcal{F}'_{j,p}(R') = G_{j-1,p+1} \mathcal{F}_{j,p}(R) G_{j-1,p+1}^{-1}. \quad (9)$$

As will become apparent in the next section, \mathcal{F} is a discrete equivalent to the curvature (field-strength) tensor of continuous gauge fields.

III. CONTINUOUS LIMIT

We now show that it is possible to choose θ , P , and Q in such a way that Eq. (1) admits a continuous limit identical to the Dirac equation for a fermion coupled to an arbitrary $U(N)$ gauge field.

In order to compute the continuous limit of Eq. (1), we first introduce a dimensionless time and space step ϵ , and consider that $\Psi_{j,p}$, $P_{j,p}$, and $Q_{j,p}$ are the values $\Psi(t_j, x_p)$, $P(t_j, x_p)$, and $Q(t_j, x_p)$ taken at spacetime point $(t_j = j\epsilon, x_p = p\epsilon)$ by a $2N$ -component wave function Ψ and two time- and spacetime-dependent matrices P and Q in $U(N)$. We then assume that Ψ , P , and Q are at least twice differentiable with respect to both space and time variables and let ϵ tend to zero.

As ϵ tends to zero, the wave functions on the left-hand side and on the right-hand side of Eq. (1) both tend towards $\Psi(t_j, x_p)$. Thus, the continuous limit of Eq. (1) can only exist if, in that limit, $B(\theta, P, Q)$ tends to unity at all points in spacetime. This is achieved by choosing an angle θ which tends to zero with ϵ and two matrices P and Q which tend to unity as ϵ goes to zero. We retain $\theta = -\epsilon m$, where m is a positive constant (as opposed to a function of t and x) which will play the role of a mass in the continuous limit. As for the matrices P and Q , we remark that $U(N)$ is a compact and connected Lie group. Thus, the exponential map generates the whole group [29], i.e., all elements $M \in U(N)$ can be written as

$$M = \exp \left(i \sum_k X_M^k \tau_k \right), \quad (10)$$

where the τ_k 's are N^2 generators of $U(N)$ and the X_M^k 's can serve as coordinates for M .

To ensure that both functions $P(t, x)$ and $Q(t, x)$ tend to unity when ϵ tends to zero, we choose $X_P^k(t, x) = \epsilon b_P^k(t, x)$ and $X_Q^k(t, x) = \epsilon b_Q^k(t, x)$, where $b_{P/Q}^k(t, x)$ are two real functions independent of ϵ .

Taylor expanding Eq. (1) at first order in ϵ and letting ϵ tend to zero then delivers

$$\begin{aligned} (\partial_0 - i b_0^k \tau_k) \psi^- - (\partial_1 - i b_1^k \tau_k) \psi^- &= -i m \psi^+, \\ (\partial_0 - i b_0^k \tau_k) \psi^+ + (\partial_1 - i b_1^k \tau_k) \psi^+ &= -i m \psi^-, \end{aligned} \quad (11)$$

where $\partial_0 = \partial_t$, $\partial_1 = \partial_x$,

$$\begin{aligned} b_0 &= (b_Q + b_P)/2, \\ b_1 &= (b_Q - b_P)/2, \end{aligned} \quad (12)$$

and summation over repeated index k is implied.

Equations (11) can be recast as

$$[i \gamma^\mu D_\mu - m] \Psi = 0, \quad (13)$$

where index μ is summed over from zero to one, with the gamma matrices $\gamma^0 = \sigma_1 \otimes \mathbf{1}_N$ and $\gamma^1 = i \sigma_2 \otimes \mathbf{1}_N$ and the covariant derivative $D_\mu = \partial_\mu - i b_\mu^k \tau_k$. Equation (13) is the flat-spacetime Dirac equation, with convention $[\eta^{\mu\nu}] = \text{diag}(+, -)$, for a spin-1/2 fermion of mass m coupled to a non-Abelian $U(N)$ potential $b_\mu^k \tau_k$ (with coupling constant

$g = -1$) [28] belonging to the Lie algebra of $U(N)$. Note that the b_μ^k 's are real-valued space- and time-dependent fields.

Taylor expanding Definition (8) for \mathcal{F} delivers

$$\mathcal{F}(t, x) = \mathbf{1}_N + 4\epsilon^2 F_{10}(t, x) + O(\epsilon^3), \quad (14)$$

where F_{10} is the only nonvanishing component of the antisymmetric curvature (field-strength) tensor $F_{\mu\nu}$ of the connection $B_\mu = b_\mu^k \tau_k$, defined by

$$F_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu - i[B_\mu, B_\nu], \quad (15)$$

with $[B_\mu, B_\nu] = B_\mu B_\nu - B_\nu B_\mu$. Note that the transformation law for $F_{\mu\nu}$ under a change of gauge reads

$$F'_{\mu\nu} = G F_{\mu\nu} G^{-1}, \quad (16)$$

which closely parallels Eq. (9).

IV. NUMERICAL SIMULATIONS OF U(2)-INVARIANT DTQWS

A. Simulated walk

As shown in the Appendix to this paper, $U(N)$ factorizes into the product of $U(1)$ and $SU(N)$. In physical terms, this means that a $U(N)$ gauge field can be seen as the superposition of a $U(1)$ Maxwell field and an $SU(N)$ gauge field. The effects of Maxwell fields on DTQWs have already been presented in several publications [20,21,24]. We want to focus on the effects of non-Abelian Yang-Mills fields and thus choose to simulate situations where the Maxwell field identically vanishes so that the $U(N)$ gauge field is then actually an $SU(N)$ gauge field. We also choose the simplest option $N = 2$. The group $SU(2)$ is compact and connected, and can thus be fully generated by the exponential map, from three generators $\bar{\tau}_k$, $k = 1, 2, 3$, belonging to its Lie algebra. We retain (see the Appendix) $\bar{\tau}_k = \sigma_k/2$ where the σ_k 's are the three Pauli matrices, and choose

$$\begin{aligned} \bar{\mathbf{X}}_0 &= (\bar{\mathbf{X}}_Q + \bar{\mathbf{X}}_P)/2 = (0, 0, 0), \\ \bar{\mathbf{X}}_1 &= (\bar{\mathbf{X}}_Q - \bar{\mathbf{X}}_P)/2 = (\epsilon E_{YM} t, 0, 0). \end{aligned} \quad (17)$$

The bar is used to distinguish the notations used for $SU(2)$ from those used for $U(N)$, including $U(2)$, in Sec. III. The boldface notation is used as a reminder that the Lie algebra of $SU(2)$ is of dimension 3. The continuous limit can be recovered by letting ϵ tend to zero (see Sec. III). In such a continuous limit, this potential, Eq. (17), generates a uniform and constant $SU(2)$ ‘‘electric’’ field E_{YM} in the $\bar{\tau}_1$ direction of the $SU(2)$ Lie algebra.

B. Convergence towards the continuous limit

To study numerically the convergence of a DTQW towards a solution of the Dirac equation, we choose an initial wave function and compare, for some given time j , its evolution $\psi_{QW}^u(t_j = \epsilon j, \cdot)$, $u \in \{+, -\}$, through the DTQW to the evolution $\psi_D^u(t_j, \cdot)$ of the same initial condition through the Dirac equation. The comparison is carried out through the following mean relative difference:

$$\delta\psi_j^u = \sqrt{\frac{\langle |\psi_D^u(t_j, \cdot) - \psi_{QW}^u(t_j, \cdot)|^2 \rangle}{\langle |\psi_D^u(t_j, \cdot)|^2 \rangle}}, \quad (18)$$




FIG. 1. Relative differences δf_j for $f = \text{Im } \psi^-$ and $f = \text{Re } \psi^-$ as functions of ϵ , at time $j = 100$, with $m = 0.1$ and $E_{YM} = 0.08$. The initial condition is given by Eq. (20) with $\sigma = 0.5$.

where

$$\langle f(t_j, \cdot) \rangle = \sum_{p=-p_{\max}(\epsilon)}^{p_{\max}(\epsilon)} f(t_j, x_p) \epsilon. \quad (19)$$

The numerical simulations are carried out over the space interval $[-x_{\max}, x_{\max}]$ with $x_{\max} = 200$, and $p_{\max}(\epsilon) \equiv x_{\max}/\epsilon$. The maximal time over which we carry out the simulations, $t_{\max} = 350$, is short enough so that the walker never reaches the spatial boundaries.

Note that $\delta\psi_j^u$ does not measure the difference between quantum states, for which phase differences are unimportant, but rather the difference between the functions ψ_D^u and ψ_{QW}^u . This is appropriate here because we want to test the convergence of a discrete scheme towards its formal continuous limit, and this convergence should be verified on both modulus and phase, i.e., on the whole complex function, and not only on the state it represents.

Since there is only a time dependence and no space dependence in the potentials (17), we can use as numerical solver for the Dirac equation standard pseudospectral methods [30,31], with resolution $2\pi/\epsilon$ in 2π -periodic boundary conditions. Time marching is performed using a second-order Runge-Kutta scheme. The original DTQW can also be simulated in spectral space using the standard translation operator in Fourier space.

Figure 1 shows that the mean relative differences δf_j for $f = \text{Im } \psi^-$ and $\text{Re } \psi^-$ scale as ϵ as expected: indeed, this scaling coincides with the theoretical expectation since, for a single time step, the discrepancy is theoretically of order ϵ^2 . Thus, after a fixed time $t = O(\epsilon^{-1})$, the discrepancy is of order $\epsilon^{-1}\epsilon^2 = \epsilon$. These results also confirm that the DTQW Eq. (1) with choice Eq. (17) can be used to simulate massive Dirac dynamics in a constant and uniform non-Abelian electric field E_{YM} .

C. Comparison with classical trajectories

Given a wave equation, it is well known [32] that the center of mass of a wave-packet solution follows classical trajectories. In the continuous-limit case described above in




FIG. 2. Time evolution of the DTQW mean trajectory $\bar{x}(t)$ for a non-Abelian coupling constant $g = 1$ and different values of E_{YM} , vs classical trajectories (black solid line). Short-time agreement between quantum and classical dynamics is shown in the ultrarelativistic range $k_0^2 = 1$ (green and blue) and the nonrelativistic range $k_0^2 = 0$ (red). The initial condition is given by Eq. (20) with $\sigma = 0.5$ (green and blue), $\sigma = 1$ (red), and $m = 0.1$.

Sec. IV A, the corresponding classical equations have been explicitly derived in [27]. We now want to investigate whether the original DTQW also reproduces classical motions of the center-of-mass of wave packets.

We consider k_0 -centered Gaussian wave packets of positive-energy eigenvector $u_+(k)$ of the two-component [i.e., without SU(2) internal degree of freedom] free Dirac Hamiltonian, tensorized with an equally weighted initial SU(2) state:

$$\Psi(x) = \int dk e^{-\frac{(k-k_0)^2}{2\sigma^2} + ixk} [u_+(k) \otimes (1,1)^T / \sqrt{2}], \quad (20)$$

where superscript T denotes the transposition, and

$$u_+(k) = \left[\frac{\sqrt{k^2 + m^2} - k}{\sqrt{(\sqrt{k^2 + m^2} - k)^2 + m^2}}, \frac{1}{\sqrt{\frac{(\sqrt{k^2 + m^2} - k)^2}{m^2} + 1}} \right]^T. \quad (21)$$

Figure 2 demonstrates the short-time agreement between solutions of classical particle trajectory equations (see [27]) and the centers of wave packets $\bar{x}(t)$ obtained from DTQW solutions, both in the nonrelativistic case, $k_0^2 = 0$, $m = 0.1$, and in the relativistic case, $k_0^2 = 1$, $m = 0.1$. When the agreement is lost, the oscillatory trajectory for $\bar{x}(t)$ is produced by the DTQW. Note that similar long-time oscillations are also found in the simple context of DTQWs corresponding to Dirac fermions coupled to electromagnetic fields [33].

V. CONCLUSION

We have introduced DTQWs on the line which exhibit an exact discrete $U(N)$ gauge invariance and the continuous limit of which coincides, when it exists, with the dynamics of Dirac fermions coupled to $U(N)$ gauge fields. We have also built a discrete generalization of the curvature tensor of the gauge fields. We have finally complemented these analytical results by numerical simulations which explore the convergence of

the DTQWs towards their continuous limit and compare the DTQWs with the dynamics of nonquantum particles in classical gauge fields. The interpretation of our results in terms of Maxwell fields superimposed to SU(N) gauge fields is presented in the Appendix. The results presented in this paper constitute a first step towards quantum simulations of generic Yang-Mills gauge theories through DTQWs. Until now, only DTQWs with two-component wave functions have been realized experimentally [34]. But experimental procedures allowing the implementation of DTQWs with wave functions having more than two components have been proposed in [35,36]. In these procedures, the DTQWs are implemented with single photons or classical light, for example in optical cavities.

Let us now mention a few avenues open to future studies. The DTQWs presented in this paper should first be extended to (1 + 2) and then to (1 + 3) spacetime dimensions. Note that DTQWs modeling Dirac fermions coupled to U(1) gauge fields have already been proposed in (1 + 1) and (1 + 2) dimensions [20–22]. Another possible extension would be the construction of DTQWs which are coupled, not only to $U(N)$ gauge fields but also to gravity. Until now, this has only been done for $N = 1$ and in (1 + 1) spacetime dimensions [24]. Also, performing full quantum simulations of Yang-Mills gauge theories will require complementing the fermionic DTQW dynamics by dynamical equations for the discrete gauge fields, i.e., for matrices $P, Q \in U(N)$ which define the DTQW. The dynamical equations for the gauge field should be a set of finite difference equations relating the discrete curvature (field-strength) tensor \mathcal{F} introduced in the present paper to a discrete gauge-invariant fermionic current associated to the DTQW. This current has already been presented in [22] for $N = 1$ in (1 + 2) spacetime dimensions, and the corresponding discrete Maxwell equations have also been written down. The procedure should now be extended to generic noncommutative discrete gauge fields. Finally, incorporating Yang-Mills fields to DTQWs defined on arbitrary graphs is certainly worth working on, if only for applications to quantum information.

ACKNOWLEDGMENTS

This work has been partially supported by the Spanish Ministerio de Educación e Innovación, MICIN-FEDER Projects No. FPA2014-54459-P and No. SEV-2014-0398, and by “Generalitat Valenciana” Grant No. GVPROMETEOII2014-087.

APPENDIX: ALTERNATIVE PHYSICAL INTERPRETATION

The Lie group $U(N)$ is the group of $N \times N$ unitary matrices, i.e., $N \times N$ matrices the determinant modulus of which equals unity. In particular, elements of $U(1)$ are complex numbers of the unit modulus, i.e., complex numbers of the form $\exp(i\beta)$, where β is an arbitrary real number. The group $U(N)$ is Abelian for $N = 1$ and non-Abelian for $N > 1$. Consider now an arbitrary element M of $U(N)$, its determinant $\det M = \exp(i\alpha)$, $\alpha \in]-\pi, +\pi]$, and we define the matrix $\bar{M} = M/\delta$ where $\delta^N = \det M$. The matrix \bar{M} has a unit determinant and is thus an element of the special unitary group $SU(N)$. The group $U(N)$ can therefore be factorized into the direct product of $U(1)$ and $SU(N)$. This

factorization is not unique because δ is not uniquely defined by the equation $\delta^N = \det M$. Indeed, this equation has the N distinct solutions $\delta_k = \exp[i(\alpha + 2k\pi)/N]$, $k = 0, \dots, N - 1$, and each solution defines a different factorization. Note also that imposing a factorization which depends continuously on M is only possible if one makes a cut along the negative real axis in the complex plane of $\det M$, i.e., if one does not define the factorization for matrices M the determinant of which corresponds to the value $\alpha = \pi$ (and is thus equal to -1).

To make all computations definite, we now choose $k = 0$ in the above definition of δ_k . This defines unambiguously a factorization of $U(N)$ into the direct product of $U(1)$ and $SU(N)$. This factorization is not continuous for matrices M with $\det M = -1$, but that should not be a practical problem when one is working on a spacetime lattice. In the continuous limit, all $U(N)$ matrices considered in this paper tend to unity. Their determinant is thus close to unity and the retained factorization is thus defined and continuous for all these matrices.

In physical terms, the existence of the factorization means that a $U(N)$ gauge field can be interpreted as the superposition of a $U(1)$ Maxwell field and an $SU(N)$ gauge field. Now, $SU(N)$ is itself a compact and connected Lie group, so the whole group is generated from the identity by the exponential map. The above factorization can thus be used to write all matrices $M \in U(N)$ as

$$M = \delta_M \bar{M} = \exp(iY_M) \exp\left(i \sum_k \bar{X}_M^k \bar{\tau}_k\right), \quad (\text{A1})$$

where the $\bar{\tau}_k$'s are the $N^2 - 1$ generators of $SU(N)$. This point of view is adopted in Sec. IV A. Note that the factorization of $U(N)$ also shows that the DTQWs presented in this paper coincide, for $N = 1$, with the DTQWs already proposed to simulate Dirac fermions coupled to arbitrary electric fields [20].

The discrete curvature \mathcal{F} also factorizes into a curvature for the Maxwell field and a curvature for the $SU(N)$ gauge field.

One finds indeed that

$$\mathcal{F}_{j,p}(R) = \mathcal{F}_{j,p}(\delta_R) \mathcal{F}_{j,p}(\bar{R}) \quad (\text{A2})$$

where $\delta_R = (e^{iY_P}, e^{iY_Q})$ and $\bar{R} = (\bar{P}, \bar{Q})$. The $SU(N)$ curvature $\mathcal{F}_{j,p}(\bar{R})$ is given by Eq. (8) and the $U(1)$ Abelian curvature reads

$$\mathcal{F}_{j,p}(\delta_R) = \exp[2i(\mathcal{I}f_{10})_{j,p}], \quad (\text{A3})$$

where

$$(f_{10})_{j,p} = (d_1 Y_0)_{j,p} - (d_0 Y_1)_{j,p}, \quad (\text{A4})$$

with

$$\begin{aligned} Y_0 &= (Y_Q + Y_P)/2, \\ Y_1 &= (Y_Q - Y_P)/2, \end{aligned} \quad (\text{A5})$$

$$d_0 = (L_0 - \Sigma_1), \quad d_1 = \Delta_1, \quad (\text{A6})$$

and

$$\begin{aligned} (L_0 K)_{j,p} &= K_{j+1,p}, \\ (\Sigma_1 K)_{j,p} &= (K_{j,p+1} + K_{j,p+1})/2, \\ (\Delta_1 K)_{j,p} &= (K_{j,p+1} - K_{j,p+1})/2, \end{aligned} \quad (\text{A7})$$

where K is an arbitrary quantity which depends on j and p . Operator \mathcal{I} is defined in terms of L_0 and L_1 by

$$\mathcal{I} = 1 + L_0^{-1} L_1^{-1}. \quad (\text{A8})$$

This form of $\mathcal{F}_{j,p}(\delta_R)$ is interesting because $(f_{10})_{j,p}$ and operators d_0 and d_1 have already been introduced in [22] in the context of DTQWs exhibiting a $U(1)$ gauge invariance.

Choosing $Y_\mu = \epsilon A_\mu$, $\bar{X}_{P/Q}^k = \epsilon \bar{b}_{P/Q}^k$ and Taylor expanding Eq. (A2) at second order in ϵ delivers

$$\mathcal{F}_{j,p}(R) = \mathbf{1}_N + 4\epsilon^2 (f_{10} \mathbf{1}_N + \bar{F}_{10}) + O(\epsilon^3), \quad (\text{A9})$$

where \bar{F}_{10} is given by Eq. (15) after substitution $B \rightarrow \bar{B}$, and $f_{10} = \partial_1 A_0 - \partial_0 A_1$ is the (10) component of the usual Abelian curvature tensor.

-
- [1] D. A. Meyer, *J. Stat. Phys.* **85**, 551 (1996).
 - [2] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, International Series in Pure and Applied Physics (McGraw-Hill, New York, 1965).
 - [3] Y. Aharonov, L. Davidovich, and N. Zagury, *Phys. Rev. A* **48**, 1687 (1993).
 - [4] H. Schmitz, R. Matjeschk, Ch. Schneider, J. Glueckert, M. Enderlein, T. Huber, and T. Schaetz, *Phys. Rev. Lett.* **103**, 090504 (2009).
 - [5] F. Zähringer, G. Kirchmair, R. Gerritsma, E. Solano, R. Blatt, and C. F. Roos, *Phys. Rev. Lett.* **104**, 100503 (2010).
 - [6] A. Schreiber, K. N. Cassemiro, V. Potoček, A. Gábris, P. J. Mosley, E. Andersson, I. Jex, and Ch. Silberhorn, *Phys. Rev. Lett.* **104**, 050502 (2010).
 - [7] M. Karski, L. Förster, J.-M. Cho, A. Steffen, W. Alt, D. Meschede, and A. Widera, *Science* **325**, 174 (2009).
 - [8] L. Sansoni, F. Sciarrino, G. Vallone, P. Mataloni, A. Crespi, R. Ramponi, and R. Osellame, *Phys. Rev. Lett.* **108**, 010502 (2012).
 - [9] B. C. Sanders, S. D. Bartlett, B. Tregenna, and P. L. Knight, *Phys. Rev. A* **67**, 042305 (2003).
 - [10] H. B. Perets, Y. Lahini, F. Pozzi, M. Sorel, R. Morandotti, and Y. Silberberg, *Phys. Rev. Lett.* **100**, 170506 (2008).
 - [11] D. Giulini, E. Joos, C. Kiefer, J. Kupsch, I.-O. Stamatescu, and H. D. Zeh, *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer-Verlag, Berlin, 1996).
 - [12] A. Ambainis, *SIAM J. Comput.* **37**, 210 (2007).
 - [13] F. Magniez, A. Nayak, J. Roland, and M. Santha, Search via Quantum Walk, *SIAM Journal on Computing: Proceedings of the Thirty-Ninth Annual ACM Symposium on Theory of Computing*, 2007 (unpublished).
 - [14] C. Aslangul, *J. Phys. A* **38**, 1 (2005).
 - [15] S. Bose, *Phys. Rev. Lett.* **91**, 207901 (2003).
 - [16] D. Burgarth, Ph.D. thesis, University College London, 2006.
 - [17] S. Bose, *Contemp. Phys.* **48**, 13 (2007).
 - [18] E. Collini, C. Y. Wong, K. E. Wilk, P. M. G. Curmi, P. Brumer, and G. D. Scholes, *Nature (London)* **463**, 644 (2010).

- [19] G. S. Engel, T. R. Calhoun, R. L. Read, T.-K. Ahn, T. Manal, Y.-C. Cheng, R. E. Blankenship, and G. R. Fleming, *Nature (London)* **446**, 782 (2007).
- [20] G. Di Molfetta and F. Debbasch, *J. Math. Phys.* **53**, 123302 (2012).
- [21] P. Arnault and F. Debbasch, *Physica A* **443**, 179 (2016).
- [22] P. Arnault and F. Debbasch, *Phys. Rev. A* **93**, 052301 (2016).
- [23] G. Di Molfetta, M. Brachet, and F. Debbasch, *Phys. Rev. A* **88**, 042301 (2013).
- [24] G. Di Molfetta, F. Debbasch, and M. Brachet, *Physica A* **397**, 157 (2014).
- [25] P. Arrighi, S. Facchini, and M. Forets, *Quant. Info. Proc.* **15**, 3467 (2016).
- [26] S. Succi, F. Fillion-Gourdeau, and S. Palpacelli, *Eur. J. Phys. Quantum Technol.* **2**, 12 (2015).
- [27] A. D. Boozer, *Am. J. Phys.* **79**, 925 (2011).
- [28] C. Quigg, *Gauge Theories of the Strong, Weak, and Electromagnetic Interactions*, 2nd ed. (Princeton University, Princeton, NJ, 2013).
- [29] J. F. Cornwell, *Group Theory in Physics* (Academic, New York, 1986), Vol. 2.
- [30] G. Ben-yu, S. Jie, and X. Cheng-long, *Adv. Comput. Math.* **19**, 35 (2003).
- [31] D. Gottlieb and S. Orszag, *Numerical Analysis of Spectral Methods* (SIAM, Philadelphia, 1977).
- [32] L. D. Landau and E. M. Lifschitz, *Quantum Mechanics*, 3rd ed., translated from Russian (Butterworth, Washington, DC, 1977).
- [33] D. Witthaut, *Phys. Rev. A* **82**, 033602 (2010).
- [34] M. Genske, W. Alt, A. Steffen, A. H. Werner, R. F. Werner, D. Meschede, and A. Alberti, *Phys. Rev. Lett.* **110**, 190601 (2013).
- [35] P. L. Knight, E. Roldán, and G. E. Sipe, *Opt. Commun.* **227**, 147 (2003).
- [36] S. K. Goyal, F. S. Roux, A. Forbes, and T. Konrad, *Phys. Rev. A* **92**, 040302(R) (2015).