

Volume monogamy of quantum steering ellipsoids for multiqubit systemsShuming Cheng,^{1,2} Antony Milne,³ Michael J. W. Hall,¹ and Howard M. Wiseman¹¹*Centre for Quantum Computation and Communication Technology (Australian Research Council), Centre for Quantum Dynamics, Griffith University, Brisbane, QLD 4111, Australia*²*Key Laboratory of Systems and Control, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, People's Republic of China*³*Controlled Quantum Dynamics Theory, Department of Physics, Imperial College London, London SW7 2AZ, United Kingdom*
(Received 13 June 2016; published 6 October 2016)

The quantum steering ellipsoid can be used to visualize 2-qubit states, and thus provides a generalization of the Bloch picture for the single qubit. Recently, a monogamy relation for the volumes of steering ellipsoids has been derived for pure 3-qubit states and shown to be stronger than the celebrated Coffman-Kundu-Wootters inequality. We first demonstrate the close connection between this volume monogamy relation and the classification of pure 3-qubit states under stochastic local operations and classical communication. We then show that this monogamy relation does not hold for general mixed 3-qubit states and derive a weaker monogamy relation that does hold for such states. We also prove a volume monogamy relation for pure 4-qubit states (further conjectured to hold for the mixed case), and generalize our 3-qubit inequality to n qubits. Finally, we study the effect of noise on the quantum steering ellipsoid and find that the volume of any 2-qubit state is nonincreasing when the state is exposed to arbitrary local noise. This implies that any volume monogamy relation for a given class of multiqubit states remains valid under the addition of local noise. We investigate this quantitatively for the experimentally relevant example of isotropic noise.

DOI: [10.1103/PhysRevA.94.042105](https://doi.org/10.1103/PhysRevA.94.042105)**I. INTRODUCTION**

Qubits play a fundamental role in quantum information processing tasks [1] and quantum measurement and control [2]. The Bloch vector faithfully captures all features of a single-qubit state. However, finding an analogous geometric representation of multiqubit states is harder. An elegant solution to this problem has recently been given for the 2-qubit case [3]. In particular, for a 2-qubit state shared between two parties, Alice and Bob say, the set of all possible Bloch vectors that Alice can steer Bob's qubit to, via all possible local measurements on her qubit, forms an ellipsoid, called the quantum steering ellipsoid. Together with Alice's and Bob's local Bloch vectors, the quantum steering ellipsoid provides a geometric representation of the shared 2-qubit state [3].

The set of 2-qubit states has a rich structure, which is mirrored, and in some cases added to, by a corresponding zoo of quantum steering ellipsoids [3]. For example, properties of steering ellipsoids reflect various features of quantum correlations, such as discord [3–5], entanglement [3,6,7], and Einstein-Podolsky-Rosen (EPR) steering [8–11]. Moreover, it appears that steering ellipsoids may be useful for characterizing correlation properties of multiqubit states, such as monogamy, in new ways. In particular, it is well known that quantum correlations cannot be freely shared between members of multipartite systems, resulting in monogamy relations for, e.g., concurrence [12,13], Bell nonlocality [14–17], and EPR-steering inequalities [18]. In this vein, Milne *et al.* have recently obtained a strong monogamy relation for the volumes of the steering ellipsoids generated by pure 3-qubit states, termed volume monogamy [6], which is strictly stronger than the well-known Coffman-Kundu-Wootters (CKW) monogamy relation for concurrence [12] in the pure regime.

The volume monogamy relation for pure 3-qubit states immediately suggests a number of questions: Does it dis-

criminate between different types of entanglement? Is it valid for the mixed case? Are there similar relations for multiqubit states? And what happens when noise, induced by inevitable interaction with the environment and imperfections of local measurements, is present? We will provide some answers to these questions in this paper, following a brief review of the quantum steering ellipsoid and its properties in Sec. II.

In Sec. III, we obtain and discuss volume monogamy relations in general scenarios ranging from pure 3-qubit states to general multiqubit states. First, in Sec. III A we discuss the known volume monogamy relation for pure 3-qubit states [6]; give an alternative proof of this relation that demonstrates a link with the quantum marginal problem [19]; review the derivation of the CKW monogamy relation from volume monogamy; and establish a close connection between properties of volume monogamy and the classification of pure 3-qubit states under stochastic local operations and classical communication (SLOCC). In Sec. III B, we show that this volume monogamy relation is violated by some mixed states, and derive a weaker volume monogamy relation that is valid for all 3-qubit states, pure or mixed. We obtain a monogamy relation of the same form for pure 4-qubit states in Sec. III C, which we conjecture is also applicable to mixed states, and give a nontrivial volume monogamy relation for general multiqubit states in Sec. III D.

In Sec. IV, we investigate the effects of noise on steering ellipsoid volumes and on volume monogamy relations. In Sec. IV A, we show that the volume of the steering ellipsoid decreases monotonically under arbitrary local operations, including under local noise channels in particular. This significantly generalizes a recent result [20], which is restricted to particular classes of states and types of noise. Moreover, it implies that any volume monogamy relation for a given set of multiqubit states remains valid under the addition of local noise. This includes, in particular, the strong volume

monogamy relation for pure 3-qubit states in Ref. [6]. Finally, in Sec. IV B we investigate the experimentally relevant case of isotropic noise acting on a family of 3-qubit W -class states. We conclude with some remarks and open questions in Sec. V.

II. QUANTUM STEERING ELLIPSOIDS FOR 2-QUBIT STATES

Any 2-qubit state ρ_{AB} can be written in the form

$$\rho_{AB} = \frac{1}{4} \left(\mathbb{1}_A \otimes \mathbb{1}_B + \mathbf{a} \cdot \boldsymbol{\sigma} \otimes \mathbb{1}_B + \mathbb{1}_A \otimes \mathbf{b} \cdot \boldsymbol{\sigma} + \sum_{j,k=1}^3 T_{jk} \sigma_j \otimes \sigma_k \right), \quad (1)$$

where $\boldsymbol{\sigma} \equiv (\sigma_1, \sigma_2, \sigma_3)$ denotes the vector of Pauli spin operators and $\mathbb{1}_A, \mathbb{1}_B$ are identity operators. Here, \mathbf{a} and \mathbf{b} are the Bloch vectors of Alice's and Bob's reduced states and T is the spin correlation matrix, i.e.,

$$\begin{aligned} a_j &:= \text{Tr}[\rho_{AB} \sigma_j \otimes \mathbb{1}_B], & b_k &:= \text{Tr}[\rho_{AB} \mathbb{1}_A \otimes \sigma_k], \\ T_{jk} &:= \text{Tr}[\rho_{AB} \sigma_j \otimes \sigma_k] \quad (j, k = 1, 2, 3). \end{aligned} \quad (2)$$

Different choices of Alice's local measurements result in different *steered* states for Bob. Specifically, each local measurement outcome for Alice corresponds to some element $E \geq 0$ of a positive operator-valued measure (POVM) describing her measurement. Thus, $E = e_0(\mathbb{1}_A + \mathbf{e} \cdot \boldsymbol{\sigma})$, with $e_0 \geq 0$ and $|\mathbf{e}| \leq 1$. This outcome is obtained with probability

$$p^E = \text{Tr}[\rho_{AB} E \otimes \mathbb{1}_B] = e_0(1 + \mathbf{a} \cdot \mathbf{e}),$$

leading to the steered state

$$\rho_B^E = \frac{\text{Tr}_A[\rho_{AB} E \otimes \mathbb{1}_B]}{p^E} = \frac{1}{2} \left[\mathbb{1}_B + \frac{(\mathbf{b} + T^\top \mathbf{e}) \cdot \boldsymbol{\sigma}}{1 + \mathbf{a} \cdot \mathbf{e}} \right] \quad (3)$$

for Bob's qubit.

Considering all possible local measurements by Alice, it follows that the corresponding set of Bob's steered states is represented by the set of Bloch vectors

$$\mathcal{E}_{B|A} = \left\{ \frac{\mathbf{b} + T^\top \mathbf{e}}{1 + \mathbf{a} \cdot \mathbf{e}} : |\mathbf{e}| \leq 1 \right\}. \quad (4)$$

While not immediately apparent from Eq. (4), this set forms a (possibly degenerate) ellipsoid, and hence is called a quantum steering ellipsoid [3]. The subscript $B|A$ indicates the steering ellipsoid for Bob that is generated by Alice's local measurements. Similarly, there is a steering ellipsoid for Alice generated by Bob's local measurements, denoted by $\mathcal{E}_{A|B}$. The ellipsoid $\mathcal{E}_{B|A}$ is fully determined by its center

$$\mathbf{c}_{B|A} = \frac{\mathbf{b} - T^\top \mathbf{a}}{1 - a^2},$$

and its orientation matrix

$$Q_{B|A} = \frac{1}{1 - a^2} (T - \mathbf{a}\mathbf{b}^\top)^\top \left(I + \frac{\mathbf{a}\mathbf{a}^\top}{1 - a^2} \right) (T - \mathbf{a}\mathbf{b}^\top),$$

where the eigenvalues and corresponding eigenvectors of $Q_{B|A}$ describe the squared lengths of the ellipsoid's semiaxes and

their orientations [3]. Here and elsewhere, we use x to denote $|\mathbf{x}|$ for the vector \mathbf{x} .

The quantum steering ellipsoid $\mathcal{E}_{B|A}$, together with the reduced Bloch vectors \mathbf{a} and \mathbf{b} , provides a faithful representation of any 2-qubit state up to local unitary operations on Alice's qubit [3]. Additionally, its various geometric properties encode useful information about the state. For example, the state is separable if and only if its steering ellipsoid can be nested in a tetrahedron that is in turn nested in the Bloch sphere [3].

The size of the steering ellipsoid is quantified by its volume [3]

$$V_{B|A} = \frac{4\pi}{3} \frac{|\det(T - \mathbf{a}\mathbf{b}^\top)|}{(1 - a^2)^2}. \quad (5)$$

It is obvious that the steering ellipsoid is constrained to lie inside the Bloch sphere, implying the volume is always no larger than $V_{\text{unit}} = \frac{4\pi}{3}$. It is therefore convenient to work with the corresponding *normalized volume* defined by

$$v_{B|A} := \frac{V_{B|A}}{V_{\text{unit}}} \leq 1. \quad (6)$$

The upper bound is achieved if and only if Alice and Bob share a pure entangled 2-qubit state [3], and hence the steering ellipsoids of such states coincide with the Bloch ball. In contrast, the normalized volume of any separable state is restricted by the nested tetrahedron condition to be no greater than $\frac{1}{27}$ [3]. Thus, the steering ellipsoid volume is, at least to some extent, connected with the entanglement of the state. This paper will explore this connection further, via volume monogamy relations.

For later reference, we note here that the quantum steering ellipsoid $\mathcal{E}_{B|A}$, and hence its volume $V_{B|A}$, is invariant under the local filtering operation on Alice's qubit defined by [6]

$$\tilde{\rho}_{AB} := [(2\rho_A)^{-1/2} \otimes \mathbb{1}_B] \rho_{AB} [(2\rho_A)^{-1/2} \otimes \mathbb{1}_B]. \quad (7)$$

The filtered state $\tilde{\rho}_{AB}$ is called the canonical form of ρ_{AB} , and has the useful properties $\tilde{\mathcal{E}}_{B|A} = \mathcal{E}_{B|A}$ and $\tilde{\mathbf{a}} = \mathbf{0}$. Thus, for example, the normalized volume of $\mathcal{E}_{B|A}$ can be rewritten in the simple form

$$v_{B|A} = \tilde{v}_{B|A} = |\det \tilde{T}_{AB}|, \quad (8)$$

via Eqs. (5) and (6), where \tilde{T}_{AB} denotes the spin correlation matrix for $\tilde{\rho}_{AB}$. Note that the canonical form is well defined whenever Alice does not have a pure qubit state, i.e., whenever $a \neq 1$. Conversely, for $a = 1$ the shared state factorizes, and hence Bob's steering ellipsoid trivially reduces to the single point $\mathcal{E}_{B|A} = \{\mathbf{b}\}$, with $v_{B|A} = 0$.

III. VOLUME MONOGAMY

A. Pure 3-qubit states

1. Monogamy, quantum marginals, and concurrence

Consider now the scenario in which Alice, Bob, and Charlie share a pure 3-qubit state, $\rho_{ABC} = |\psi_{ABC}\rangle\langle\psi_{ABC}|$, and let $\mathcal{E}_{B|A}$ and $\mathcal{E}_{C|A}$ denote the steering ellipsoids for Bob and Charlie, respectively, generated by local measurements on Alice's qubit. Milne *et al.* have derived the strong volume monogamy relation [6,21]

$$\sqrt{v_{B|A}} + \sqrt{v_{C|A}} \leq 1 \quad (9)$$




FIG. 1. The volume monogamy relation [Eq. (9)] for pure states. The x axis and y axis refer to the normalized volumes $v_{B|A}$ and $v_{C|A}$, respectively. Both fully factorizable states and bipartite entangled states $|\psi_A\rangle|\psi_{BC}\rangle$ are mapped to the origin. Two red points (1,0) and (0,1) correspond to the other two classes of bipartite entangled states $|\psi_{AB}\rangle|\psi_C\rangle$ and $|\psi_{AC}\rangle|\psi_B\rangle$. The blue solid line is $\sqrt{v_{A|B}} + \sqrt{v_{C|B}} = 1$ and represents the W -class states, except for the red points (1,0) and (0,1). The shaded region, including the two axes, represents the GHZ-class states.

for the corresponding normalized volumes.

This relation immediately implies that Alice cannot steer both Bob and Charlie to a large set of states. For example, if Alice is able to steer Bob to the whole Bloch sphere (i.e., they share a pure entangled state), then Charlie’s steering ellipsoid has zero volume (and indeed reduces to a single point). The volume monogamy relation (9) is depicted in Fig. 1. It is nontrivially saturated if and only if $|\psi_{ABC}\rangle$ is a W -class state [6,21] (see also Sec. III A 2 below).

We note here an alternative proof of Eq. (9) to that given in Ref. [21], which does not require consideration of extremal ellipsoid volumes, and which provides an interesting connection between volume monogamy and the quantum marginal problem [19,22]. For $a = 1$ the proof is trivial: Alice’s state is pure and hence the shared state factorizes, implying that $v_{B|A} = v_{C|A} = 0$. Otherwise, for $a \neq 1$ we can apply a local filtering operation similarly to Eq. (7), with $\mathbb{1}_B$ replaced by $\mathbb{1}_{BC}$ and ρ_{AB} replaced by ρ_{ABC} , to obtain the corresponding canonical state $\tilde{\rho}_{ABC}$ with $\tilde{\mathbf{a}} = \mathbf{0}$. Taking partial traces, the normalized volumes of Bob’s and Charlie’s steering ellipsoids follow via Eq. (8) as

$$v_{B|A} = \tilde{v}_{B|A} = |\det \tilde{T}_{AB}| = \tilde{c}^2, \tag{10}$$

$$v_{C|A} = \tilde{v}_{C|A} = |\det \tilde{T}_{AC}| = \tilde{b}^2, \tag{11}$$

respectively, where the final equalities may be proved for pure canonical states by direct calculation, or via partial transposition properties as per Ref. [21]. Finally, we apply

the polygon inequality [19,22]

$$b + c \leq 1 + a, \tag{12}$$

for the Bloch vectors of any pure 3-qubit state, to the canonical state $\tilde{\rho}_{ABC}$, yielding

$$\sqrt{v_{B|A}} + \sqrt{v_{C|A}} = \tilde{c} + \tilde{b} \leq 1 + \tilde{a} = 1$$

as required.

Remarkably, the volume monogamy relation (9) also has a deep connection with entanglement monogamy. In particular, the concurrence of a bipartite state ρ_{AB} has the upper bound [6]

$$C^2(\rho_{AB}) \leq (1 - a^2)\sqrt{v_{B|A}}, \tag{13}$$

which combined with Eq. (9) immediately yields the celebrated CKW inequality [12]

$$C^2(\rho_{AB}) + C^2(\rho_{AC}) \leq (1 - a^2) = 4 \det \rho_A. \tag{14}$$

Thus, volume monogamy is a mathematically stronger condition than the monogamy of concurrence. Note, however, the latter is valid for any 3-qubit state, including mixed states. This is not the case for Eq. (9), as we will show in Sec. III B.

2. Connection to SLOCC classes and 3-tangle

It has been shown in Ref. [23] that any pure 3-qubit state can be classified into one of six entanglement classes, with all members of any one class being interconvertible under SLOCC transformations, i.e., under local operations and classical communication with some nonzero probability. Here, we show that these classes map onto corresponding regions of Fig. 1, thus relating these classes to properties of steering ellipsoid volumes.

We first consider the four classes having no tripartite entanglement. The first of these comprises all fully factorizable states, i.e., those of the form $|\psi_{ABC}\rangle = |\psi_A\rangle|\psi_B\rangle|\psi_C\rangle$. All steering ellipsoids reduce to single points for this class, with zero volumes, and thus it is mapped to the origin in Fig. 1. Next are the three classes of bipartite entangled states, with the corresponding forms $|\psi_A\rangle|\psi_{BC}\rangle$, $|\psi_{AB}\rangle|\psi_C\rangle$, and $|\psi_{AC}\rangle|\psi_B\rangle$. For the first of these, Alice’s qubit is uncorrelated with Bob’s and Charlie’s qubits, and hence the steering ellipsoids $\mathcal{E}_{B|A}$ and $\mathcal{E}_{C|A}$ again have zero volumes, corresponding to the origin in Fig. 1. For the other two bipartite classes, Alice can steer one of Bob and Charlie’s qubits to the entire Bloch sphere, and the other to a single point. Thus, these two classes correspond to the two red dots in Fig. 1, and trivially saturate volume monogamy relation (9).

The remaining two entanglement classes comprise genuine 3-party entangled states. They correspond to states which are convertible, under SLOCC transformations, to either the W state

$$|\psi_{ABC}\rangle = \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle) \tag{15}$$

or to the Greenberger-Horne-Zeilinger (GHZ) state

$$|\psi_{ABC}\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle), \tag{16}$$

and are called the W class and the GHZ class, respectively [23]. These two classes are distinguished by having 3-tangles

$\tau(\rho_{ABC}) = 0$ and $\tau(\rho_{ABC}) > 0$, respectively [23], where for pure states [12]

$$\tau(\rho_{ABC}) := 4 \det \rho_A - C^2(\rho_{AB}) - C^2(\rho_{AC}). \quad (17)$$

As indicated in Fig. 1, and demonstrated below, W -class states are precisely those states which nontrivially saturate the volume monogamy relation, while the GHZ-class states correspond to the entire region below this saturating curve.

It has been shown by Milne *et al.* that the volume monogamy relation is saturated if and only if the state is a ‘‘maximum volume’’ state, with the canonical form

$$|\tilde{\psi}_{ABC}\rangle = (|100\rangle + \cos\theta|010\rangle + \sin\theta|001\rangle)/\sqrt{2} \quad (18)$$

up to local unitary transformations, where $\theta \in [0, \pi/2]$ [6,21]. It is straightforward to check that such states are bipartite entangled states for $\theta = 0, \pi/2$, corresponding to the points $(0,1)$ and $(1,0)$ in Fig. 1, and are W -class states otherwise [23]. Hence, all states nontrivially saturating the volume monogamy relation are W -class states. Conversely, noting that W -class states have zero 3-tangle and $a < 1$ [23], the inequality

$$\tau(\rho_{ABC}) \geq (1 - a^2)(1 - \sqrt{v_{B|A}} - \sqrt{v_{C|A}})$$

following from Eqs. (13) and (17) immediately implies that every W -class state saturates the monogamy relation.

It follows from the above that all GHZ-class states must correspond to points in the region below the saturating curve in Fig. 1. We show that indeed every point in this region, including the axes, corresponds to such a state. In particular, we give a family of canonical GHZ-class states for which the normalized volumes are mapped to every point (x,y) in the shaded area in Fig. 1, including the axes. The explicit form of these states is defined via two real free parameters:

$$|\tilde{\psi}_{ABC}\rangle = \frac{1}{\sqrt{2}}(\sin\alpha|100\rangle + \sin\beta|010\rangle + \cos\beta|001\rangle + \cos\alpha|111\rangle), \quad (19)$$

where $\alpha, \beta \in (0, \pi/2)$, which immediately maps to the coordinates $x = v_{B|A} = \frac{1}{4}(\cos 2\alpha + \cos 2\beta)^2$ and $y = v_{C|A} = \frac{1}{4}(\cos 2\alpha - \cos 2\beta)^2$ via Eqs. (10) and (11). These fill the shaded area because any point (x,y) therein corresponds to either $2\alpha = \arccos(\sqrt{x} + \sqrt{y})$ and $2\beta = \arccos(\sqrt{x} - \sqrt{y})$ or $2\alpha = \arccos(\sqrt{x} - \sqrt{y})$ and $2\beta = \arccos(\sqrt{x} + \sqrt{y})$.

B. Mixed 3-qubit states

A natural question to consider is whether the volume monogamy relation (9) is also valid for a general mixed 3-qubit state. Unfortunately, the answer is negative. A counterexample is given by

$$\rho_{ABC} = \frac{1}{2}(|\chi_1\rangle\langle\chi_1| + |\chi_2\rangle\langle\chi_2|), \quad (20)$$

with

$$|\chi_1\rangle = \frac{1}{\sqrt{6}}(|101\rangle - 2|011\rangle + |110\rangle),$$

$$|\chi_2\rangle = \frac{1}{\sqrt{6}}(|010\rangle - 2|100\rangle + |001\rangle).$$

The 2-qubit reduced states ρ_{AB} and ρ_{AC} are then identical Werner states, of the form

$$\frac{2}{3}|\psi^-\rangle\langle\psi^-| + \frac{1}{3}\frac{\mathbb{1}}{2} \otimes \frac{\mathbb{1}}{2}, \quad (21)$$

where $|\psi^-\rangle$ denotes the singlet state $(|01\rangle - |10\rangle)/\sqrt{2}$. It is easy to verify that the corresponding steering ellipsoids are spheres of radius $\frac{2}{3}$ [3]. Hence, $\sqrt{v_{B|A}} + \sqrt{v_{C|A}} = 2\sqrt{8/27} = 1.0888 > 1$, implying that the volume monogamy relation (9) does not hold for all mixed 3-qubit states.

In Sec. IV, we will show that Eq. (9) does remain valid for the particular case of mixed states obtained via local operations on pure 3-qubit states. Here, however, we will derive a volume monogamy relation that is valid for *all* 3-qubit states:

$$(v_{B|A})^{2/3} + (v_{C|A})^{2/3} \leq 1. \quad (22)$$

This monogamy relation is clearly weaker than Eq. (9) for pure states. However, it is saturated in some cases: for example, when $\rho_{ABC} = \rho_{AB} \otimes \rho_C$ where ρ_{AB} is a pure entangled state, in which case $v_{B|A} = 1$ and $v_{C|A} = 0$.

Our derivation of Eq. (22) is based on a relatively strong tradeoff relation for the pairwise spin correlations of any 3-qubit state ρ_{ABC} ,

$$\text{Tr}[T_{AB}^\top T_{AB}] + \text{Tr}[T_{AC}^\top T_{AC}] + \text{Tr}[T_{BC}^\top T_{BC}] \leq 3 \quad (23)$$

(with equality for pure states), obtained as follows. First, consider a pure state ρ_{ABC} . From the Schmidt decomposition, the purities of any bipartition of such a state satisfy $\text{Tr}[\rho_{AB}^2] = \text{Tr}[\rho_C^2]$, $\text{Tr}[\rho_{AC}^2] = \text{Tr}[\rho_B^2]$, and $\text{Tr}[\rho_{BC}^2] = \text{Tr}[\rho_A^2]$, which from the definitions in Eq. (2) are equivalent to

$$\begin{aligned} \text{Tr}[T_{AB}^\top T_{AB}] + a^2 + b^2 &= 1 + 2c^2, \\ \text{Tr}[T_{AC}^\top T_{AC}] + a^2 + c^2 &= 1 + 2b^2, \\ \text{Tr}[T_{BC}^\top T_{BC}] + b^2 + c^2 &= 1 + 2a^2. \end{aligned}$$

Summing these three equations immediately leads to the identity

$$\text{Tr}[T_{AB}^\top T_{AB}] + \text{Tr}[T_{AC}^\top T_{AC}] + \text{Tr}[T_{BC}^\top T_{BC}] = 3 \quad (24)$$

for pure 3-qubit states. Second, expressing a mixed state ρ_{ABC} as a convex combination of pure states, $\rho_{ABC} = \sum_m p_m |\varphi_m\rangle\langle\varphi_m|$, and letting T_{AB}^m denote the spin correlation matrix of $\text{Tr}_C |\varphi_m\rangle\langle\varphi_m|$, we have

$$\begin{aligned} \text{Tr}[T_{AB}^\top T_{AB}] &= \sum_{m,n} p_m p_n \text{Tr}[(T_{AB}^m)^\top T_{AB}^n] \\ &\leq \sum_{m,n} p_m p_n \{ \text{Tr}[(T_{AB}^m)^\top T_{AB}^m] \text{Tr}[(T_{AB}^n)^\top T_{AB}^n] \}^{1/2} \\ &\leq \frac{1}{2} \sum_{m,n} p_m p_n \{ \text{Tr}[(T_{AB}^m)^\top T_{AB}^m] + \text{Tr}[(T_{AB}^n)^\top T_{AB}^n] \} \\ &= \sum_m p_m \text{Tr}[(T_{AB}^m)^\top T_{AB}^m]. \end{aligned} \quad (25)$$

Here, the first inequality follows from the Schwarz inequality, and the second from the geometric mean being no greater than the arithmetic mean. One has similar relations for $\text{Tr}[T_{AC}^\top T_{AC}]$ and $\text{Tr}[T_{BC}^\top T_{BC}]$. Summing these and using identity (24) for pure states then yields Eq. (23) as required.

We note that Eq. (23) can itself be considered as a monogamy relation, for the strengths of the pairwise spin correlations, and subsumes the known monogamy relation for pairwise Bell nonlocality [24].

Similarly to the proof of Eq. (9) in the previous section, we now consider the canonical state $\tilde{\rho}_{ABC}$ for $a \neq 1$ [since Eq. (22) is similarly trivially satisfied when $a = 1$]. Equation (8) then yields

$$\begin{aligned} & (v_{B|A})^{2/3} + (v_{C|A})^{2/3} \\ &= |\det[(\tilde{T}_{AB})^\top \tilde{T}_{AB}]|^{1/3} + |\det[(\tilde{T}_{AC})^\top \tilde{T}_{AC}]|^{1/3} \\ &\leq \frac{1}{3} \text{Tr}[(\tilde{T}_{AB})^\top \tilde{T}_{AB}] + \frac{1}{3} \text{Tr}[(\tilde{T}_{AC})^\top \tilde{T}_{AC}] \\ &\leq 1 \end{aligned} \tag{26}$$

as desired. Here, the first inequality follows from the arithmetic-geometric mean inequality, applied to the eigenvalues of $(\tilde{T}_{AB})^\top \tilde{T}_{AB}$ and $(\tilde{T}_{AC})^\top \tilde{T}_{AC}$, and the second inequality from the tradeoff relation in Eq. (23).

In the next section, we show that a monogamy relation of the same form (involving $\frac{2}{3}$ powers) also holds for pure 4-qubit states.

C. 4-qubit states

We first remark that the strong monogamy relation (9) for pure 3-qubit states cannot be generalized to a similar form for pure n -qubit states, for any $n \geq 4$. In particular, by purifying the counterexample in Eq. (20), we can construct the pure 4-qubit state

$$|\psi\rangle_{ABCD} = \frac{1}{\sqrt{2}}(|\chi_1\rangle_{ABC}|0\rangle_D + |\chi_2\rangle_{ABC}|1\rangle_D), \tag{27}$$

implying, similarly to the counterexample, that $\sqrt{v_{B|A}} + \sqrt{v_{C|A}} + \sqrt{v_{D|A}} = 2\sqrt{8/27} + \sqrt{v_{D|A}} > 1$. It follows more generally, by considering an n -qubit pure state with factor $|\psi\rangle_{ABCD}$, that the form of Eq. (9) does not generalize to any $n \geq 4$.

We will show here, however, that the volume monogamy relation

$$(v_{B|A})^{2/3} + (v_{C|A})^{2/3} + (v_{D|A})^{2/3} \leq 1 \tag{28}$$

is valid for any pure 4-qubit state ρ_{ABCD} , and give numerical evidence strongly supporting its validity for mixed 4-qubit states.

To prove result (28), we adapt the techniques used in the proof of Eq. (22) in Sec. III B. First, using the equalities of purities of any bipartition of a pure state, we have

$$a^2 + b^2 + \text{Tr}[T_{AB}^\top T_{AB}] = c^2 + d^2 + \text{Tr}[T_{CD}^\top T_{CD}], \tag{29}$$

$$a^2 + c^2 + \text{Tr}[T_{AC}^\top T_{AC}] = b^2 + d^2 + \text{Tr}[T_{BD}^\top T_{BD}], \tag{30}$$

$$a^2 + d^2 + \text{Tr}[T_{AD}^\top T_{AD}] = b^2 + c^2 + \text{Tr}[T_{BC}^\top T_{BC}], \tag{31}$$

with respect to the bipartitions (AB, CD) , (AC, BD) , and (AD, BC) . We also have, for the bipartition (A, BCD) ,

$$\begin{aligned} & b^2 + c^2 + d^2 + \text{Tr}[T_{BC}^\top T_{BC} + T_{BD}^\top T_{BD} + T_{CD}^\top T_{CD}] \\ &+ L_{BCD} = 3 + 4a^2, \end{aligned} \tag{32}$$

where

$$L_{BCD} := \sum_{l,m,n} (\text{Tr}[\mathbb{1} \otimes \sigma_l \otimes \sigma_m \otimes \sigma_n \rho_{ABCD}])^2 \tag{33}$$

is a measure of the tripartite spin correlation strength between Bob, Charlie, and Dianne. Summing Eqs. (29)–(32) yields

$$\begin{aligned} & \text{Tr}[T_{AB}^\top T_{AB}] + \text{Tr}[T_{AC}^\top T_{AC}] + \text{Tr}[T_{AD}^\top T_{AD}] \\ &= 3 + a^2 - L_{BCD} \leq 3 + a^2. \end{aligned} \tag{34}$$

Applying a local filtering operation similarly to Eq. (7), with $\mathbb{1}_B$ replaced by $\mathbb{1}_{BCD}$ and ρ_{AB} replaced by ρ_{ABCD} , to obtain the corresponding canonical state $\tilde{\rho}_{ABCD}$, we have $\tilde{\mathbf{a}} = \mathbf{0}$ and hence that

$$\text{Tr}[(\tilde{T}_{AB})^\top \tilde{T}_{AB}] + \text{Tr}[(\tilde{T}_{AC})^\top \tilde{T}_{AC}] + \text{Tr}[(\tilde{T}_{AD})^\top \tilde{T}_{AD}] \leq 3. \tag{35}$$

Equation (28) then follows via the same arguments used in the derivation of Eq. (26).

Finally, we conjecture that inequality (35) generalizes to

$$\text{Tr}[T_{AB}^\top T_{AB}] + \text{Tr}[T_{AC}^\top T_{AC}] + \text{Tr}[T_{AD}^\top T_{AD}] \leq 3 \tag{36}$$

for all pure 4-qubit states. We have employed numerical simulations to generate 2×10^5 random pure states and find no violation of inequality (36). The validity of this conjecture would imply, using the same techniques as above, that monogamy relation (28) in fact holds for *all* 4-qubit states.

D. Multiqubit states

We now obtain a general volume monogamy relation for n -qubit states, pure or mixed, based on Eq. (22) for 3-qubit states. In particular, for an n -qubit state $\rho_{ABCD\dots}$ consider the normalized volumes $v_{B|A}, v_{C|A}, v_{D|A}, \dots$ of the steering ellipsoids generated by Alice's local measurements. The steered parties B, C, D, \dots can be grouped into $\frac{1}{2}(n-1)$ $(n-2)$ distinct pairs, with each pair satisfying a volume monogamy relation as per Eq. (22). Summing these relations over all such pairs and rearranging terms then yields the general monogamy relation

$$(v_{B|A})^{2/3} + (v_{C|A})^{2/3} + (v_{D|A})^{2/3} + \dots \leq \frac{n-1}{2}. \tag{37}$$

This reduces to the 3-qubit relation for $n = 3$, and in general places a nontrivial constraint on the degree to which Alice can steer the states of the other parties. For example, noting that $v \leq v^{2/3}$ for $v \leq 1$, it follows that the average volume of the $n-1$ ellipsoids to which Alice can steer the other parties is bounded by

$$\bar{v}_{|A} := \frac{v_{B|A} + v_{C|A} + v_{D|A} + \dots}{n-1} \leq \frac{1}{2}. \tag{38}$$

For the 4-qubit case, the upper bound in Eq. (37) is $\frac{3}{2}$. While this is weaker than the upper bound of 1 in Eq. (28) for pure 4-qubit states, it has the advantage of also being valid for the mixed case.

IV. VOLUME MONOGAMY AND NOISE

A. Local noise and steering ellipsoids

Taking into account the imperfections of any experiment, including in state preparation and measurement, the quantum state is inevitably exposed to all kinds of noise. Such noise processes can be modeled as a noisy channel acting on an ideal initial state. We are interested in the problem of how such channels affect the desired properties of the initial state, and in particular the steering ellipsoid.

Mathematically, a noisy channel acting on a bipartite state ρ_{AB} is equivalent to a completely positive and trace-preserving (CPTP) map, Φ , mapping the initial state ρ_{AB} to some ρ'_{AB} . Here, we consider the case that the noise acts locally on each subsystem. Thus, $\Phi = \phi_A \otimes \phi_B$ where ϕ_A and ϕ_B are CPTP maps acting on A and B , respectively, and

$$\rho'_{AB} = \Phi(\rho_{AB}) = (\phi_A \otimes \phi_B)(\rho_{AB}). \quad (39)$$

The set of reduced states generated by Alice's local measurements on ρ'_{AB} follows from Eq. (3) of Sec. II as

$$\begin{aligned} \{\rho'_B{}^E\} &= \left\{ \frac{\text{Tr}_A[(\phi_A \otimes \phi_B)(\rho_{AB}) E \otimes \mathbb{1}_B]}{\text{Tr}[(\phi_A \otimes \phi_B)(\rho_{AB}) E \otimes \mathbb{1}_B]} \right\} \\ &= \left\{ \frac{\text{Tr}_A[(I_A \otimes \phi_B)(\rho_{AB}) \phi_A^*(E) \otimes \mathbb{1}_B]}{\text{Tr}[(I_A \otimes \phi_B)(\rho_{AB}) \phi_A^*(E) \otimes \mathbb{1}_B]} \right\}, \end{aligned}$$

where E ranges over all positive operators, I denotes the identity map, and the dual map ϕ^* of any CP map ϕ is defined by $\text{Tr}[\phi^*(X)Y] := \text{Tr}[X\phi(Y)]$. Noting ϕ_B is trace preserving and that ϕ_A^* maps positive operators to positive operators then yields

$$\begin{aligned} \{\rho'_B{}^E\} &= \left\{ \frac{\phi_B(\text{Tr}_A[\rho_{AB} \phi_A^*(E) \otimes \mathbb{1}_B])}{\text{Tr}[\rho_{AB} \phi_A^*(E) \otimes \mathbb{1}_B]} \right\} \\ &\subseteq \left\{ \frac{\phi_B(\text{Tr}_A[\rho_{AB} E \otimes \mathbb{1}_B])}{\text{Tr}[\rho_{AB} E \otimes \mathbb{1}_B]} \right\} \\ &= \left\{ \phi_B \left[\frac{\text{Tr}_A[\rho_{AB} E \otimes \mathbb{1}_B]}{\text{Tr}[\rho_{AB} E \otimes \mathbb{1}_B]} \right] \right\}. \end{aligned}$$

Hence, the steering ellipsoids of ρ'_{AB} and ρ_{AB} are related by

$$\mathcal{E}'_{B|A} \subseteq \phi_B(\mathcal{E}_{B|A}). \quad (40)$$

To determine how the volumes of the steering ellipsoids are related, note that the trace distance between two states contracts under any CPTP map ϕ [1]. Moreover, for qubits, the trace distance is proportional to the Euclidean distance in the Bloch ball [1]. As a consequence, the volume of any set of qubit Bloch vectors contracts under CPTP maps, yielding the inequality chain

$$V(\mathcal{E}'_{B|A}) \leq V(\phi_B(\mathcal{E}_{B|A})) \leq V(\mathcal{E}_{B|A}) \quad (41)$$

via Eq. (40). Thus, local noise never increases the volume of the steering ellipsoid.

An immediate consequence of this result is that any volume monogamy relation, for a given set of multiqubit states, will remain valid under the addition of local noise. For example, it follows via Eq. (9) that

$$\sqrt{v'_{B|A}} + \sqrt{v'_{C|A}} \leq 1 \quad (42)$$




FIG. 2. The effect of local noise on the left-hand side of the volume monogamy relation [Eq. (9)] for the family of states in Eq. (43). Here, the noise on each qubit is a depolarizing channel with strength $\varepsilon = 0$ (black solid curve), 0.001 (red solid curve), 0.005 (blue dashed curve), 0.01 (purple dotted curve).

for any state obtained by adding local noise to a pure 3-qubit state, i.e., for any 3-qubit state of the form $\rho'_{ABC} = (\phi_A \otimes \phi_B \otimes \phi_C)(|\psi_{ABC}\rangle\langle\psi_{ABC}|)$.

B. An example: Local isotropic noise

We consider the set of states

$$|\varphi_{ABC}\rangle = p|100\rangle + \sqrt{\frac{1-p^2}{2}}|010\rangle + \sqrt{\frac{1-p^2}{2}}|001\rangle, \quad (43)$$

with $p \in (0,1)$. These states are symmetric with respect to Bob and Charlie, so that $v_{B|A} = v_{C|A}$. Moreover, it is easy to verify that these are W -class states [23]. Thus, they saturate the volume monogamy relation in Eq. (9) (see Sec. III B), making them of experimental interest.

However, the inevitable presence of noise in any experiment means that in practice these states cannot be perfectly generated and measured. We therefore investigate the robustness of these states under a simple noise model. In particular, we consider the addition of local isotropic noise

$$\rho'_{ABC} = \Phi_\varepsilon(\rho_{ABC}) := (\phi_\varepsilon \otimes \phi_\varepsilon \otimes \phi_\varepsilon)(\rho_{ABC}), \quad (44)$$

where

$$\phi_\varepsilon : \rho \rightarrow \frac{\varepsilon}{2}\mathbb{1} + (1-\varepsilon)\rho \quad (45)$$

corresponds to adding isotropic noise of strength $\varepsilon \in [0,1]$. Thus, for each qubit, $\varepsilon = 0$ corresponds to no noise, while $\varepsilon = 1$ corresponds to noise so strong that the state becomes completely mixed. The Bloch vector of each qubit is scaled by $1-\varepsilon$.

The noisy channel Φ_ε preserves the symmetry of the state ρ_{ABC} with respect to Bob and Charlie, and the volumes of the steering ellipsoid may be analytically calculated for the states in Eq. (43) as

$$v'_{B|A} = v'_{C|A} = \frac{4p^4(1-p^2)^2(1-\varepsilon)^6}{[1-(1-\varepsilon)^2(1-2p^2)]^2}. \quad (46)$$

The corresponding sensitivity of the volume monogamy relation (9) to noise is depicted in Fig. 2, for a range of

experimentally relevant noise strengths. It is seen that while the relation is no longer saturated for $\varepsilon > 0$, those states with p taking values in the mid-range of the unit interval are relatively robust.

V. CONCLUSION

We have studied volume monogamy relations for multiqubit systems. We have demonstrated a close connection between the volume monogamy relation (9) for pure 3-qubit states and the SLOCC classification of such states. A counterexample (20) was constructed to show this relation does not generalize to all 3-qubit states, and a suitable universal volume monogamy relation (22) was obtained for the general case. A similar relation was obtained in Eq. (28) for pure 4-qubit states, and conjectured to also hold for mixed states. Furthermore, we have found a generalized volume monogamy relation valid for all multiqubit states. Finally, we studied the effects of noise on the quantum steering ellipsoid and showed that local noise channels do not invalidate the volume monogamy relation, as such noise decreases the volume of steering ellipsoids.

More generally, it is remarkable that the simple concept of the steering ellipsoid, i.e., the set of Bob's local states that Alice can prepare by local measurements on her system, can geometrically capture many important aspects of quantum correlations and information processing tasks. Properties of steering ellipsoids are not only strongly connected to quantum monogamy, as investigated here, but have also been closely

linked with quantum communication protocols based on, for example, Bell nonlocality [7], teleportation [7], and Einstein-Podolsky-Rosen steering [8,11].

It is hoped that further investigation of steering ellipsoids (and their generalizations to higher dimensions) will cement their relevance to understanding the properties and usefulness of quantum correlations. This goes well beyond properties of ellipsoid volumes: for example, while local dissipation reduces volumes as per Sec. IV, it is also known that such noise can enhance teleportation fidelity for some states [25]. Hence, a geometric characterization of such states will require consideration of aspects other than volume (such as semiaxis lengths [7]).

Many open questions remain for future work even within the confines of volume monogamy relations. For example, can stronger monogamy relations than Eqs. (22) and (28) be obtained? Is the 4-qubit conjecture in Eq. (36) valid? Is there some underlying connection between volume monogamy relations and that of other types of entanglement [26–28]? How close might we get to these bounds with experiments? Finally, can volume monogamy be generalized to higher-dimensional systems?

ACKNOWLEDGMENTS

We thank J. Chen for his assistance in finding the counterexample (20). S.C. thanks L. Li for helpful discussions. This work was supported by the ARC Centre of Excellence Grant No. CE110001027. A.M. is supported by the EPSRC.

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2010).
 - [2] H. M. Wiseman and G. J. Milburn, *Quantum Measurement and Control* (Cambridge University Press, Cambridge, UK, 2009).
 - [3] S. Jevtic, M. Pusey, D. Jennings, and T. Rudolph, *Phys. Rev. Lett.* **113**, 020402 (2014).
 - [4] M. Shi, F. Jiang, C. Sun, and J. Du, *New J. Phys.* **13**, 073016 (2011).
 - [5] M. Shi, C. Sun, F. Jiang, X. Yan, and J. Du, *Phys. Rev. A* **85**, 064104 (2012).
 - [6] A. Milne, S. Jevtic, D. Jennings, H. Wiseman, and T. Rudolph, *New J. Phys.* **16**, 083017 (2014).
 - [7] A. Milne, D. Jennings, S. Jevtic, and T. Rudolph, *Phys. Rev. A* **90**, 024302 (2014).
 - [8] S. Jevtic, M. J. W. Hall, M. R. Anderson, M. Zwierz, and H. M. Wiseman, *J. Opt. Soc. Am. B* **32**, A40 (2015).
 - [9] Q. Quan, H. Zhu, S.-Y. Liu, S.-M. Fei, H. Fan, and W.-L. Yang, *Sci. Rep.* **6**, 22025 (2016).
 - [10] H. C. Nguyen and T. Vu, *Phys. Rev. A* **94**, 012114 (2016).
 - [11] H. C. Nguyen and T. Vu, *Europhys. Lett.* **115**, 10003 (2016).
 - [12] V. Coffman, J. Kundu, and W. K. Wootters, *Phys. Rev. A* **61**, 052306 (2000).
 - [13] T. J. Osborne and F. Verstraete, *Phys. Rev. Lett.* **96**, 220503 (2006).
 - [14] V. Scarani and N. Gisin, *Phys. Rev. Lett.* **87**, 117901 (2001).
 - [15] B. Toner and F. Verstraete, [arXiv:quant-ph/0611001](https://arxiv.org/abs/quant-ph/0611001)
 - [16] B. Toner, *Proc. R. Soc. London A* **465**, 59 (2009).
 - [17] P. Kurzyński, T. Paterek, R. Ramanathan, W. Laskowski, and D. Kaszlikowski, *Phys. Rev. Lett.* **106**, 180402 (2011).
 - [18] M. D. Reid, *Phys. Rev. A* **88**, 062108 (2013).
 - [19] A. Klyachko, [arXiv:quant-ph/0409113](https://arxiv.org/abs/quant-ph/0409113).
 - [20] X. Hu and H. Fan, *Phys. Rev. A* **91**, 022301 (2015).
 - [21] A. Milne, S. Jevtic, D. Jennings, H. Wiseman, and T. Rudolph, *New J. Phys.* **17**, 019501 (2015).
 - [22] A. Higuchi, A. Sudbery, and J. Szulc, *Phys. Rev. Lett.* **90**, 107902 (2003).
 - [23] W. Dür, G. Vidal, and J. I. Cirac, *Phys. Rev. A* **62**, 062314 (2000).
 - [24] H.-H. Qin, S.-M. Fei, and X. Li-Jost, *Phys. Rev. A* **92**, 062339 (2015).
 - [25] P. Badziąg, M. Horodecki, P. Horodecki, and R. Horodecki, *Phys. Rev. A* **62**, 012311 (2000).
 - [26] B. Regula, S. Di Martino, S. Lee, and G. Adesso, *Phys. Rev. Lett.* **113**, 110501 (2014).
 - [27] Y.-K. Bai, Y.-F. Xu, and Z. D. Wang, *Phys. Rev. Lett.* **113**, 100503 (2014).
 - [28] C. Eltschka and J. Siewert, *Phys. Rev. Lett.* **114**, 140402 (2015).