

Spectral theory of Liouvillians for dissipative phase transitions

Fabrizio Minganti, Alberto Biella, Nicola Bartolo, and Cristiano Ciuti*

Laboratoire Matériaux et Phénomènes Quantiques, Université Paris Diderot, CNRS-UMR7162, 75013 Paris, France



(Received 30 April 2018; revised manuscript received 20 July 2018; published 17 October 2018)

A state of an open quantum system is described by a density matrix, whose dynamics is governed by a Liouvillian superoperator. Within a general framework, we explore fundamental properties of both first-order dissipative phase transitions and second-order dissipative phase transitions associated with a symmetry breaking. In the critical region, we determine the general form of the steady-state density matrix and of the Liouvillian eigenmatrix whose eigenvalue defines the Liouvillian spectral gap. We illustrate our exact results by studying some paradigmatic quantum optical models exhibiting critical behavior.

DOI: [10.1103/PhysRevA.98.042118](https://doi.org/10.1103/PhysRevA.98.042118)

I. INTRODUCTION AND MOTIVATION

In classical physics, phase transitions are driven by a competition between the value of the system energy and the entropy produced by its thermal fluctuations [1]. A quantum system at zero temperature has zero entropy and is in its ground state, which is the state minimizing the system energy [2]. However, critical phenomena can occur in the thermodynamic limit as a result of the competition between noncommuting terms of the Hamiltonian.

Driven-dissipative systems have an intrinsic nonequilibrium nature and the properties of the stationary state of the system cannot be determined via a free energy analysis [3–7]. The statistical mechanics of such systems can be remarkably rich. For example, classical systems [8] can display long-range order in 2D [9], since their driven-diffusive nature can violate the Mermin-Wagner theorem [10], which is valid at equilibrium. At a quantum level, by properly designing the coupling with the environment and the driving mechanisms, it is possible to stabilize phases without an equilibrium counterpart [11,12].

The impressive experimental advances of the last decade provide the opportunity to explore nonequilibrium critical phenomena on a variety of platforms. Lattices of superconducting resonators [13,14], Rydberg atoms in optical lattices [15,16], optomechanical systems [17,18], and exciton-polariton condensates [19,20] provide a highly controllable playground in which to study the emergence of dissipative phase transitions. In the thermodynamic limit, the competition between Hamiltonian evolution, pumping, and dissipation processes can trigger a nonanalytical change in the steady state [21]. The engineering of complex many-body phases has been deeply explored in different contexts [22,23]. Dissipative phase transitions have been discussed theoretically for photonic systems [24–36], lossy polariton condensates [37–39], and spin models [11,12,21,40–45].

The interplay between classical and quantum fluctuations in triggering nonequilibrium phase transitions has been

addressed by different methods, including renormalization group approaches based on the Keldysh formalism [37,46,47] and via extensive numerical analysis of lattice systems [43,48,49]. Very recently, the critical properties have been investigated also experimentally in single superconducting cavities [50], semiconductor micropillars [51,52], and large arrays of microwave cavities [14]. Our understanding of criticality in such systems, however, is still in its infancy, since their description cannot be traced back to the traditional framework of equilibrium statistical mechanics.

One of the first investigations in this direction was reported by Kessler *et al.* [21]. They considered a specific spin model, conjecturing some general properties of dissipative phase transitions. However, a general theory connecting Liouvillian spectral properties and dissipative phase transitions is still lacking. In this work, we wish to provide a common theoretical framework to describe the emergence of critical behavior in Markovian open quantum systems, analyzing both first- and second-order phase transitions. We show the general form of the steady-state density matrix in the vicinity of the critical point. We determine also the form of the eigenmatrix of the Liouvillian superoperator corresponding to the nonzero eigenvalue with the smallest modulus of the real part (the so-called Liouvillian spectral gap or asymptotic decay rate). When the transition is of the first order, we show that the gap closes only at the critical point, where the stationary state is bimodal. Concerning second-order phase transitions associated with a symmetry breaking, we provide a general spectral description proving that the Liouvillian gap remains closed in the whole region of broken symmetry. In this context, we highlight the connection between the structure of the eigenmatrices and the symmetry properties of the Lindblad master equation. Note that according to the theory presented in this work, when the Liouvillian gap closes, also the imaginary part of the corresponding eigenvalue must vanish. This is a more stringent constraint with respect to that discussed in [21], where only the real part is assumed to vanish. Particular attention is devoted to the connection between our results and their relation to mean-field solutions, bistability phenomena, and metastability. We bring under a common paradigm apparently different phenomena related to dissipative phase transitions

*cristiano.ciuti@univ-paris-diderot.fr

which have been observed experimentally [14,50,52] and predicted theoretically [12,48,49,53] for specific models. We remark that our results are model independent. One of the goals of the present work is to identify a general spectral mechanism which can explain these phenomena regardless of the nature of the system (bosons, fermions, or spins) and dimensionality.

In order to better illustrate our general results, we analyze some specific paradigmatic cases of linearly [54] and quadratically driven [29,55,56] Kerr resonators. Those models are known to undergo a phase transition (of first [31] and second order [29], respectively) in the thermodynamic limit of large excitation numbers.

This paper is structured as follows. In Sec. II we introduce the theoretical framework, pointing out some general key properties of the Liouvillian superoperator and of density matrices. In Sec. III and Sec. IV we consider, respectively, first- and second-order dissipative phase transitions. Section V is devoted to the numerical study of the two paradigmatic examples mentioned above. Finally, in Sec. VI we draw our conclusions and discuss possible perspectives for future studies. In Appendix A we include the proofs of some useful lemmas, while in Appendix B we consider an exactly solvable model presenting a nondiagonalizable Liouvillian.

II. THEORETICAL FRAMEWORK

In this work, we will consider open quantum systems where the coupling to a reservoir leads to a Markovian dynamics for the system density matrix $\hat{\rho}(t)$, described by a master equation in the Lindblad form [57]

$$\partial_t \hat{\rho}(t) = -\frac{i}{\hbar} [\hat{H}, \hat{\rho}(t)] + \sum_i \frac{\gamma_i}{2} \mathcal{D}[\hat{\Gamma}_i] \hat{\rho}(t), \quad (1)$$

where \hat{H} is the Hamiltonian describing the unitary evolution of the system, while the dissipation superoperators $\mathcal{D}[\hat{\Gamma}_i]$ are defined as

$$\mathcal{D}[\hat{\Gamma}_i] \hat{\rho}(t) = 2\hat{\Gamma}_i \hat{\rho}(t) \hat{\Gamma}_i^\dagger - \hat{\Gamma}_i^\dagger \hat{\Gamma}_i \hat{\rho}(t) - \hat{\rho}(t) \hat{\Gamma}_i^\dagger \hat{\Gamma}_i. \quad (2)$$

Each quantum jump operator $\hat{\Gamma}_i$ is associated with a dissipation channel occurring at the rate γ_i . In the following, we will consider the case where \hat{H} , $\hat{\Gamma}_i$, and γ_i are time independent (for each i). This kind of master equation can be applied for example to photonic quantum systems (see, for example, Refs. [13,14,50–52]).

Since the Lindblad master equation (1) is linear in $\hat{\rho}$, it is possible to associate with it the so-called Liouvillian superoperator \mathcal{L} , defined as

$$\partial_t \hat{\rho}(t) = \mathcal{L} \hat{\rho}(t). \quad (3)$$

The superoperator \mathcal{L} of the Lindblad master equation is trace preserving and generates a completely positive map $e^{\mathcal{L}t}$ describing the time evolution of the system [3–5,58]. For a time-independent Liouvillian, there is at least one steady state (if the dimension of the Hilbert space is finite [58]), i.e., a matrix such that

$$\mathcal{L} \hat{\rho}_{ss} = 0. \quad (4)$$

This equation means that the steady-state density matrix is an eigenmatrix of the superoperator \mathcal{L} corresponding to the zero eigenvalue. Moreover, under quite general conditions (see Refs. [59,60] and Appendix A 1), the steady state is unique. As we will see below, dissipative phase transitions are strictly related to the violation of this unicity condition.

Let us call \mathcal{H} the Hilbert space of the system. A density matrix $\hat{\rho}$, as any other operator $\hat{\xi}$, belongs to the operator space $\mathcal{H} \otimes \mathcal{H}$. The Liouvillian superoperator, instead, is $\mathcal{L} \in L = (\mathcal{H} \otimes \mathcal{H})^* \otimes (\mathcal{H} \otimes \mathcal{H})$, where L is the Liouville space. In this article, we will systematically adopt the following notation: operators will be denoted by hats (e.g., \hat{A}), superoperators will be written in calligraphic characters (e.g., \mathcal{A}), and states and their duals will be expressed in the Dirac notation ($|a\rangle$ and $\langle a|$). A vectorized representation of an operator \hat{A} will be denoted by \vec{A} , while the matrix representing a superoperator \mathcal{A} is indicated by $\vec{\vec{A}}$. In particular, the matrix representation of the Liouvillian is

$$\vec{\vec{L}} = -i[(\hat{H} \otimes \mathbb{1}) - (\mathbb{1} \otimes \hat{H}^{\text{TR}})] + \frac{\gamma}{2}(2\hat{\Gamma} \otimes \hat{\Gamma}^* - \hat{\Gamma}^\dagger \hat{\Gamma} \otimes \mathbb{1} - \mathbb{1} \otimes \hat{\Gamma}^{\text{TR}} \hat{\Gamma}^*), \quad (5)$$

where the superscript TR denotes the transposition. Moreover, we will introduce the Hilbert-Schmidt inner product

$$\langle \hat{A}, \hat{B} \rangle = \text{Tr}[\hat{A}^\dagger \hat{B}], \quad (6)$$

which, in the vectorized representation, takes the intuitive form $\vec{A} \cdot \vec{B} = \sum_{\mu,\nu} (\vec{A})_{\mu,\nu} (\vec{B})_{\mu,\nu}$. The definition of the norm naturally follows as

$$\|\hat{A}\|^2 = \text{Tr}[\hat{A}^\dagger \hat{A}] = \vec{A} \cdot \vec{A} = \sum_{\mu,\nu} |\hat{A}_{\mu,\nu}|^2. \quad (7)$$

A. Spectral properties of Liouvillian superoperators

To fully determine the dynamics of the system, the knowledge of the steady-state density matrix $\hat{\rho}_{ss}$ is not enough. Indeed, one has to know all the spectrum of the Liouvillian superoperator \mathcal{L} , whose eigenmatrices and eigenvalues are defined via the relation

$$\mathcal{L} \hat{\rho}_i = \lambda_i \hat{\rho}_i. \quad (8)$$

Equivalently, in the vector representation, $\vec{\rho}_i$ is a right eigenvector of the superoperator matrix $\vec{\vec{L}}$. Having introduced a norm, we require the eigenstates to be normalized: $\|\hat{\rho}_i\|^2 = 1$ [61]. Since \mathcal{L} is not Hermitian, its eigenvectors are, in general, not orthogonal: $\vec{\rho}_i \cdot \vec{\rho}_j \neq 0$. If the Liouvillian is diagonalizable, we can conveniently use the eigenstates of \mathcal{L} as a basis of the Liouville space, apart from some exceptional points (see Appendix A 1 and Ref. [62]). Under this hypothesis, for any operator \hat{A} there exists a unique decomposition

$$\hat{A} = \sum_i c_i \hat{\rho}_i. \quad (9)$$

It can be proved [57,58] that $\text{Re}[\lambda_i] \leq 0, \forall i$. Since the real part of the eigenvalues is responsible for the relaxation to the steady state, $\hat{\rho}_{ss} = \lim_{t \rightarrow +\infty} e^{\mathcal{L}t} \rho(0)$. For convenience, we sort the eigenvalues in such a way that $|\text{Re}[\lambda_0]| < |\text{Re}[\lambda_1]| < \dots < |\text{Re}[\lambda_n]|$. From this definition it follows that $\lambda_0 = 0$ and $\hat{\rho}_{ss} = \hat{\rho}_0 / \text{Tr}[\hat{\rho}_0]$. We can also identify another relevant

quantity: the Liouvillian gap $\lambda = |\text{Re}[\lambda_1]|$, which is also called the asymptotic decay rate [21], determining the slowest relaxation dynamics in the long-time limit.

For any Liouvillian, the following lemmas hold (for a detailed proof of Lemmas 3 and 4 see Appendix A 2):

Lemma 1. Given Eq. (8), $e^{\mathcal{L}t} \hat{\rho}_i = e^{\lambda_i t} \hat{\rho}_i$.

Lemma 2. $\text{Tr}[\hat{\rho}_i] = 0$ if $\text{Re}[\lambda_i] \neq 0$. Indeed, the Liouvillian evolution conserves the trace [3–5,58] and if $\text{Re}[\lambda_i] \neq 0$ for $t \rightarrow +\infty$ we have $e^{\mathcal{L}t} \hat{\rho}_i = e^{\lambda_i t} \hat{\rho}_i \rightarrow 0$.

Lemma 3. If $\mathcal{L}\hat{\rho}_i = \lambda_i \hat{\rho}_i$ then $\mathcal{L}\hat{\rho}_i^\dagger = \lambda_i^* \hat{\rho}_i^\dagger$. This implies that, if $\hat{\rho}_i$ is Hermitian, then λ_i has to be real. Conversely, if λ_i is real and of degeneracy 1, $\hat{\rho}_i$ is Hermitian. If λ_i has geometric multiplicity n and \mathcal{L} is diagonalizable, it is always possible to construct n Hermitian eigenmatrices of \mathcal{L} with eigenvalue λ_i [63].

Lemma 4. If $\lambda_i = 0$ has degeneracy n , then there exist n independent eigenvectors of the Liouvillian (the algebraic multiplicity is identical to the geometrical one). Therefore, there exist n different steady states towards which the system can evolve, depending on the initial condition.

B. Spectral decomposition of density matrices

Let us consider a system admitting a unique steady state. To be physical, its $\hat{\rho}(t)$ must be a Hermitian, positive-definite matrix with trace equal to one. Hence, from Lemma 2, to ensure $\text{Tr}[\hat{\rho}(t)] = 1$ at every time, we must have

$$\hat{\rho}(t) = \frac{\hat{\rho}_0}{\text{Tr}[\hat{\rho}_0]} + \sum_{i \neq 0} c_i(t) \hat{\rho}_i = \hat{\rho}_{ss} + \sum_{i \neq 0} c_i(0) e^{\lambda_i t} \hat{\rho}_i. \quad (10)$$

1. The case of a real Liouvillian eigenvalue λ_i

When λ_i is real, $\hat{\rho}_i$ can be constructed to be Hermitian (see Lemma 3 of Sec. II A). Thus, it can be diagonalized, obtaining the spectral decomposition [58]

$$\hat{\rho}_i = \sum_n p_n^{(i)} |\psi_n^{(i)}\rangle \langle \psi_n^{(i)}|, \quad (11)$$

where all the $p_n^{(i)}$ must be real and $\langle \psi_n^{(i)} | \psi_m^{(i)} \rangle = \delta_{n,m}$. Moreover, since $\hat{\rho}_i$ is traceless (see Lemma 2 of Sec. II A), some of the $p_n^{(i)}$ must be positive and the others negative. We can order them in such a way to have $p_n^{(i)} > 0$ for $n \leq \bar{n}$ and $p_n^{(i)} < 0$ for $n > \bar{n}$. Thus, we have

$$\hat{\rho}_i \propto \hat{\rho}_i^+ - \hat{\rho}_i^-, \quad (12)$$

where

$$\begin{aligned} \hat{\rho}_i^+ &= \sum_{n \leq \bar{n}} p_n^{(i)} |\psi_n^{(i)}\rangle \langle \psi_n^{(i)}|, \\ \hat{\rho}_i^- &= - \sum_{n > \bar{n}} p_n^{(i)} |\psi_n^{(i)}\rangle \langle \psi_n^{(i)}|, \end{aligned} \quad (13)$$

and where the $\{p_n\}$ have been normalized to ensure $\text{Tr}[\hat{\rho}_i^+] = \text{Tr}[\hat{\rho}_i^-] = 1$. With this definition, $\hat{\rho}_i^\pm$ are density matrices. Consequently, a state of the form $\hat{\rho}(0) = \hat{\rho}_{ss} + A \hat{\rho}_i$ will evolve in time as [62]

$$\hat{\rho}(t) = \hat{\rho}_{ss} + A e^{\lambda_i t} (\hat{\rho}_i^+ - \hat{\rho}_i^-). \quad (14)$$

2. The case of a complex Liouvillian eigenvalue λ_i

Let us now consider an eigenmatrix $\hat{\rho}_i$ with a complex eigenvalue λ_i . As it stems from Eq. (10), to ensure a Hermitian $\hat{\rho}(t)$ such an eigenmatrix must always appear in combination with its Hermitian conjugate $\hat{\rho}_i^\dagger$, which is also an eigenmatrix of \mathcal{L} (Lemma 3 of Sec. II A). Thus, one can simply consider the Hermitian combinations $\hat{\rho}_i + \hat{\rho}_i^\dagger$ and $i(\hat{\rho}_i - \hat{\rho}_i^\dagger)$. For example, given an initial condition $\hat{\rho}(0) = \hat{\rho}_{ss} + A(\hat{\rho}_i + \hat{\rho}_i^\dagger)$ with A real, one has [62]

$$\begin{aligned} \hat{\rho}(t) &= \hat{\rho}_{ss} + A(e^{\lambda_i t} \hat{\rho}_i + e^{\lambda_i^* t} \hat{\rho}_i^\dagger) = \hat{\rho}_{ss} + A e^{\text{Re}[\lambda_i]t} (\hat{\rho}_i e^{i\text{Im}[\lambda_i]t} \\ &\quad + \hat{\rho}_i^\dagger e^{-i\text{Im}[\lambda_i]t}) \\ &= \hat{\rho}_{ss} + 2A e^{\text{Re}[\lambda_i]t} [(\hat{\rho}_i + \hat{\rho}_i^\dagger) \cos(\text{Im}[\lambda_i]t) \\ &\quad + i(\hat{\rho}_i - \hat{\rho}_i^\dagger) \sin(\text{Im}[\lambda_i]t)]. \end{aligned} \quad (15)$$

C. Definition of dissipative phase transitions

Let us consider a system where a thermodynamic limit is obtained when a parameter $N \rightarrow +\infty$. For example, in a lattice of spins, N would be the number of lattice sites. For any finite N , the system always admits a unique steady-state solution. In the thermodynamic limit $N \rightarrow +\infty$, a transition between two different phases is characterized by the nonanalytical behavior of some ζ -independent observable \hat{o} when the parameter ζ tends to the critical value ζ_c . Formally, we say that there is a phase transition of order M if

$$\lim_{\zeta \rightarrow \zeta_c} \left| \frac{\partial^M}{\partial \zeta^M} \lim_{N \rightarrow +\infty} \text{Tr}[\hat{\rho}_{ss}(\zeta, N) \hat{o}] \right| = +\infty. \quad (16)$$

Since \hat{o} does not depend on ζ , the discontinuity in Eq. (16) is due to a discontinuous behavior in $\hat{\rho}_{ss}(\zeta, N \rightarrow \infty)$. As proved in [64], a discontinuity of an eigenmatrix is to be associated with a level crossing in the spectrum of the Liouvillian. Since $\hat{\rho}_{ss}$ is associated with $\lambda_0 = 0$, the phase transition must coincide with the closure of the Liouvillian gap [21,65] (indeed, in this case, is more correct to talk about *level touching*). Therefore, dissipative phase transitions are intimately connected to the emergence of multiple steady states in the thermodynamic limit $N \rightarrow +\infty$.

III. FIRST-ORDER PHASE TRANSITION

In this section we consider the emergence of a first-order dissipative phase transition at $\zeta = \zeta_c$ in the thermodynamic limit $N \rightarrow +\infty$. Such a transition must be associated with the existence of two different steady states, one for $\zeta < \zeta_c$ and the other for $\zeta > \zeta_c$, which implies that

$$\lambda_1(\zeta, N \rightarrow +\infty) \neq \lambda_0 = 0 \quad \text{for } \zeta \neq \zeta_c. \quad (17)$$

According to our definition, a first-order dissipative phase transition occurs when Eq. (16) is satisfied for $M = 1$, which also corresponds to

$$\lim_{\zeta \rightarrow \zeta_c^+} \lim_{N \rightarrow +\infty} \hat{\rho}_{ss}(\zeta, N) \equiv \hat{\rho}^+ \neq \hat{\rho}^- \equiv \lim_{\zeta \rightarrow \zeta_c^-} \lim_{N \rightarrow +\infty} \hat{\rho}_{ss}(\zeta, N), \quad (18)$$

which defines ρ^+ (ρ^-) as the steady state in the thermodynamic limit right after (before) the critical point. From Eq. (18) we can write that $\hat{\rho}_{ss}(\zeta) = \theta(\zeta - \zeta_c) \hat{\rho}^+ + \theta(\zeta_c -$

$\zeta)\hat{\rho}^-$ for $\zeta \neq \zeta_c$, where $\theta(x)$ is the Heaviside step function. Assuming the continuity of the Liouvillian, we can state that $\mathcal{L}(\zeta_c)\hat{\rho}^\pm = 0$ (we drop the explicit dependence on N when assuming the thermodynamic limit). This implies that $\lambda_1(\zeta_c) = \lambda_0 = 0$ and hence $\hat{\rho}_{ss}(\zeta_c)$ and $\hat{\rho}_1(\zeta_c)$ belong to the kernel spanned by $\hat{\rho}^\pm$. It is worth stressing that in the thermodynamic limit and for $\zeta = \zeta_c$, both the real and imaginary part of λ_1 must vanish. Furthermore, in a first-order dissipative phase transition, the condition $\text{Im}[\lambda_1] = 0$ must hold in a finite domain around ζ_c , as a consequence of Lemma 3 [66].

Lemma 2 (Sec. II) ensures that $\text{Tr}[\hat{\rho}_1(\zeta)] = 0$ if $\lambda_1(\zeta) \neq 0$ (i.e., $\zeta \neq \zeta_c$). Moreover, as discussed in [64], $\lambda_1(\zeta)$ must be continuous in a domain of the parameter space around $\zeta = \zeta_c$ (cf. Fig. 1). By analogy, we want also $\hat{\rho}_1(\zeta)$ to be continuous around ζ_c , and to extend the zero-trace condition we must set

$$\hat{\rho}_1(\zeta_c) \propto \hat{\rho}^+ - \hat{\rho}^-. \quad (19)$$

The above equation allows the identification of the states $\hat{\rho}_1^\pm$ obtained with the eigendecomposition (12) with the two phases $\hat{\rho}^\pm$ [Eq. (18)] emerging in the thermodynamic limit. Together with the continuity requirement, this allows us to interpret $\hat{\rho}_1^\pm(\zeta) \simeq \hat{\rho}^\pm$ in a domain around $\zeta = \zeta_c$. In this region, since the Liouvillian gap is finite, we also have $\hat{\rho}_0(\zeta) \propto \hat{\rho}_{ss}(\zeta)$. Using that $\theta(0) = 1/2$, we can infer

$$\hat{\rho}_0(\zeta_c) \propto \hat{\rho}^+ + \hat{\rho}^-. \quad (20)$$

Accordingly, $\hat{\rho}_0(\zeta_c)$ and $\hat{\rho}_1(\zeta_c)$ are orthogonal, since $\langle \hat{\rho}_0(\zeta_c), \hat{\rho}_1(\zeta_c) \rangle \propto \text{Tr}[(\rho^+)^2] - \text{Tr}[(\rho^-)^2] = 0$.




FIG. 1. Sketch depicting the paradigm of a first-order dissipative phase transition, formally described in Sec. III. In the thermodynamic limit, the Liouvillian gap $\lambda = |\text{Re}[\lambda_1]|$ closes when the parameter ζ of the Liouvillian assumes the critical value ζ_c . We note that for $\zeta \simeq \zeta_c$, we must also have $\text{Im}[\lambda_1] = 0$. Right before (after) the critical point, the steady-state density matrix $\hat{\rho}_{ss} \simeq \hat{\rho}_1^-$ ($\hat{\rho}_{ss} \simeq \hat{\rho}_1^+$), which represents one of the two different phases of the system. At the critical point $\zeta = \zeta_c$, $\hat{\rho}_{ss}$ is bimodal: the steady state is a statistical mixture of $\hat{\rho}_1^+$ and $\hat{\rho}_1^-$.

For large but finite N , provided that $|\text{Re}[\lambda_2]| \gg |\text{Re}[\lambda_1]| > 0$, Eqs. (19) and (20) are asymptotic good approximations and, since $\hat{\rho}^\pm = \hat{\rho}_1^\pm(\zeta_c, N)$, we get the asymptotic expression

$$\hat{\rho}_{ss}(\zeta_c, N) \simeq \frac{\hat{\rho}_1^+(\zeta_c, N) + \hat{\rho}_1^-(\zeta_c, N)}{2}, \quad (21)$$

which ensures Hermiticity and unit trace of the $\hat{\rho}_{ss}(\zeta_c, N)$. Let us note that Eq. (21) has a clear physical interpretation: at the critical point, for a finite-size system, the steady state is the equiprobable mixture of the two phases, which are encoded in the spectral decomposition of $\hat{\rho}_1(\zeta_c, N)$. Remarkably, in a small region on the left (right) of the critical point, $\hat{\rho}_1^+$ ($\hat{\rho}_1^-$) is metastable. This means that if the system is initialized in one of these two states it will remain stuck, for a time proportional to $1/\lambda$, before reaching the steady state [62]. This can give rise to hysterical behavior, typical of first-order phase transitions [51].

Conversely, if $\lambda_1 = 0$ in a point, one has to have a first-order phase transition. A proof can be found in Appendix A 3.

IV. SECOND-ORDER PHASE TRANSITIONS WITH SYMMETRY BREAKING

In this section, we will consider second-order dissipative phase transitions associated with a symmetry breaking. A symmetry of an open quantum system is described by a unitary superoperator $\mathcal{U} = \hat{V} \bullet \hat{V}^{-1}$ (where \hat{V} is a unitary operator and the \bullet in the previous definition means that, upon the action of the superoperator \mathcal{U} on a generic operator, the latter has to be inserted in between \hat{V} and \hat{V}^{-1}) [67], such that

$$\mathcal{U}^{-1} \mathcal{L} \mathcal{U} = \mathcal{L}, \quad (22)$$

or, equivalently, $[\mathcal{L}, \mathcal{U}] = 0$. It follows that the matrix representations $\bar{\mathcal{U}}$ of \mathcal{U} and $\bar{\mathcal{L}}$ of \mathcal{L} can be simultaneously diagonalized. From now on, we will call the symmetry sector L_u the subspace of the Liouville space L spanned by the eigenmatrices of \mathcal{U} with eigenvalue u . The existence of a symmetry means that the Lindblad master equation cannot mix different symmetry sectors. Therefore $\bar{\mathcal{L}}$ can be cast in a block-diagonal form:

$$\bar{\mathcal{L}} = \begin{bmatrix} \bar{\mathcal{L}}_{u_0} & 0 & \dots & 0 \\ 0 & \bar{\mathcal{L}}_{u_1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \bar{\mathcal{L}}_{u_n} \end{bmatrix}. \quad (23)$$

Consider an arbitrary density matrix $\hat{\rho}$ which is an eigenmatrix of \mathcal{U} : $\mathcal{U}\hat{\rho} = u\hat{\rho}$. Taking the trace of both sides of the previous identity, and given the form of \mathcal{U} , one finds $u = 1$. If $\hat{\rho}_{ss}$ is the only eigenmatrix with zero eigenvalue of \mathcal{L} (unique steady state), it must also be an eigenmatrix of \mathcal{U} . From a physical perspective, this tells us that the symmetry sector to which $\hat{\rho}_{ss}$ (and therefore $\hat{\rho}_0$) belongs is *always* $L_{u=1}$.

A symmetry-breaking dissipative phase transition is associated with the emergence of multiple eigenmatrices of \mathcal{L} with $\lambda_i = 0$, each of them belonging to a different symmetry sector L_{u_i} . The structure imposed by Eq. (23) is preserved and the previous considerations still hold. Therefore, $\hat{\rho}_0$ (belonging to the symmetry sector $L_{u=1}$) is still the only eigenmatrix of \mathcal{L} with nonzero trace.

The block-diagonal structure of the Liouvillian [see Eq. (23)], together with the previous observations, can play a fundamental role in reducing the complexity of the problem. Indeed, by properly exploiting spatial and/or internal symmetries, one can explicitly construct the reduced subspace in which the steady-state density matrix belongs. This can give a substantial speed-up for algorithms based on Monte Carlo strategies [68], cluster expansions [69,70], corner methods [71], and the tensor-network ansatz [72–74].

A. \mathcal{Z}_2 symmetry

Let us consider first a system which has a discrete Z_2 symmetry represented by the superoperator $\mathcal{Z}_2 = \hat{Z}_2 \bullet \hat{Z}_2^\dagger$. Later, we will deal with the general case of a Z_n symmetry. The symmetry superoperator \mathcal{Z}_2 admits two eigenvalues, namely ± 1 . For $\zeta < \zeta_c$ (ζ being the critical parameter) there exists a unique steady state associated with the eigenvalue $\lambda_0 = 0$ and $\mathcal{Z}_2 \hat{\rho}_{ss} = \hat{\rho}_{ss}$. For $\zeta \geq \zeta_c$, a phase transition with a symmetry breaking takes place. Consequently, $\lambda_0 = \lambda_1 = 0$ while $\hat{\rho}_0$ and $\hat{\rho}_1$ belong to two different symmetry sectors (cf. Fig. 2). From these properties, it follows that $\hat{\rho}_0$ and $\hat{\rho}_1$ are orthogonal,

since

$$\langle \hat{\rho}_0 | \hat{\rho}_1 \rangle = \langle \mathcal{Z}_2 \hat{\rho}_0 | \hat{\rho}_1 \rangle = \langle \hat{\rho}_0 | \mathcal{Z}_2 \hat{\rho}_1 \rangle = -\langle \hat{\rho}_0 | \hat{\rho}_1 \rangle, \quad (24)$$

where we exploited the Hermiticity of \mathcal{Z}_2 . Similarly, $\langle \hat{\rho}_0^\dagger | \hat{\rho}_1 \rangle = 0$. Since $\lambda_1 = 0$ is real and $\hat{\rho}_1^\dagger \neq \hat{\rho}_0$, the eigenmatrix $\hat{\rho}_1$ is Hermitian (lemmas of Sec. II A). Hence, the density matrices

$$\hat{\rho}^\pm = \frac{\hat{\rho}_0 \pm \hat{\rho}_1}{\text{Tr}[\hat{\rho}_0]} \quad (25)$$

are steady states of the master equation breaking the symmetry, as $\mathcal{Z}_2 \hat{\rho}^\pm = \hat{\rho}^\mp$. From Eq. (24) it follows that $\hat{\rho}^+$ and $\hat{\rho}^-$ are orthogonal as well. So we have

$$\hat{\rho}_0 \propto \hat{\rho}^+ + \hat{\rho}^-, \quad (26a)$$

$$\hat{\rho}_1 \propto \hat{\rho}^+ - \hat{\rho}^-. \quad (26b)$$

Thus, we can conclude that the two symmetry-broken states $\hat{\rho}^\pm$ are the two matrices stemming from the spectral decomposition of $\hat{\rho}_1$, i.e., $\hat{\rho}_1^\pm$ [cf. Eq. (12)]. For a finite-size system, where the steady state is unique,

$$\hat{\rho}_{ss}(\zeta \geq \zeta_c, N) \simeq \frac{\hat{\rho}_1^+(\zeta_c, N) + \hat{\rho}_1^-(\zeta_c, N)}{2}. \quad (27)$$

Since we are considering a second-order phase transition, we must ensure that the unique steady state in ζ_c^- coincides with both the symmetry-breaking steady states in ζ_c^+ : $\hat{\rho}_{ss}(\zeta_c^-) = \hat{\rho}^+(\zeta_c^+) = \hat{\rho}^-(\zeta_c^+)$. Consequently, according to this discussion, $\hat{\rho}_1(\zeta_c) = 0$. Therefore, a second-order phase transition is characterized by the coalescence of two eigenvectors of the Liouvillian, which may give rise to a Jordan form of the Liouvillian (see Appendix B). In order to unveil the symmetry breaking in a finite-size system (where the symmetry is always preserved) one can resort to different strategies. To identify the critical point, one can use an external weak probe which breaks the symmetry (see for example Refs. [43,69]) and look for divergences in the associated susceptibility. To characterize the existence of the two (or more) metastable states which individually break the symmetry, one can also resort to a quantum trajectory protocol [7]. Indeed, the dynamics of a single trajectory can explicitly break the symmetry, even if once the average over many trajectories is taken, such a symmetry is restored [49,56].

B. \mathcal{Z}_n symmetry

Consider now a generic symmetry superoperator $\mathcal{Z}_n = \hat{Z}_n \bullet \hat{Z}_n^\dagger$. In this case, the Liouvillian can be partitioned into n blocks, each characterized by an eigenvalue $z_j = \exp[2i\pi j/n]$, with $j = 0, 1, \dots, n-1$ (i.e., the eigenvalues must satisfy the equation $z_j^n = 1$). In the symmetry-broken phase, in each of those blocks there exists an eigenmatrix $\hat{\rho}_j$ such that $\mathcal{L}\hat{\rho}_j = 0$ and $\mathcal{Z}_n \hat{\rho}_j = z_j \hat{\rho}_j$. Lemma 3 of Sec. II A imposes $\mathcal{L}\hat{\rho}_j^\dagger = 0$. Moreover, $\hat{\rho}_j^\dagger$ is also an eigenmatrix of \mathcal{Z}_n of eigenvalue z_j^* , since

$$\mathcal{Z}_n \hat{\rho}_j^\dagger = (\hat{Z}_n \hat{\rho}_j^\dagger \hat{Z}_n^\dagger) = (\hat{Z}_n \hat{\rho}_j \hat{Z}_n^\dagger)^\dagger = z_j^* \hat{\rho}_j^\dagger. \quad (28)$$




FIG. 2. Sketch depicting the paradigm of a second-order dissipative phase transition (cf. Sec. IV), associated with the breaking of a Z_n symmetry (in the sketch $n = 5$). In the thermodynamic limit, the Liouvillian gap λ closes over the whole region $\zeta \geq \zeta_c$, ζ being the critical parameter triggering the transition. Moreover, one has that $\lambda_0, \dots, \lambda_{n-1} = 0$ for $\zeta \geq \zeta_c$. When $\lambda \neq 0$ (here for $\zeta < \zeta_c$), the steady-state density matrix $\hat{\rho}_{ss}$ is monomodal. In the symmetry-broken phase ($\lambda = 0$ and $\zeta \geq \zeta_c$), $\hat{\rho}_{ss}$ is an n -modal statistical mixture of density matrices $\hat{\rho}_j$, which are mapped one onto the other under the action of the symmetry superoperator \mathcal{Z}_n .

Note that by definition $z_j^* = z_{n-j}$, and hence $\hat{\rho}_j^\dagger = \hat{\rho}_{n-j}$. As a particular case, if $z_j = z_j^*$ then $\hat{\rho}_j = \hat{\rho}_j^\dagger$.

To construct a basis of the degenerate subspace made of density matrices, consider the operator

$$\hat{\rho}_0 = \sum_{j=0}^{n-1} \frac{\hat{\rho}_j}{\text{Tr}[\hat{\rho}_0]} = \sum_{j=0}^{n-1} \frac{\hat{\rho}_j + \hat{\rho}_{n-j}}{2 \text{Tr}[\hat{\rho}_0]} = \sum_{j=0}^{n-1} \frac{\hat{\rho}_j + \hat{\rho}_j^\dagger}{2 \text{Tr}[\hat{\rho}_0]}. \quad (29)$$

With this choice, $\hat{\rho}_0$ is a density matrix, since it is Hermitian and it has trace 1 ($\text{Tr}[\hat{\rho}_j] = \text{Tr}[\hat{\rho}_0] \delta_{j,0}$). For $\hat{\rho}_1 = \mathcal{Z}_n \hat{\rho}_0$, one has

$$\hat{\rho}_1 = \sum_{j=0}^{n-1} \frac{z_j \hat{\rho}_j + z_j^* \hat{\rho}_j^\dagger}{2 \text{Tr}[\hat{\rho}_0]}, \quad (30)$$

which is still Hermitian and of unitary trace, and therefore a density matrix. By iterative application of the symmetry operator \mathcal{Z}_n , and since $\hat{\rho}_i \neq \hat{\rho}_j$ for $i \neq j$, one obtains a basis $\{\hat{\rho}_i\}$ of density matrices, with $i = 0, \dots, n-1$. In compact notation, one has

$$\hat{\rho}_l = \mathcal{Z}_n^l \sum_{j=0}^{n-1} \frac{\hat{\rho}_j}{\text{Tr}[\hat{\rho}_0]} = \sum_{j=0}^{n-1} \frac{z_j^l(i) \hat{\rho}_j}{\text{Tr}[\hat{\rho}_0]}. \quad (31)$$

Equation (31) can be inverted to obtain $\hat{\rho}_k$ as a function of $\hat{\rho}_l$:

$$\begin{aligned} \sum_{l=0}^{n-1} (z_k^*)^l \hat{\rho}_l &= \sum_{l=0}^{n-1} \sum_{j=0}^{n-1} \frac{(z_k^* z_j)^l \hat{\rho}_j}{\text{Tr}[\hat{\rho}_0]} = \sum_{l=0}^{n-1} \sum_{j=0}^{n-1} \frac{z_{j-k}^l \hat{\rho}_j}{\text{Tr}[\hat{\rho}_0]} \\ &= \frac{n}{\text{Tr}[\hat{\rho}_0]} \sum_{j=0}^{n-1} \delta_{j,k} \hat{\rho}_j = \frac{n}{\text{Tr}[\hat{\rho}_0]} \hat{\rho}_k, \end{aligned} \quad (32)$$

where we used the identity

$$\sum_{l=0}^{n-1} z_{j-k}^l = \sum_{l=0}^{n-1} \left(e^{\frac{2i\pi(j-k)}{n}} \right)^l = n \delta_{k,j}. \quad (33)$$

We conclude that

$$\hat{\rho}_k \propto \sum_{l=0}^{n-1} \frac{(z_k^*)^l \hat{\rho}_l}{n}. \quad (34)$$

Summarizing, we have constructed a basis of $\{\hat{\rho}_i\}$ of the kernel of the Liouvillian made of density matrices such that $\mathcal{Z}_n \hat{\rho}_i = \hat{\rho}_{\text{mod}(i+1,n)}$, as depicted in Fig. 2. This construction ensures that $\mathcal{Z}_n \hat{\rho}_0 = \hat{\rho}_0$. Again, for large enough but finite N , where the steady state is unique also for $\zeta \geq \zeta_c$, we get the asymptotic expression

$$\hat{\rho}_{ss}(\zeta \geq \zeta_c, N) \simeq \sum_{l=0}^{n-1} \frac{\hat{\rho}_l(\zeta \geq \zeta_c, N)}{n}. \quad (35)$$

V. APPLICATIONS TO SPECIFIC MODELS

In the following, we will explore some specific models exhibiting dissipative phase transitions in a thermodynamic limit. We will show that in the finite-size case, our theory predicts with high fidelity the form of $\hat{\rho}_{ss}$ in the vicinity of the critical point. In particular, we will analyze some systems

for which a brute force diagonalization of the Liouvillian supermatrix $\bar{\mathcal{L}}$ is possible.

A. The driven-dissipative Kerr resonator

The first example which we discuss is the general model of a single driven-dissipative Kerr nonlinear resonator, for which an exact solution of $\hat{\rho}_{ss}$ exists [54]. In a reference frame rotating at the coherent pump frequency ω_p , the Hamiltonian of this system is

$$\hat{H} = -\Delta \hat{a}^\dagger \hat{a} + \frac{U}{2} \hat{a}^\dagger \hat{a}^\dagger \hat{a} \hat{a} + F(\hat{a}^\dagger + \hat{a}), \quad (36)$$

where $\Delta = \omega_p - \omega_c$ is the pump-cavity detuning, F is the driving amplitude, and U quantifies the Kerr nonlinearity. The operators \hat{a}^\dagger and \hat{a} are the bosonic creation and annihilation operators, respectively. The corresponding Lindblad master equation reads

$$\partial_t \hat{\rho}(t) = -i[\hat{H}, \hat{\rho}(t)] + \frac{\gamma}{2} \mathcal{D}[\hat{a}] \hat{\rho}(t), \quad (37)$$

where γ is the dissipation rate of the cavity mode. The properties of this model, and the emergence of a first-order phase transition, have been extensively discussed in Refs. [28,29,31]. A well-defined thermodynamic limit is obtained for $|F| \rightarrow +\infty$ while keeping $U|F|^2$ constant [31]. This is equivalent to expressing nonlinearity and driving amplitude in the following form:

$$U = \tilde{U}/N, \quad F = \tilde{F}\sqrt{N}, \quad (38)$$

and letting $N \rightarrow +\infty$. In Fig. 3 we study numerically the emergence of the first-order phase transition by increasing N . The top panel shows the mean value of $\langle \hat{a}^\dagger \hat{a} \rangle / N = \text{Tr}[\hat{\rho}_{ss} \hat{a}^\dagger \hat{a}] / N$ as a function of \tilde{F}/γ . The middle panel shows the rescaled Liouvillian gap $-\text{Re}[\lambda_1/\gamma]$ as a function of the rescaled driving amplitude. Such Liouvillian gap tends to zero in the thermodynamic limit $N \rightarrow +\infty$, while $\text{Im}[\lambda_1] = 0$ around the critical point also for finite N . The bottom panel of Fig. 3 presents a study of the fidelity between the steady state $\hat{\rho}_{ss}$ and the matrix $\hat{\xi} = (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$, obtained by the spectral decomposition of $\hat{\rho}_1$ [Eq. (12)]. We recall that the fidelity is defined as $f(\hat{\rho}, \hat{\xi}) = \text{Tr}[\sqrt{\sqrt{\hat{\rho}} \hat{\xi} \sqrt{\hat{\rho}}}]$. A fidelity equal to 1 indicates that the two states are identical. As the thermodynamical parameter N increases, we notice two important effects: (i) in the region in which the Liouvillian gap is minimal the fidelity is maximal; (ii) the region in which $\hat{\rho}_{ss}$ and $\hat{\xi}$ are close becomes narrower and narrower. This is consistent with our general results which are exact in the thermodynamic limit.

It is interesting now to connect our findings with the results predicted by mean-field theories. A Gross-Pitaevskii-like mean-field approximation for the driven-dissipative Kerr model is known to exhibit bistability, while the full quantum solution is always unique [54]. In the same way, a Gutzwiller mean-field theory predicts multiple solutions [33,75]. In Fig. 4, we investigate the properties of the exact steady state $\hat{\rho}_{ss}$ and of the density matrices $\hat{\rho}_1^+$ and $\hat{\rho}_1^-$ for a system with $N = 10$ as a function of the rescaled driving amplitude \tilde{F}/γ . In the top panel, we plot the mean photon density $\langle \hat{a}^\dagger \hat{a} \rangle / N = \text{Tr}[\hat{a}^\dagger \hat{a} \hat{\chi}] / N$, for $\hat{\chi} = \hat{\rho}_{ss}, \hat{\rho}_1^+, \hat{\rho}_1^-$ as indicated in the legend.




FIG. 3. Numerical results for the driven-dissipative Kerr model. Top panel: Rescaled number of photons $\langle \hat{a}^\dagger \hat{a} \rangle / N$ as a function of the rescaled driving \tilde{F}/γ for different values of N . Middle panel: $-\text{Re}[\lambda_1/\gamma]$ (Liouvillian gap) for different values of N . In the selected range of parameters we find that $\text{Im}[\lambda_1]$ is zero within the numerical error. Bottom panel: The error $1 - f$, where f is the fidelity between the steady-state density matrix $\hat{\rho}_{ss}$ and the one reconstructed via the eigendecomposition of the first eigenstate $\hat{\xi} = (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$. Parameters: $\Delta/\gamma = 10$, $\tilde{U}/\gamma = 10$.

To further characterize the nature of $\hat{\rho}_1^\pm$ in the phase transition, in the bottom panel we plot the fidelity between $\hat{\rho}_{ss}$ and $\hat{\chi} = \hat{\xi}$, $\hat{\rho}_1^+$, $\hat{\rho}_1^-$ [where $\hat{\xi} = (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$]. For $\tilde{F} < \tilde{F}_c$, $\hat{\rho}_{ss}$ is almost exactly $\hat{\rho}_1^-$. Around the critical point $\tilde{F} \simeq \tilde{F}_c$, $\hat{\rho}_{ss}$ becomes an equal mixture of $\hat{\rho}_1^+$ and $\hat{\rho}_1^-$. The maximal mixed character occurs for $\tilde{F} = \tilde{F}_c$. Finally, for $\tilde{F} > \tilde{F}_c$, the density matrix becomes very close to $\hat{\rho}_1^+$. This analysis allows us to interpret the two stable solutions predicted by the mean-field approach in terms of the metastable states which compose $\hat{\rho}_1$.

B. The driven-dissipative resonator with two-photon pumping

As an example of second-order dissipative phase transition with symmetry breaking, we will consider the driven-dissipative Kerr model with two-photon pumping and losses. In a reference frame rotating at the parametric pump frequency, the Hamiltonian of this system is [29,56]

$$\hat{H} = -\Delta \hat{a}^\dagger \hat{a} + \frac{U}{2} \hat{a}^\dagger \hat{a}^\dagger \hat{a} \hat{a} + \frac{G}{2} (\hat{a}^\dagger \hat{a}^\dagger + \hat{a} \hat{a}), \quad (39)$$

where G is the two-photon driving amplitude. This time, in addition to the Hamiltonian and to the one-photon dissipa-




FIG. 4. Top panel: Average number of photons according to $\hat{\rho}_{ss}$ and $\hat{\rho}^\pm$ as a function of the rescaled driving \tilde{F}/γ for $N = 10$ (see definitions in the main text). The dotted line indicates the Gross-Pitaevskii prediction. Bottom panel: Fidelity f between the steady-state density matrix $\hat{\rho}_{ss}$ and a density matrix $\hat{\chi} = \hat{\rho}^+$, $\hat{\rho}^-$, $\hat{\xi}$ [for $\hat{\xi} = (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$] as a function of the rescaled driving \tilde{F}/γ . The transition can be seen as a switching from a region where $\hat{\rho}_1^-$ describes the system to one where the physics is dominated by $\hat{\rho}_1^+$. Even if the region of phase coexistence in $\hat{\rho}_{ss}$ is very narrow, $\hat{\rho}_1$ describes the physics in a larger region. Parameters are set as in Fig. 3.

tion superoperator $\mathcal{D}[\hat{a}]$, we will consider also a two-photon dissipation channel with rate η . The corresponding Lindblad master equation reads

$$\partial_t \rho(t) = -i[\hat{H}, \rho(t)] + \frac{\gamma}{2} \mathcal{D}[\hat{a}]\rho(t) + \frac{\eta}{2} \mathcal{D}[\hat{a}^2]\rho(t). \quad (40)$$

The analytical solution of the steady state of this model has been provided in Ref. [55], and the emergence of first- and second-order phase transitions (according to the value of Δ) has been discussed in Ref. [29]. The emergence of a similar symmetry breaking has been also observed in an equivalent classical system [76]. The thermodynamic limit of this model is obtained by expressing U and η as

$$U = \tilde{U}/N, \quad \eta = \tilde{\eta}/N, \quad (41)$$

and considering the limit $N \rightarrow +\infty$. In this way the ratio U/η is kept constant. This model has a discrete \mathbb{Z}_2 symmetry, resulting from the invariance under the transformation $\hat{a} \rightarrow -\hat{a}$. The corresponding superoperator \mathcal{Z}_2 is

$$\mathcal{Z}_2 = e^{i\pi \hat{a}^\dagger \hat{a}} \bullet e^{-i\pi \hat{a}^\dagger \hat{a}}, \quad (42)$$

with $\mathcal{Z}_2 \hat{\rho}_{ss} = \hat{\rho}_{ss}$.

In Fig. 5 we show the emergence of a second-order phase transition by increasing the value of N . The top panel shows $\langle \hat{a}^\dagger \hat{a} \rangle / N = \text{Tr}[\hat{\rho}_{ss} \hat{a}^\dagger \hat{a}] / N$ as a function of G/γ . In the middle panel we show the rescaled Liouvillian gap $-\text{Re}[\lambda_1/\gamma]$ as a function of the rescaled pump amplitude. The abrupt




FIG. 5. Numerical results for the driven-dissipative two-photon Kerr model. Top panel: Rescaled number of photons $\langle \hat{a}^\dagger \hat{a} \rangle / N$ as a function of the rescaled driving G/γ for different values of N . Middle panel: $-\text{Re}[\lambda_1/\gamma]$ (Liouvillian gap) for different values of N . Bottom panel: The error $1 - f$, where f is the fidelity between the steady-state density matrix $\hat{\rho}_{ss}$ and $\hat{\xi} = (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$. The curves are shown in the region where λ_1 is purely real. Parameters: $\Delta/\gamma = -10$, $\tilde{U}/\gamma = 10$, $\tilde{\eta}/\gamma = 1.0$.

change in the behavior of λ indicates the onset of the phase transition. In the whole region of broken symmetry, the gap is much smaller than γ and λ_1 is real, while $\hat{\rho}_1$ is a traceless Hermitian matrix which belongs to the odd symmetry sector of \mathcal{Z}_2 ($\mathcal{Z}_2 \hat{\rho}_1 = -\hat{\rho}_1$). The states $\hat{\rho}_1^+$ and $\hat{\rho}_1^-$ obtained via the spectral decomposition of $\hat{\rho}_1$ are such that $\mathcal{Z}_2 \hat{\rho}_1^+ = \hat{\rho}_1^-$. As has been shown in Sec. IV, in the symmetry-broken region, $\hat{\rho}_{ss}$ can be constructed as a symmetric mixture of $\hat{\rho}_1^+$ and $\hat{\rho}_1^-$. As shown in the bottom panel of Fig. 5, this gives an excellent approximation for the finite-sized systems considered here. Remarkably, this expression for $\hat{\rho}_{ss}$ remains very accurate even quite far from the thermodynamic limit.

In order to characterize the abrupt change in the behavior of λ_1 , which becomes discontinuous for $N \rightarrow +\infty$, we plot part of the full spectrum of $\tilde{\mathcal{L}}$ for $N = 20$ across the critical point. In the top panel of Fig. 6, we show the real part of the spectrum, while the bottom one reports the imaginary part. Starting from the imaginary part, we clearly see that there is a point in which two complex-conjugate eigenvalues (highlighted by the red line) become real. We call $G_B(N)$ the point at which this bifurcation happens. Looking at the top panel, this merging is associated with a change in the behavior of the real part of those eigenvalues, which split and bifurcate. The one approaching zero is responsible for the phase transi-




FIG. 6. Liouvillian spectrum in the two-photon Kerr model for $N = 20$. Top and bottom panels: Real and imaginary part of the eigenvalues of $\tilde{\mathcal{L}}$. The dots represent the 10 smallest-modulus eigenvalues obtained by numerical diagonalization. The red lines are a guide for the eye indicating the two eigenvalues which merge into λ_1 for $G \gg \gamma$. Inset: A log-log plot of $\Delta G = G_B(N) - G_c$ (defined in the text) as a function of the parameter N , showing the power-law behavior $\Delta G = A N^{-\eta}$ with $A = 21.1 \pm 0.2$ and $\eta = 0.881 \pm 0.006$. In the thermodynamic limit, the bifurcation point $G_B(N)$ and the critical point G_c coincide. Same parameters as in Fig. 5.

tion and its associated eigenvector becomes $\hat{\rho}_1 = \hat{\rho}_1^+ - \hat{\rho}_1^-$ for $G > G_B(N)$. As we saw in Fig 5, it is not clear where the gap starts to close, but one might guess that it happens when the two eigenvalues bifurcate. To test this conjecture, in the inset we plot, as a function of N , the scaling of the bifurcation point $\Delta G = G_B(N) - G_c$, where G_c is the critical point extrapolated via the study of the analytic solution for $N = 1000$. Indeed, the clear power-law decay of this quantity demonstrates that the onset of this transition can be understood in terms of a merging of two eigenvalues. The emergence of criticality is thus to be associated with a *touching* of two eigenvalues in the complex plane. This fact, together with the emergence of a discontinuity in λ_1 for $N \rightarrow \infty$, implies that at the bifurcation point, the Liouvillian becomes nondiagonalizable, resulting in a Jordan structure. This leads to a nonexponential relaxation dynamics at criticality. To better understand this behavior, in Appendix B we study an exactly solvable two-level system which admits a Jordan block structure for a specific choice of parameters.

Up to now, we considered the case in which an eigenvalue of the symmetry sector L_{-1} approaches zero, which gives rise to a symmetry breaking without inducing first-order discontinuities in $\hat{\rho}_{ss}$. The two-photon Kerr model is known to present also a first-order phase transition with symmetry breaking for $\Delta > 0$ [29]. Indeed, together with the emerging of a zero in L_{-1} , the symmetry sector L_1 acquires two zero eigenvalues of \mathcal{L} : one associated with $\hat{\rho}_{ss}$, the other with an eigenmatrix whose eigenvalue touches zero only at the critical point.




FIG. 7. Study of the first-order phase transition with symmetry breaking in the two-photon Kerr resonator. Top panel: Rescaled number of photons $\langle \hat{a}^\dagger \hat{a} \rangle / N$ as a function of the rescaled driving G/γ for different values of N . Middle panel: Real part of $\lambda_{1,2}$ rescaled by γ for $N = 10$. The two branches of Liouvillian eigenvalues lead to the first-order phase transition (red) and a symmetry breaking (blue). Bottom panel: Average number of photons according to $\hat{\rho}_{ss}$ and $\hat{\rho}_2^\pm$. Parameters: $\Delta/\gamma = 10$, $U/\gamma = 10$, $\eta/\gamma = 1.0$.

This allows a discontinuous behavior of $\hat{\rho}_{ss}$ with symmetry breaking. In Fig. 7 we plot the behavior of the system in such regime. The top panel shows the emergence of a first-order phase transition in the rescaled density. In the middle panel, we plot the real part of the two eigenvalues of the Liouvillian with the smallest modulus. One presents the phenomenology we expect from a symmetry breaking: $-\text{Re}[\lambda_1] \ll \gamma$ in the symmetry-broken phase $G \geq G_c(N)$. The other is responsible for the discontinuous first-order behavior: $-\text{Re}[\lambda_2] \ll \gamma$ only for $G \simeq G_c(N)$. Indeed, we tested that $\hat{\rho}_1$ (associated with λ_1) satisfies $\mathcal{Z}_2 \hat{\rho}_1 = -\hat{\rho}_1$. Moreover, $\mathcal{Z}_2 \hat{\rho}_1^\pm = \hat{\rho}_1^\mp$ and $\hat{\rho}_{ss} \simeq (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$. As for $\hat{\rho}_2$, $\mathcal{Z}_2 \hat{\rho}_2 = \hat{\rho}_2$ and it cannot be associated with a symmetry breaking. In the bottom panel we test the structure of $\hat{\rho}_{ss}$ in connection to the spectral decomposition of $\hat{\rho}_2$: the first-order phase transition can be interpreted as a switch between $\hat{\rho}_2^-$ and $\hat{\rho}_2^+$. The symmetry breaking emerges in the fact that $\hat{\rho}_2^+ \simeq (\hat{\rho}_1^+ + \hat{\rho}_1^-)/2$. In conclusion, in these specific numerical examples we recover all the features predicted by our general theory.

VI. CONCLUSIONS

In this article, we have presented theoretical results for first- and second-order dissipative phase transitions. Within

a general formalism, we have determined the structure of the density matrix in the vicinity of a critical point. In particular, due to the closure of the Liouvillian gap at the critical point, we have shown how the steady-state density matrix is directly related to the eigenmatrix of the Liouvillian superoperator corresponding to the eigenvalue λ_1 (the one with the smallest absolute value of the real part). We have illustrated our general results by considering two specific quantum optical models, where the emergence of a dissipative phase transition can be studied analytically and numerically. Our work provides a general insight into dissipative phase transitions. Moreover, it gives precise constraints for variational methods [25,77] to describe critical phenomena in open quantum systems, whose corresponding ansatz matrices must satisfy the relations derived in this work.

ACKNOWLEDGMENTS

We acknowledge discussions with V. Albert, G. Orso, D. Rossini, R. Rota, and M. Vogel. We acknowledge support from ERC (via Consolidator Grant CORPHO No. 616233) and ANR (via the grant UNIQ ANR-16-CE24-0029).

APPENDIX A: PROOFS

1. Remarks on the Liouvillian diagonalizability

Generally, the Liouvillian is a non-Hermitian superoperator with a holomorphic dependence on the system parameter(s) ζ . Therefore, there might exist values of ζ for which $\mathcal{L}(\zeta)$ is not diagonalizable: this implies the existence of a degenerate eigensubspace.

The eigenvalues $\lambda_i(\zeta)$ of $\mathcal{L}(\zeta)$ can be obtained via the resolution of the characteristic equation $\det[\tilde{\mathcal{L}}(\zeta) - \lambda_i(\zeta)\mathbb{I}] = 0$. A well-known result of function theory [64] guarantees that the roots of this equation are branches of analytic functions of ζ with, at most, algebraic singularities. Therefore, the number s of distinct eigenvalues of $\mathcal{L}(\zeta)$ is a constant except in a countable number of points. This ensures that if the Liouvillian has a simple spectrum on a finite region of the parameter space, it will be diagonalizable for any ζ , except the countable exceptional points. For all the systems considered in this work, this condition is fulfilled far from the thermodynamic limit.

2. Proofs of the lemmas in Sec. II A

Lemma 3. If $\mathcal{L}\hat{\rho}_i = \lambda_i\hat{\rho}_i$ then $\mathcal{L}\hat{\rho}_i^\dagger = \lambda_i^*\hat{\rho}_i^\dagger$. This implies that, if $\hat{\rho}_i$ is Hermitian then λ_i has to be real. Conversely, if λ_i is real and of degeneracy 1, $\hat{\rho}_i$ is Hermitian. If λ_i has geometric multiplicity n and \mathcal{L} is diagonalizable, it is always possible to construct n Hermitian eigenmatrices of \mathcal{L} with eigenvalue λ_i .

Proof. Thanks to the master equation we have

$$\begin{aligned} \mathcal{L}\hat{\rho}_i^\dagger &= -i[\hat{H}, \hat{\rho}_i^\dagger] + \frac{\gamma}{2}(2\hat{a}\hat{\rho}_i^\dagger\hat{a}^\dagger - \hat{a}^\dagger\hat{a}\hat{\rho}_i^\dagger - \hat{\rho}_i^\dagger\hat{a}^\dagger\hat{a}) \\ &= \left\{ -i[\hat{H}, \hat{\rho}_i] + \frac{\gamma}{2}(2\hat{a}\hat{\rho}_i\hat{a}^\dagger - \hat{a}^\dagger\hat{a}\hat{\rho}_i - \hat{\rho}_i\hat{a}^\dagger\hat{a}) \right\}^\dagger \\ &= (\mathcal{L}\hat{\rho}_i)^\dagger = \lambda_i^*\hat{\rho}_i^\dagger. \end{aligned} \quad (\text{A1})$$

If $\hat{\rho}_i$ is Hermitian, we have $\lambda_i\hat{\rho}_i = \mathcal{L}\hat{\rho}_i = \mathcal{L}\hat{\rho}_i^\dagger = \lambda_i^*\hat{\rho}_i^\dagger = \lambda_i^*\hat{\rho}_i$. Thus, we can conclude $\lambda_i = \lambda_i^*$. Conversely, in the case

in which $\lambda_i \in \mathbb{R}$ is a simple eigenvalue (i.e., with degeneracy 1), we can conclude that $\hat{\rho}_i = \hat{\rho}_i^\dagger$, and thus $\hat{\rho}_i$ is Hermitian. If the eigenvalues have geometric multiplicity n , it may happen that for some eigenmatrices $\hat{\rho}_i^\dagger \neq \hat{\rho}_i$. From Eq. (A1) it follows $\mathcal{L}\hat{\rho}_i^\dagger = \lambda_i\hat{\rho}_i^\dagger$. In this case, we can consider the matrices $(\hat{\rho}_i + \hat{\rho}_i^\dagger)/2$ and $i(\hat{\rho}_i - \hat{\rho}_i^\dagger)/2$, which are Hermitian by construction, and whose eigenvalue is λ_i .

Lemma 4. If $\lambda_i = 0$ has degeneracy n , then there exist n independent eigenvectors of the Liouvillian (the algebraic multiplicity is identical to the geometrical one). Therefore, there exist n different steady states towards which the system can evolve, depending on the initial condition.

Proof. We will prove this lemma by contradiction. Let us suppose that the algebraic multiplicity is greater than the geometrical one (see [63]). Since the dimension of the reduced space is n , we can write the Liouvillian as a matrix acting on a basis of vector in this reduced space; i.e., the invariant space of λ_0 has a finite dimension. Since we can write the Liouvillian as a matrix, this means that we can put in its canonical Jordan form. In other words the Liouvillian acting on the vectors of this subspace can be decomposed in a diagonal part $\bar{\Lambda}_0$ and a nilpotent matrix \bar{N} via a similarity transformation S :

$$\bar{\mathcal{L}}_{\lambda_0} = S^{-1}(\bar{\Lambda}_0 + \bar{N})$$

$$S = S^{-1} \begin{bmatrix} \lambda_0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & \lambda_0 & 1 & \cdots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & \lambda_0 \end{bmatrix} S. \quad (\text{A2})$$

Of course, the new basis of vectors obtained by the nonunitary transformation S may not be orthonormal. The time evolution of the system is given by $e^{\bar{\mathcal{L}}t}$, and since $\bar{\Lambda}_0$ and \bar{N} commute, one has

$$e^{\bar{\mathcal{L}}_{\lambda_0}t} = S^{-1}e^{\bar{\Lambda}_0 t}e^{\bar{N}t}S$$

$$= S^{-1}e^{\lambda_0 t} \left(\mathbb{1} + \bar{N}t + \frac{(\bar{N}t)^2}{2} + \cdots + \frac{(\bar{N}t)^n}{n!} \right) S$$

$$= S^{-1}e^{\lambda_0 t} \begin{bmatrix} 1 & t & \frac{t^2}{2} & \cdots & \frac{t^{n-1}}{(n-1)!} & \frac{t^n}{n!} \\ 0 & 1 & t & \cdots & \frac{t^{n-2}}{(n-2)!} & \frac{t^{n-1}}{(n-1)!} \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ 0 & 0 & 0 & \cdots & 1 & t \\ 1 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix} S. \quad (\text{A3})$$

Since $\lambda_0 = 0$, the previous expression clearly will cause the dynamics to diverge, proving the absurd. We stress that this reasoning cannot be directly extended to $\lambda_i \neq 0$ nor to infinite degeneracies $n \rightarrow \infty$.

3. Vanishing of λ_1 associated with a first-order phase transition

In Sec. III we proved that if there is a jump in one observable at the critical point $\zeta = \zeta_c$, then $\lambda_1 = 0$. Here, we prove

that the last condition is also sufficient; i.e., $\lim_{\zeta \rightarrow \zeta_c} \lambda_1(\zeta) = 0$ implies a first-order phase transition.

We will prove this statement by contradiction. Let us suppose that even if $\lambda_1 = 0$ there is no phase transition. From the definition (16), we deduce that for any operator \hat{o} in $\mathcal{H} \otimes \mathcal{H}$, $\langle \hat{o}(\zeta) \rangle$ is continuous in ζ_c . Hence, we have that also $\hat{\rho}_0(\zeta)$ is continuous. From Lemmas 3 and 4 of Sec. II A, the eigenstate $\hat{\rho}_1(\zeta_c)$, being associated with $\lambda_1(\zeta_c) = 0$, exists and is Hermitian. By exploiting its spectral decomposition, we can write $\hat{\rho}_1(\zeta_c) = [\hat{\rho}_1^+(\zeta_c) - \hat{\rho}_1^-(\zeta_c)]/\sqrt{2}$ [we stress that here we have $\|\hat{\rho}_1^\pm(\zeta_c)\| = 1$, and $\langle \hat{\rho}_1^+(\zeta_c), \hat{\rho}_1^-(\zeta_c) \rangle = 0$ by construction].

The first part of the proof is to show that $\hat{\rho}_0(\zeta_c) = [\hat{\rho}_1^+(\zeta_c) + \hat{\rho}_1^-(\zeta_c)]/\sqrt{2}$. Indeed, $\|\hat{\rho}_1(\zeta_c)\| = 1$ and $e^{\mathcal{L}t}\hat{\rho}_1 = \hat{\rho}_1$. Thus, exploiting the triangular inequality, we have

$$1 = \|\hat{\rho}_1(\zeta_c)\|^2 = \|e^{\mathcal{L}t}\hat{\rho}_1(\zeta_c)\|^2 = \left\| e^{\mathcal{L}t} \frac{\hat{\rho}_1^+(\zeta_c) - \hat{\rho}_1^-(\zeta_c)}{\sqrt{2}} \right\|^2$$

$$\leq \frac{\|e^{\mathcal{L}t}\hat{\rho}_1^+(\zeta_c)\|^2 + \|e^{\mathcal{L}t}\hat{\rho}_1^-(\zeta_c)\|^2}{2} \leq 1. \quad (\text{A4})$$

It follows that $\|e^{\mathcal{L}t}\hat{\rho}_1^\pm(\zeta_c)\| = 1$ for every time t . Hence, $\hat{\rho}_1^\pm(\zeta_c)$ must be a linear superposition of eigematrices of the Liouvillian with zero eigenvalue. Considering that $\hat{\rho}_1(\zeta_c) = [\hat{\rho}_1^+(\zeta_c) - \hat{\rho}_1^-(\zeta_c)]/\sqrt{2}$, $\|\hat{\rho}_0(\zeta_c)\| = 1$ and $\langle \hat{\rho}_1^+(\zeta_c), \hat{\rho}_1^-(\zeta_c) \rangle = 0$, we obtain $\hat{\rho}_0(\zeta_c) = [\hat{\rho}_1^+(\zeta_c) + \hat{\rho}_1^-(\zeta_c)]/\sqrt{2}$.

Having proved the first part, let us consider the eigendecomposition of $\hat{\rho}_1(\zeta)$ around ζ_c . Except at the critical point, we have $\lim_{t \rightarrow \infty} e^{\mathcal{L}t}\hat{\rho}_1^\pm(\zeta) = \hat{\rho}_0(\zeta)/\text{Tr}[\hat{\rho}_1^\pm]$. But, by hypothesis, all function are continuous; hence,

$$\hat{\rho}_1^\pm(\zeta_c) = \lim_{\zeta \rightarrow \zeta_c} \lim_{t \rightarrow \infty} e^{\mathcal{L}t}\hat{\rho}_1^\pm(\zeta) = \lim_{\zeta \rightarrow \zeta_c} \hat{\rho}_0(\zeta)/\text{Tr}[\hat{\rho}_1^\pm]$$

$$= \frac{\hat{\rho}_1^+(\zeta_c) + \hat{\rho}_1^-(\zeta_c)}{2}. \quad (\text{A5})$$

Consequently, we find that at the critical point, $\hat{\rho}_1^+(\zeta_c) = \hat{\rho}_1^-(\zeta_c)$. This statement would require that at $\zeta = \zeta_c$ $\hat{\rho}_1(\zeta_c) = 0$. This statement is absurd, since Lemma 4 of Sec. II A guarantees that $\hat{\rho}_1(\zeta_c)$ is a well-defined eigenvector of the Liouvillian. Therefore, by the absurd, we deduce that the function $\hat{\rho}_0(\zeta)$ cannot be continuous at $\zeta = \zeta_c$.

APPENDIX B: NONEXPONENTIAL DECAYS AND JORDAN BLOCKS

In this appendix, we provide a simple analytic example of a nonexponential decay associated with a Jordan block structure of the Liouvillian. Let us consider a spin-1/2 subjected to the action of two competing decay channels whose evolution obeys ($\hbar = 1$) [78]

$$\partial_t \hat{\rho}(t) = \mathcal{L}\hat{\rho}(t) = -i[\hat{H}, \hat{\rho}(t)] + \frac{\epsilon}{2}\mathcal{D}[\hat{\sigma}^-]\hat{\rho}(t)$$

$$+ \frac{\gamma}{2}\mathcal{D}[\hat{\sigma}^x]\hat{\rho}(t), \quad (\text{B1})$$

where $\hat{H} = \frac{\omega}{2}\hat{\sigma}^z$. The steady-state density matrix can be obtained as

$$\hat{\rho}_{ss} = \frac{1}{2\gamma + \epsilon} \begin{pmatrix} \gamma & 0 \\ 0 & \gamma + \epsilon \end{pmatrix}. \quad (\text{B2})$$

The eigenvalues of \mathcal{L} are

$$\begin{aligned}\lambda_0 &= 0, \\ \lambda_1 &= -\gamma - \frac{\epsilon}{2} + \sqrt{\gamma^2 - \omega^2}, \\ \lambda_2 &= -\gamma - \frac{\epsilon}{2} - \sqrt{\gamma^2 - \omega^2}, \\ \lambda_3 &= -2\gamma - \epsilon,\end{aligned}\tag{B3}$$

which are associated with the following (unnormalized) eigenmatrices:

$$\begin{aligned}\hat{\rho}_0 \propto \hat{\rho}_{ss} &= \frac{1}{2\gamma + \epsilon} \begin{pmatrix} \gamma & 0 \\ 0 & \gamma + \epsilon \end{pmatrix}, \quad \hat{\rho}_1 \propto \begin{pmatrix} 0 & \sqrt{\gamma^2 - \omega^2 - i\omega} \\ 1 & \gamma \end{pmatrix}, \\ \hat{\rho}_2 \propto \begin{pmatrix} 0 & -\frac{i\omega + \sqrt{\gamma^2 - \omega^2}}{\gamma} \\ 1 & 0 \end{pmatrix}, \quad \hat{\rho}_3 \propto \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.\end{aligned}\tag{B4}$$

The eigenmatrices $\hat{\rho}_{1,2}$ describe the decay of the $\hat{\sigma}^{x,y}$ components with rate $\lambda_{1,2}$, while $\hat{\rho}_3$ is associated with $\hat{\sigma}^z$ and λ_3 .

This simple model is particularly interesting since, according to the values of the couplings, it can display different relaxation dynamics toward the steady state:

(1) If $\gamma > \omega$ the Liouvillian has 4 real distinct eigenvalues (it is diagonalizable). In this case, the decay at long times will be exponential. The asymptotic decay rate is λ_1 .

(2) If $\gamma < \omega$ the Liouvillian has 4 distinct eigenvalues (it is diagonalizable), 2 of which are complex conjugate ($\lambda_1 = \lambda_2^*$). In this case, the decay at long times will be an exponential decay of magnitude $\text{Re}[\lambda_1] = \text{Re}[\lambda_2]$ multiplied by an oscillating term given by $\text{Im}[\lambda_1]$.

(3) If $\lambda = \omega$ we have $\lambda_1 = \lambda_2$ and $\hat{\rho}_1 = \hat{\rho}_2$: the Liouvillian is not diagonalizable but it can be written in a Jordan form.

The presence of a Jordan form has strong consequences for the long-time dynamics. Indeed, given a general initial state

$$\hat{\rho}(0) = \begin{pmatrix} a & b \\ b^* & 1 - a \end{pmatrix},\tag{B5}$$

the decay of the observables $\hat{\sigma}^{x,y}$ is given by

$$\begin{aligned}\text{Tr}[\hat{\sigma}^x \hat{\rho}(t)] &= 2e^{-\lambda_1 t} \{t\omega(\text{Re}[b] + \text{Im}[b]) + \text{Re}[b]\}, \\ \text{Tr}[\hat{\sigma}^y \hat{\rho}(t)] &= 2e^{-\lambda_1 t} \{t\omega(\text{Re}[b] + \text{Im}[b]) - \text{Re}[b]\},\end{aligned}\tag{B6}$$

hence not exponential. However, we stress that the asymptotic decay rate is λ_3 for $\hat{\sigma}^z$ (purely exponential decay). We also remark on the strong similarity between the behavior of $\lambda_{1,2}$ and that of the eigenvalues characterizing a second-order phase transition. In both cases, a pair of two complex-conjugate eigenvalues becomes real in proximity to an exceptional point.

-
- [1] L. D. Landau and E. M. Lifshitz, *Statistical Physics* (Pergamon Press, Oxford, 1969).
- [2] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 2001).
- [3] H. J. Carmichael, *Statistical Methods in Quantum Optics I: Master Equations and Fokker-Planck Equations*, Physics and Astronomy Online Library (Springer-Verlag, Berlin, Heidelberg, 1999).
- [4] C. Gardiner and P. Zoller, *Quantum Noise: A Handbook of Markovian and Non-Markovian Quantum Stochastic Methods with Applications to Quantum Optics*, Springer Series in Synergetics (Springer-Verlag, Berlin, Heidelberg, 2004).
- [5] S. Haroche and J. M. Raimond, *Exploring the Quantum: Atoms, Cavities, and Photons*, Oxford Graduate Texts (Oxford University Press, Oxford, 2006).
- [6] U. Weiss, *Quantum Dissipative Systems* (World Scientific, Singapore, 2008).
- [7] A. J. Daley, Quantum trajectories and open many-body quantum systems, *Adv. Phys.* **63**, 77 (2014).
- [8] B. Schmittmann and R. K. P. Zia, *Statistical Mechanics of Driven Diffusive Systems*, Phase Transitions and Critical Phenomena Vol. 17, edited by C. Domb and J. L. Lebowitz (Academic Press, London, 1995).
- [9] J. Toner and Y. Tu, Flocks, herds, and schools: A quantitative theory of flocking, *Phys. Rev. E* **58**, 4828 (1998).
- [10] N. D. Mermin and H. Wagner, Absence of Ferromagnetism or Antiferromagnetism in One- or Two-Dimensional Isotropic Heisenberg Models, *Phys. Rev. Lett.* **17**, 1133 (1966).
- [11] T. E. Lee, S. Gopalakrishnan, and M. D. Lukin, Unconventional Magnetism Via Optical Pumping of Interacting Spin Systems, *Phys. Rev. Lett.* **110**, 257204 (2013).
- [12] J. Jin, A. Biella, O. Viyuela, L. Mazza, J. Keeling, R. Fazio, and D. Rossini, Cluster Mean-Field Approach to the Steady-State Phase Diagram of Dissipative Spin Systems, *Phys. Rev. X* **6**, 031011 (2016).
- [13] A. A. Houck, H. E. Tureci, and J. Koch, On-chip quantum simulation with superconducting circuits, *Nat. Phys.* **8**, 292 (2012).
- [14] M. Fitzpatrick, N. M. Sundaresan, A. C. Y. Li, J. Koch, and A. A. Houck, Observation of a Dissipative Phase Transition in a One-Dimensional Circuit QED Lattice, *Phys. Rev. X* **7**, 011016 (2017).
- [15] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, Engineered open systems and quantum simulations with atoms and ions, *Adv. At. Mol. Opt. Phys.* **61**, 1 (2012).
- [16] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, Probing many-body dynamics on a 51-atom quantum simulator, *Nature (London)* **551**, 579 (2017).
- [17] M. Aspelmeyer, T. J. Kippenberg, and F. Marquardt, Cavity optomechanics, *Rev. Mod. Phys.* **86**, 1391 (2014).
- [18] E. Gil-Santos, M. Labousse, C. Baker, A. Goetschy, W. Hease, C. Gomez, A. Lemaître, G. Leo, C. Ciuti, and I. Favero, Light-Mediated Cascaded Locking of Multiple Nano-Optomechanical Oscillators, *Phys. Rev. Lett.* **118**, 063605 (2017).

- [19] J. Kasprzak, M. Richard, S. Kundermann, A. Baas, P. Jeambrun, J. M. J. Keeling, F. M. Marchetti, M. H. Szymanska, R. Andre, J. L. Staehli, V. Savona, P. B. Littlewood, B. Deveaud, and Le Si Dang, Bose-Einstein condensation of exciton polaritons, *Nature (London)* **443**, 409 (2006).
- [20] I. Carusotto and C. Ciuti, Quantum fluids of light, *Rev. Mod. Phys.* **85**, 299 (2013).
- [21] E. M. Kessler, G. Giedke, A. Imamoglu, S. F. Yelin, M. D. Lukin, and J. I. Cirac, Dissipative phase transition in a central spin system, *Phys. Rev. A* **86**, 012116 (2012).
- [22] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. P. Büchler, and P. Zoller, Quantum states and phases in driven open quantum systems with cold atoms, *Nat. Phys.* **4**, 878 (2008).
- [23] F. Verstraete, M. M. Wolf, and J. I. Cirac, Quantum computation and quantum-state engineering driven by dissipation, *Nat. Phys.* **5**, 633 (2009).
- [24] H. J. Carmichael, Breakdown of Photon Blockade: A Dissipative Quantum Phase Transition in Zero Dimensions, *Phys. Rev. X* **5**, 031028 (2015).
- [25] H. Weimer, Variational Principle for Steady States of Dissipative Quantum Many-Body Systems, *Phys. Rev. Lett.* **114**, 040402 (2015).
- [26] M. Benito, C. Sánchez Muñoz, and C. Navarrete-Benlloch, Degenerate parametric oscillation in quantum membrane optomechanics, *Phys. Rev. A* **93**, 023846 (2016).
- [27] J. J. Mendoza-Arenas, S. R. Clark, S. Felicetti, G. Romero, E. Solano, D. G. Angelakis, and D. Jaksch, Beyond mean-field bistability in driven-dissipative lattices: Bunching-antibunching transition and quantum simulation, *Phys. Rev. A* **93**, 023821 (2016).
- [28] W. Casteels, F. Storme, A. Le Boité, and C. Ciuti, Power laws in the dynamic hysteresis of quantum nonlinear photonic resonators, *Phys. Rev. A* **93**, 033824 (2016).
- [29] N. Bartolo, F. Minganti, W. Casteels, and C. Ciuti, Exact steady state of a Kerr resonator with one- and two-photon driving and dissipation: Controllable Wigner-function multimodality and dissipative phase transitions, *Phys. Rev. A* **94**, 033841 (2016).
- [30] W. Casteels and C. Ciuti, Quantum entanglement in the spatial-symmetry-breaking phase transition of a driven-dissipative Bose-Hubbard dimer, *Phys. Rev. A* **95**, 013812 (2017).
- [31] W. Casteels, R. Fazio, and C. Ciuti, Critical dynamical properties of a first-order dissipative phase transition, *Phys. Rev. A* **95**, 012128 (2017).
- [32] M. Foss-Feig, P. Niroula, J. T. Young, M. Hafezi, A. V. Gorshkov, R. M. Wilson, and M. F. Maghrebi, Emergent equilibrium in many-body optical bistability, *Phys. Rev. A* **95**, 043826 (2017).
- [33] M. Biondi, G. Blatter, H. E. Türeci, and S. Schmidt, Nonequilibrium gas-liquid transition in the driven-dissipative photonic lattice, *Phys. Rev. A* **96**, 043809 (2017).
- [34] A. Biella, F. Storme, J. Lebreuilly, D. Rossini, R. Fazio, I. Carusotto, and C. Ciuti, Phase diagram of incoherently driven strongly correlated photonic lattices, *Phys. Rev. A* **96**, 023839 (2017).
- [35] V. Savona, Spontaneous symmetry breaking in a quadratically driven nonlinear photonic lattice, *Phys. Rev. A* **96**, 033826 (2017).
- [36] C. Sánchez Muñoz, A. Lara, J. Puebla, and F. Nori, Hybrid Systems for the Dissipative Preparation of Mechanical Schrödinger Cats, *Phys. Rev. Lett.* **121**, 123604 (2018).
- [37] L. M. Sieberer, S. D. Huber, E. Altman, and S. Diehl, Dynamical Critical Phenomena in Driven-Dissipative Systems, *Phys. Rev. Lett.* **110**, 195301 (2013).
- [38] L. M. Sieberer, S. D. Huber, E. Altman, and S. Diehl, Nonequilibrium functional renormalization for driven-dissipative Bose-Einstein condensation, *Phys. Rev. B* **89**, 134310 (2014).
- [39] E. Altman, L. M. Sieberer, L. Chen, S. Diehl, and J. Toner, Two-Dimensional Superfluidity of Exciton Polaritons Requires Strong Anisotropy, *Phys. Rev. X* **5**, 011017 (2015).
- [40] T. E. Lee, H. Häffner, and M. C. Cross, Antiferromagnetic phase transition in a nonequilibrium lattice of Rydberg atoms, *Phys. Rev. A* **84**, 031402 (2011).
- [41] C. K. Chan, T. E. Lee, and S. Gopalakrishnan, Limit-cycle phase in driven-dissipative spin systems, *Phys. Rev. A* **91**, 051601(R) (2015).
- [42] M. F. Maghrebi and A. V. Gorshkov, Nonequilibrium many-body steady states via Keldysh formalism, *Phys. Rev. B* **93**, 014307 (2016).
- [43] R. Rota, F. Storme, N. Bartolo, R. Fazio, and C. Ciuti, Critical behavior of dissipative two-dimensional spin lattices, *Phys. Rev. B* **95**, 134431 (2017).
- [44] V. R. Overbeck, M. F. Maghrebi, A. V. Gorshkov, and H. Weimer, Multicritical behavior in dissipative Ising models, *Phys. Rev. A* **95**, 042133 (2017).
- [45] R. Dietrich, S. Diehl, and M. Buchhold, Phenomenology of a first-order dark state phase transition, [arXiv:1803.08514](https://arxiv.org/abs/1803.08514).
- [46] E. G. Dalla Torre, E. Demler, T. Giamarchi, and E. Altman, Dynamics and universality in noise-driven dissipative systems, *Phys. Rev. B* **85**, 184302 (2012).
- [47] J. Marino and S. Diehl, Driven Markovian Quantum Criticality, *Phys. Rev. Lett.* **116**, 070407 (2016).
- [48] F. Vicentini, F. Minganti, R. Rota, G. Orso, and C. Ciuti, Critical slowing down in driven-dissipative Bose-Hubbard lattices, *Phys. Rev. A* **97**, 013853 (2018).
- [49] R. Rota, F. Minganti, A. Biella, and C. Ciuti, Dynamical properties of dissipative XYZ Heisenberg lattices, *New J. Phys.* **20**, 045003 (2018).
- [50] J. M. Fink, A. Dombi, A. Vukics, A. Wallraff, and P. Domokos, Observation of the Photon-Blockade Breakdown Phase Transition, *Phys. Rev. X* **7**, 011012 (2017).
- [51] S. R. K. Rodriguez, W. Casteels, F. Storme, N. Carlon Zambon, I. Sagnes, L. Le Gratiet, E. Galopin, A. Lemaître, A. Amo, C. Ciuti, and J. Bloch, Probing a Dissipative Phase Transition Via Dynamical Optical Hysteresis, *Phys. Rev. Lett.* **118**, 247402 (2017).
- [52] T. Fink, A. Schade, S. Höfling, C. Schneider, and A. Imamoglu, Signatures of a dissipative phase transition in photon correlation measurements, *Nat. Phys.* **14**, 365 (2018).
- [53] R. Gutierrez-Jauregui and H. J. Carmichael, Dissipative quantum phase transition of light in a generalized Jaynes-Cummings-Rabi model, *Phys. Rev. A* **98**, 023804 (2018).
- [54] P. D. Drummond and D. F. Walls, Quantum theory of optical bistability. I. Nonlinear polarisability model, *J. Phys. A: Math. Gen.* **13**, 725 (1980).
- [55] F. Minganti, N. Bartolo, J. Lolli, W. Casteels, and C. Ciuti, Exact results for Schrödinger cats in driven-dissipative systems and their feedback control, *Sci. Rep.* **6**, 26987 (2016).
- [56] N. Bartolo, F. Minganti, J. Lolli, and C. Ciuti, Homodyne versus photon-counting quantum trajectories for dissipative Kerr

- resonators with two-photon driving, *Eur. Phys. J.: Spec. Top.* **226**, 2705 (2017).
- [57] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2007).
- [58] Á. Rivas and S. F. Huelga, *Open Quantum Systems: An Introduction*, Springer Briefs in Physics (Springer, Berlin, 2011).
- [59] V. V. Albert and L. Jiang, Symmetries and conserved quantities in Lindblad master equations, *Phys. Rev. A* **89**, 022118 (2014).
- [60] D. Nigro, On the uniqueness of the steady-state solution of the Lindblad-Gorini-Kossakowski-Sudarshan equation, [arXiv:1803.06279](https://arxiv.org/abs/1803.06279).
- [61] The steady-state density matrix $\hat{\rho}_{ss}$ is thus proportional to the eigenstate of \mathcal{L} whose eigenvalue is zero, since $\hat{\rho}_{ss}$ must satisfy $\text{Tr}[\hat{\rho}_{ss}] = 1$, which may not correspond to a state with norm one.
- [62] K. Macieszczak, Mădălin Guță, I. Lesanovsky, and J. P. Garrahan, Towards a Theory of Metastability in Open Quantum Dynamics, *Phys. Rev. Lett.* **116**, 240404 (2016).
- [63] The algebraic multiplicity of λ is defined as the number of times λ appears as a root of the characteristic equation. The geometric multiplicity, instead, is the maximum number of linearly independent eigenvectors associated with λ .
- [64] T. Kato, *Perturbation Theory for Linear Operators*, Classics in Mathematics (Springer-Verlag, Berlin, Heidelberg, 1995).
- [65] B. Horstmann, J. I. Cirac, and G. Giedke, Noise-driven dynamics and phase transitions in fermionic systems, *Phys. Rev. A* **87**, 012108 (2013).
- [66] Suppose that for $\zeta \simeq \zeta_c$, $\text{Im}[\lambda_1(\zeta)] \neq 0$. In virtue of Lemma 3, also $\lambda_2(\zeta) = \lambda_1^*(\zeta)$ belongs to the spectrum of \mathcal{L} . If, in the thermodynamic limit, $\text{Im}[\lambda_1(\zeta)]$ goes to zero only at $\zeta = \zeta_c$, so does $\text{Im}[\lambda_2(\zeta)]$. This would result in having three zeros in the Liouvillian spectrum at $\zeta = \zeta_c$. This is in contrast with the present theory of first-order dissipative phase transitions, which predicts a double degeneracy. Hence, the imaginary part of λ_1 must be zero in a finite domain around ζ_c .
- [67] B. Baumgartner and H. Narnhofer, Analysis of quantum semi-groups with GKS-Lindblad generators: II. General, *J. Phys. A: Math. Theor.* **41**, 395303 (2008).
- [68] A. Nagy and V. Savona, A driven-dissipative quantum Monte Carlo method for open quantum systems, *Phys. Rev. A* **97**, 052129 (2018).
- [69] A. Biella, J. Jin, O. Viyuela, C. Ciuti, R. Fazio, and D. Rossini, Linked cluster expansions for open quantum systems on a lattice, *Phys. Rev. B* **97**, 035103 (2018).
- [70] M. Biondi, S. Lienhard, G. Blatter, H. E. Türeci, and S. Schmidt, Spatial correlations in driven-dissipative photonic lattices, *New J. Phys.* **19**, 125016 (2017).
- [71] S. Finazzi, A. Le Boité, F. Storme, A. Baksic, and C. Ciuti, Corner-Space Renormalization Method for Driven-Dissipative Two-Dimensional Correlated Systems, *Phys. Rev. Lett.* **115**, 080604 (2015).
- [72] A. Kshetrimayum, H. Weimer, and R. Orús, A simple tensor network algorithm for two-dimensional steady states, *Nat. Commun.* **8**, 1291 (2017).
- [73] G. Vidal, Classical Simulation of Infinite-Size Quantum Lattice Systems in One Spatial Dimension, *Phys. Rev. Lett.* **98**, 070201 (2007).
- [74] M. Zwolek and G. Vidal, Mixed-State Dynamics in One-Dimensional Quantum Lattice Systems: A Time-Dependent Superoperator Renormalization Algorithm, *Phys. Rev. Lett.* **93**, 207205 (2004).
- [75] A. Le Boité, G. Orso, and C. Ciuti, Steady-State Phases and Tunneling-Induced Instabilities in the Driven Dissipative Bose-Hubbard Model, *Phys. Rev. Lett.* **110**, 233601 (2013).
- [76] A. Leuch, L. Papariello, O. Zilberberg, C. L. Degen, R. Chitra, and A. Eichler, Parametric Symmetry Breaking in a Nonlinear Resonator, *Phys. Rev. Lett.* **117**, 214101 (2016).
- [77] J. Cui, J. I. Cirac, and M. C. Bañuls, Variational Matrix Product Operators for the Steady State of Dissipative Quantum Systems, *Phys. Rev. Lett.* **114**, 220601 (2015).
- [78] M. S. Sarandy and D. A. Lidar, Adiabatic approximation in open quantum systems, *Phys. Rev. A* **71**, 012331 (2005).