



Deconfined criticality and a gapless \mathbb{Z}_2 spin liquid in the square-lattice antiferromagnetHenry Shackleton ¹, Alex Thomson,^{2,3,4} and Subir Sachdev ¹¹*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*²*Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA*³*Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, California 91125, USA*⁴*Department of Physics, California Institute of Technology, Pasadena, California 91125, USA*

(Received 29 April 2021; revised 23 June 2021; accepted 23 June 2021; published 6 July 2021)

The theory for the vanishing of Néel order in the spin $S = 1/2$ square lattice antiferromagnet has been the focus of attention for many decades. A consensus appears to have emerged in recent numerical studies on the antiferromagnet with first and second neighbor exchange interactions (the J_1 - J_2 model): A gapless spin liquid is present for a narrow window of parameters between the vanishing of the Néel order and the onset of a gapped valence bond solid state. We propose a deconfined critical SU(2) gauge theory for a transition into a stable \mathbb{Z}_2 spin liquid with massless Dirac spinon excitations; on the other side of the critical point, the SU(2) spin liquid (the ‘ π -flux’ phase) is presumed to be unstable to confinement to the Néel phase. We identify a dangerously irrelevant coupling in the critical SU(2) gauge theory, which contributes a logarithm-squared renormalization. This critical theory is also not Lorentz invariant and weakly breaks the SO(5) symmetry which rotates between the Néel and valence bond solid order parameters. We also propose a distinct deconfined critical U(1) gauge theory for a transition into the same gapless \mathbb{Z}_2 spin liquid; on the other side of the critical point, the U(1) spin liquid (the ‘staggered flux’ phase) is presumed to be unstable to confinement to the valence bond solid. This critical theory has no dangerously irrelevant coupling, dynamic critical exponent $z \neq 1$, and no SO(5) symmetry. All of these phases and critical points are unified in a SU(2) gauge theory with Higgs fields and fermionic spinons which can naturally realize the observed sequence of phases with increasing J_2/J_1 : Néel, gapless \mathbb{Z}_2 spin liquid, and valence bond solid.

DOI: [10.1103/PhysRevB.104.045110](https://doi.org/10.1103/PhysRevB.104.045110)**I. INTRODUCTION**

Antiferromagnetism on the square lattice became a topic of intense study soon after the discovery of high temperature superconductivity in the cuprates, and it continues to be a wellspring of interesting experimental and theoretical physics. It was established early on that the insulating antiferromagnet with $S = 1/2$ spins on each site and only nearest neighbor antiferromagnetic exchange interactions (J_1) has long-range Néel order in its ground state i.e., global SU(2) spin rotation symmetry was broken with the spin expectation value $\langle S_i \rangle = \eta_i N_0$ where S_i is the spin operator on site i , $\eta_i = \pm 1$ on the two checkerboard sublattices, and N_0 is the antiferromagnetic moment. Much attention has since been lavished on the insulating J_1 - J_2 antiferromagnet [1–6], which also has a second-neighbor antiferromagnetic exchange interaction J_2 . The key questions are the nature of the quantum phases of the model, and of the quantum phase transitions between them, as a function of increasing J_2/J_1 after the Néel order vanishes at a critical value of J_2/J_1 . These questions are also the focus of our attention here.

An early proposal [5,6,12,13] was that there was a direct transition from the Néel state to a valence bond solid (VBS) (see Fig. 1) which restores spin rotation symmetry but breaks lattice symmetries (followed by a first order transition at larger J_2/J_1 to a ‘columnar’ state which breaks spin rotation sym-

metry and which we do not address in the present paper). A theory of ‘deconfined criticality’ was developed [14–16] showing that a continuous Néel-VBS transition was possible, even though it was not allowed in the Landau-Ginzburg-Wilson framework because distinct symmetries were broken in the two phases. Evidence has since accumulated for the presence of a VBS phase in the J_1 - J_2 model, but the nature of the Néel-VBS transition in this model has remained a question of significant debate. However, in the past year, a consensus appears to have emerged [11] among the groups investigating this question by different numerical methods [7–10] and is summarized in Fig. 1: There is a narrow window with a gapless spin liquid phase between the Néel and VBS states. This gapless phase has been identified [8,17–20] as a \mathbb{Z}_2 spin liquid [5,6,21,22] with gapless, fermionic, $S = 1/2$ spinon excitations with a Dirac-like dispersion [18,23–26], labeled Z2Az13 in Wen’s classification [24].

The starting point of our analysis is the fermionic spinon dual [27–30] of the bosonic spinon CP¹ model used earlier [12,13,15,16] to describe the Néel-VBS transition. This fermionic dual is a relativistic SU(2) gauge theory of two flavors of two-component, massless Dirac fermions carrying fundamental gauge charges: This formulation is preferred over the bosonic spinons because the massless Dirac fermions connect naturally to the gapless fermionic spinons in the \mathbb{Z}_2 spin liquid. Recent studies [31–34] have indicated that the




FIG. 1. Phases of the $S = 1/2$ J_1 - J_2 antiferromagnet on the square lattice, from the numerical results of Refs. [7–10], all of which agree that the spin liquid is gapless. Each ellipse in the valence bond solid (VBS) represents a singlet pair of electrons. Lower part of figure adapted from Ref. [11].

two fermion flavor $SU(2)$ gauge theory does not ultimately describe a conformal field theory needed for Néel-VBS criticality but exhibits a ‘pseudocriticality’ associated with a proximate fixed point at complex coupling [27,35–37]. Reference [28] used connections to bosonic spinon theories to argue that the two fermion flavor $SU(2)$ gauge theory was ultimately unstable to confinement and symmetry breaking leading to the appearance of Néel order. We assume this is the case, and we can then describe the transition to the \mathbb{Z}_2 spin liquid by the condensation of Higgs fields which break the $SU(2)$ gauge symmetry down to \mathbb{Z}_2 : see Fig. 2. The Néel- \mathbb{Z}_2 spin liquid transition is a confinement-Higgs transition, and the critical theory is proposed to be a two-flavor $SU(2)$ gauge theory with critical Higgs fields [28]. We note that a similar critical theory was proposed in Ref. [38] for a continuous transition from the Néel state to a different gapless state with a \mathbb{Z}_2 gauge field (the ‘orthogonal semimetal’), and this scenario was supported there by quantum Monte Carlo simulations. Evidently, it is possible that critical Higgs fields can stabilize a scale-invariant critical point of the two-flavor $SU(2)$ gauge theory at the boundary of a Higgs phase where the $SU(2)$ gauge symmetry is broken down to \mathbb{Z}_2 .

As we will see below, an important difference between our critical Higgs $SU(2)$ gauge theory and that of Ref. [38] is that our theory does not preserve Lorentz invariance. The Lorentz symmetry is broken by the Yukawa couplings between the Higgs fields and fermions. The Yukawa couplings also do not preserve the $SO(5)$ flavor symmetry of the $SU(2)$ gauge theory with only fermionic matter; this symmetry rotates between the Néel and VBS states. Both these features have important




FIG. 2. Mean field phase diagram of our low energy theory obtained by minimization of the Higgs potential in Eq. (3.26). Dashed (solid red) lines indicate second (first) order transitions in mean field theory. We assume the $SU(2)$ π -flux gauge theory confines to a Néel state, the $U(1)$ staggered flux gauge theory confines to a VBS state, except at their deconfined critical boundaries to Wen’s stable, gapless \mathbb{Z}_2 spin liquid Z2Azz13. The dotted blue line indicates a possible trajectory of the square lattice antiferromagnet with increasing J_2/J_1 . However, as we discuss in Sec. VI, we cannot rule out interchanging the assignments of the confining states between the $SU(2)$ and $U(1)$ spin liquids, in which case the orientation of the blue arrow would be reversed. The critical $SU(2)$ gauge theory has a dangerously irrelevant coupling, but the critical $U(1)$ gauge theory does not. The mean-field analysis was performed with $w = u = 1$, $v = -1$, $\tilde{u} = 0.75$, and $v_4 = 0.5$ in Eq. (3.26). We use the ansatz $\Phi_1^a = c_1 \delta_{ax}$, $\Phi_2^a = c_1 \delta_{ay}$, and $\Phi_3^a = c_2 \delta_{az}$, so the terms in $V(\Phi)$ proportional to v_1, v_3 are automatically zero.

consequences for the Néel- \mathbb{Z}_2 spin liquid critical point and lead to predictions described below which can be tested by numerical studies.

In earlier work, Ran and Wen [39,40] had considered the two-flavor $SU(2)$ gauge theory as the description of an extended gapless phase on the square lattice—also called the π -flux phase [41]. They proposed a theory for a transition from the π -flux phase to the Z2Azz13 spin liquid by the condensation of a pair of adjoint Higgs fields, which we denote $\vec{\Phi}_{1,2}$ (the vector symbol implies gauge $SU(2)$ adjoint index). In light of our arguments above on the confining instability of the π -flux phase to the Néel state, the critical Higgs theory of Ran and Wen [39,40] can serve as the deconfined critical theory for the Néel to Z2Azz13 spin liquid transition. However, as we shall see in Sec. IV, additional ‘dangerously irrelevant’ terms are needed to fully define the critical theory in a $1/N_f$ expansion, and these contribute a logarithm-squared renormalization.

The mean-field phase diagram of the $SU(2)$ gauge theory with adjoint Higgs fields describing the π flux to Z2Azz13 transition turns out to naturally acquire an additional phase, as explained in Sec. III D—this is the $U(1)$ staggered flux spin

liquid [41]. We show that the adjoint Higgs field

$$\vec{\Phi}_3 \sim \vec{\Phi}_1 \times \vec{\Phi}_2 \quad (1.1)$$

[see Eq. (3.30)] is precisely that required to go from the SU(2) π -flux phase to the U(1) staggered flux phase. Specifically, starting from the π -flux phase, if both $\vec{\Phi}_{1,2}$ condense with $\langle \vec{\Phi}_1 \rangle \times \langle \vec{\Phi}_2 \rangle \neq 0$, we obtain the gapless \mathbb{Z}_2 spin liquid (the simultaneous condensation of $\vec{\Phi}_1$ and $\vec{\Phi}_2$ does not require fine tuning because of symmetry constraints that we will describe). On the other hand, Eq. (1.1) implies that if only the composite field $\vec{\Phi}_1 \times \vec{\Phi}_2$ condenses, but the individual fields $\vec{\Phi}_{1,2}$ do not, then the π -flux phase turns into the U(1) staggered flux phase. Speaking imprecisely, starting from the parent π -flux phase, the Higgs condensate for the gapless \mathbb{Z}_2 spin liquid is the ‘square root’ of the Higgs condensate for the staggered flux phase. (Let us also note that Song *et al.* [29] proposed that a trivial monopole would drive the staggered flux state into the π -flux state: so the Higgs field $\vec{\Phi}_3$ can be viewed as a ‘dual’ description of the trivial monopole and induces a transition in the opposite direction. Four-fermion terms have also been proposed as a route to reducing the emergent symmetry of the staggered flux state to that of the π -flux state [42].) The phase diagram of the Higgs fields $\vec{\Phi}_{1,2,3}$ is computed in Sec. III D and shown in Fig. 2. We propose here that the transition from the gapless \mathbb{Z}_2 spin liquid to the VBS state is described by the deconfined critical theory appearing at the onset of the U(1) spin liquid. Other works [25,29,30] have discussed the possible instability of this U(1) spin liquid to either Néel or VBS order via monopole proliferation. The critical U(1) gauge theory is described briefly in Sec. V, where we show that it does *not* contain the dangerously irrelevant terms found in the critical SU(2) theory.

We will review the derivation of the Ran-Wen theory and discuss its symmetry properties in some detail in Sec. II and Appendix A. A continuum SU(2) gauge theory coupled to three adjoint Higgs fields and gapless Majorana fermions will be obtained in Sec. III. The critical SU(2) gauge theory for the onset of the gapless \mathbb{Z}_2 spin liquid phase from the π -flux phase will be presented in Sec. IV, along with an analysis of its properties in a $1/N_f$ expansion. The critical U(1) gauge theory for the onset of the same gapless \mathbb{Z}_2 spin liquid from the staggered flux phase appears in Sec. V.

II. GAPLESS \mathbb{Z}_2 SPIN LIQUID GAPLESS \mathbb{Z}_2 SPIN LIQUID

The fermionic spinon theory of \mathbb{Z}_2 spin liquids proceeds by re-expressing the spin operators in terms of spinons $f_{i\alpha}$, $\alpha = \uparrow, \downarrow$ at site $\mathbf{i} = (i_x, i_y)$ of the square lattice using

$$\mathbf{S}_i = \frac{1}{2} \sum_{\alpha, \beta} f_{i\alpha}^\dagger \boldsymbol{\sigma}_{\alpha\beta} f_{i\beta}. \quad (2.1)$$

We write down a Bogoliubov Hamiltonian for the $f_{i\alpha}$ to obtain a \mathbb{Z}_2 spin liquid. Following Wen’s notation [24], we introduce the Nambu spinor

$$\psi_i = \begin{pmatrix} f_{i\uparrow} \\ f_{i\downarrow}^\dagger \end{pmatrix}, \quad (2.2)$$

resulting in the Bogoliubov Hamiltonian

$$H = - \sum_{ij} \psi_i^\dagger u_{ij} \psi_j. \quad (2.3)$$

Here,

$$u_{ij} = iu_{ij}^0 + u_{ij}^x \tau^x + u_{ij}^y \tau^y + u_{ij}^z \tau^z, \quad (2.4)$$

with τ^a Pauli matrices acting on the Nambu indices of ψ_i . Invariance under global SU(2)_s spin rotation requires that the u_{ij}^μ are all real numbers obeying

$$u_{ji}^0 = -u_{ij}^0, \quad u_{ji}^x = u_{ij}^x, \quad u_{ji}^y = u_{ij}^y, \quad u_{ji}^z = u_{ij}^z. \quad (2.5)$$

This fermionic spinon representation has a SU(2)_g gauge symmetry, under which

$$\text{SU}(2)_g : \psi_i \rightarrow U_{g,i} \psi_i \quad (2.6)$$

and a corresponding transformation for u_{ij} .

We will provide three different ansatzes for the u_{ij} in the Z2Azz13 spin liquid, each suited for different purposes. The three ansatzes are, of course, related to each other by SU(2)_g gauge transformations. Wen’s ansatz for the Z2Azz13 spin liquid is given in Appendix A, where the continuum Lagrangian describing the different spin liquid phases is deduced from symmetry fractionalization considerations. In the main text, we obtain the continuum theory directly from the lattice model, for which the ansatz given in Eq. (2.11) will be most useful. To derive this ansatz, we first describe the Z2Azz13 spin liquid by starting from the familiar staggered flux phase with U(1) gauge symmetry [41] and perturbing it with d_{xy} pairing. Explicitly, the ansatz is

$$\begin{aligned} \tilde{u}_{i,i+\hat{x}} &= \begin{pmatrix} te^{-i\phi} & 0 \\ 0 & -te^{i\phi} \end{pmatrix}, & i_x + i_y \text{ even} \\ \tilde{u}_{i,i+\hat{x}} &= \begin{pmatrix} te^{i\phi} & 0 \\ 0 & -te^{-i\phi} \end{pmatrix}, & i_x + i_y \text{ odd} \\ \tilde{u}_{i,i+\hat{y}} &= \begin{pmatrix} -te^{i\phi} & 0 \\ 0 & te^{-i\phi} \end{pmatrix}, & i_x + i_y \text{ even} \\ \tilde{u}_{i,i+\hat{y}} &= \begin{pmatrix} -te^{-i\phi} & 0 \\ 0 & te^{i\phi} \end{pmatrix}, & i_x + i_y \text{ odd} \\ \tilde{u}_{i,i+\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & -(\gamma_1 - i\gamma_2) \\ -(\gamma_1 + i\gamma_2) & 0 \end{pmatrix} \\ \tilde{u}_{i,i-\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & (\gamma_1 - i\gamma_2) \\ (\gamma_1 + i\gamma_2) & 0 \end{pmatrix}. \end{aligned} \quad (2.7)$$

The first four terms in (2.7) represent the fermion hopping, which is sketched in Fig. 3, and the last two terms are the d_{xy} pairing. With this ansatz, three distinct spin liquids may be described depending on the choice of parameters. These spin liquids are shown in Fig. 2, and we list them below:

(i) The π -flux phase with SU(2) gauge symmetry corresponds to $\phi = \pi/4$, and no fermion pairing $\gamma_{1,2} = 0$.

(ii) The ‘staggered flux’ U(1) spin liquid is obtained for general ϕ , and no fermion pairing $\gamma_{1,2} = 0$. The U(1) gauge field corresponds to a nearly spatially uniform modulation in the phases of the fermion hopping terms.

(iii) The Z2Azz13 spin liquid is obtained when the d_{xy} pairing $\gamma_1 + i\gamma_2$ is present and breaks the U(1) down to \mathbb{Z}_2 .




FIG. 3. Nearest-neighbor fermionic spinon hopping showing the A ($i_x + i_y$ even) and B ($i_x + i_y$ odd) sublattices.

Note that we have d_{xy} pairing in the \mathbb{Z}_2 spin liquid only, with opposite signs on the two sublattices.

In momentum space, we choose the A and B checkerboard sublattices as the basis sites (shown in Fig. 3), and the Hamiltonian acting on $(f_{A,k\uparrow}, f_{B,k\uparrow}, f_{A,-k,\downarrow}^\dagger, f_{B,-k,\downarrow}^\dagger)^T$ in the gauge

of Eq. (2.7) is

$$H = \begin{pmatrix} 0 & C_k & D_k & 0 \\ C_k^* & 0 & 0 & D_k \\ D_k^* & 0 & 0 & -C_k^* \\ 0 & D_k^* & -C_k & 0 \end{pmatrix} \quad (2.8)$$

where

$$\begin{aligned} C_k &= -2t(e^{-i\phi} \cos(k_x) - e^{i\phi} \cos(k_y)), \\ D_k &= 4(\gamma_1 - i\gamma_2) \sin(k_x) \sin(k_y). \end{aligned} \quad (2.9)$$

The eigenvalues of (2.8) are

$$\varepsilon_k = \pm([\text{Re}(C_k)]^2 + [\text{Im}(C_k) \pm |D_k|]^2)^{1/2} \quad (2.10)$$

and these coincide with those obtained in Wen's gauge in (A3). Note that the dispersion depends only on $|\gamma_1 + i\gamma_2|$ and not on $\gamma_{1,2}$ separately. This is natural in the staggered flux gauge, where U(1) the gauge transformation acts simply as $f_{i\alpha} \rightarrow f_{i\alpha} e^{i\phi_i}$, and so the d_{xy} pairing acts like a charge 2 Higgs field: A simple identification of the charge 2 Higgs field is the advantage of the present gauge. This dispersion is plotted in Fig. 4. The staggered flux phase has Dirac nodal points at $(\pm\pi/2, \pm\pi/2)$. Introducing d_{xy} pairing does not gap these nodal points but moves them away from these high symmetry points. Although the dispersion does not have full square lattice symmetry, all gauge-invariant observables do, and this is verified by the analysis in Appendix A.

A. Majorana gauge

For the remainder of the analysis in the body of the paper we map (2.7) onto the gauge used by Wang *et al.* [27] for the π -flux phase, which is convenient for eventual representation in Majorana fermions and making the gauge and spin rotation symmetries manifest. In this gauge, the ansatz of the Z2Azz13 spin liquid [which is gauge equivalent to Eq. (2.7)] is

$$\begin{aligned} \bar{u}_{i,i+\hat{x}} &= \begin{pmatrix} ite^{-4i\phi} & 0 \\ 0 & ite^{4i\phi} \end{pmatrix}, & i_x + i_y \text{ even} \\ \bar{u}_{i,i+\hat{x}} &= \begin{pmatrix} ite^{4i\phi} & 0 \\ 0 & ite^{-4i\phi} \end{pmatrix}, & i_x + i_y \text{ odd} \\ \bar{u}_{i,i+\hat{y}} &= (-1)^{i_x} \begin{pmatrix} te^{2i\phi} & 0 \\ 0 & -te^{-2i\phi} \end{pmatrix}, & i_x + i_y \text{ odd} \\ \bar{u}_{i,i+\hat{y}} &= (-1)^{i_x} \begin{pmatrix} -te^{-2i\phi} & 0 \\ 0 & te^{2i\phi} \end{pmatrix}, & i_x + i_y \text{ even} \\ \bar{u}_{i,i+\hat{x}+\hat{y}} = \bar{u}_{i,i-\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & (\gamma_1 - i\gamma_2)e^{-2i\phi} \\ (\gamma_1 + i\gamma_2)e^{2i\phi} & 0 \end{pmatrix}, & i_x + i_y \text{ even} \\ \bar{u}_{i,i+\hat{x}+\hat{y}} = \bar{u}_{i,i-\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & (-\gamma_1 + i\gamma_2)e^{4i\phi} \\ (-\gamma_1 - i\gamma_2)e^{-4i\phi} & 0 \end{pmatrix}, & i_x + i_y \text{ odd.} \end{aligned} \quad (2.11)$$

As in the previous gauge, the π -flux phase is obtained when $\phi = \pi/4$ while the staggered-flux phase corresponds to general ϕ .

III. CONTINUUM THEORY FOR HIGGS TRANSITION FROM SU(2) TO \mathbb{Z}_2

A. π -flux state with SO(5) symmetry

We begin by working out the continuum SU(2) gauge theory with the two-flavor massless Dirac fermion from the mean-field ansatz for the π -flux phase, using the Majorana gauge given in Eq. (2.11). In this gauge, we replace the Nambu

spinor in Eq. (2.2) by the matrix operator

$$\mathcal{X}_i = \begin{pmatrix} f_{i\uparrow} & -f_{i\downarrow}^\dagger \\ f_{i\downarrow} & f_{i\uparrow}^\dagger \end{pmatrix}. \quad (3.1)$$

The spinon SU(2) gauge symmetry of Eq. (2.6) now acts on \mathcal{X}_i as

$$\text{SU}(2)_g : \mathcal{X}_i \rightarrow \mathcal{X}_i U_{g,i}^\dagger. \quad (3.2)$$




FIG. 4. Plot of the dispersion, ε_k , of the fermionic spinons of the \mathbb{Z}_2 spin liquid Z2AZz13. The eigenvalues of the spinon Hamiltonian are $\pm\varepsilon_k$. All gauge invariant observables are invariant under the square lattice space group, although the spinon dispersion is not. The plot is of Eq. (2.10) for $t = 1.118$, $\phi = 0.464$, $\gamma_1 = 0.5$, $\gamma_2 = 0$.

The physical spin symmetry acts on \mathcal{X}_i on the left:

$$\text{SU}(2)_s : \mathcal{X}_i \rightarrow U_s \mathcal{X}_i. \quad (3.3)$$

We write the Bogoliubov Hamiltonian Eq. (2.3) as

$$H_{MF} = \sum_{\langle ij \rangle} [i\alpha_{ij} \text{Tr}(\mathcal{X}_i^\dagger \mathcal{X}_j) + \beta_{ij}^a \text{Tr}(\sigma^a \mathcal{X}_i^\dagger \mathcal{X}_j) + i\gamma_{ij} \text{Tr}(\sigma^a \mathcal{X}_i^\dagger \sigma^a \mathcal{X}_j)]. \quad (3.4)$$

The correspondence with the notation in Eq. (2.3) is

$$u_{ij} = i\alpha_{ij}\tau^0 + \beta_{ij}^a \tau^a. \quad (3.5)$$

The additional γ_{ij} hoppings involve projective realizations of the spin rotation symmetry and will not be relevant. The degrees of freedom in this Hamiltonian can be represented by four Majorana fermions,

$$\mathcal{X}_i = \frac{1}{\sqrt{2}}(\chi_0 + i\chi_a \sigma^a). \quad (3.6)$$

The $\text{SU}(2)$ -invariant π -flux state comes from the hoppings $\beta^a = 0$ and

$$\alpha_{ij} = -\alpha_{ji} \quad \alpha_{i+\hat{x},i} = 2t \quad \alpha_{i+\hat{y},i} = (-1)^{i_x} 2t. \quad (3.7)$$

The low-energy behavior of this mean-field ansatz is described by an $\text{SU}(2)$ gauge theory with an emergent $\text{SO}(5)$ symmetry. To work out the dispersion relation of this Hamiltonian, we increase our unit cell by one lattice site in the x direction and so χ acquires an additional sublattice index $m = A, B$. Note that this unit cell differs slightly from the one used in the staggered flux gauge. In momentum space, we then have

$$H = \sum_k \chi_{-k}^\dagger H(k) \chi_k, \quad (3.8)$$

$$H(k) = -2t[\sin(k_y)\rho^z + \sin(k_x)\rho^x].$$

ρ^i are Pauli operators acting on the sublattice space, $m = A, B$. This Hamiltonian is diagonal in the $0, a$ indices in Eq. (3.6), and the gauge was chosen to have this feature. The Hamiltonian in Eq. (3.8) has Dirac points at $k_y = 0, \pi$, $k_x = 0$. Labelling these Dirac points by another index $v = 1, 2$ and expanding around these two points, we decompose our Majorana operator as

$$\chi_{m,i} \sim \rho^x \chi_{m,v=1}(x) + (-1)^{i_y} \chi_{m,v=2}(x). \quad (3.9)$$

With this, the Hamiltonian reduces to

$$H \approx 2it \sum_{v=1,2} \chi_v^\dagger (\rho^x \partial_x - \rho^z \partial_y) \chi_v, \quad (3.10)$$

with the sublattice and $0, a$ indices implicit. This gives the continuum Lagrangian

$$\mathcal{L}_{MF} = 2it \bar{\chi}_v \gamma^\mu \partial_\mu \chi_v \quad (3.11)$$

where $\gamma^0 = \rho^y$, $\gamma^x = i\rho^z$, $\gamma^y = i\rho^x$, and $\bar{\chi} \equiv \chi^T \gamma^0$. Here we have chosen to express \mathcal{L}_{MF} in the Minkowski metric (+, −, −); we ultimately move to the Euclidean metric below to perform calculations.

We now define the 4×2 matrix operator

$$X_{\alpha,v;\beta} = \frac{1}{\sqrt{2}}(\chi_{0,v} \delta_{\alpha\beta} + i\chi_{a,v} \sigma_{\alpha\beta}^a) \quad (3.12)$$

and $\bar{X} = X^\dagger \gamma^0$, where the sublattice/Dirac index m is left implicit. This lets us write our Lagrangian as

$$\mathcal{L}_{MF} = i\text{Tr}(\bar{X} \gamma^\mu \partial_\mu X), \quad (3.13)$$

where we set $t = 1/2$ from now on. In this form, the Hamiltonian describes eight massless Majorana fermions (these are two-component ‘relativistic’ Majorana fermions with an additional sublattice index). The $\text{SU}(2)$ gauge symmetry acts on the right index [β in Eq. (3.12)] of X , and the gradient in \mathcal{L}_{MF} must be replaced by the appropriate covariant gradient when the gauge field is included. Global spin rotations act of the left index [α in Eq. (3.12)] of X , and global valley rotations act of the v index. These global rotations combine to yield an




FIG. 5. Shown are the leading-order perturbations that away from the $\text{SU}(2)$ π -flux state, in the Majorana gauge given by Eq. (2.11). Note that the unit cell, with sublattice sites A and B , differs from the gauge illustrated in Fig. 3. (Left) The perturbation that shifts the π -flux state to the staggered flux state, with hoppings proportional to τ^z . Thickness of the line denotes strength (weaker in the y direction) and solid/dashed indicates positive/negative sign. (Right) The d_{xy} pairing that breaks the $\text{U}(1)$ gauge symmetry to \mathbb{Z}_2 , with pairing $\gamma_1 \tau^y - \gamma_2 \tau^x$ on solid lines and $\gamma_1 \tau^x + \gamma_2 \tau^y$ on dashed lines.

emergent, low energy $\text{Sp}(4)/\mathbb{Z}_2 \equiv \text{SO}(5)$ global symmetry in the π -flux phase [27,39].

In the following subsections, we derive the continuum form of the perturbations given in Eq. (2.11), which break the π -flux state down to either the staggered flux state or the Z2Azz13 spin liquid. We do so by rewriting these perturbations in terms of the low-energy modes given in Eq. (3.9) and keeping only the lowest order gradient terms. These perturbations are coupled to adjoint Higgs fields, and the transition of the π -flux state to either the staggered flux state or Z2Azz13 spin liquid is obtained by condensing the corresponding Higgs fields. An alternative derivation of these continuum perturbations based on symmetry fractionalization is provided in Appendix A and agrees with the following analysis.

B. From π flux to staggered flux

We obtain the continuum version of the perturbations to the staggered flux phase by expanding the mean field param-

eters \bar{u}_{ij} defined in Eq. (2.11) in powers of $\phi = \pi/4 + \delta\phi$. We subsequently employ Eq. (3.5), which in turn yields additional hopping parameters to the Hamiltonian of the form

$$\beta_{i,i+\hat{x}}^z = -4\delta\phi(-1)^{i_x+i_y}, \quad \beta_{i,i+\hat{y}}^z = 2\delta\phi(-1)^{i_y}. \quad (3.14)$$

These terms are illustrated in Fig. 5. If we look at the components of the Majorana fermions [as defined in Eq. (3.6), with $(0, a)$, $a = x, y, z$], we see that these new terms introduce hopping between the $0 \leftrightarrow z$ and $x \leftrightarrow y$ Majorana fermions. For simplicity, we focus on the $0 \leftrightarrow z$ hoppings, as the $x \leftrightarrow y$ hoppings will be identical. We start with the hoppings in the x direction, expand our Majorana operators in terms of low-energy modes, and keep only the lowest-order gradient terms. As in Eq. (3.12), the two indices on χ correspond to $(0, x, y, z)$ and valley, respectively, with the sublattice index implicit.

$$\begin{aligned} \delta H &= -4\delta\phi \sum_i [\chi_{0,1}^T(x_i)\rho^x + (-1)^{i_y}\chi_{0,2}^T(x_i)](-1)^{i_y}\rho^x[\rho^x\chi_{z,1}(x_i) + (-1)^{i_y}\chi_{z,2}(x_i)] \\ &\quad + 2\delta\phi \sum_i [\chi_{0,1}^T(x_i)\rho^x + (-1)^{i_y}\chi_{0,2}^T(x_i)](-1)^{i_y}(\rho^x - i\rho^y)[\rho^x\chi_{z,1}(x_{i+\hat{x}}) + (-1)^{i_y}\chi_{z,2}(x_{i+\hat{x}})] \\ &\quad + 2\delta\phi \sum_i [\chi_{0,1}^T(x_i)\rho^x + (-1)^{i_y}\chi_{0,2}^T(x_i)](-1)^{i_y}(\rho^x + i\rho^y)[\rho^x\chi_{z,1}(x_{i-\hat{x}}) + (-1)^{i_y}\chi_{z,2}(x_{i-\hat{x}})] \\ &\approx 2\delta\phi \int d^2x [\chi_{0,1}^T\rho^z\partial_x\chi_{z,2} - \chi_{0,2}^T\rho^z\partial_x\chi_{z,1}] \\ \Rightarrow \delta\mathcal{L} &= -2i\delta\phi\text{Tr}(\sigma^z\bar{X}\mu^y\gamma^y\partial_x X) \end{aligned} \quad (3.15)$$

In our final term, we have reintroduced the $x \leftrightarrow y$ hoppings. For the hoppings in the y direction,

$$\begin{aligned} \delta H &= 2\delta\phi \sum_i [\chi_{0,1}^T(x_i)\rho^x + (-1)^{i_y}\chi_{0,2}^T(x_i)](-1)^{i_y}[\rho^x\chi_{z,1}(x_{i+\hat{y}}) - (-1)^{i_y}\chi_{z,2}(x_{i+\hat{y}})] \\ &\quad - 2\delta\phi \sum_i [\chi_{0,1}^T(x_i)\rho^x + (-1)^{i_y}\chi_{0,2}^T(x_i)](-1)^{i_y}[\rho^x\chi_{z,1}(x_{i-\hat{y}}) - (-1)^{i_y}\chi_{z,2}(x_{i-\hat{y}})] \\ &\approx -2\delta\phi \int d^2x [\chi_{0,1}^T\rho^x\partial_y\chi_{z,2} - \chi_{0,2}^T\rho^x\partial_y\chi_{z,1}] \\ \Rightarrow \delta\mathcal{L} &= -2i\delta\phi\text{Tr}(\sigma^z\bar{X}\mu^y\gamma^x\partial_y X). \end{aligned} \quad (3.16)$$

Note that here and in Eq. (3.15) the Pauli matrix σ^z is acted on by the $\text{SU}(2)$ gauge symmetry of the π -flux phase. Gauge invariance requires there exist nearly identical continuum model bilinears containing instead σ^x and σ^y Pauli matrices. It is therefore useful to express the perturbation in a gauge independent fashion using an adjoint Higgs field Φ_3^a , where $a = x, y, z$ is a $\text{SU}(2)$ gauge index:

$$\delta\mathcal{L} = \Phi_3^a\text{Tr}[\sigma^a\bar{X}\mu^y(\gamma^y i\partial_x + \gamma^x i\partial_y)X]. \quad (3.17)$$

(Our choice of subscript ‘‘3’’ will be clear shortly.) This Higgs field mediates the onset of the staggered flux phase, and in this present gauge we have the identification

$$\Phi_3^z \sim \delta\phi. \quad (3.18)$$

Condensing the Higgs field with $\langle\Phi_3^a\rangle$ leads to a transition from the $\text{SU}(2)$ π -flux state to the $\text{U}(1)$ staggered flux state. For concreteness, we continue to work in the gauge where Φ_3^a condenses in the z direction, as implied by Eqs. (3.15) and (3.16).

C. From π flux to Z2Az13

We now evaluate the effects of a nonzero $\gamma_{1,2}$ in the π -flux phase, using the Majorana gauge as given in Eq. (2.11). We first consider turning on the perturbation

$$\bar{u}_{i,i+\hat{x}+\hat{y}} = \bar{u}_{i,i-\hat{x}+\hat{y}} = \gamma_1 \tau^y - \gamma_2 \tau^x, \quad i_x + i_y \text{ even.} \quad (3.19)$$

Recall that in the Majorana basis, terms proportional to τ^x (τ^y) correspond to hoppings between the $0 \leftrightarrow x(y)$ and $z \leftrightarrow y(x)$ Majorana fermions. Focusing on the γ_1 term, we expand in low-energy modes

$$\begin{aligned} \delta H &= \gamma_1 \sum_i [\chi_{0,1}^T(x_i) \rho^x + (-1)^{i_y} \chi_{0,2}^T(x_i)] [\rho^x + (-1)^{i_y} \rho^y] [\rho^x \chi_{y,1}(x_{i+\hat{x}+\hat{y}}) - (-1)^{i_y} \chi_{y,2}(x_{i+\hat{x}+\hat{y}})] \\ &\approx \gamma_1 \sum_i \chi_{0,1}^T(x_i) \rho^x \chi_{y,1}(x_i) - \chi_{0,2}^T(x_i) \rho^x \chi_{y,2}(x_i) + \chi_{0,1}^T(x_i) \rho^z \chi_{y,2}(x_i) + \chi_{0,2}^T(x_i) \rho^z \chi_{y,1}(x_i) \\ \Rightarrow \delta \mathcal{L} &= \gamma_1 \text{Tr}[\sigma^y \bar{X}(\mu^z \gamma^x + \mu^x \gamma^y) X]. \end{aligned} \quad (3.20)$$

The perturbation is identical for the γ_2 term but with $\sigma^y \Rightarrow -\sigma^x$.

As in the previous section, the addition of the hopping parameters of Eq. (3.19) can be formulated in a gauge-invariant fashion by coupling the bilinear above to an adjoint Higgs field Φ_1^a , $a = x, y, z$ (the bar on the “1” will be apparent below). In particular, when a term $\Phi_1^a \text{Tr}[\sigma^a \bar{X}(\mu^z \gamma^x + \mu^x \gamma^y) X]$ is added to the Lagrangian, we reproduce the continuum version of Eq. (3.19) we just derived when Φ_1^a condenses as

$$\langle \Phi_1^x \rangle \sim \gamma_2, \quad \langle \Phi_1^y \rangle \sim \gamma_1. \quad (3.21)$$

We perform the same analysis for the second term proportional to $\gamma_{1,2}$:

$$u_{i,i+\hat{x}+\hat{y}} = u_{i,i-\hat{x}+\hat{y}} = \gamma_1 \tau^x + \gamma_2 \tau^y, \quad i_x + i_y \text{ odd.} \quad (3.22)$$

The continuum derivation of this is essentially identical to as before, yielding

$$\delta \mathcal{L} = \text{Tr}[(\gamma_1 \sigma^x + \gamma_2 \sigma^y) \bar{X}(\mu^z \gamma^x - \mu^x \gamma^y) X], \quad (3.23)$$

prompting us to introduce $\Phi_2^a \text{Tr}[\sigma^a \bar{X}(\mu^z \gamma^x - \mu^x \gamma^y) X]$. The continuum version of Eq. (3.19) is obtained through the condensation Φ_2^a such that $\langle \Phi_2^a \rangle = \gamma_1 \delta_{ax} + \gamma_2 \delta_{ay}$.

D. Majorana-Higgs Lagrangian

We now combine the results of Secs. III A, III B, and III C to obtain the low energy Lagrangian for the Majorana field X , and three real, adjoint Higgs scalars, which we now identify as $\Phi_1^a, \Phi_2^a, \Phi_3^a$ ($\Phi_{1,2}^a$ are rotations of $\Phi_{1,2}^a$ in the 1,2 plane). We do not explicitly write out the coupling to the SU(2) gauge field in this subsection, which can be included by the usual requirements of minimal coupling.

The Lagrangian is

$$\begin{aligned} \mathcal{L} &= i \text{Tr}(\bar{X} \gamma^\mu \partial_\mu X) + \Phi_1^a \text{Tr}(\sigma^a \bar{X} \mu^z \gamma^x X) + \Phi_2^a \text{Tr}(\sigma^a \bar{X} \mu^x \gamma^y X) \\ &\quad + \Phi_3^a \text{Tr}(\sigma^a \bar{X} \mu^y (\gamma^y i \partial_x + \gamma^x i \partial_y) X) + V(\Phi). \end{aligned} \quad (3.24)$$

The staggered flux state is obtained when $\langle \Phi_3 \rangle \propto (0, 0, \delta\phi)$. The Z2Az13 state follows from $\langle \Phi_1 \rangle \propto (\gamma_1 - \gamma_2, \gamma_1 + \gamma_2, 0)$ and $\langle \Phi_2 \rangle \propto (-\gamma_1 - \gamma_2, \gamma_1 - \gamma_2, 0)$.

The Higgs potential $V(\Phi)$ arises from integrating out the high energy spinon degrees of freedom. We deduce its form by carefully considering the symmetry properties of the theory, which are described in some detail in Appendix A. Here, we

note that the theory should respect time reversal and the lattice symmetries,

$$\begin{aligned} T_x : (i_x, i_y) &\mapsto (i_x + 1, i_y), & T_y : (i_x, i_y) &\mapsto (i_x, i_y + 1), \\ P_x : (i_x, i_y) &\mapsto (-i_x, i_y), & P_y : (i_x, i_y) &\mapsto (i_x, -i_y), \\ R_{\pi/2} : (i_x, i_y) &\mapsto (-i_y, i_x), \end{aligned} \quad (3.25)$$

and we summarize the transformations of the Higgs fields here:

	T_x	T_y	P_x	P_y	\mathcal{T}	$R_{\pi/2}$
Φ_1^a	−	+	−	−	−	− Φ_2^a
Φ_2^a	+	−	−	−	−	− Φ_1^a
Φ_3^a	−	−	+	+	+	−

From this, we can deduce that the following gauge-invariant terms are allowed to quartic order in the Higgs potential

$$\begin{aligned} V(\Phi) &= s(\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a) + \tilde{s} \Phi_3^a \Phi_3^a + w \epsilon_{abc} \Phi_1^a \Phi_2^b \Phi_3^c \\ &\quad + u(\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a)^2 + \tilde{u}(\Phi_3^a \Phi_3^a)^2 + v_1(\Phi_1^a \Phi_2^a)^2 \\ &\quad + v_2(\Phi_1^a \Phi_1^a)(\Phi_2^b \Phi_2^b) + v_3[(\Phi_1^a \Phi_3^a)^2 + (\Phi_2^a \Phi_3^a)^2] \\ &\quad + v_4(\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a)(\Phi_3^b \Phi_3^b), \end{aligned} \quad (3.26)$$

where ϵ_{abc} is the antisymmetric unit tensor.

An important feature of $V(\Phi)$ is the cubic term proportional to w . This term implies that if any two of the Higgs fields are condensed, then so must the third. It also shows that even if we were only considering the transition from the SU(2) π -flux phase to the gapless \mathbb{Z}_2 spin liquid by the condensation of $\Phi_{1,2}^a$, we would be forced to include Φ_3^a in our theory and hence the additional possibility of a U(1) staggered flux phase. The symmetry transformations show that Φ_3^a is the unique adjoint Higgs field that can be made from the tensor product of the Higgs fields needed to describe the gapless \mathbb{Z}_2 spin liquid, Φ_1^a and Φ_2^a : so the staggered flux phase is a natural partner of this gapless \mathbb{Z}_2 spin liquid and the π -flux phase.

We can perform a mean-field minimization of Eq. (3.26), and typical results are shown in Fig. 2. There are three phases

as a function of the tuning parameters s and \tilde{s} , which correspond to exactly those obtained in the lattice mean-field theory described in Sec. II. The presence of the w term implies that there is a first order transition line near the point where the three phases meet [43], as shown in Fig. 2. We summarize and re-express the lattice theory results in terms of the continuum model parameters below.

1. SU(2) π -flux phase

Here, there is no Higgs condensate $\langle \Phi_{1,2}^a \rangle = 0$, $\langle \Phi_3^a \rangle = 0$, and the system lies in the red region on the top right of Fig. 2: the SU(2) π -flux phase. The continuum model possesses an SU(2) gauge symmetry, along with the corresponding gauge bosons. The theory is believed to confine to the Néel or VBS phase—as discussed in Secs. I and VI, we view the Néel phase to be more likely.

2. U(1) Staggered flux phase

This state as $\langle \Phi_3^a \rangle$ nonzero, while $\langle \Phi_{1,2}^a \rangle = 0$, resulting in the U(1) staggered flux phase represented on the top left of Fig. 2. Making contact with the lattice ansatz, we have

$$\langle \Phi_3^a \rangle \propto (0, 0, \phi - \pi/4) \neq 0. \quad (3.27)$$

Again, the theory has a continuous unbroken gauge degree of freedom, now with only a U(1) symmetry. There is a single gauge boson, which we nevertheless assume triggers confinement. As argued, the most likely fate of the theory is the VBS state, but we cannot preclude the Néel phase.

3. \mathbb{Z}_2 spin liquid Z2Azz13

The \mathbb{Z}_2 spin liquid Z2Azz13 corresponds to a Higgs condensate satisfying $\langle \Phi_{1,2}^a \rangle \neq 0$; it is shown in the lower half of the phase diagram of Fig. 2. The symmetry transformations imply that Φ_1^a and Φ_2^a have the same mass, so only a single tuning parameter s is required to make them condense from the SU(2) π -flux phase. From the symmetry transformations, we also see that the absence of a broken symmetry requires that the gauge-invariant bilinears obey

$$\langle \Phi_1^a \Phi_1^a \rangle = \langle \Phi_2^a \Phi_2^a \rangle > 0, \quad \langle \Phi_1^a \Phi_2^a \rangle = 0. \quad (3.28)$$

Such saddle points are obtained from the Higgs potential for a range of v_1 positive and v_2 negative. Moreover, such a saddle point is indeed present in the lattice ansatz of the previous section where

$$\begin{aligned} \langle \Phi_{1a} \rangle &\propto (-\gamma_1 - \gamma_2, \gamma_1 - \gamma_2, 0), \\ \langle \Phi_{2a} \rangle &\propto (\gamma_1 - \gamma_2, \gamma_1 + \gamma_2, 0). \end{aligned} \quad (3.29)$$

We note that this implies $\langle \Phi_1 \rangle \perp \langle \Phi_2 \rangle$ and $|\langle \Phi_1 \rangle| = |\langle \Phi_2 \rangle|$, where we use a vector shorthand for the indices $a = x, y, z$ of the Higgs fields. By minimizing the potential $V(\Phi)$ in Eq. (3.26), we see that this \mathbb{Z}_2 spin liquid also implies the condensation of the remaining Higgs field:

$$\langle \Phi_3^a \rangle \propto w \epsilon_{abc} \langle \Phi_1^b \rangle \langle \Phi_2^c \rangle. \quad (3.30)$$

It follows that \tilde{s} can change sign within this phase without any phase transition.

E. Visons

The \mathbb{Z}_2 spin liquid is obtained from the theory in Eq. (3.24) + SU(2) gauge fields [which is Eq. (4.2) below] by condensing $\Phi_{1,2}^a$. This spin liquid has gapless fermionic spinon excitations, whose low energy dispersion can also be determined from the continuum theory. However, as in all \mathbb{Z}_2 spin liquids, there must also be vison excitations, which are mutual semions with respect to the spinons. In the theory in Eq. (4.2), the vison is a finite energy excitation associated with vortexlike saddle point in which the Higgs fields $\Phi_{1,2}^a$ undergo a topologically nontrivial SO(3) rotation, associated with $\pi_1(\text{SO}(3)) = \mathbb{Z}_2$, around the core of the vortex: see Ref. [44] for an explicit solution in a theory without the fermionic spinons. Given that the vison appears in a lattice model with a background spinon density of one spinon per site, we expect the vison transforms projectively under translational symmetries with $T_x T_y = -T_y T_x$, where T_α is translation by one lattice spacing in the α direction [23,45–48]. For the case of gapped spinons, this fact now has a modern interpretation in the theory of symmetry fractionalization in topological phases [49–53]. We expect that a similar result applies in the present gapless spinon case, but this has not been explicitly established. For the case of gapped spinons, the vison projective transformation can be derived from a parent U(1) gauge theory (which is Higgsed down to \mathbb{Z}_2) in which the monopoles carry Berry phases [45,46,48]. Such monopole Berry phases are in turn related to a SO(5) Wess-Zumino-Witten term in an effective theory the Néel and VBS order parameters [54,55]. Notably, this SO(5) WZW term is also linked to an anomaly of the Majorana theory in Eq. (4.2) [27]. It would therefore be interesting to establish $T_x T_y = -T_y T_x$ for gapped visons in the presence of gapless spinons starting directly from Eq. (4.2) and condensing the Higgs fields: We leave such an analysis for future work.

IV. RENORMALIZED PERTURBATION EXPANSION FOR THE CRITICAL SU(2) GAUGE THEORY

This section will present an analysis of the transition obtained by tuning the Higgs ‘mass’ s in Eq. (3.26) across a quantum critical point at $s = s_c$, for $\tilde{s} > 0$ in Fig. 2, between the SU(2) and \mathbb{Z}_2 spin liquids. We have $\langle \Phi_{1,2,3}^a \rangle = 0$ for $s > s_c$, yielding the π -flux spin liquid. For $s < s_c$, we have $\langle \Phi_{1,2}^a \rangle \neq 0$ yielding the \mathbb{Z}_2 spin liquid Z2Azz13. As we noted below Eq. (3.29), $\langle \Phi_3^a \rangle$ will also be nonzero once both $\langle \Phi_{1,2}^a \rangle$ are nonzero. However, as $\langle \Phi_3^a \rangle$ is quadratic in $\langle \Phi_{1,2}^a \rangle$ [see Eq. (3.30)], it is not a primary order parameter for the transition. So we can entirely neglect Φ_3^a in the analysis of the criticality in the present section.

It is also convenient to write the theory in terms of two flavors of complex Dirac fermions which also carry a fundamental SU(2) gauge charge, $\psi_{\alpha,v}$; here α is the SU(2) gauge index, $v = 1, 2$ is the valley index, and the Dirac/sublattice index is suppressed. The global SU(2) spin symmetry is not manifest in this formalism, unlike in the earlier Majorana formalism. Since the Lagrangian in Eq. (3.24) does not contain terms that act on the physical SU(2) spin, our Lagrangian nevertheless has a simple form in terms of these Dirac fermions, although a more careful analysis will be required to calculate

the behavior of the Néel order parameter, which does involve the physical SU(2) spin. Explicitly, the relationship between the Dirac and Majorana fermions is

$$\psi_{\alpha,v} = i\sigma_{\alpha,\beta}^y X_{1,v;\beta}. \quad (4.1)$$

Applying this change of variables to Eq. (3.24), and including the SU(2) gauge field A_μ^a , we obtain the Lagrangian for ψ and the $\Phi_{1,2}^a$ Higgs fields

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_\psi + \mathcal{L}_\Phi + \mathcal{L}_{\Phi\psi} \\ \mathcal{L}_\psi &= i \sum_v \bar{\psi}_v \gamma^\mu (\partial_\mu - iA_\mu^a \sigma^a) \psi_v. \\ \frac{\mathcal{L}_\Phi}{N_f} &= \frac{K}{2} [(\partial_x \Phi_1^a - 2\epsilon_{abc} A_x^b \Phi_1^c)^2 + (\partial_y \Phi_2^a - 2\epsilon_{abc} A_y^b \Phi_2^c)^2] \\ &\quad + \frac{s}{2} (\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a) + u (\Phi_1^a \Phi_1^a + \Phi_2^a \Phi_2^a)^2 \\ &\quad + v_1 (\Phi_1^a \Phi_2^a)^2 + v_2 (\Phi_1^a \Phi_1^a) (\Phi_2^b \Phi_2^b) \\ \mathcal{L}_{\Phi\psi} &= \lambda (\Phi_1^a \bar{\psi} \mu^z \gamma^x \sigma^a \psi + \Phi_2^a \bar{\psi} \mu^x \gamma^y \sigma^a \psi). \end{aligned} \quad (4.2)$$

We will henceforth work in Euclidean signature, with $(\gamma^\mu)^2 = 1$ for all μ . This Lagrangian includes an important new term not present in Eq. (3.24): a bare spatial gradient term for the Higgs field proportional to the coupling K (we will define N_f shortly). This coupling is allowed by symmetry and will turn out to be ‘dangerously irrelevant,’ i.e., under renormalization, K flows to zero, but it cannot be set to zero at the outset because of some singular effects that we will describe below. In contrast, the quartic couplings $u, v_{1,2}$ are genuinely irrelevant at the critical point and will not be considered further.

The theory \mathcal{L} is invariant under SU(2) gauge, SU(2) spin rotation, time-reversal, and space group transformations, as it must be, because these are symmetries of the underlying Hamiltonian and its parton representation. However, the Yukawa coupling λ breaks both the emergent Lorentz and SO(5) symmetries of the fermion kinetic term. As we will show below, λ is not an irrelevant perturbation, and so the absence of these emergent symmetries will be apparent in the critical correlation functions.

We will analyze the critical properties of Eq. (4.2) by the $1/N_f$ expansion used in earlier treatments of Dirac fermions coupled to scalar fields by Yukawa couplings which break relativistic invariance [56]. For this purpose, we will endow




FIG. 6. The leading order effective propagators for the Higgs (left) and gauge field (right) are generated by the one-loop contributions from N fermions.

the fermions with an additional flavor index (not shown explicitly) which ranges over N_f values. Combined with the v index, there are a total of $2N_f$ flavors and two colors of two-component Dirac fermions. The physical case of interest to us is $N_f = 1$.

As in Ref. [56], we will compute the renormalization constants of the theory \mathcal{L} in a $1/N_f$ expansion. The most important of these will be the renormalization of the Fermi velocity, which has been implicitly set to unity above: This is nonzero because of the lack of the Lorentz invariance in the Yukawa coupling. The renormalization of the Fermi velocity in turn defines a dynamic critical exponent z : We will compute z to order $1/N_f$ and find it to be a universal number at this order. Next, we shall examine the renormalization of the field scales. As in Ref. [56], a convenient choice, as we explain below, is to renormalize the boson field scale Φ so that the Yukawa coupling $\lambda = 1$; we will assume $\lambda = 1$ below. As usual, the renormalization of the fermion field Z_ψ is determined from the fermion self energy, which then determines a fermion anomalous dimension η_ψ . Here we will find an unusual phenomenon, which is one of our main results: The value of η_ψ is not universal at order $1/N_f$ but has a logarithmic dependence upon the irrelevant coupling K . Finally, we will also compute the renormalization of the fermion bilinears associated with the Néel and VBS order parameters: These are not equal to each other because the SO(5) symmetry is explicitly broken.

A. Boson propagators

The first step in the large N_f expansion is to integrate out the large number of fermions ψ , which allows us to determine the propagators of the bosons: the Higgs fields and the gauge fields. To leading order in $1/N_f$, we have to evaluate the diagrams in Fig. 6, and this leads to an effective quadratic action of the following form

$$\frac{S_b}{N_f} = \int_k \frac{1}{2} (s + Kk_x^2 + \Gamma_1(k)) \Phi_1^a(k) \Phi_1^a(-k) + \frac{1}{2} (s + Kk_y^2 + \Gamma_2(k)) \Phi_2^a(k) \Phi_2^a(-k) + \frac{\Gamma_A(k)}{2} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) A_\mu^a(k) A_\nu^a(-k).$$

We work in the Euclidean time signature, and k is a 3-momentum.

We first calculate the one-loop corrections to the Higgs propagators. The correction to the Φ_1 propagator is shown in the first diagram in Fig. 6 and is

$$\begin{aligned} \Gamma_1(k) \delta_{ab} &= \lambda^2 \text{Tr} \int \frac{d^3 p}{(2\pi)^3} [\mu^z \gamma^x \sigma^a] \left[\frac{\not{p}}{p^2} \right] [\mu^z \gamma^x \sigma^b] \left[\frac{\not{p} + \not{k}}{(k+p)^2} \right] \\ &= -8\lambda^2 \delta_{ab} \int \frac{d^3 p}{(2\pi)^3} \frac{p_0(p_0 + k_0) - p_x(p_x + k_x) + p_y(p_y + k_y)}{p^2(k+p)^2} = \frac{\lambda^2 \delta_{ab}}{4} \frac{(k_0^2 + k_y^2)}{\sqrt{k^2}}. \end{aligned} \quad (4.3)$$




FIG. 7. The two leading order contributions to the fermion self-energy, arising from Higgs (left) and gauge boson (right) couplings. To leading order in $1/N_f$, both the Higgs and gauge boson propagators are generated by the fermions.

We have omitted a constant term, which will be tuned to zero at the critical point.

The correction to the Φ_2 propagator is identical to the Φ_1 correction, with $k_x \leftrightarrow k_y$.

$$\Gamma_2(k) = \frac{\lambda^2 (k_0^2 + k_x^2)}{4 \sqrt{k^2}} \quad (4.4)$$

The reader should now notice some key features. As in Ref. [56], the overall scaling in momentum is $\Gamma_{1,2} \sim |k|$. So, this fermion-induced contribution to the Φ propagators is more important at low momenta than the k^2 terms which would be present in the bare theory. In general, the bare boson k^2 terms are irrelevant, and this is why we choose to set the field scale of Φ with the renormalization condition $\lambda = 1$. However, unlike Ref. [56], we will see below in some detail that we cannot entirely ignore the bare k^2 term. The expression for Γ_1 (Γ_2) is not an increasing function of k_x (k_y) when it is larger than the other momentum components, and this will lead to infrared singularities at first order in $1/N_f$. Specifically, the integral over the propagator $1/\Gamma_1$ ($1/\Gamma_2$) has an infrared divergence in the k_0, k_y (k_0, k_x) plane. Consequently, we do need to include the *dangerously* irrelevant Kk_x^2 (Kk_y^2) term in the bare action for Φ_1^a (Φ_2^a), as we have anticipated in Eqs. (4.2) and (4.3).

The $\mathcal{O}(N_f)$ propagator for the gauge field is obtained from

$$\begin{aligned} & \Gamma_A(k)(k^2 \delta^\mu \delta^\nu - k^\mu k^\nu) \\ &= -\text{Tr} \int \frac{d^3 p}{(2\pi)^3} \gamma^\mu \left[\frac{\not{p}}{p^2} \right] \gamma^\nu \left[\frac{(\not{k} + \not{p})}{(k+p)^2} \right] \\ &= \frac{1}{4\sqrt{k^2}} (k^2 \delta^\mu \delta^\nu - k^\mu k^\nu) + \mathcal{O}(k^2). \end{aligned} \quad (4.5)$$

This is relativistically invariant, as expected.

B. Fermion self-energy

We first calculate the one-loop corrections to the fermion self-energy, which will determine the anomalous dimension of the fermion operators as well as the dynamical critical exponent z . Although the anomalous dimension of the fermion is not a gauge-invariant observable, it will be needed to calculate the critical behavior of the gauge-invariant SO(5) order parameter. The three contributions to the fermion self-energy, as shown in Fig. 7, come from the two Higgs bosons and the gauge field, $\Sigma = \Sigma_1 + \Sigma_2 + \Sigma_A$.

$$\Sigma_1(k) = \frac{3}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^x \left[\frac{\not{p} + \not{k}}{(p+k)^2} \right] \gamma^x \frac{1}{\Gamma_1(p) + Kp_x^2} \quad (4.6)$$

$$\Sigma_2(k) = \frac{3}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^y \left[\frac{\not{p} + \not{k}}{(p+k)^2} \right] \gamma^y \frac{1}{\Gamma_2(p) + Kp_y^2} \quad (4.7)$$

$$\Sigma_A(k) = \frac{3}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^\mu \left[\frac{\not{p} + \not{k}}{(p+k)^2} \right] \gamma^\nu \frac{\eta^{\mu\nu} - (1-\xi) \frac{p^\mu p^\nu}{p^2}}{\Gamma_A(p) p^2} \quad (4.8)$$

We have introduced ξ as a gauge-fixing parameter to obtain the gauge boson propagator.

Focusing on the Higgs corrections [Eqs. (4.6) and (4.7)], we analyze the behavior at small external momenta k_i . Note that the self-energy integrals are fully regulated by the presence of K and a nonzero external momenta. Since Σ_1 (Σ_2) is invariant under $k_y \leftrightarrow k_0$ ($k_x \leftrightarrow k_0$), and the two transform into each other under a 90 degree spatial rotation, there are two distinct types of contributions for small external momenta. The first is proportional to $k_x \gamma^x$ for Σ_1 , and $k_y \gamma^y$ for Σ_2 . The second type includes all other possible choices of momenta, such as $k_0 \gamma^0$.

As we shall justify below and in Appendix B, we can focus on the regime $|p_x| \gg |p_0|, |p_y|$ for graphs with a Φ_1 propagator. In this limit, we can approximate the Φ_1 propagator as

$$\frac{4|p_x|}{p_0^2 + p_y^2 + 4K|p_x|^3}. \quad (4.9)$$

At $K = 0$, this propagator has an infrared divergence when integrated over the $p_{0,y}$ plane, so K is needed an infrared regulator. With this, we extract the γ^x correction to the self-energy from the Φ_1 propagator by considering the $k_0 = k_y = 0$ limit:

$$\begin{aligned} \gamma^x \Sigma_1(k_x) &= \frac{12}{N_f} \int^\Lambda \frac{d^3 p}{(2\pi)^3} \frac{(p_x + k_x)}{(p_x + k_x)^2 + p_0^2 + p_y^2} \\ &\times \frac{|p_x|}{p_0^2 + p_y^2 + 4K|p_x|^3}. \end{aligned} \quad (4.10)$$

We have indicated a cutoff Λ to regulate the theory at large momenta, and this is needed in conformal gauge theories in 2+1 dimensions. However, with our inclusion of the irrelevant K to control the infrared singularity, we find that the integrand vanishes faster at large momenta. It is not difficult to see that for $K \neq 0$ Eq. (4.10) is finite as $\Lambda \rightarrow \infty$, and we will take this limit in the present section. The theory with a finite Λ will be examined in Appendix B in a renormalization group computation.

We will now show that Eq. (4.10) has a leading $k_x \ln^2(k_x)$ contribution. One factor of $\ln(k_x)$ is the usual one: It follows from the fact that at $K = 0$ the integrand divided by k_x is a homogeneous function of momenta of dimension -3 . The other comes from the infrared divergence regulated by K noted below Eq. (4.9).

Extracting the coefficient of the $k_x \ln^2(k_x)$ contribution requires a number of approximations. To understand the values of p that dominate the integral in Eq. (4.10), it is useful to

perform the integral over p_0 and p_y :

$$\gamma^x \Sigma_1(k_x) = \frac{12}{N_f} \int \frac{dp_x}{8\pi^2} \frac{|p_x|(p_x+k_x) \ln[(p_x+k_x)^2/(4K|p_x|^3)]}{(p_x+k_x)^2 - 4K|p_x|^3}. \quad (4.11)$$

By examining the form of the integrals in Eqs. (4.10) and (4.11), one can verify that the dominant term at small k_x and K is proportional to $k_x \ln^2(Kk_x)$ and arises from the integration regime

$$[K|p_x|^3]^{1/2} \ll [p_0^2 + p_y^2]^{1/2} \ll |p_x| \ll \frac{1}{K}. \quad (4.12)$$

The scale K appears both as an ultraviolet cutoff and in defining the infrared bound. For future calculations, this integration regime will prove to be the relevant one in isolating similar \log^2 contributions in other diagrams, although in principle one must still carry out an explicit calculation like in Eq. (4.11) to verify that no other integration regimes give comparable contributions. We provide these calculations in Appendix C in addition to numerical evaluations of the one-loop integrals which confirm the validity of our approximations and simply evaluate the one-loop integrals in the Eq. (4.12) limit in the main text.

We can extract the coefficient of this \log^2 term by performing the integral in this regime,

$$\begin{aligned} \gamma^x \Sigma_1(k_x) &\approx \frac{12}{N_f} \int_{-1/K}^{1/K} \frac{dp_x}{2\pi} \frac{|p_x|}{(p_x+k_x)} \\ &\times \int_{(K|p_x|^3)^{1/2}}^{|p_x|} \frac{dp_y dp_0}{4\pi^2} \frac{1}{p_0^2 + p_y^2} \\ &\approx \frac{12}{N_f} \int_{-1/K}^{1/K} \frac{dp_x}{2\pi} \frac{|p_x|}{(p_x+k_x)} \frac{1}{4\pi} \ln(1/(K|p_x|)) \\ &\approx -\frac{12}{N_f} \frac{k_x}{8\pi^2} [\ln(Kk_x)]^2. \end{aligned} \quad (4.13)$$

Another discussion of the origin of the $k_x \ln^2(k_x)$ is presented in Appendix B using a renormalization group analysis.

We now calculate the form of the second type of corrections using the limits in Eq. (4.12), evaluating the Φ_1 contribution to the self-energy with external momentum k_0 for concreteness.

$$\begin{aligned} \gamma^0 \Sigma_1(k_0) &\approx -\frac{12k_0}{N_f} \int_{-1/K}^{1/K} \frac{dp_x}{2\pi} \frac{1}{|p_x|} \int_{(K|p_x|^3)^{1/2}}^{|p_x|} \frac{dp_y dp_0}{(2\pi)^2} \frac{1}{p_0^2 + p_y^2} \\ &\approx -\frac{12k_0}{N_f} \int_{-1/K}^{1/K} \frac{dp_x}{2\pi} \frac{1}{|p_x|} \frac{1}{4\pi} \ln(1/(K|p_x|)) \\ &\approx -\frac{12}{N_f} \frac{k_0}{8\pi^2} [\ln(Kk_0)]^2 \end{aligned} \quad (4.14)$$

Combining the corrections from both Higgs propagators, we obtain the full expression for the self-energy for small external momenta at \log^2 order,

$$\begin{aligned} \Sigma(k) &\approx -\frac{3}{\pi^2 N_f} [k_0 \ln^2(Kk_0) \gamma^0 + k_x \ln^2(Kk_x) \gamma^x \\ &+ k_y \ln^2(Kk_y) \gamma^y]. \end{aligned} \quad (4.15)$$

In principle, the dependence on external momenta inside the logarithms could be more complicated for general k , i.e.,

$\Sigma(k) \gamma^0 \sim k_0 \ln^2(Kf(k_0, k_x, k_y))$, but since we have verified that $f(k_0, 0, 0) = k_0$, corrections to this are subleading.

These divergent corrections are absorbed into the renormalization of the fermion field, $\psi = \sqrt{Z_\psi} \psi_R$, with

$$Z_\psi = 1 - \frac{3}{\pi^2 N_f} \ln^2(K\mu), \quad (4.16)$$

where we have renormalized the theory at some momentum scale μ . This counterterm only cures the divergence at \log^2 order, since the renormalized self-energy at some other momentum scale k will scale as

$$\ln^2(K\mu) - \ln^2(Kk) = \ln(\mu/k) \ln(K^2 k \mu). \quad (4.17)$$

This, along with the RG analysis in Appendix B, indicates that the subleading single-logarithm corrections will generically give nonuniversal behavior. However, these \log^2 corrections to the self-energy are Lorentz invariant and do not affect the renormalization of the dynamical critical exponent z . Therefore, the subleading single-logarithm correction to the velocity anisotropy will lead to a universal correction to the dynamical critical exponent. To extract the subleading correction to z using K and the external momenta as a regulator, we start with the expression

$$\begin{aligned} \frac{\partial \Sigma}{\partial k_0} \gamma^0 - \frac{\partial \Sigma}{\partial k_x} \gamma^x \\ = -\frac{12}{N_f} \int \frac{d^3 p}{(2\pi)^3} \frac{2(p_y + k_y)^2}{(p+k)^4} \frac{|p|}{4K p_y^2 |p| + p_0^2 + p_y^2}. \end{aligned} \quad (4.18)$$

To leading order in k , we set $k = 0$ inside the integrand and simply use it as an IR cutoff, which gives

$$\frac{\partial \Sigma}{\partial k_0} \gamma^0 - \frac{\partial \Sigma}{\partial k_x} \gamma^x \approx \frac{6}{N_f \pi^2} \ln(Kk). \quad (4.19)$$

This result can be obtained analytically by approximating the integration region $k \leq |p_x| \leq 1/K$ and can be verified by a numerical evaluation of Eq. (4.18). This implies a renormalization of the Fermi velocity, $v_F = Z_v v_{F,R}$

$$Z_v = 1 + \frac{6}{\pi^2 N_f} \ln(K\mu). \quad (4.20)$$

The logarithmic derivative with respect to $1/K$ determines the renormalization of the dynamical critical exponent,

$$z = 1 + \frac{6}{\pi^2 N_f}. \quad (4.21)$$

The one-loop calculation defined in Eq. (4.18) is actually well defined when $K = 0$ and can be regulated via more standard approaches, such as dimensional regularization, as shown in Appendix D. The same value of z is also obtained in a renormalization group computation in Appendix B.

C. SO(5) order parameter

In the absence of the Higgs fields, our theory possesses an emergent SO(5) symmetry corresponding to rotations between Néel and VBS order parameters. This SO(5) symmetry is broken by the critical Higgs fields, and as a result, the scaling behavior of Néel and VBS order parameters will differ. In




FIG. 8. The $\mathcal{O}(N_f^{-1})$ vertex corrections which contribute to the renormalization of the SO(5) order parameter. The order parameter receives corrections at one-loop order from the Higgs fields (left) and the gauge boson (center), although only the former gives a \log^2 correction. An additional two-loop $\mathcal{O}(N_f^{-1})$ contribution (right) is possible; we show in Appendix F that it does not contain any \log^2 divergences.

terms of Dirac fermions, the fermion bilinears corresponding to the two-component VBS order parameter—determined by the action of the square lattice symmetries on the bilinears—may be written as

$$V^i = \bar{\psi} \Gamma^i \psi, \quad \Gamma^i = \{\mu^x, \mu^z\}. \quad (4.22)$$

$$\frac{\mu^z \sigma^a \mu^i \sigma^a \mu^z}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^x \frac{\not{p} - \not{k}_1}{(p - k_1)^2} \frac{\not{p} - \not{k}_2}{(p - k_2)^2} \gamma^x \frac{1}{\Gamma_1(p) + K p_x^2} + \frac{\mu^x \sigma^a \mu^i \sigma^a \mu^x}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^y \frac{\not{p} - \not{k}_1}{(p - k_1)^2} \frac{\not{p} - \not{k}_2}{(p - k_2)^2} \gamma^y \frac{1}{\Gamma_2(p) + K p_y^2}, \quad (4.24)$$

where the first and second terms arise from interactions with Φ_1^a and Φ_2^a , respectively. The gauge field correction does not break SO(5) symmetry and does not contribute to the renormalization at \log^2 order, so we will focus on the Higgs corrections. Additionally, there is a possible two-loop diagram shown in Fig. 8 that contributes at $\mathcal{O}(N_f^{-1})$, but we show explicitly in Appendix F that these corrections also do not contribute to the renormalization at \log^2 order. At zero external momenta, the \log^2 Higgs corrections to the VBS order parameter ($\mu^i = \mu^x, \mu^z$) drops out entirely, leaving only Higgs corrections coming from the fermion renormalization.

We focus on vertex corrections to the Néel order parameter ($\mu^i = \mu^y$). As is the case in the fermion self-energy, the spatial anisotropy in the Higgs propagators gives rise to \log^2 divergences in their corrections to the SO(5) vertex. We isolate \log^2 divergences in the Higgs correction to the SO(5) order parameter by including an external momenta $2k_x$ to the order parameter, which is distributed symmetrically between the two fermion fields. We calculate this for the Φ_1 propagator—approximating the Higgs propagator as $4|p_x|/(p_0^2 + p_y^2 + 4K|p_x|^3)$ as in the previous section and taking the limit in Eq. (4.12), the one-loop correction is

$$\begin{aligned} & \frac{\mu^z \sigma^a \mu^y \sigma^a \mu^z}{N_f} \int \frac{d^3 p}{(2\pi)^3} \frac{p_x^2 - k_x^2 + p_y^2 + p_0^2}{[(p_x + k_x)^2 + p_y^2 + p_0^2][(p_x - k_x)^2 + p_y^2 + p_0^2]} \frac{4|p_x|}{p_0^2 + p_y^2 + 4K|p_x|^3} \\ & \approx -\frac{12\mu^y}{N_f} \int_{-1/K}^{1/K} \frac{dp_x}{2\pi} \frac{(p_x^2 - k_x^2)|p_x|}{(p_x + k_x)^2(p_x - k_x)^2} \int_{(K|p_x|^3)^{1/2}}^{|p_x|} \frac{dp_y dp_0}{4\pi^2} \frac{1}{p_0^2 + p_y^2} \\ & \approx -\frac{12\mu^y}{N_f} \int_{1/K}^{1/K} \frac{dp_x}{8\pi^2} \frac{(p_x^2 - k_x^2)|p_x|}{(p_x + k_x)^2(p_x - k_x)^2} \ln(1/(K|p_x|)) \approx -\frac{3\mu^y}{2N_f \pi^2} \ln^2(Kk_x). \end{aligned} \quad (4.25)$$

The Φ_2 propagator gives an identical correction. Since the external momenta only play the role of an IR cutoff to leading order, we generalize this result to an arbitrary external momentum and obtain the composite operator renormalizations [57]

$$Z_{\text{VBS}} = 1 \quad Z_{\text{Néel}} = 1 + \frac{3}{N_f \pi^2} \ln^2(K\mu). \quad (4.26)$$

We can state these results in terms of the perturbative corrections to the two-point correlator of the order parameters, $\langle \bar{\psi} \Gamma^i \psi(k) \bar{\psi} \Gamma^j \psi(-k) \rangle$, i.e., the corresponding susceptibili-

The three-component Néel order parameter has a less concise expression in terms of Dirac fermions; this is due to the fact that the Dirac fermion representation obfuscates the action of the physical SU(2) spin rotation symmetry. In terms of the Majorana field X , the order parameter is $\text{Tr}(\bar{X} \mu^y \sigma^a X)$, $a = x, y, z$. In order to calculate corrections to the Néel order parameter, we focus on the σ^z component, which happens to be simply expressible in terms of a Dirac fermion bilinear:

$$N^z = \bar{\psi} \mu^y \psi. \quad (4.23)$$

Because the Higgs couplings preserve the physical SU(2) spin rotation symmetry, the other components must have the same corrections, and this has been confirmed by an explicit calculation in terms of the Majorana fermions. To compute the corrections to the scaling dimensions of these composite operators, we couple the fermion bilinear $n^i = \bar{\psi} \mu^i \psi$ to a source field J_i and compute the $\mathcal{O}(N_f^{-1})$ vertex corrections in Fig. 8. Aside from the corrections coming from the renormalization of the fermion self-energy, the $\mathcal{O}(N_f^{-1})$ corrections that we will be interested in come from one-loop corrections of the Higgs fields with external momenta $k_{1,2}$

ties χ_{VBS} and $\chi_{\text{Néel}}$; these combine the consequences of the composite operator renormalizations in Eq. (4.26), and Z_ψ in Eq. (4.16), to yield

$$\begin{aligned} \chi_{\text{VBS}}(k) & \sim -|k| \left(\frac{Z_\psi}{Z_{\text{VBS}}} \right)^2 = -|k| \left[1 - \frac{6}{N_f \pi^2} \ln^2(K|k|) \right] \\ \chi_{\text{Néel}}(k) & \sim -|k| \left(\frac{Z_\psi}{Z_{\text{Néel}}} \right)^2 = -|k| \left[1 - \frac{12}{N_f \pi^2} \ln^2(K|k|) \right]. \end{aligned} \quad (4.27)$$

After a Fourier transform to real space, these correlators are

$$\begin{aligned}\chi_{\text{VBS}}(r) &\sim \frac{1}{|r|^4} \left[1 - \frac{6}{N_f \pi^2} \ln^2(|r|/K) \right] \\ \chi_{\text{Néel}}(r) &\sim \frac{1}{|r|^4} \left[1 - \frac{12}{N_f \pi^2} \ln^2(|r|/K) \right].\end{aligned}\quad (4.28)$$

The renormalization group analysis in Appendix B shows how the above results may be renormalized to large r ; we find

$$\begin{aligned}\chi_{\text{VBS}}(r) &\sim \frac{1}{|r|^a} \exp\left(-\frac{6}{N_f \pi^2} \ln^2(|r|/K)\right) \\ \chi_{\text{Néel}}(r) &\sim \frac{1}{|r|^b} \exp\left(-\frac{12}{N_f \pi^2} \ln^2(|r|/K)\right),\end{aligned}\quad (4.29)$$

where the exponents of the prefactors, a and b , are *nonuniversal* numbers.

Leading logarithm-squared corrections have appeared earlier in a few other problems in quantum many-body theory. They appear in the theory of weakly disordered two-dimensional metals with Coulomb interactions [58–60]. More recently, \log^2 terms have also been found in computations of the density of states of clean bilayer graphene with Coulomb interactions [61,62]. Renormalization group analyses of these cases [59,62] also yield an exponentiation similar to that in Eq. (4.29).

As an aside, we note that the one-loop vertex corrections to the bilinear $\bar{\psi}\psi$, whose symmetry properties identify it as the scalar spin chirality [63], have the same magnitude and opposite sign as the Néel order parameter. Because of this, the \log^2 divergence is in fact canceled by the fermion self-energy. As shown in Appendix F, the two-loop corrections coming from the Higgs fields vanish, meaning that correlations of the scalar spin chirality should have power law decay at $\mathcal{O}(N_f^{-1})$. Since this power law decay is slower than the Néel and VBS correlations, this may indicate proximity to a chiral spin liquid.

V. TRANSITION FROM U(1) STAGGERED FLUX TO GAPLESS \mathbb{Z}_2 SPIN LIQUID

This section discusses the critical U(1) gauge theory for the transition between the U(1) staggered flux spin liquid and the gapless \mathbb{Z}_2 spin liquid Z2Az13 in Fig. 2. A similar theory has been considered earlier [25] for the Néel- \mathbb{Z}_2 spin liquid transition.

Both phases have the Higgs field $\langle \Phi_3^a \rangle \neq 0$. So let us fix $\Phi_3^a = \delta_{az} \Phi$, with Φ a nonzero constant, which will turn into a coupling constant in the low energy theory below. In this situation, the SU(2) gauge symmetry is broken down to U(1), and we need only consider a U(1) gauge theory with the U(1) gauge field $A_\mu \equiv A_\mu^z$. Also important is the consequence of the w term in the Higgs potential Eq. (3.26):

$$V(\Phi) = \dots + w \Phi (\Phi_1^x \Phi_2^y - \Phi_1^y \Phi_2^x) + \dots \quad (5.1)$$

Choosing a gauge with $w \Phi < 0$, and diagonalizing the quadratic form of the Higgs potential for $\Phi_{1,2}^{x,y}$, we deduce that we need only focus on a single low energy complex Higgs field near the critical point

$$\mathcal{H} = \frac{1}{2}(\Phi_1^x + \Phi_2^y + i(\Phi_1^y - \Phi_2^x)). \quad (5.2)$$

It can now be checked that \mathcal{H} transforms as a charge 2 Higgs field under the unbroken U(1) gauge symmetry. Other linear combinations of $\Phi_{1,2}^{x,y}$ can be ignored for the critical theory.

We can now obtain the critical theory for the fermions ψ , the complex Higgs field \mathcal{H} , and the U(1) gauge field A_μ from Eq. (3.24):

$$\begin{aligned}\mathcal{L}_{\text{sf}} &= \mathcal{L}_\psi + \mathcal{L}_\mathcal{H} + \mathcal{L}_{\mathcal{H}\psi} \\ \mathcal{L}_\psi &= i \sum_v \bar{\psi}_v \gamma^\mu D_\mu \psi_v + \Phi \bar{\psi} \mu^y \sigma^z (\gamma^y D_x + \gamma^x D_y) \psi. \\ \frac{\mathcal{L}_\mathcal{H}}{N_f} &= s |\mathcal{H}|^2 + u |\mathcal{H}|^4 \\ \mathcal{L}_{\mathcal{H}\psi} &= \lambda (\mathcal{H} \bar{\psi} (\mu^z \gamma^x + i \mu^x \gamma^y) \sigma^- \psi \\ &\quad + \mathcal{H}^* \bar{\psi} (\mu^z \gamma^x - i \mu^x \gamma^y) \sigma^+ \psi).\end{aligned}\quad (5.3)$$

We define the covariant derivative $D_\mu = \partial_\mu - i A_\mu \sigma^z$ and operators $\sigma^\pm = (\sigma^x \pm i \sigma^y)/2$. Note that Φ is a marginal coupling constant here, not a fluctuating field. A crucial feature of \mathcal{L}_{sf} is that it does not contain the K gradient terms: These terms are now truly irrelevant. This can be seen in the large N_f expansion: Upon integrating the fermions, we obtain, in place of Eq. (4.3),

$$\begin{aligned}\frac{S_b}{N_f} &= \int_k \left(s + \frac{\Gamma_1(k) + \Gamma_2(k)}{2} \right) |\mathcal{H}(k)|^2 \\ &\quad + \frac{\Gamma_A(k)}{2} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) A_\mu(k) A_\nu(-k),\end{aligned}\quad (5.4)$$

where $\Gamma_{1,2}(k)$ are specified in Eqs. (4.3) and (4.4) for $\Phi = 0$. In general, the sum $\Gamma_1(k) + \Gamma_2(k)$ has the rotational symmetry of the square lattice, and its inverse does not contain the infrared singularities we encountered earlier. Consequently, there is no logarithmic violation of scaling by a dangerously irrelevant K here, and the $1/N_f$ expansion of \mathcal{L}_{sf} should proceed along more conventional lines.

The $1/N_f$ expansion of the theory \mathcal{L}_ψ was presented in Refs. [63,64]: They found a stable Lorentz invariant fixed point with $\Phi = 0$ at the fixed point. In our case, for \mathcal{L}_{sf} we expect a critical theory with dynamic scaling with an exponent $z \neq 1$, SO(5) symmetry broken by $\mathcal{L}_{\mathcal{H}\psi}$, and a spatial anisotropy in the fermion velocities at the Dirac nodes determined by the fixed point value of Φ . Note that even for $\Phi = 0$ we do not expect Lorentz invariance with $z = 1$, because the relevant Yukawa couplings in $\mathcal{L}_{\mathcal{H}\psi}$ are not Lorentz invariant, and consequently $\Gamma_1(k) + \Gamma_2(k)$ is not Lorentz invariant.

VI. CONCLUSIONS

Building upon the results of recent numerical studies [7–10], we have proposed resolutions of long-standing controversies connected to theories of the cuprates: the phases of the frustrated square lattice spin $S = 1/2$ antiferromagnets and the nature of deconfined criticality in such models. Deconfined criticality expresses the low energy physics in terms of fractionalized degrees of freedom and emergent gauge fields, which can enter various confining states with possible broken symmetries on either or both sides of the critical point. Although there are several well-established examples, the transition between Néel and VBS states in square lattice

antiferromagnets [5,6,12,13] has been of particular interest. One formulation of this deconfined critical point is a version of QCD₃, quantum chromodynamics in 2+1 dimensions: a SU(2) gauge theory with two flavors of two-component massless Dirac fermions, each carrying a fundamental color charge. This theory is dual to a SO(5) nonlinear sigma model with a Wess-Zumino-Witten term [27,54,55]. There is now significant numerical evidence that such a conformal field theory (CFT) does not exist, although there is likely a nearby ‘complex’ CFT [31–37]. This leaves open the fate of a physical model with a Hermitian Hamiltonian, such as the J_1 - J_2 antiferromagnet on the square lattice, between the Néel and VBS states. Here we have presented a theory in which the putative QCD₃ CFT is resolved into an intermediate stable gapless phase with \mathbb{Z}_2 topological order and gapless Dirac fermions [23,24,26]. The intermediate \mathbb{Z}_2 spin liquid is flanked by two proposed deconfined critical points, neither of which is a CFT, or even invariant under Lorentz transformations. The absence of Lorentz symmetry permits several novel phenomena, including the appearance of dangerously irrelevant couplings and logarithm-squared renormalizations, which can be tested in numerical studies. All of these phases and critical points are described by extending QCD₃ with three real adjoint Higgs fields. The couplings of these Higgs fields are tightly constrained by the transformations of QCD₃ under the symmetries of the underlying square lattice antiferromagnet, and an analysis of these symmetries occupy a significant portion of this paper.

Our main results can be summarized in the context of the mean-field phase diagram in Fig. 2 obtained from the SU(2) gauge theory with three adjoint Higgs field $\Phi_{1,2,3}^a$ in Eq. (3.24). This mean field theory yields three spin liquids, with deconfined SU(2), U(1), and \mathbb{Z}_2 gauge fields. We assume that the spin liquids with continuous gauge symmetries confine, except at possible deconfined critical transitions to the \mathbb{Z}_2 spin liquid. This phase diagram maps onto the J_1 - J_2 model along the trajectory of the dotted blue line, and our proposed deconfined critical theories are at the boundaries between the mean field SU(2) and \mathbb{Z}_2 spin liquids, and the U(1) and \mathbb{Z}_2 spin liquids.

The numerical evidence for the confinement of the SU(2) π -flux spin liquid was reviewed in Sec. I. This confining state should have either Néel or VBS order [27], and Ref. [28] argued by comparing to bosonic spinon theories that it should be the Néel state. The structure of the critical theory from such a confining state to the gapless \mathbb{Z}_2 spin liquid was presented in Sec. IV, and we found some unusual \log^2 corrections to both the Néel and VBS critical correlators. From the geometry of the mean field phase diagrams in Fig. 2, and the numerical studies on the square lattice antiferromagnet noted in Fig. 1, it is then natural to propose that the U(1) staggered flux spin liquid confines to the VBS state. The critical U(1) gauge theory for the boundary between the U(1) and \mathbb{Z}_2 spin liquid was presented in Sec. V, and this has no \log^2 terms. We also note that the \log^2 correlators in Eqs. (4.28) and (4.29) show a faster decay of the Néel order than the VBS order, which might be evidence that the SU(2) critical theory is proximate to the VBS state rather than the Néel state, which would reverse the direction of the arrow in Fig. 2.

Irrespective of the assignment of the Néel or VBS confining states to the SU(2) or U(1) spin liquids in Fig. 2, we expect any direct phase boundary between the Néel and VBS states to be a first order transition. This follows from the numerical studies [31,32,34] noted in Sec. I.

Our critical SU(2) gauge theory for the $S = 1/2$ square lattice antiferromagnet has massless two-component Dirac fermions with two flavors and two colors, and real critical Higgs fields with two flavors and three colors, and is shown in Eq. (4.2). This derives from a theory for the π -flux to gapless \mathbb{Z}_2 spin liquid transition proposed by Ran and Wen [39,40] and includes an additional ‘dangerously irrelevant’ coupling K , which is the coefficient of a spatial gradient term in the Higgs fields. We analyzed this theory along the lines of the $1/N_f$ expansion of Ref. [56] (the case of interest to us here is $N_f = 1$). We found that the theory with $K = 0$ has infrared divergencies that arise from the highly anisotropic spatial structure of the Higgs correlations, which is in turn a consequence of the non-Lorentz invariant Yukawa couplings between the Higgs fields and the fermions. So even though the coupling K is formally irrelevant, it must be included to understand the long-distance and long-time behavior of the theory, i.e., the coupling K is dangerously irrelevant. We found that the coupling K leads to leading logarithm-squared corrections to various correlators, such as those in Eqs. (4.27) and (4.28) for the correlations of the Néel and VBS order parameters; Appendix B showed how these corrections are exponentiated in a renormalization group analysis, leading to Eq. (4.29). We also note that the logarithm-squared term was absent in the contributions to the dynamic critical exponent z , and we computed a non-Lorentz-invariant value for z in Eq. (4.21).

The critical U(1) gauge theory for the $S = 1/2$ square lattice antiferromagnet was discussed in Sec. V. It has massless two-component Dirac fermions with four flavors and ± 1 U(1) gauge charges, and a single complex critical Higgs fields with ± 2 U(1) gauge charge. We found that K was not dangerously irrelevant in this theory. The critical theory is not Lorentz invariant and so has dynamic critical exponent $z \neq 1$. The critical theory also does not have the SO(5) symmetry between the Néel and VBS order parameters. A full analysis of this theory requires a study of the role of anisotropies in the Dirac fermion velocities [associated with the coupling Φ in Eq. (5.3)], and we leave this for future work.

It would be useful to examine numerical studies of the square lattice antiferromagnet for logarithmic violations of scaling, Lorentz invariance, and SO(5) symmetry and compare to our predictions. In particular, we note the violations of scaling observed in Ref. [65], although for a different square lattice antiferromagnet. Finally, we note that gapless \mathbb{Z}_2 spin liquid studied is an attractive candidate for the ancilla model of doped antiferromagnets [66–68], as it can realize a stable state in the second ancilla layer for the pseudogap state.

Recently, we became aware of some related work:

(i) Superconductivity has been observed [69,70] in the doped J_1 - J_2 model; doping the gapless \mathbb{Z}_2 spin liquid is a known to be a natural route to d -wave superconductivity [18,25].

(ii) Yang *et al.* [71] have detected a gapless spin liquid phase next to the Néel phase on the Shastry-Sutherland model,

which is obtained from the J_1 - J_2 model by removing 3/4 of the J_2 bonds.

ACKNOWLEDGMENTS

We thank Federico Becca and Anders Sandvik for enlightening discussions which stimulated this work. We are also grateful to Masatoshi Imada, Steve Kivelson, Patrick Ledwith, Ying Ran, T. Senthil, Ashvin Vishwanath, Chong Wang, and Cenke Xu for valuable discussions. This research was supported by the National Science Foundation under Grant No. DMR-2002850. This work was also supported by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651440, S.S.).

APPENDIX A: PROJECTIVE SYMMETRY ANALYSIS

This Appendix will present a detailed analysis of the projective symmetry group (PSG) of the Z2Azz13 spin liquid and its neighboring phases. Here, we will employ the gauge used by Wen [24]. Wen described the Z2Azz13 spin liquid by the Bogoliubov Hamiltonian in Eq. (2.3) with the ansatz

$$\begin{aligned} u_{i,i+\hat{x}} &= \chi \tau^x - \eta \tau^y \\ u_{i,i+\hat{y}} &= \chi \tau^x + \eta \tau^y \\ u_{i,i+\hat{x}+\hat{y}} &= -\gamma_1 \tau^x \\ u_{i,i-\hat{x}+\hat{y}} &= \gamma_1 \tau^x. \end{aligned} \quad (\text{A1})$$

In terms of the spinons $f_{i\alpha}$, this can be written as

$$\begin{aligned} H = - \sum_{\mathbf{k}} [2\chi(\cos(k_x) + \cos(k_y)) - i2\eta(\cos(k_x) - \cos(k_y)) \\ + 4\gamma_1 \sin(k_x) \sin(k_y)] f_{-\mathbf{k}\downarrow} f_{\mathbf{k}\uparrow} + \text{H.c.} \end{aligned} \quad (\text{A2})$$

So in this gauge, the Z2Azz13 spin liquid has both $d_{x^2-y^2} + is$ and d_{xy} pairing and no hopping, and the fermion dispersion relation is

$$\begin{aligned} \varepsilon_{\mathbf{k}}^2 &= [2\chi(\cos(k_x) + \cos(k_y)) + 4\gamma_1 \sin(k_x) \sin(k_y)]^2 \\ &+ [2\eta(\cos(k_x) - \cos(k_y))]^2. \end{aligned} \quad (\text{A3})$$

In the ansatz in Eq. (A1), the three spin liquids are

(i) The π -flux phase with SU(2) gauge symmetry corresponds to $\chi = \eta \neq 0$, $\gamma_1 = 0$.

(ii) The ‘staggered flux’ U(1) spin liquid is obtained for $\chi \neq 0$, $\gamma_1 = 0$, $\eta \neq 0$ with $\chi \neq \eta$.

(iii) The Z2Azz13 spin liquid is obtained when the d_{xy} pairing γ_1 breaks the U(1) down to \mathbb{Z}_2 .

For our purposes, and in general, a complex Higgs field is needed to break U(1) down to \mathbb{Z}_2 . We have characterized the d_{xy} pairing above by a real parameter γ_1 , and we need to generalize this to a complex parameter. From the analysis in Sec. II, we deduce that this is obtained by taking a complex d_{xy} order parameter which has opposite phases on the two sublattices, i.e.,

$$\begin{aligned} u_{i,i+\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & -(\gamma_1 - i\gamma_2) \\ -(\gamma_1 + i\gamma_2) & 0 \end{pmatrix}, \quad \mathbf{i}_x + \mathbf{i}_y \text{ even} \\ u_{i,i+\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & -(\gamma_1 + i\gamma_2) \\ -(\gamma_1 - i\gamma_2) & 0 \end{pmatrix}, \quad \mathbf{i}_x + \mathbf{i}_y \text{ odd} \end{aligned}$$

$$\begin{aligned} u_{i,i-\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & (\gamma_1 - i\gamma_2) \\ (\gamma_1 + i\gamma_2) & 0 \end{pmatrix}, \quad \mathbf{i}_x + \mathbf{i}_y \text{ even} \\ u_{i,i-\hat{x}+\hat{y}} &= \begin{pmatrix} 0 & (\gamma_1 + i\gamma_2) \\ (\gamma_1 - i\gamma_2) & 0 \end{pmatrix}, \quad \mathbf{i}_x + \mathbf{i}_y \text{ odd.} \end{aligned} \quad (\text{A4})$$

1. Lattice PSGs

We first recall the spin liquid classification scheme of Ref. [24]. If u_{ij} is the mean field ansatz for a spin liquid symmetric under the group action G , it transforms as

$$PG: u_{ij} \rightarrow W_g^\dagger(\mathbf{i}) u_{G(\mathbf{i}, G(\mathbf{j}))} W(\mathbf{j}), \quad (\text{A5})$$

where $W_G(\mathbf{i})$ is a gauge transform. In addition to the symmetries, these gauge transformations characterize the spin liquid, yielding the projective symmetry group (PSG) [24].

Using the notation from Ref. [24], the spin liquid Z2Azz13 is defined by the PSG

$$\begin{aligned} W_{tx}(\mathbf{i}) &= \tau^0, \quad W_{px}(\mathbf{i}) = (-)^{i_x+i_y} i\tau^z, \quad W_{pxy}(\mathbf{i}) = i\tau^x, \\ W_{ty}(\mathbf{i}) &= \tau^0, \quad W_{py}(\mathbf{i}) = (-)^{i_x+i_y} i\tau^z, \quad W_t(\mathbf{i}) = i\tau^z \end{aligned} \quad (\text{A6})$$

while the PSG of U1Cn01n (the staggered flux phase) is

$$\begin{aligned} W_{tx}(\mathbf{i}) &= g_3(\theta_x) i\tau^x, \quad W_{px}(\mathbf{i}) = (-)^{i_x} g_3(\theta_x) i\tau^x, \\ W_{pxy}(\mathbf{i}) &= g_3(\theta_{pxy}) i\tau^x, \\ W_{ty}(\mathbf{i}) &= g_3(\theta_y) i\tau^x, \quad W_{py}(\mathbf{i}) = (-)^{i_y} g_3(\theta_y), \\ W_t(\mathbf{i}) &= (-)^{i_x+i_y} g_3(\theta_t), \end{aligned} \quad (\text{A7})$$

where $g_3(\theta) = e^{i\theta\tau^z}$. From these PSGs we can extract the symmetry fractionalization through the group relations given in the Appendix of Ref. [28] [Eq. (B8)]. These are provided in Table I. Note that instead of $P_{xy} : (i_x, i_y) \rightarrow (i_y, i_x)$, we consider the 90° rotation $R_{\pi/2} = P_{xy} P_y$. Similarly, P_x is related to the other symmetries through $R_{\pi/2} P_y R_{\pi/2}^{-1}$.

2. Identification of staggered flux in continuum model

The staggered flux state (U1Cn01n) can be obtained by coupling a Higgs field to the bilinear

$$\mathcal{O}_3^a = \text{tr}(\sigma^a \bar{X} \mu^y (\gamma^x i\partial_y + \gamma^y i\partial_x) X) \quad (\text{A8})$$

giving something like

$$\mathcal{L} = \text{tr}(\bar{X} \gamma^\mu i\partial_\mu X) + \Phi_3^a \text{tr}(\sigma^a \bar{X} M X). \quad (\text{A9})$$

The U(1) spin liquid U1Cn01n is then obtained upon condensing one component of Φ_3 . This was determined by considering the symmetry fractionalization of the U(1) spin liquid obtained by condensing the z component of Φ_3 : $\langle \Phi_3^z \rangle \neq 0$. Based on the symmetry transformations outlined in Table II, this condensate has a corresponding continuum PSG

$$\begin{aligned} V_{tx} &= g_3(\phi_x) i\sigma^x, \quad V_{px} = g_3(\phi_{px}), \quad V_r = g_3(\phi_r) i\sigma^x, \\ V_{ty} &= g_3(\phi_y) i\sigma^x, \quad V_{py} = g_3(\phi_{py}), \quad V_t = g_3(\phi_t), \end{aligned} \quad (\text{A10})$$

where $g_3(\phi) = e^{i\phi\sigma^z}$ is an arbitrary gauge transformation. Importantly, in the U(1) spin liquid, the phases ϕ can take any value. When these phases are rewritten in terms of the U(1)

TABLE I. Symmetry fractionalization. In keeping with the conventions of Ref. [28], the gauge is chosen such that group relation 7 is fixed to equal -1 for the \mathbb{Z}_2 spin liquid.

	Group relations	Z2Azz13	U1Cn01n lattice	U1Cn01n cont
1	$T_y^{-1}T_xT_yT_x^{-1}$	1	$e^{-2i(\theta_x - \theta_y)\tau^z}$	$-e^{-2i(\phi_x - \phi_y)\sigma^z}$
2	$P_y^{-1}T_xP_yT_x^{-1}$	-1	$e^{-2i\theta_{py}\tau^z}$	$e^{2i\phi_{py}\sigma^z}$
3	$P_y^{-1}T_yP_yT_y$	-1	$e^{-2i\theta_{py}\tau^z}$	$e^{2i\phi_{py}\sigma^z}$
4	P_y^2	-1	$e^{2i\theta_{py}\tau^z}$	$e^{-2i\phi_{py}\sigma^z}$
5	$P_y^{-1}R_{\pi/2}P_yR_{\pi/2}$	1	$-e^{-2i\theta_{py}\tau^z}$	$-e^{2i\phi_{py}\sigma^z}$
6	$R_{\pi/2}^4$	1	$\mathbb{1}$	$\mathbb{1}$
7	$R_{\pi/2}^{-1}T_xR_{\pi/2}T_y$	-1	$e^{i(2[\theta_{pxy} + \theta_{py}] - [\theta_x + \theta_y])\tau^z}$	$e^{i(2\phi_r - \phi_x - \phi_y)\sigma^z}$
8	$R_{\pi/2}^{-1}T_xR_{\pi/2}T_x^{-1}$	-1	$e^{i(2[\theta_{pxy} + \theta_{py}] - [\theta_x + \theta_y])\tau^z}$	$e^{i(2\phi_r - \phi_x - \phi_y)\sigma^z}$
9	$\mathcal{T}^{-1}R_{\pi/2}^{-1}\mathcal{T}R_{\pi/2}$	-1	$e^{-2i\theta_t\tau^z}$	$e^{-2i\phi_t\sigma^z}$
10	$\mathcal{T}^{-1}P_y^{-1}\mathcal{T}P_y$	1	$\mathbb{1}$	$\mathbb{1}$
11	$\mathcal{T}^{-1}T_x^{-1}\mathcal{T}T_x$	1	$-e^{-2i\theta_t\tau^z}$	$-e^{-2i\phi_t\sigma^z}$
12	$\mathcal{T}^{-1}T_y^{-1}\mathcal{T}T_y$	1	$-e^{-2i\theta_t\tau^z}$	$-e^{-2i\phi_t\sigma^z}$
13	\mathcal{T}^2	-1	$e^{2i\theta_t\sigma^z}$	$e^{2i\phi_t\sigma^z}$

phases from Eq. (A7), θ_G , according to

$$(\phi_x, \phi_y, \phi_{py}, \phi_t, \phi_r) = \left(\theta_x + \frac{\pi}{4}, \theta_y - \frac{\pi}{4}, -\theta_y, \theta_t, \theta_{pxy} + \theta_{py} \right) \quad (\text{A11})$$

the symmetry fractionalizations given in columns 4 and 5 of Table I are identical. It is possible that two distinct spin liquids (as defined by having distinct PSGs) could nevertheless have identical symmetry fractionalization. This seems unlikely in this situation and is, moreover, proven false by the explicit derivation of the continuum action from the lattice model.

3. Identification of Z2Azz13 in continuum model

The spin liquid Z2Azz13 is proximate to U1Cn01n in that the PSG of Eq. (A6) may be obtained through gauge transformations and judicious choices of the angles θ_G in Eq. (A7). It is, however, simpler to determine the U(1) transformations (i.e., the angles ϕ_μ) that map the symmetry fractionalization of U1Cn01n to the symmetry fractionalization of Z2Azz13. That is, we find that the assignment

$$\begin{aligned} &(\phi_x, \phi_y, \phi_{py}, \phi_t, \phi_r) \\ &= \left(\theta + \frac{\pi}{4}, \theta - \frac{\pi}{4} + n_y\pi, (2n_{py} + 1)\frac{\pi}{2}, (2n_t + 1)\frac{\pi}{2}, \theta \right. \\ &\quad \left. + (2n_r + n_y + 1)\frac{\pi}{2} \right), \quad n_\mu \in \mathbb{Z} \end{aligned} \quad (\text{A12})$$

transforms the fifth column of Table I into a set of ± 1 s that match the third column. Inserting these ϕ_μ s into the PSG

defined in Eq. (A10) and selecting $n_\mu = 0$, $\mu = y, py, t, r$ and $\theta = \pi/4$, we obtain the \mathbb{Z}_2 continuum PSG [72]

$$\begin{aligned} V_{tx} &= -i\sigma^y, & V_{px} &= \pm i\sigma^z \\ V_r &= -\frac{i}{\sqrt{2}}(\sigma^x - \sigma^y) \\ V_{ty} &= -i\sigma^x, & V_{py} &= -i\sigma^z, & V_t &= i\sigma^z. \end{aligned} \quad (\text{A13})$$

We can now ask what form of operator needs to couple to a new Higgs field in order to realize this PSG and hence the \mathbb{Z}_2 spin liquid Z2Azz13. Firstly, it's clear that the σ^x or σ^y components of the Higgs field must condense—condensing in the σ^z channel, $\sim \langle \Phi^z \rangle \text{tr}(\sigma^z \bar{X} \bar{M} X)$, would not break the U(1) symmetry. However, in considering condensates in x or y , we see that the gauge transformations corresponding to the translations T_x and T_y are different and, further, the rotation $R_{\pi/2}$ exchanges σ^x and σ^y , meaning that both must be present in a symmetric spin liquid.

Based on the symmetry relations documented in Ref. [28], we find that the operators $\mathcal{O}_{1,2}$,

$$\mathcal{O}_1^a = \text{tr}(\sigma^a \bar{X} \mu^z \gamma^x X), \quad \mathcal{O}_2^a = \text{tr}(\sigma^a \bar{X} \mu^x \gamma^y X), \quad (\text{A14})$$

induce the PSG of Eq. (A13) provided they couple to Higgs fields that condense in perpendicular directions. The symmetry transformation properties of $\mathcal{O}_{1,2}^a$ are given in Table II. That is, given a Lagrangian:

$$\begin{aligned} \mathcal{L}' &= \text{tr}(\bar{X} \gamma^\mu i \partial_\mu X) + \Phi_1^a \text{tr}(\sigma^a \bar{X} \mu^z \gamma^x X) + \Phi_2^a \text{tr}(\sigma^a \bar{X} \mu^x \gamma^y X) \\ &\quad + \Phi_3^a \text{tr}(\sigma^a \bar{X} \mu^y (\gamma^x i \partial_y + \gamma^y i \partial_x) X), \end{aligned} \quad (\text{A15})$$

TABLE II. Symmetry transformations of the three fermion bilinears which must be coupled to SU(2) adjoint Higgs fields in order to realize the PSG of the \mathbb{Z}_2 spin liquid Z2Azz13. The symmetry transformations of these bilinears were previously worked out in Ref. [28].

	Operators	\mathcal{T}	P_x	P_y	T_x	T_y	$R_{\pi/2}$
\mathcal{O}_1^a	$\text{tr}(\sigma^a \bar{X} \mu^z \gamma^x X)$	-	-	-	-	+	$-\mathcal{O}_2^a$
\mathcal{O}_2^a	$\text{tr}(\sigma^a \bar{X} \mu^x \gamma^y X)$	-	-	-	+	-	$-\mathcal{O}_1^a$
\mathcal{O}_3^a	$\text{tr}(\sigma^a \bar{X} \mu^y (\gamma^x i \partial_y + \gamma^y i \partial_x) X)$	+	+	+	-	-	-

the PSG in Eq. (A13) is obtained when $\langle \Phi_1 \rangle = (\alpha, 0, 0)$, $\langle \Phi_2 \rangle = (0, \alpha, 0)$, and $\langle \Phi_3 \rangle = (0, 0, \beta)$, for $\alpha, \beta \in \mathbb{R}$. This agrees with conclusions reached in Sec. III D.

APPENDIX B: RENORMALIZATION GROUP ANALYSIS OF THE SU(2) GAUGE THEORY

In this Appendix, we describe the origin of the \log^2 terms in the critical SU(2) gauge theory in a renormalization group (RG) framework. Integrating the RG equations will lead to an exponentiated prediction for the correlators.

We start with the expression in Eq. (4.10), keep the full Higgs propagator as in Eq. (4.6), and perform a standard momentum shell RG in the window $\Lambda - d\Lambda < (p_0^2 + p_x^2 + p_y^2)^{1/2} < \Lambda$

$$\delta\gamma^x \Sigma_1(k_x) = \frac{12}{N_f} \int_{\Lambda-d\Lambda}^{\Lambda} \frac{d^3 p}{(2\pi)^3} \frac{(p_x + k_x)}{(p_x + k_x)^2 + p_0^2 + p_y^2} \times \frac{|p|}{p_0^2 + p_y^2 + 4K|p|p_x^2}. \quad (\text{B1})$$

$$\delta\gamma^0 \Sigma_1(k_0) = -\frac{12}{N_f} \int_{\Lambda-d\Lambda}^{\Lambda} \frac{d^3 p}{(2\pi)^3} \frac{(p_0 + k_0)}{(p_0 + k_0)^2 + p_x^2 + p_y^2} \frac{|p|}{p_0^2 + p_y^2 + 4K|p|p_x^2}. \quad (\text{B5})$$

In spherical coordinates this simplifies to

$$\delta\gamma^0 \Sigma_1(k_0) = -\frac{6k_0}{N_f \pi^2} \frac{d\Lambda}{\Lambda} \int_0^1 d\mu (1 - (1 - \mu^2)) \times \frac{1}{1 - \mu^2 + 4K\Lambda\mu^2} = -\frac{3k_0}{N_f \pi^2} \frac{d\Lambda}{\Lambda} \left[\ln\left(\frac{1}{K\Lambda}\right) - 2 + \mathcal{O}(K\Lambda) \right]. \quad (\text{B6})$$

The expression for $\delta\gamma^y \Sigma_1(k_y)$ is the same as $\delta\gamma^0 \Sigma_1(k_0)$, after mapping $k_0 \Rightarrow k_y$.

We can also examine the vertex correction for the SO(5) order parameter in a similar manner. From Eq. (4.25) at zero external momentum, we note that the vertex correction needs the integral

$$\delta V = \frac{1}{N_f} \int_{\Lambda-d\Lambda}^{\Lambda} \frac{d^3 p}{(2\pi)^3} \frac{1}{p_x^2 + p_y^2 + p_0^2} \frac{4|p|}{p_0^2 + p_y^2 + 4K|p|p_x^2} = \frac{1}{2N_f \pi^2} \frac{d\Lambda}{\Lambda} \int_0^1 d\mu \frac{4}{1 - \mu^2 + 4K\Lambda\mu^2} = \frac{1}{N_f \pi^2} \frac{d\Lambda}{\Lambda} \left[\ln\left(\frac{1}{K\Lambda}\right) + \mathcal{O}(K\Lambda) \right]. \quad (\text{B7})$$

We now proceed as usual to obtain the RG equations from the momentum shell results under the rescaling

$$x' = xe^{-\ell} y' = ye^{-\ell} \tau' = \tau \exp\left(-\int_0^\ell d\ell' z(\ell')\right). \quad (\text{B8})$$

Importantly, we note the flow of the irrelevant coupling K under this transformation

$$\frac{dK}{d\ell} = -K. \quad (\text{B9})$$

Expanding to linear order in k_x , using spherical coordinates with

$$(p_0, p_y, p_x) = \Lambda(\sin\theta \cos\phi, \sin\theta \sin\phi, \cos\theta), \quad (\text{B2})$$

and setting $\mu = \cos\theta$, we obtain

$$\delta\gamma^x \Sigma_1(k_x) = \frac{6k_x}{N_f \pi^2} \frac{d\Lambda}{\Lambda} \int_0^1 d\mu (1 - 2\mu^2) \frac{1}{1 - \mu^2 + 4K\Lambda\mu^2}. \quad (\text{B3})$$

Under normal circumstances, the μ integral would be a finite numerical constant, and the coefficient of $d\Lambda/\Lambda$ with the usual RG log which would then contribute (in this case) to the exponent η_ψ . However, that is not the case here, because of the logarithmic divergence of the μ integral near $\mu = 1$. Evaluating the μ integral, we obtain

$$\delta\gamma^x \Sigma_1(k_x) = -\frac{3k_x}{N_f \pi^2} \frac{d\Lambda}{\Lambda} \left[\ln\left(\frac{1}{K\Lambda}\right) - 4 + \mathcal{O}(K\Lambda) \right]. \quad (\text{B4})$$

In a similar manner, we obtain for the frequency dependence of the self energy

For the fermion field we define

$$\psi' = \psi \exp\left(\int_0^\ell d\ell' \frac{1 + z(\ell') + \eta_\psi(\ell')}{2}\right). \quad (\text{B10})$$




FIG. 9. Denoting the integrands of the two types of self-energy contributions in Eqs. (4.10) and (C1) with $K = 1$ as $f_A(k_x)$ and $f_B(k_0)$, we plot a numerical evaluation of $8\pi^2 f_i(k)/(k \ln^3(k))$ vs $1/\ln(k)$. The form of this expression is designed to isolate the \log^2 contribution at small momenta, and agreement with our analytic predictions should give a straight line with a slope of 1. The approximation of the Higgs propagator as Eq. (4.9) allows for greater numerical precision, as the dimensionality of the integral can be reduced by performing portions of the integral analytically. These numerical evaluations give good agreement with analytic predictions as well as calculations using the full form of the Higgs propagator.

The field ψ is not gauge invariant, and neither is its anomalous dimension η_ψ . However, the leading \log^2 term we shall find shortly is gauge invariant. In the presence of the \log^2 term, we will also see that the usual logarithm terms have a nonuniversal coefficient. So we ignore the gauge field contributions here (the gauge field induced renormalizations have been computed in Refs. [39,40]), because they only contribute logarithm terms which become part of overall terms which are nonuniversal.

Matching Eqs. (B4) and (B6) to Eqs. (B8) and (B10) we obtain

$$\eta(\ell) = \frac{6}{N_f \pi^2} \left[\ln \left(\frac{1}{K(\ell)\Lambda} \right) - 3 \right]$$

$$z(\ell) = 1 + \frac{6}{N_f \pi^2}. \quad (\text{B11})$$

Assuming a bare value $K(0) = K_0$, integrating Eq. (B9) to obtain $K(\ell) = K_0 e^{-\ell}$, and then integrating Eq. (B11) we obtain

$$\int_0^\ell \eta(\ell') d\ell' = \frac{6}{N_f \pi^2} \left[\frac{\ell^2}{2} - (\ln(K_0 \Lambda) + 3)\ell \right]. \quad (\text{B12})$$

We first analyze the Φ_1 contribution to the fermion self-energy with external momenta k_0 . This is equivalent to the k_y external momenta, as well as the Φ_2 contribution with external momenta k_0, k_x .

$$\begin{aligned} \gamma^0 \Sigma(k_0) &\approx -\frac{12}{N_f} \int \frac{d^3 p}{(2\pi)^3} \frac{p_0 + k_0}{(p_0 + k_0)^2 + p_x^2 + p_y^2} \frac{|p_x|}{p_0^2 + p_y^2 + 4K|p_x|^3} \\ &= -\frac{12}{N_f} \int \frac{dp_x}{8\pi^2} \frac{|p_x|(p_x^2 - 4K|p_x|^3 - k_0^2)}{k_0 \sqrt{-(4K)^2 |p_x|^6 + 8K|p_x|^3(p_x^2 - k_0^2) - (p_x^2 + k_0^2)^2}} \\ &\quad \times \left[\tan^{-1} \left(\frac{-p_x^2 + 4K|p_x|^3 - k_0^2}{\sqrt{-(4K)^2 |p_x|^6 + 8K|p_x|^3(p_x^2 - k_0^2) - (p_x^2 + k_0^2)^2}} \right) \right. \\ &\quad \left. + \tan^{-1} \left(\frac{-p_x^2 + 4K|p_x|^3 + k_0^2}{\sqrt{-(4K)^2 |p_x|^6 + 8K|p_x|^3(p_x^2 - k_0^2) - (p_x^2 + k_0^2)^2}} \right) \right] - \frac{12}{N_f} \int \frac{dp_x}{8\pi^2} \frac{|p_x|}{2k_0} \ln \left(\frac{p_x^2}{4K|p_x|^3} \right). \quad (\text{C1}) \end{aligned}$$

We see that the dominant term is proportional to $k_0 \log^2(Kk_0)$, arising in the same limit as in Eq. (4.12). This \log^2 dependence comes from the inverse tangents, since $2 \tan^{-1}(x) = i \log(\frac{1+ix}{1-ix})$. Assuming $K|p_x|^3 \ll p_x^2$, the expression in the denominator of the inverse tangent argument is $\approx ip_x^2$, so our integrand $\approx i \tan^{-1}(i(1 + 4K|p_x|))$. If we further assume $K|p_x| \ll 1$, we get an integrand that scales like $k_0/p_x \ln(Kp_x)$, and hence the full expression scales as $k_0 \log^2(Kk_0)$. These \log^2 contributions are verified by numerically evaluating Eqs. (4.10) and (C1) and analyzing the behavior at small k , as shown in Fig. 9.

For the Higgs vertex correction to the SO(5) order parameter, we can also isolate a \log^2 divergence. Calculating the Φ_1 correction to the vertex, we regulate the integral by including an external momenta $2k_x$ evenly distributed between the two outgoing fermions. The vertex correction is

$$\begin{aligned} &\frac{\mu^z \sigma^a \Gamma^i \sigma^a \mu^z}{N_f} \int \frac{d^3 p}{(2\pi)^3} \frac{p_x^2 - k_x^2 + p_y^2 + p_0^2}{[(p_x + k_x)^2 + p_y^2 + p_0^2][(p_x - k_x)^2 + p_y^2 + p_0^2]} \frac{4|p_x|}{p_0^2 + p_y^2 + 4K|p_x|^3} \\ &= \frac{\mu^z \sigma^a \Gamma^i \sigma^a \mu^z}{N_f} \int \frac{rdrdp_x}{(2\pi)^2} \frac{p_x^2 - k_x^2 + r^2}{[(p_x + k_x)^2 + r^2][(p_x - k_x)^2 + r^2]} \frac{4|p_x|}{r^2 + 4K|p_x|^2} \\ &= \frac{\mu^z \sigma^a \Gamma^i \sigma^a \mu^z K}{N_f} \int \frac{dp_x}{4\pi^2} \frac{|p_x|}{k_x p_x [(p_x - x_x)^2 - 4K|p_x|^2][(p_x + k_x)^2 - 4K|p_x|^2]} \\ &\quad \times \left[\left((p_x^2 - k_x^2) |p_x|^2 + \frac{(p_x + k_x)^2 (p_x - k_x)^2}{4K} \right) \ln \left(\frac{(p_x + k_x)^2}{(p_x - k_x)^2} \right) \right] \end{aligned}$$

We can now obtain the momentum dependence of physical observables by evaluating them at a scale $\ell = \ell^* = \ln(\Lambda/|p|)$. Note that the coefficient of ℓ^* involves the bare value of K_0 , and hence the coefficient of the logarithm term is nonuniversal, as claimed earlier. The leading term is \log^2 , and its coefficient is universal and agrees with that in Eq. (4.16); similarly, Eq. (B7) agrees with Eq. (4.26). Inserting the integral Eq. (B12) into Eq. (B10), we obtain results of the form in Eq. (4.29).

APPENDIX C: ISOLATION OF LOGARITHM-SQUARED DIVERGENCES IN ONE-LOOP CORRECTIONS

We state in the main text that logarithm-squared divergences in the critical SU(2) gauge theory arise in the one-loop diagrams in a certain parameter regime, given by Eq. (4.12). This is shown in the main text for the simplest one-loop calculation, which is the Φ_1 (Φ_2) contribution to the fermion self-energy with external momenta k_x (k_y). Here, we provide more general calculations for other cases.

$$\begin{aligned}
 & + (p_x + k_x)^2 \left[|p_x|^3 \ln \left(\frac{4K|p_x|^3}{(p_x + k_x)^2} \right) + \frac{(k_x^2 - p_x^2)}{4K} \ln \left(\frac{4K|p_x|^3}{(p_x - k_x)^2} \right) \right] \\
 & - (p_x - k_x)^2 \left[|p_x|^3 \ln \left(\frac{4K|p_x|^3}{(p_x - k_x)^2} \right) + \frac{(k_x^2 - p_x^2)}{4K} \ln \left(\frac{4K|p_x|^3}{(p_x + k_x)^2} \right) \right].
 \end{aligned} \tag{C2}$$

We can obtain a \log^2 from the second logarithm in the brackets in the limit given by Eq. (4.12).

APPENDIX D: ALTERNATE COMPUTATION OF Z IN THE SU(2) GAUGE THEORY

In the main text we emphasize that although irrelevant terms in the Higgs propagator turn out to strongly influence the renormalization of the critical theory, these effects cancel in the dynamical critical exponent, and its value can be computed through more standard methods. In this section, we compute z via dimensional regularization after explicitly setting the irrelevant Higgs terms to zero and show that it gives the same answer for z as in the main text. For the calculation of z , we are interested in the counterterms generated by

$$\gamma^0 \frac{\partial \Sigma}{\partial k_0} \Big|_{k=0} - \gamma^x \frac{\partial \Sigma}{\partial k_x} \Big|_{k=0} = -\frac{6}{N_f} \int \frac{d^3 p}{(2\pi)^3} \left[\frac{p_y^2}{p^4 \Gamma_1(p)} - \frac{p_0^2 - p_x^2}{p^4 \Gamma_2(p)} \right]. \tag{D1}$$

This integrand is well behaved for $p^2 \neq 0, \infty$, and one can see that the second term in brackets vanishes, since the Γ_2 propagator is invariant under $p_x \leftrightarrow p_0$.

The integrals over p_0, p_y can be performed exactly in radial coordinates, which gives

$$-\frac{3}{\pi^2 N_f} \int_{-\infty}^{\infty} dp_x \frac{1}{|p_x|} = -\frac{6}{\pi^2 N_f} \int_0^{\infty} \frac{dp_x}{p_x}. \tag{D2}$$

Continuing the p_x integral to $1 - \epsilon$ dimensions and imposing a UV cutoff Λ yields

$$-\frac{6}{\pi^2 N_f} \mu^\epsilon \int_0^\Lambda \frac{dp_0}{p_0^{1+\epsilon}} = -\frac{6}{\pi^2 N_f \epsilon} \mu^\epsilon \Lambda^{-\epsilon} = -\frac{6}{\pi^2 N_f} \left(\frac{1}{\epsilon} + \ln \left(\frac{\mu}{\Lambda} \right) + \mathcal{O}(\epsilon) \right), \tag{D3}$$

which gives the same answer for z as when the irrelevant Higgs terms were used to regulate the divergences in the self-energy.

APPENDIX E: HIGGS FIELD RENORMALIZATION IN THE SU(2) GAUGE THEORY

For completeness, we compute the \log^2 corrections in the critical SU(2) gauge theory to the Yukawa couplings at one-loop level, since these determine the renormalization of the Higgs fields. The calculations are nearly identical to those of the SO(5) order parameter.

The correction to the Φ_1 Yukawa coupling is given by the integral

$$\begin{aligned}
 & \frac{(\mu^z \sigma^b)(\mu^z \sigma^a)(\mu^z \sigma^b)}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^x \frac{\not{p} - \not{k}_1}{(p - k_1)^2} \gamma^x \frac{\not{p} - \not{k}_2}{(p - k_2)^2} \gamma^x \frac{1}{\Gamma_1(p) + K p_x^2} \\
 & + \frac{(\mu^x \sigma^b)(\mu^z \sigma^a)(\mu^x \sigma^b)}{N_f} \int \frac{d^3 p}{(2\pi)^3} \gamma^y \frac{\not{p} - \not{k}_1}{(p - k_1)^2} \gamma^x \frac{\not{p} - \not{k}_2}{(p - k_2)^2} \gamma^y \frac{1}{\Gamma_2(p) + K p_y^2}.
 \end{aligned} \tag{E1}$$

Evaluating the first term in the limit in Eq. (4.12), we set the external momenta to zero and use it as an IR cutoff k , which gives to \log^2 order

$$\frac{(\mu^z \sigma^b)(\mu^z \sigma^a)(\mu^z \sigma^b)}{2\pi^2 N_f} \ln^2(Kk). \tag{E2}$$

This coefficient is identical to the SO(5) correction, as the two integrals are the same to leading order in k . The contribution is the same for the second term in Eq. (E1), giving the final Yukawa correction

$$-\frac{\mu_z \sigma^a}{\pi^2 N_f} \ln^2(Kk). \tag{E3}$$

The correction to the Φ_2 Yukawa term is identical, as the two are related by a spatial rotation.

We now renormalize the $\Phi_{1,2}$ fields so that the Yukawa coupling remains invariant, as in Ref. [56]. Hence, the Higgs fields are renormalized at \log^2 order, $\Phi_i = \sqrt{Z_\Phi} \Phi_{i,R}$, with corrections from Z_ψ and (E3)

$$\begin{aligned}
 Z_\Phi & = 1 + \frac{6}{\pi^2 N_f} \ln^2(K\mu) + \frac{2}{\pi^2 N_f} \ln^2(K\mu) \\
 & = 1 + \frac{8}{\pi^2 N_f} \ln^2(K\mu).
 \end{aligned} \tag{E4}$$

APPENDIX F: EVALUATION OF TWO-LOOP SO(5) ORDER PARAMETER CORRECTIONS

In this Appendix, we evaluate the $\mathcal{O}(N_f^{-1})$ two-loop correction to the SO(5) order parameter, shown in the main text




FIG. 10. The $\mathcal{O}(N_f^{-1})$ two-loop correction to the SO(5) order parameter. We set all external momenta to zero. Shown is one of four possible diagrams; the other three can be obtained by either exchanging $\Phi_1 \leftrightarrow \Phi_2$, crossing the lines of the Higgs propagators, or both. All give the same contribution at zero external momenta.

and displayed here in Fig. 10 with internal momenta labeled. The diagram shown is one of four possible contributions -

additional diagrams can be generated by either exchanging $\Phi^1 \leftrightarrow \Phi^2$ or crossing the propagators of the Higgs bosons, but all give the same correction for zero external momenta. The main conclusion of this Appendix is that this contribution is well behaved upon setting the dangerously irrelevant operators to zero and only contributes standard logarithm divergences, which we argue in the main text and in Appendix B give nonuniversal corrections to the order parameter scaling. Intuitively, this may be thought of as related to the fact that these two-loop diagrams require *both* types of Higgs $\Phi_{1,2}$, as they vanish trivially when both Higgs propagators are of the same type. As the \log^2 divergences are connected to the rotational symmetry breaking in the $\mathcal{O}(N_f)$ effective action for the Higgs propagators, it is natural—although still a nontrivial fact—that these two-loop diagrams which respect rotational symmetry only contribute single logarithm divergences.

This two-loop correction vanishes for the VBS order parameter, so we focus on the Néel order parameter, where the source vertex contributes a factor of $\mu^y \sigma^a$. We first evaluate the fermion loop integral,

$$\begin{aligned}
 & (-1)N_f \text{Tr} \sigma^b \sigma^c \mu^y \mu^x \mu^z \int \frac{d^3 p}{(2\pi)^3} \frac{\not{p}}{p^2} \gamma^x \frac{\not{p} + \not{q}}{(p+q)^2} \gamma^y \frac{\not{p}}{p^2} \\
 &= -4N_f \delta_{bc} \text{Tr} [\gamma^\mu \gamma^x \gamma^y \gamma^v \gamma^\sigma] \int \frac{d^3 p}{(2\pi)^3} \frac{p_\mu (p+q)_\nu p_\sigma}{p^4 (p+q)^2} \\
 &= -8iN_f \delta_{bc} (\delta^{\mu x} \epsilon^{y\nu\sigma} + \delta^{y\nu} \epsilon^{x\sigma\mu} + \delta^{\sigma y} \epsilon^{x\nu\mu} - \delta^{\nu\sigma} \delta^{\mu 0}) \int \frac{d^3 p}{(2\pi)^3} \frac{p_\mu (p+q)_\nu p_\sigma}{p^4 (p+q)^2}. \tag{F1}
 \end{aligned}$$

The integral over p yields

$$\int \frac{d^3 p}{(2\pi)^3} \frac{p_\mu (p+q)_\nu p_\sigma}{p^4 (p+q)^2} = \frac{1}{128} \frac{1}{|q|} \left[3\delta^{\mu\sigma} q_\nu - \delta^{\sigma\nu} q_\mu - \delta^{\mu\nu} q_\sigma + \frac{q_\mu q_\nu q_\sigma}{q^2} \right] \tag{F2}$$

and contracting with the tensors in Eq. (F1) gives the final contribution of the fermion loop

$$\frac{iN_f}{2} \frac{q_0}{|q|} \delta_{bc}. \tag{F3}$$

We combine this with the remaining loop integral, setting the coefficient K of the irrelevant operators to zero, to give

$$\frac{i}{2N_f} \delta_{bc} \mu^z \mu^x \sigma^b \sigma^c \int \frac{d^3 q}{(2\pi)^3} \frac{q_0}{|q|} \gamma^x \frac{(-\not{q})}{q^2} \gamma^y \frac{16q^2}{(q_0^2 + q_x^2)(q_0^2 + q_y^2)} = -\frac{16\mu^y \sigma^a}{N_f} \int \frac{d^3 q}{(2\pi)^3} \frac{q_0^2}{|q|} \frac{1}{(q_0^2 + q_x^2)(q_0^2 + q_y^2)}. \tag{F4}$$

Focusing on the integrand, we can compute this by converting to radial coordinates,

$$\int \frac{dz d\theta r dr}{(2\pi)^3} \frac{z^2}{\sqrt{z^2 + r^2}} \frac{1}{(z^2 + r^2 \cos^2 \theta)(z^2 + r^2 \sin^2 \theta)} = \frac{1}{2\pi^2} \int dz dr \frac{|z|r}{(z^2 + r^2)(2z^2 + r^2)} = \frac{\ln 2}{4\pi^2} \int dz \frac{1}{|z|}. \tag{F5}$$

Hence, this two-loop contribution only contributes a standard logarithm divergence and is subleading in comparison to the one-loop Higgs corrections.

We also analyze the two-loop corrections to the $\bar{\psi} \psi$ bilinear, whose symmetry properties correspond to the scalar spin chirality. This is motivated by the fact that \log^2 terms in the $\mathcal{O}(1/N_f)$ one-loop corrections exactly cancel the \log^2 self-energy terms. Hence, if two-loop corrections only contributed standard logarithm divergences, then the scalar spin chirality would have a power law decay at $\mathcal{O}(1/N_f)$. In fact, the

two-loop corrections involving two Higgs propagators vanish exactly. If the Higgs propagators are different, as was the case for the Néel corrections, then the trace over μ in the fermion loop vanishes. If the Higgs propagators are the same, then the trace over γ vanishes, since

$$\begin{aligned}
 & \text{Tr} [\gamma^\mu \gamma^x \gamma^y \gamma^v \gamma^x \gamma^\sigma] p_\mu (p+q)_\nu p_\sigma \\
 &= \text{Tr} [\gamma^\mu \gamma^y \gamma^v \gamma^y \gamma^\sigma] p_\mu (p+q)_\nu p_\sigma = 0. \tag{F6}
 \end{aligned}$$

- [1] L. B. Ioffe and A. I. Larkin, Effective action of a two-dimensional antiferromagnet, *Int. J. Mod. Phys. B* **02**, 203 (1988).
- [2] M. P. Gelfand, R. R. P. Singh, and D. A. Huse, Zero-temperature ordering in two-dimensional frustrated quantum Heisenberg antiferromagnets, *Phys. Rev. B* **40**, 10801 (1989).
- [3] E. Dagotto and A. Moreo, Phase Diagram of the Frustrated Spin-1/2 Heisenberg Antiferromagnet in 2 Dimensions, *Phys. Rev. Lett.* **63**, 2148 (1989).
- [4] P. Chandra, P. Coleman, and A. I. Larkin, Ising Transition in Frustrated Heisenberg Models, *Phys. Rev. Lett.* **64**, 88 (1990).
- [5] N. Read and S. Sachdev, Large- N Expansion for Frustrated Quantum Antiferromagnets, *Phys. Rev. Lett.* **66**, 1773 (1991).
- [6] S. Sachdev and N. Read, Large N Expansion for Frustrated and Doped Quantum Antiferromagnets, *Int. J. Mod. Phys. B* **5**, 219 (1991).
- [7] L. Wang and A. W. Sandvik, Critical Level Crossings and Gapless Spin Liquid in the Square-Lattice Spin-1/2 J_1 - J_2 Heisenberg Antiferromagnet, *Phys. Rev. Lett.* **121**, 107202 (2018).
- [8] F. Ferrari and F. Becca, Gapless spin liquid and valence-bond solid in the J_1 - J_2 Heisenberg model on the square lattice: Insights from singlet and triplet excitations, *Phys. Rev. B* **102**, 014417 (2020).
- [9] Y. Nomura and M. Imada, Dirac-type nodal spin liquid revealed by machine learning, [arXiv:2005.14142](https://arxiv.org/abs/2005.14142).
- [10] W.-Y. Liu, S.-S. Gong, Y.-B. Li, D. Poilblanc, W.-Q. Chen, and Z.-C. Gu, Gapless quantum spin liquid and global phase diagram of the spin-1/2 J_1 - J_2 square antiferromagnetic Heisenberg model, [arXiv:2009.01821](https://arxiv.org/abs/2009.01821).
- [11] F. Becca, Talk at KITP, University of California, Santa Barbara, November 5, Correlated Systems with Multicomponent Local Hilbert Spaces (2020).
- [12] N. Read and S. Sachdev, Valence-Bond and Spin-Peierls Ground States of Low-Dimensional Quantum Antiferromagnets, *Phys. Rev. Lett.* **62**, 1694 (1989).
- [13] N. Read and S. Sachdev, Spin-Peierls, valence-bond solid, and Néel ground states of low-dimensional quantum antiferromagnets, *Phys. Rev. B* **42**, 4568 (1990).
- [14] O. I. Motrunich and A. Vishwanath, Emergent photons and transitions in the O(3) sigma model with hedgehog suppression, *Phys. Rev. B* **70**, 075104 (2004).
- [15] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, Deconfined Quantum Critical Points, *Science* **303**, 1490 (2004).
- [16] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, Quantum criticality beyond the Landau-Ginzburg-Wilson paradigm, *Phys. Rev. B* **70**, 144407 (2004).
- [17] L. Capriotti, F. Becca, A. Parola, and S. Sorella, Resonating Valence Bond Wave Functions for Strongly Frustrated Spin Systems, *Phys. Rev. Lett.* **87**, 097201 (2001).
- [18] D. A. Ivanov and T. Senthil, Projected wave functions for fractionalized phases of quantum spin systems, *Phys. Rev. B* **66**, 115111 (2002).
- [19] W.-J. Hu, F. Becca, A. Parola, and S. Sorella, Direct evidence for a gapless \mathbb{Z}_2 spin liquid by frustrating Néel antiferromagnetism, *Phys. Rev. B* **88**, 060402(R) (2013).
- [20] F. Ferrari and F. Becca, Spectral signatures of fractionalization in the frustrated Heisenberg model on the square lattice, *Phys. Rev. B* **98**, 100405(R) (2018).
- [21] X. G. Wen, Mean-field theory of spin-liquid states with finite energy gap and topological orders, *Phys. Rev. B* **44**, 2664 (1991).
- [22] A. Kitaev, Fault tolerant quantum computation by anyons, *Ann. Phys.* **303**, 2 (2003).
- [23] T. Senthil and M. P. A. Fisher, \mathbb{Z}_2 gauge theory of electron fractionalization in strongly correlated systems, *Phys. Rev. B* **62**, 7850 (2000).
- [24] X.-G. Wen, Quantum orders and symmetric spin liquids, *Phys. Rev. B* **65**, 165113 (2002).
- [25] T. Senthil and P. A. Lee, Cuprates as doped U(1) spin liquids, *Phys. Rev. B* **71**, 174515 (2005).
- [26] A. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys.* **321**, 2 (2006).
- [27] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, Deconfined Quantum Critical Points: Symmetries and Dualities, *Phys. Rev. X* **7**, 031051 (2017).
- [28] A. Thomson and S. Sachdev, Fermionic Spinon Theory of Square Lattice Spin Liquids near the Néel State, *Phys. Rev. X* **8**, 011012 (2018).
- [29] X.-Y. Song, Y.-C. He, A. Vishwanath, and C. Wang, From Spinon Band Topology to the Symmetry Quantum Numbers of Monopoles in Dirac Spin Liquids, *Phys. Rev. X* **10**, 011033 (2020).
- [30] X.-Y. Song, C. Wang, A. Vishwanath, and Y.-C. He, Unifying Description of Competing Orders in Two Dimensional Quantum Magnets, *Nat. Commun.* **10**, 4254 (2019).
- [31] R. Ma and C. Wang, Theory of deconfined pseudocriticality, *Phys. Rev. B* **102**, 020407(R) (2020).
- [32] A. Nahum, Note on Wess-Zumino-Witten models and quasiuniversality in 2+1 dimensions, *Phys. Rev. B* **102**, 201116(R) (2020).
- [33] Z. Wang, M. P. Zaletel, R. S. K. Mong, and F. F. Assaad, Phases of the (2+1) Dimensional SO(5) Nonlinear Sigma Model with Topological Term, *Phys. Rev. Lett.* **126**, 045701 (2021).
- [34] Y.-C. He, J. Rong, and N. Su, Non-Wilson-Fisher kinks of $O(N)$ numerical bootstrap: from the deconfined phase transition to a putative new family of CFTs, *SciPost Phys.* **10**, 115 (2021).
- [35] V. Gorbenko, S. Rychkov, and B. Zan, Walking, Weak first-order transitions, and Complex CFTs, *J. High Energy Phys.* **10** (2018) 108.
- [36] V. Gorbenko, S. Rychkov, and B. Zan, Walking, Weak first-order transitions, and Complex CFTs II. Two-dimensional Potts model at $Q > 4$, *SciPost Phys.* **5**, 050 (2018).
- [37] H. Ma and Y.-C. He, Shadow of complex fixed point: Approximate conformality of $Q > 4$ Potts model, *Phys. Rev. B* **99**, 195130 (2019).
- [38] S. Gazit, F. F. Assaad, S. Sachdev, A. Vishwanath, and C. Wang, Confinement transition of \mathbb{Z}_2 gauge theories coupled to massless fermions: Emergent quantum chromodynamics and SO(5) symmetry, *Proc. Natl. Acad. Sci.* **115**, E6987 (2018).
- [39] Y. Ran and X.-G. Wen, Continuous quantum phase transitions beyond Landau's paradigm in a large- N spin model, [arXiv:cond-mat/0609620](https://arxiv.org/abs/cond-mat/0609620).
- [40] Y. Ran, Spin liquids, exotic phases and phase transitions, Ph.D. thesis, MIT Physics Department, Cambridge, MA, 2007.
- [41] I. Affleck and J. B. Marston, Large- n limit of the Heisenberg-Hubbard model: Implications for high- T_c superconductors, *Phys. Rev. B* **37**, 3774 (1988).

- [42] C. Xu and S. Sachdev, Square-Lattice Algebraic Spin Liquid with SO(5) Symmetry, *Phys. Rev. Lett.* **100**, 137201 (2008).
- [43] S. Sachdev and K. Park, Ground States of Quantum Antiferromagnets in Two Dimensions, *Ann. Phys.* **298**, 58 (2002).
- [44] S. Sachdev, H. D. Scammell, M. S. Scheurer, and G. Tarnopolsky, Gauge theory for the cuprates near optimal doping, *Phys. Rev. B* **99**, 054516 (2019).
- [45] R. A. Jalabert and S. Sachdev, Spontaneous alignment of frustrated bonds in an anisotropic, three-dimensional Ising model, *Phys. Rev. B* **44**, 686 (1991).
- [46] S. Sachdev and M. Vojta, Translational symmetry breaking in two-dimensional antiferromagnets and superconductors, *J. Phys. Soc. Jpn* **69**, Supp. B, 1 (1999).
- [47] Y. Huh, M. Punk, and S. Sachdev, Vison states and confinement transitions of Z_2 spin liquids on the kagome lattice, *Phys. Rev. B* **84**, 094419 (2011).
- [48] S. Sachdev, Topological order, emergent gauge fields, and Fermi surface reconstruction, *Rept. Prog. Phys.* **82**, 014001 (2019).
- [49] M. P. Zaletel and A. Vishwanath, Constraints on Topological Order in Mott Insulators, *Phys. Rev. Lett.* **114**, 077201 (2015).
- [50] M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, Translational Symmetry and Microscopic Constraints on Symmetry-Enriched Topological Phases: A View from the Surface, *Phys. Rev. X* **6**, 041068 (2016).
- [51] P. Bonderson, M. Cheng, K. Patel, and E. Plamadeala, Topological Enrichment of Luttinger's Theorem, [arXiv:1601.07902](https://arxiv.org/abs/1601.07902).
- [52] M. A. Metlitski and R. Thorngren, Intrinsic and emergent anomalies at deconfined critical points, *Phys. Rev. B* **98**, 085140 (2018).
- [53] D. V. Else and R. Thorngren, Topological theory of Lieb-Schultz-Mattis theorems in quantum spin systems, *Phys. Rev. B* **101**, 224437 (2020).
- [54] A. Tanaka and X. Hu, Many-Body Spin Berry Phases Emerging from the π -Flux State: Competition between Antiferromagnetism and the Valence-Bond-Solid State, *Phys. Rev. Lett.* **95**, 036402 (2005).
- [55] T. Senthil and M. P. A. Fisher, Competing orders, nonlinear sigma models, and topological terms in quantum magnets, *Phys. Rev. B* **74**, 064405 (2006).
- [56] Y. Huh and S. Sachdev, Renormalization group theory of nematic ordering in d -wave superconductors, *Phys. Rev. B* **78**, 064512 (2008).
- [57] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, International series of monographs on physics (Clarendon Press, Oxford, 2002).
- [58] B. L. Altshuler, A. G. Aronov, and P. A. Lee, Interaction Effects in Disordered Fermi Systems in Two Dimensions, *Phys. Rev. Lett.* **44**, 1288 (1980).
- [59] A. M. Finkelstein, Influence of Coulomb interaction on the properties of disordered metals, *Sov. Phys. JETP* **57**, 97 (1983) [*Zh. Eksp. Teor. Fiz.* **84**, 168 (1983)].
- [60] P. A. Lee and T. V. Ramakrishnan, Disordered electronic systems, *Rev. Mod. Phys.* **57**, 287 (1985).
- [61] Y. Barlas and K. Yang, Non-Fermi-liquid behavior in neutral bilayer graphene, *Phys. Rev. B* **80**, 161408(R) (2009).
- [62] R. Nandkishore and L. Levitov, Electron interactions in bilayer graphene: Marginal Fermi liquid and zero-bias anomaly, *Phys. Rev. B* **82**, 115431 (2010).
- [63] M. Hermele, T. Senthil, and M. P. A. Fisher, Algebraic spin liquid as the mother of many competing orders, *Phys. Rev. B* **72**, 104404 (2005).
- [64] W. Rantner and X.-G. Wen, Spin correlations in the algebraic spin liquid: Implications for high- T_c superconductors, *Phys. Rev. B* **66**, 144501 (2002).
- [65] H. Shao, W. Guo, and A. W. Sandvik, Quantum criticality with two length scales, *Science* **352**, 213 (2016).
- [66] Y.-H. Zhang and S. Sachdev, From the pseudogap metal to the Fermi liquid using ancilla qubits, *Phys. Rev. Research* **2**, 023172 (2020).
- [67] Y.-H. Zhang and S. Sachdev, Deconfined criticality and ghost Fermi surfaces at the onset of antiferromagnetism in a metal, *Phys. Rev. B* **102**, 155124 (2020).
- [68] A. Nikolaenko, M. Tikhonovskaya, S. Sachdev, and Y.-H. Zhang, Small to large Fermi surface transition in a single band model, using randomly coupled ancillas, *Phys. Rev. B* **103**, 235138 (2021).
- [69] H.-C. Jiang and S. A. Kivelson, High Temperature Superconductivity in a Lightly Doped Quantum Spin Liquid, [arXiv:2104.01485](https://arxiv.org/abs/2104.01485).
- [70] S. Gong, W. Zhu, and D. N. Sheng, Robust d-wave superconductivity in the square-lattice t - J model, [arXiv:2104.03758](https://arxiv.org/abs/2104.03758) [cond-mat.str-el].
- [71] J. Yang, A. W. Sandvik, and L. Wang, Quantum Criticality and Spin Liquid Phase in the Shastry-Sutherland model, [arXiv:2104.08887](https://arxiv.org/abs/2104.08887) [cond-mat.str-el].
- [72] The symmetry P_x is related to the other symmetries through $R_{\pi/2} P_y R_{\pi/2}^{-1}$.