



Fracton-elasticity duality in twisted moiré superlattices

Jonas Gaa ¹, Grgur Palle,¹ Rafael M. Fernandes,² and Jörg Schmalian ^{1,3}

¹*Institute for Theory of Condensed Matter, Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany*

²*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

³*Institute for Quantum Materials and Technologies, Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany*



(Received 7 May 2021; revised 26 July 2021; accepted 2 August 2021; published 13 August 2021)

We formulate a fracton-elasticity duality for twisted moiré superlattices, taking into account that they are incommensurate crystals with dissipative phason dynamics. From a dual tensor-gauge formulation, as compared to standard crystals, we identify twice the number of conserved charges that describe topological lattice defects, namely, disclinations and another type of defects that we dub *discompressions*. The key implication of these conservation laws is that both glide and climb motions of lattice dislocations are suppressed, indicating that dislocation networks may become exceptionally stable. We also generalize our results to other planar incommensurate crystals and quasicrystals.

DOI: [10.1103/PhysRevB.104.064109](https://doi.org/10.1103/PhysRevB.104.064109)

I. INTRODUCTION

Twisted bilayer graphene (TBG) [1–4] forms an incommensurate moiré lattice. The description of its electronic degrees of freedom, to a very good approximation, can be done using concepts of periodic crystals, due to the weak scattering between opposite valleys [5–8]. However, structurally, there is no periodicity in the system if the twist angle takes a generic value. As was demonstrated by Ochoa [9], this has profound implications for the elasticity theory of TBG. Phason modes $\mathbf{w}(\mathbf{x}, t)$, which correspond to acoustic branches of the incommensurate lattice, dominate lattice vibrations on the scale of the moiré period and give rise to an additional rotational stiffness κ in the elastic energy

$$E_{\text{el}} \rightarrow E_{\text{el}} + \frac{\kappa}{8} \int d^2x [\partial_x w_y - \partial_y w_x]^2. \quad (1)$$

While in standard elasticity theory such a term is not allowed by rotational symmetry [10], in the case of TBG the adhesive potential between the two layers gives rise to $\kappa > 0$. Notice that the elasticity theory of planar quasicrystals [11–13] can also be formulated in terms of Eq. (1). The influence of the rotational term and phason excitations on electron-lattice couplings was discussed in Refs. [9,14], while their role in the context of electronic nematicity was analyzed in Ref. [15].

Phasons in TBG have a very straightforward interpretation. Ignoring the structural relaxation of the lattice, they have a simple relationship with the relative displacement between layers [9]. To illustrate a twist of the phason mode, we therefore consider a moiré superlattice with sixfold rotational symmetry, described in terms of the density profile

$$\varrho_\theta(\mathbf{r}) = \sum_{\mathbf{G}} |\rho_{\mathbf{G}}| e^{i(\mathbf{G} \cdot \mathbf{r} - \phi_{\mathbf{G}})}. \quad (2)$$

\mathbf{G} are reciprocal lattice vectors of the moiré superlattice and $\phi_{\mathbf{G}} = \mathbf{w} \cdot \mathbf{G}$. In Fig. 1, we show the density profile with

$\mathbf{w} = \theta \hat{\mathbf{z}} \times \mathbf{r}$ for $\theta = 0$ and with an excited phason, i.e., for a finite rotation θ of the moiré pattern. For simplicity, we consider only the leading harmonics, \mathbf{G}_1 , \mathbf{G}_2 , and $\mathbf{G}_3 \equiv -\mathbf{G}_1 - \mathbf{G}_2$, with $|\rho_{\mathbf{G}_i}| \equiv \rho$ and $\mathbf{G}_{1,2}$ denoting the standard primitive vectors of the triangular lattice.

The dynamics of phasons in incommensurate crystals also differs from that of standard acoustic phonons by the fact that it is strongly damped at long times [16,17]. This is due to the friction that opposes the relative motion of the incommensurate mass-density wave. In usual elasticity, this damping does not occur because the displacement couples to the generator of translations, i.e. the momentum density. In TBG, however, anharmonic terms of the adhesion potential between the layers give rise to a finite damping [18].

In this paper, we discuss the impact of the rotational stiffness and phason damping on inevitable defects of the moiré lattice such as dislocations and disclinations. To this end, we extend the recently formulated fracton formulation of standard elasticity [19–21] to incommensurate crystals with overdamped dynamics by developing an appropriate tensor gauge field theory. The analogy to a tensor version of electromagnetism allows us to draw general conclusions about the lattice defect dynamics of TBG. The finite rotational stiffness of Eq. (1) increases the degrees of freedom compared to standard elasticity theory, giving rise to a type of defect that we call *discompression*. As a result, defects of the moiré lattice, which can be described in terms of fractons, are governed by an additional conservation law that dramatically affects the mobility of lattice defects. In particular, we find that in an incommensurate crystal with phason excitations, both glide and climb of dislocations are forbidden and the dynamics of defects is always subdiffusive. This implies that dislocation networks in twisted bilayer graphene are expected to be exceptionally stable. We also show that our theory directly applies to other incommensurate systems such as planar quasicrystals.




FIG. 1. Density profile for the triangular moiré superlattice, as given by Eq. (2), in the cases without rotation [panel (a), $\varrho_{\theta=0}(\mathbf{r})$] and with a rotation of $\theta/(2\pi) = 0.03$ [panel (b), $\varrho_{\theta}(\mathbf{r})$]. The phason excitation is illustrated by the difference in density profiles, $\delta\varrho_{\theta}(\mathbf{r}) = \varrho_{\theta}(\mathbf{r}) - \varrho_{\theta=0}(\mathbf{r})$, which is plotted in panel (c), and whose absolute value is plotted in panel (d). The underlying triangular lattice is a guide to the eye.

The impact of topological lattice defects on the mechanical and electronic properties of graphene and bilayer graphene has been widely discussed [22–24]. Lattice disorder has also been recognized as a key ingredient in TBG, with experiments reporting sharp variations of the twist angle and of uniaxial heterostrain over relatively short length scales, which are manifested in local electronic properties [25,26]. Here, we focus on the topological defects of the emergent moiré incommensurate lattice in TBG. While its elastic properties derive from those of the coupled graphene layers, we adopt a coarse-grained low-energy description, as in Ref. [9], that does not require mapping the defects of the moiré lattice onto individual carbon atoms.

Defects of two-dimensional lattices can be efficiently captured in terms of a dual description of elasticity theory, where disclinations emerge as effective charges, dislocations as dipoles of these charges, and elastic forces between them are transmitted by gauge fields [27–32]. This approach readily demonstrates why dislocations in commensurate lattices can glide in the direction of the Burgers vector \mathbf{b} , but not climb perpendicular to it [31]. While defects in aperiodic crystals were discussed in Ref. [33–35], their dynamics remains an open problem.

A very elegant formulation of this duality was recently achieved in Refs. [19–21,36–39]. It was recognized that the dual formulation of the usual elasticity theory is, in fact, a fracton field theory. Fracton phases describe quantum phases of matter with excitations of restricted mobility [40]. If considered in isolation, a fracton is immobile, either along certain directions or as a whole. It can only move collectively, via interactions with other degrees of freedom. Fractons were initially discussed in the context of nonergodic quantum glass models [41] and stabilizer codes for self-correcting quantum memory [42]. More recently, it was recognized that fractons can be efficiently described in terms of tensor gauge theories

[43]. The restricted mobility of fractons emerges in terms of dipole conservation laws and gives rise to anomalous, sub-diffusive hydrodynamics [44–47].

II. ELASTICITY THEORY OF TWISTED BILAYER GRAPHENE CRYSTALS

We start with the action of elasticity theory [10,29], supplemented by the rotational stiffness of Eq. (1). To include dissipative dynamics, we use the Schwinger-Keldysh formalism [48]:

$$S = \frac{1}{2} \int_{\mathbf{x}, t, t' \in \mathcal{C}} w_i(\mathbf{x}, t) D^{-1}(t - t') w_i(\mathbf{x}, t') - \frac{1}{2} \int_{\mathbf{x}, t \in \mathcal{C}} [C_{ij,kl} w_{ij}(\mathbf{x}, t) w_{kl}(\mathbf{x}, t) + \kappa \vartheta_w^2(\mathbf{x}, t)]. \quad (3)$$

Here, $\int_{\mathbf{x}, t \in \mathcal{C}} = \int_{\mathcal{C}} dt \int_V d^2x$ indicates that the time integration has to be performed on the round-trip Keldysh contour, while the spatial integration goes over the crystal volume V . $w_{ij} \equiv \frac{1}{2}(\partial_i w_j + \partial_j w_i)$ is the symmetric phason strain, the elastic constants $C_{ij,kl}$ are summarized in Ref. [9], and the bond angle variation is $\vartheta_w \equiv \frac{1}{2}\epsilon_{ij}\partial_i w_j$. Damping enters through the nonlocal-in-time contribution

$$D^{-1}(t - t') = -\delta(t - t')\partial_t^2 + \gamma(t - t'), \quad (4)$$

where we assume Ohmic damping $\text{Im}\gamma^R(\omega) = \gamma_0\omega$ for the retarded self-energy. Without damping, the equation of motion become elastic wave equations, while the dynamics becomes diffusive at large γ_0 . For further details on the Keldysh formalism, see the Appendix.

We shall use a notation that will prove efficient in two dimensions. In it, we always start with lower index vectors or tensors A_i . The raising of their indices we define as a contraction with the Levi-Civita symbol:

$$A^i \equiv \epsilon_{ij} A_j. \quad (5)$$

From $A_i = (A_x, A_y)$ and $A^i \equiv \epsilon_{ij} A_j$ we see that the upper index vector $A^i = (A_y, -A_x)$ is the lower index vector rotated in the clockwise direction by $\pi/2$. This ensures that $\mathbf{A} \cdot \mathbf{B} = A_i B_i = A^i B^i$. On the other hand, $(\mathbf{A} \times \mathbf{B})_z = A_i B^i = -A^i B_i$ and in particular $A_i A^i = 0$. The divergence and two-dimensional curl may likewise be expressed in this notation as

$$\nabla \cdot \mathbf{A} = \partial_i A_i, \quad (\nabla \times \mathbf{A})_z = \partial_i A^i = -\partial^i A_i. \quad (6)$$

For single-valued functions f , $\partial_i \partial^i f = 0$. For multivalued functions, the partial derivative do not commute at the branch cuts. The most important benefit of this notation, reminiscent of the van der Waerden spinor notation [49], is that the correspondence between elasticity and tensor electromagnetism [19–21,36] amounts to simply raising all the indices.

A. Dual gauge theory

Building on the approach of Refs. [19–21], the first step in formulating a dual description is to introduce the fields $\partial_i w_i \rightarrow \pi_i$, $C_{ij,kl} w_{kl} \rightarrow \sigma_{ij}$, and $\kappa \vartheta_w \rightarrow M$ through a series

of Hubbard-Stratonovich transformations. They can be associated with momentum, stress, and torque, respectively. This yields the real-time action

$$S = \frac{1}{2} \int_{\mathbf{x}, t \in \mathcal{C}} [C_{ij,kl}^{-1} \sigma_{ij}(\mathbf{x}, t) \sigma_{kl}(\mathbf{x}, t) + \kappa^{-1} M^2(\mathbf{x}, t)] - \frac{1}{2} \int_{\mathbf{x}, t, t' \in \mathcal{C}} \pi_i(\mathbf{x}, t) \Gamma(t - t') \pi_i(\mathbf{x}, t') + S_w, \quad (7)$$

where the retarded version of Γ is $\Gamma^R(\omega) = \omega^2 / (\omega^2 + i\gamma_0\omega)$. The last term

$$S_w = \int_{\mathbf{x}, t \in \mathcal{C}} [\pi_i(\partial_t w_i) - \sigma_{ij} w_{ij} - M \vartheta_w] \quad (8)$$

still contains the original phason field.

The next step is to decompose $w_i = \tilde{w}_i + w_i^{(s)}$ into a smooth part \tilde{w}_i and a singular part $w_i^{(s)}$ due to topological defects. After integrating out the smooth part, we obtain the constraint

$$\partial_t \pi_j = \partial_i \Sigma_{ij}, \quad (9)$$

which resembles Newton's second law with a non-symmetric stress

$$\Sigma_{ij} = \sigma_{ij} + \frac{1}{2} \epsilon_{ij} M, \quad (10)$$

consisting of the symmetric stress tensor σ_{ij} and the torque M . Whereas σ_{ij} also appears in ordinary crystals, the torque only arises due to the finite rotational stiffness.

The last step is to introduce (tensorial) gauge potentials that enforce the constraint (9):

$$\begin{aligned} \Sigma_{ij} &= -\partial_t A_{ij} - \partial^i \phi_j, \\ \pi_j &= -\partial_t A_{ij}. \end{aligned} \quad (11)$$

These gauge potentials are invariant under the gauge transformations

$$\begin{aligned} A_{ij} &\mapsto A_{ij} + \partial^i \Lambda_j, \\ \phi_j &\mapsto \phi_j - \partial_t \Lambda_j. \end{aligned} \quad (12)$$

The stress-strain coupling Eq. (8) now takes the familiar form of a charge/current-potential coupling:

$$S_w = \int_{\mathbf{x}, t \in \mathcal{C}} (A_{ij} J_{ij} - \phi_j Q_j). \quad (13)$$

Due to the tensorial nature of the gauge fields, the charge density

$$Q_j = \partial^i \partial_i w_j^{(s)} \quad (14)$$

is a vector and the corresponding current density,

$$J_{ij} = (\partial_i \partial_t - \partial_t \partial_i) w_j^{(s)}, \quad (15)$$

a tensor. Demanding the invariance of Eq. (13) under the gauge transformations yields the continuity equation:

$$\partial_t Q_j + \partial^i J_{ij} = 0. \quad (16)$$

B. Electromagnetic analogy

Let us develop, in analogy to Refs. [19–21,36] for standard elasticity, a physically illuminating electromagnetic analogy

of the conserved vector charge Q_j . Consider a single charge

$$Q_j(\mathbf{x}, t) = q_j \delta(\mathbf{x} - \mathbf{r}(t)) \quad (17)$$

at position $\mathbf{r}(t)$ that moves with velocity $\mathbf{v}(t) = \dot{\mathbf{r}}(t)$. Equation (16) is then fulfilled by

$$J_{ij}(\mathbf{x}, t) = v^i q_j \delta(\mathbf{x} - \mathbf{r}(t)). \quad (18)$$

Inserting this into the action (13) yields the Lorentz force

$$F_i = \delta S_w / \delta r_i = -(\Sigma^i_j + v^i \pi_j) q_j. \quad (19)$$

The problem behaves like the electrodynamics of non-symmetric tensor electric fields Σ^i_j and vector magnetic fields π_i . This analogy also helps us understand why the overdamped dynamics of the phasons does not spoil the entire gauge description. The dissipative propagator $\Gamma(t - t')$ for $\pi_i(t)$ in Eq. (7) translates, in standard electromagnetism, to a frequency-dependent magnetic permeability $\Gamma(\omega)$, which clearly does not violate the gauge description of electromagnetism.

Besides the effects of dissipation, a crucial new aspect of the incommensurate moiré lattice elasticity is the vector continuity equation (16) that can be rephrased in terms of two conserved scalar densities:

$$\begin{aligned} \rho^{(\ell)} &= \partial_j Q_j, \\ \rho^{(t)} &= \partial^j Q_j, \end{aligned} \quad (20)$$

with continuity equations

$$\begin{aligned} \partial_t \rho^{(\ell)} + \partial_j \partial^j J_{ij} &= 0, \\ \partial_t \rho^{(t)} + \partial^j \partial^j J_{ij} &= 0. \end{aligned} \quad (21)$$

The transverse density $\rho^{(t)}$ also appears in standard elasticity, where it was identified as the density of disclinations, with Q_i denoting the Burgers vector density [19,30]. The longitudinal density $\rho^{(\ell)}$ is the new conserved charge present only in incommensurate crystals.

C. Interpretation of the charges

The charges of the elastic gauge theory are topological lattice defects that make the displacement field w_i multivalued [32]. We can write the singular part of a dislocation displacement field as

$$\mathbf{w}^{(s,Q)}(\mathbf{x}) = \frac{\text{Im} \text{Log}(z)}{2\pi} \mathbf{b}, \quad (22)$$

where $z = x_1 + ix_2$, $\text{Log}(z)$ is the principal branch of the logarithm with a branch cut along the negative x_1 axis, and $\mathbf{b} = b_1 \hat{\mathbf{e}}_1 + b_2 \hat{\mathbf{e}}_2$ is the Burgers vector. At the origin the partial derivatives do not commute, yielding $Q_j(\mathbf{x}) = b_j \delta(\mathbf{x})$, which confirms that the vector charge is the Burgers vector density.

Next, we construct a disclination from a Dirac string of dislocations. By integrating $\int_{-\infty}^0 dx'_1 \mathbf{w}^{(s,Q)}(x_1 - x'_1, x_2)$, ignoring all except the singular parts, and setting $b_1 = 0$ and $b_2 \neq 0$, we obtain

$$\mathbf{w}^{(s,t)}(\mathbf{x}) = -b_2 \frac{\text{Im}[z \text{Log}(z)]}{2\pi} \hat{\mathbf{e}}_2, \quad (23)$$

with $Q_j = b_2 \delta_{2j} \Theta(-x_1) \delta(x_2)$, $\rho^{(\ell)} = b_2 \Theta(-x_1) \delta'(x_2)$, and $\rho^{(t)} = b_2 \delta(x)$. This confirms that $\rho^{(t)}$ is the density of

This gives

$$\partial_t C_{ii,kl}^{-1} \sigma_{kl} = \partial_i(\Gamma^R \pi_i) - J_{ii}, \quad (27)$$

$$\frac{2}{\kappa} \partial_t M = \partial_i(\Gamma^R \pi^i) - J_i^i, \quad (28)$$

where

$$(\Gamma^R \pi_j)(x, t) = \int dt' \Gamma^R(t-t') \pi_j(x, t'). \quad (29)$$

Now we use the conservation laws (21), partially integrate the quadrupolar moments twice, and then use the equations of motion. This yields

$$\begin{aligned} \frac{d}{dt} \int_V |\mathbf{x}|^2 \rho^{(t)} &\stackrel{(21)}{=} - \int_V x_k x_k \partial_j \partial^j J_{ij} \\ &= -2 \int_V J_{ii} + \int_V \partial_j (2x_i J^{ji} - |\mathbf{x}|^2 \partial_i J^{ij}) \\ &\stackrel{(27)}{=} 2 \int_V \partial_i C_{ii,kl}^{-1} \sigma_{kl} \\ &\quad + \int_V \partial_j (2x_i J^{ji} - |\mathbf{x}|^2 \partial_i J^{ij} - 2\Gamma^R \pi_j) \end{aligned} \quad (30)$$

and

$$\begin{aligned} \frac{d}{dt} \int_V |\mathbf{x}|^2 \rho^{(\ell)} &\stackrel{(21)}{=} - \int_V x_k x_k \partial_j \partial^j J_{ij} \\ &= -2 \int_V J_i^i + \int_V \partial_j (|\mathbf{x}|^2 \partial_i J^j_i - 2x_i J^j_i) \\ &\stackrel{(28)}{=} \frac{4}{\kappa} \int_V \partial_t M \\ &\quad + \int_V \partial_j (|\mathbf{x}|^2 \partial_i J^j_i - 2x_i J^j_i - 2\Gamma^R \pi^j). \end{aligned} \quad (31)$$

Finally, we rewrite the sources by exploiting the Euler-Lagrange equations of the action (7), $C_{ij,kl}^{-1} \sigma_{kl} = w_{ij}$ and $M = \kappa \vartheta_w$, and thus obtain

$$\frac{d}{dt} \int d^2x |\mathbf{x}|^2 \rho^{(\ell,t)} = \frac{d}{dt} S^{(\ell,t)}. \quad (32)$$

The source terms for disclination quadrupoles is

$$S^{(t)} = 2 \int d^2x \partial_i w_i \quad (33)$$

and corresponds to a costly volume change of the system. In turn,

$$S^{(\ell)} = 4 \int d^2x \vartheta_w \quad (34)$$

for discompression quadrupoles corresponds to a global twist. These quadrupoles can easily be interpreted if one considers an isolated dislocation $Q_j(\mathbf{x}) = b_j \delta(\mathbf{x} - \mathbf{r}(t))$ whose $\int d^2x |\mathbf{x}|^2 \rho^{(t)} = 2b^j r_j(t)$ and $\int d^2x |\mathbf{x}|^2 \rho^{(\ell)} = -2b_j r_j(t)$. Thus the quadrupolar moments are the components of the dislocation position parallel and perpendicular to the Burgers vector. Equation (32) was obtained up to boundary terms. The boundary terms vanish or, more physically, cannot be changed by local operations that act in the bulk of the medium. Thus Eq. (32) expresses how in the bulk the sources

of changing quadrupolar moments are the energetically costly compression and disorientation of the crystal.

In our analysis we freely used expressions that are valid only on-shell, such as the conservation laws (16), (21), (25), (32), equations of motion (A8), (26), $C_{ij,kl}^{-1} \sigma_{kl} = \partial_i w_j$, $M = \kappa \vartheta_w$, expressions for the sources $S^{(t)}$ and $S^{(\ell)}$, etc. They can all be formulated as rigorous statements that apply to the respective path-integral averages. This is proved by using the Schwinger-Dyson equations.

From Eq. (32) we may therefore conclude that a climb, i.e., the motion perpendicular to b_j of a dislocation, is accompanied by an energetically costly change in the volume of the crystal. In addition, a glide, i.e., the motion parallel to b_j of a dislocation, is accompanied by an energetically costly change of the orientation of the crystal. While the first result holds in any crystal, the second is due to the nonzero rotational stiffness of incommensurate ones. In its presence the motion of dislocations is forbidden entirely instead of just being restricted to one direction.

F. Sub-diffusive hydrodynamics

Thus far we have analyzed the motion of isolated dipoles and quadrupoles. The situation becomes more complicated for higher-order multipoles. However, using the hydrodynamic description of fractons [44–47] one readily finds that there cannot be a Fick's law where gradients of the vector charges yield currents (i.e., $J_{ij} \propto \partial_i Q_j$ or $\propto \partial^i Q_j$), leading to ordinary diffusion of all charges. If this were true, the mere existence of scalar charges would induce a current $J_{ii} \sim \rho^{(\ell,t)}$, which is not allowed. Expanding in gradients, the leading symmetry-allowed term connects J_{ij} with a third spatial derivative of Q_j . This gives rise to a coupled dynamics of the two scalar densities that form two subdiffusive modes with dispersions $\omega = iB_{\pm} k^4$.

III. GENERALIZATION TO GENERIC PLANAR QUASICRYSTALS AND INCOMMENSURATE CRYSTALS

Finally, we demonstrate that our results are not restricted to TBG and apply directly to planar quasicrystals and incommensurate crystals with two length scales. To this end we relate the elastic gauge theory with rotational term considered in the main text to the elasticity theory of two-dimensional quasicrystals. In these systems, the low-energy elastic variables are doubled compared to periodic crystals [11–13]. In addition to the phonons \mathbf{u} , there are phasons \mathbf{w} that can be shown to have a rotational stiffness like in Eq. (1). The defects of the system are characterized by two Burgers vectors $\mathbf{b}_u = \oint d\mathbf{u}$ and $\mathbf{b}_w = \oint d\mathbf{w}$ [12]. Our results imply that pure displacement dislocations can glide, whereas those that involve \mathbf{b}_w cannot. This remains true even in the presence of phonon-phason coupling.

A symmetry analysis of the problem yields the following result for the elastic energy [11–13]:

$$\mathcal{E}_{\text{el}} = \frac{1}{2} C_{ij,kl} u_{ij} u_{kl} + R_{ij,kl} u_{ij} w_{kl} + \frac{1}{2} K_{ij,kl} w_{ij} w_{kl}, \quad (35)$$

where $u_{ij} = \frac{1}{2}(\partial_i u_j + \partial_j u_i)$ is the usual symmetric strain tensor describing phonons and $w_{ij} = \partial_i w_j$ is the non-symmetric phason tensor.

For the moment, let us focus on the case of a 12-fold symmetric planar quasicrystals that has the special property of no phonon-phason mixing, i.e., $R_{ij,kl} = 0$ [13]. Because of the high rotational symmetry, the elastic constants $C_{ij,kl}$ have the same form as those of isotropic media:

$$C_{ij,kl} = \lambda \delta_{ij} \delta_{kl} + \mu (\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}). \quad (36)$$

On the other hand,

$$K_{ij,kl} = K_1 \delta_{ik} \delta_{jl} + K_2 (\delta_{ij} \delta_{kl} - \delta_{il} \delta_{jk}) + \tilde{K}_3 \epsilon_{ij}^S \epsilon_{kl}^S, \quad (37)$$

where $\epsilon_{11}^S = \epsilon_{22}^S = 0$ and $\epsilon_{12}^S = \epsilon_{21}^S = 1$. We note that $K_{ij,kl}$ satisfies only the major index symmetry $K_{ij,kl} = K_{kl,ij}$, but not the minor ones $K_{ij,kl} \neq K_{ji,kl}$ and $K_{ij,kl} \neq K_{ij,lk}$.

To make contact with our system (3), we merely have to recognize that $K_{ij,kl}$ can be rewritten as

$$K_{ij,kl} = \tilde{C}_{ij,kl} + \frac{\kappa}{4} \epsilon_{ij} \epsilon_{kl}, \quad (38)$$

where

$$\tilde{C}_{ij,kl} = \tilde{\lambda} \delta_{ij} \delta_{kl} + \tilde{\mu} (\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}) + \tilde{K}_3 \epsilon_{ij}^S \epsilon_{kl}^S \quad (39)$$

is the minor-index-symmetric part that satisfies $\tilde{C}_{ij,kl} = \tilde{C}_{ji,kl} = \tilde{C}_{ij,lk}$ and has $\tilde{\lambda} = K_2$ and $\tilde{\mu} = \frac{1}{2}(K_1 - K_2)$. The minor-index-antisymmetric part plays the role of a rotational term with stiffness

$$\kappa = 2(K_1 + K_2). \quad (40)$$

Thus the phonon displacement field u_i behaves like in the elasticity theory of a conventional crystal, while the phason w_i is governed by an elastic energy identical to the one of twisted bilayer graphene, given in Eq. (1) or (3). Hence all our conclusions about the mobility of defects of the phason modes carry over.

The situation is more complicated for quasicrystals with fivefold or eightfold symmetry. In these cases, $C_{ij,kl}$ is still isotropic, as in Eq. (36), and $K_{ij,kl}$ still has the form (37), although with $\tilde{K}_3 = 0$ in the case of fivefold symmetry, and can therefore be decomposed according to Eq. (38). Most significantly, a phonon-phason coupling of the form

$$R_{ij,kl} = R(\delta_{i1} - \delta_{i2})[\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}] \quad (41)$$

is allowed. Consequently, one has to redo the whole gauge formulation.

The end results, however, turn out to be insensitive to phonon-phason coupling. Specifically, for the phonon field one finds the usual results [19–21]:

$$\partial_t \rho_u^{(t)} + \partial^j \partial^i J_{uij}^S = 0, \quad (42)$$

$$\frac{d}{dt} \int d^2x x_k \rho_u^{(t)} = 0, \quad (43)$$

$$\frac{d}{dt} \int d^2x |\mathbf{x}|^2 \rho_u^{(t)} = \frac{d}{dt} S_u^{(t)}, \quad (44)$$

where $Q_{uj} = \partial^i \partial_i u_j^{(s)}$, $J_{uij} = (\partial_i \partial_t - \partial_t \partial_i) u_j^{(s)}$, $\rho_u^{(t)} = \partial^j Q_{uj}$, $J_{uij}^S = \frac{1}{2}(J_{uij} + J_{uji})$, and $S_u^{(t)} = 2 \int d^2x \partial_i u_i$. For the phason fields one similarly finds that Eqs. (16), (21), (25), and (32) continue to hold, but with the appropriate $Q_{wj} = \partial^i \partial_i w_j^{(s)}$, $J_{wij} = (\partial_i \partial_t - \partial_t \partial_i) w_j^{(s)}$, $\rho_w^{(t)} = \partial_j Q_{wj}$, $\rho_w^{(t)} = \partial^j Q_{wj}$, $S_w^{(t)} =$

$2 \int d^2x \partial_i w_i$, and $S_w^{(t)} = 4 \int d^2x \vartheta_w$. Therefore, the glide of defects that have a finite phason Burgers vectors $\mathbf{b}_w = \oint d\mathbf{w}$ is suppressed, and the parameter that controls the suppression is the effective rotational stiffness (40).

Details of the phonon-phason coupled gauge theory: It is convenient to introduce an additional index μ , $\nu \in \{u, w\}$ that differentiates the phonon and phason displacement fields. That way, $u_{ui} = u_i$, $u_{wi} = w_i$, $u_{uij} = u_{ij}$, $u_{wij} = w_{ij}$, and

$$C_{uij,ukl} = C_{ij,kl}, \quad C_{wij,wkl} = K_{ij,kl}, \quad (45)$$

$$C_{uij,wkl} = C_{wkl,uij} = R_{ij,kl}. \quad (46)$$

The appropriate generalization of the action (3) we may now write as

$$S = \frac{1}{2} \int_{x,t,t' \in \mathcal{C}} u_{\mu i}(t) D_{\mu}^{-1}(t-t') u_{\mu i}(t') - \frac{1}{2} \int_{x,t \in \mathcal{C}} C_{\mu ij, \nu kl} u_{\mu ij} u_{\nu kl}. \quad (47)$$

Next, we introduce the Hubbard-Stratonovich fields $\partial_t u_{\mu i} \rightarrow \pi_{\mu i}$ and $C_{\mu ij, \nu kl} u_{\nu kl} \rightarrow \Sigma_{\mu ij}$, integrate out the regular parts of $u_{\mu i}$, and then enforce the constraints $\partial_t \pi_{\mu j} = \partial_t \Sigma_{\mu ij}$ through gauge potentials:

$$\Sigma_{uij} = -\partial_t A_{uij}^S - \partial^i \partial^j \phi_u, \quad (48)$$

$$\pi_{uj} = -\partial_t A_{uj}^S, \quad (48)$$

$$\Sigma_{wij} = -\partial_t A_{wij} - \partial^i \phi_{wj}, \quad (49)$$

$$\pi_{wj} = -\partial_t A_{wj},$$

where $A_{uij}^S = A_{uji}^S$ is symmetric. This yields the action

$$S = -\frac{1}{2} \int_{x,t,t' \in \mathcal{C}} \pi_{\mu i}(t) \Gamma_{\mu}(t-t') \pi_{\mu i}(t') + \frac{1}{2} \int_{x,t \in \mathcal{C}} C_{\mu ij, \nu kl}^{-1} \Sigma_{\mu ij} \Sigma_{\nu kl} + S_{uw}, \quad (50)$$

where $\Gamma_{\mu}(t-t') = \overleftarrow{\partial}_t D_{\mu}(t-t') \overrightarrow{\partial}_{t'}$ and

$$S_{uw} = \int_{x,t \in \mathcal{C}} [A_{uij}^S J_{uij}^S + \phi_u \rho_u^{(t)} + A_{vij} J_{vij} - \phi_{vj} Q_{vj}], \quad (51)$$

with the previously defined charges and current densities.

In this action we see why the phonon-phason coupling does not affect the argument leading to the quadrupolar conservation laws. On the one hand, the charge conservation laws are a consequence of the local gauge symmetry and from the form of the source term (51) we immediately see that they are unaffected by $C_{\mu ij, \nu kl}^{-1}$ or $R_{ij,kl}$. On the other hand, let us take a look at the Euler-Lagrange equations of $A_{\mu ij}$ that allowed for an additional partial integration in Eqs. (30) and (31):

$$\partial_t \tilde{\Sigma}_{\mu ij}(t) = \int dt' \Gamma_{\mu}^R(t-t') \partial_t \pi_{\mu j}(t') - J_{\mu ij}(t). \quad (52)$$

Once one expresses $\tilde{\Sigma}_{\mu ij} = C_{\mu ij, \nu kl}^{-1} \Sigma_{\nu kl}$ in terms of the original fields $u_{\mu ij}$, one finds that $\tilde{\Sigma}_{\mu ij} = u_{\mu ij}$ according to the Euler-Lagrange equations of $\Sigma_{\mu ij}$. Thus $C_{\mu ij, \nu kl}$ again drop out of the argument. Therefore, $C_{\mu ij, \nu kl}$ formally influence

the argument only through the symmetry properties that they entail for $\Sigma_{\mu ij}$. Physically, however, $C_{\mu ij, \nu k \ell}$ also set the energy scales of glide and climb suppressions.

IV. CONCLUSIONS

The dual formulation of elasticity theory that exploits the concept of fractons allows for rather general insights into the mobility of dissipative incommensurate crystals, such as twisted bilayer graphene. In particular, we find that these systems have, in distinction to the usual crystals, completely immobile dislocations in the low defect density limit that become subdiffusive in the high-density limit. Hence, while the electronic properties of graphene can, to a very good approximation, be treated using concepts of periodic crystals [5–8], the incommensurate nature of the material is much more visible in its mechanical properties. In order to estimate whether the effects discussed here are quantitatively relevant, we follow Ref. [9]. Here, κ is shown to be of the order of $10\text{eV/nm}^2 \sin(\Theta/2)$. For twist angles $\Theta \sim 1^\circ$ this yields $\kappa \sim 0.1\text{eV/nm}^2$. Hence, on the length scale of the moiré crystal $\sim 10\text{nm}$, this stiffness is clearly relevant in a wide temperature regime.

Our results are not unique to twisted bilayer graphene. As we demonstrate in Sec. III, they can be generalized to generic planar quasicrystals and incommensurate crystals. The lack of mobility of defects stabilizes networks of defects, such as the soliton network discussed in Ref. [9]. Another implication is that these incommensurate systems should mechanically be rather brittle.

The power of the formalism that led to the results of this work is drawn primarily from the deep intuition that follows from the analogy to electromagnetism, including multipole expansions or the electromagnetism of dissipative media. An interesting open question is how the properties of these defects, and particularly the discompressions, impact the global mechanical properties of TBG, as well as its local electronic spectrum, where stable defect configurations on the electronic spectrum are expected to lead to localized bound state formation. For isolated disclinations and discompressions one might even expect topologically protected bound states in the electronic spectrum [50].

Note added. Recently, we became aware of interesting related works concerned with a fracton description [51] and topologically protected defect motion [52] of quasicrystals.

ACKNOWLEDGMENTS

We thank H. Ochoa for fruitful discussions. J.G. was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project No. SCHM 1031/12-1. G.P. and J.S. were supported by the DFG - TRR 288 - 422213477 Elasto-Q-Mat (Project No. A07). R.M.F. was supported by the US Department of Energy, Office of Science, Basic Energy Sciences, under Award No. DE-SC0020045. R.M.F. also acknowledges a Mercator Fellowship (TRR 288 - 422213477) from the German Research Foundation (DFG).

APPENDIX: SCHWINGER-KELDYSH FORMULATION OF THE ELASTICITY THEORY

In order to include damping of fractons, we perform the analysis on the Keldysh contour [48]. We consider the generating functional:

$$W[h_i] = \int \mathcal{D}w e^{iS[w] - i \int_C h_i(t) w_i(t)}. \quad (\text{A1})$$

Here, $\int_{x,t \in C} \dots = \int_C dt \int_V d^2x \dots$ indicates that the time integration has to be performed on the round-trip Keldysh contour, while the spatial integration goes over the volume V . The action of the problem is given in Eq. (3), where the damping term enters through the nonlocal-in-time contribution

$$D^{-1}(t-t') = -\delta(t-t')\partial_t^2 + \gamma(t-t'), \quad (\text{A2})$$

with the friction selfenergy term $\gamma(t)$. Let $w_i^+(t)$ and $w_i^-(t)$ refer to the phason modes on the upper and lower contour, respectively. Transforming the Keldysh degrees of freedom to quantum and classical fields

$$w_i^{c,q}(t) = \frac{1}{\sqrt{2}}(w_i^+(t) \pm w_i^-(t)), \quad (\text{A3})$$

the self-energy as function of frequency takes the form

$$\gamma(\omega) = \begin{pmatrix} 0 & \gamma^R(\omega) \\ \gamma^A(\omega) & \gamma^K(\omega) \end{pmatrix}. \quad (\text{A4})$$

We assume Ohmic damping

$$\text{Im}\gamma^R(\omega) = \gamma_0 \omega e^{-|\omega|/\omega_c} \quad (\text{A5})$$

with upper cutoff ω_c . The real part of $\gamma^R(\omega)$ is determined via a Kramers-Kronig transformation, where constant terms $\gamma^R(0)$ have to be subtracted. In addition,

$$\gamma^K(\omega) = -2 \coth\left(\frac{\omega}{2T}\right) \text{Im}\gamma^R(\omega) \quad (\text{A6})$$

follows from the fluctuation-dissipation theorem. Hence, the retarded Fourier transform is at low energies given by

$$D^R(\omega) = \frac{1}{\omega^2 + i\gamma_0\omega}. \quad (\text{A7})$$

At low frequencies, the damping term is the dominant one. The equations of motion for the displacement fields w_i are

$$\begin{aligned} \frac{\partial^2 w_j}{\partial t^2} &= \int dt' \gamma^R(t-t') w_j(t') \\ &+ \partial_i [C_{ij, k\ell} w_{k\ell} + \frac{1}{2} \epsilon_{ij\kappa} \vartheta_w]. \end{aligned} \quad (\text{A8})$$

For strong damping and long times $\int dt' \gamma^R(t-t') \rightarrow -\gamma_0 \partial_t$, and the dynamics is diffusive. When we perform the Hubbard-Stratonovich transformations, we obtain Eq. (7) with

$$\Gamma(t-t') = \overleftarrow{\partial}_t D(t-t') \overrightarrow{\partial}_{t'}. \quad (\text{A9})$$

The retarded version of Γ becomes

$$\Gamma^R(\omega) = \frac{\omega^2}{\omega^2 + i\gamma_0\omega}. \quad (\text{A10})$$

In the limit $\gamma_0 \rightarrow 0$, $\Gamma^R(\omega) \rightarrow 1$.

- [1] Y. Cao, V. Fatemi, A. Demir, S. Fang, S. L. Tomarken, J. Y. Luo, J. D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras, R. C. Ashoori, and P. Jarillo-Herrero, Correlated insulator behaviour at half-filling in magic-angle graphene superlattices, *Nature (London)* **556**, 80 (2018).
- [2] Y. Cao, V. Fatemi, S. Fang, K. Watanabe, T. Taniguchi, E. Kaxiras, and P. Jarillo-Herrero, Unconventional superconductivity in magic-angle graphene superlattices, *Nature (London)* **556**, 43 (2018).
- [3] L. Balents, C. R. Dean, D. K. Efetov, and A. Young, Superconductivity and strong correlations in moiré flat bands, *Nat. Phys.* **16**, 725 (2020).
- [4] E. Y. Andrei and A. H. MacDonald, Graphene bilayers with a twist, *Nat. Mater.* **19**, 1265 (2020).
- [5] J. M. B. Lopes dos Santos, N. M. R. Peres, and A. H. Castro Neto, Graphene Bilayer with a Twist: Electronic Structure, *Phys. Rev. Lett.* **99**, 256802 (2007).
- [6] G. Trambly de Laissardière, D. Mayou, and L. Magaud, Localization of Dirac electrons in rotated graphene bilayers, *Nano Lett.* **10**, 804 (2010).
- [7] E. Suárez Morell, J. D. Correa, P. Vargas, M. Pacheco, and Z. Barticevic, Flat bands in slightly twisted bilayer graphene: Tight-binding calculations, *Phys. Rev. B* **82**, 121407(R) (2010).
- [8] R. Bistritzer and A. H. MacDonald, Moiré bands in twisted double-layer graphene, *Proc. Natl. Acad. Sci. USA* **108**, 12233 (2011).
- [9] H. Ochoa, Moiré-pattern fluctuations and electron-phonon coupling in twisted bilayer graphene, *Phys. Rev. B* **100**, 155426 (2019).
- [10] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, 2000).
- [11] D. Levine, T. C. Lubensky, S. Ostlund, S. Ramaswamy, P. J. Steinhardt, and J. Toner, Elasticity and Dislocations in Pentagonal and Icosahedral Quasicrystals, *Phys. Rev. Lett.* **54**, 1520 (1985).
- [12] P. De and R. A. Pelcovits, Linear elasticity theory of pentagonal quasicrystals, *Phys. Rev. B* **35**, 8609 (1987).
- [13] D-h. Ding, W. Yang, C. Hu, and R. Wang, Generalized elasticity theory of quasicrystals, *Phys. Rev. B* **48**, 7003 (1993).
- [14] B. Lian, Z. Wang, and B. A. Bernevig, Twisted Bilayer Graphene: A Phonon-Driven Superconductor, *Phys. Rev. Lett.* **122**, 257002 (2019).
- [15] R. M. Fernandes and J. W. F. Venderbos, Nematicity with a twist: Rotational symmetry breaking in a moiré superlattice, *Sci. Adv.* **6**, eaba8834 (2020).
- [16] T. C. Lubensky, S. Ramaswamy, and J. Toner, Hydrodynamics of icosahedral quasicrystals, *Phys. Rev. B* **32**, 7444 (1985).
- [17] M. Baggioli and M. Landry, Effective field theory for quasicrystals and phasons dynamics, *SciPost Phys.* **9**, 062 (2020).
- [18] H. Ochoa and R. M. Fernandes (unpublished).
- [19] M. Pretko and L. Radzihovsky, Fracton-Elasticity Duality, *Phys. Rev. Lett.* **120**, 195301 (2018).
- [20] S. Pai and M. Pretko, Fractonic line excitations: An inroad from three-dimensional elasticity theory, *Phys. Rev. B* **97**, 235102 (2018).
- [21] M. Pretko, Z. Zhai, and L. Radzihovsky, Crystal-to-fracton tensor gauge theory dualities, *Phys. Rev. B* **100**, 134113 (2019).
- [22] O. V. Yazyev and S. G. Louie, Topological defects in graphene: Dislocations and grain boundaries, *Phys. Rev. B* **81**, 195420 (2010).
- [23] J. H. Warner, E. R. Margine, M. Mukai, A. W. Robertson, F. Giustino, and A. I. Kirkland, Dislocation-Driven Deformations in Graphene, *Science* **337**, 209 (2012).
- [24] B. Butz, C. Dolle, F. Niekkel, K. Weber, D. Waldmann, H. B. Weber, B. Meyer, and E. Spiecker, Dislocations in bilayer graphene, *Nature (London)* **505**, 533 (2014).
- [25] A. Uri, S. Grover, Y. Cao, J. A. Crosse, K. Bagani, D. Rodan-Legrain, Y. Myasoedov, K. Watanabe, T. Taniguchi, P. Moon *et al.*, Mapping the twist-angle disorder and Landau levels in magic-angle graphene, *Nature (London)* **581**, 47 (2020).
- [26] N. P. Kazmierczak, M. Van Winkle, C. Ophus, K. C. Bustillo, S. Carr, H. G. Brown, J. Ciston, T. Taniguchi, K. Watanabe, and D. K. Bediako, Strain fields in twisted bilayer graphene, *Nat. Mater.* **20**, 956 (2021).
- [27] D. Nelson and B. I. Halperin, Dislocation-mediated melting in two dimensions, *Phys. Rev. B* **19**, 2457 (1979).
- [28] A. P. Young, Melting and the vector Coulomb gas in two dimensions, *Phys. Rev. B* **19**, 1855 (1979).
- [29] H. Kleinert, *Gauge fields in Condensed Matter Physics, Stresses and Defects, Differential Geometry, Crystal Defects* (World Scientific, Singapore, 1989).
- [30] J. Zaanen, Z. Nussinov, and S. I. Mukhin, Duality in 2 + 1D quantum elasticity: superconductivity and quantum nematic order, *Ann. Phys.* **310**, 181 (2004).
- [31] V. Cvetković, Z. Nussinov, and J. Zaanen, Topological kinematic constraints: dislocations and the glide principle, *Philos. Mag.* **86**, 2995 (2007).
- [32] H. Kleinert, *Multivalued Fields In Condensed Matter, Electromagnetism, and Gravitation* (World Scientific, Singapore, 2008).
- [33] J. Bohsung and H.-R. Trebin, Disclinations in Quasicrystals, *Phys. Rev. Lett.* **58**, 1204 (1987); **58**, 2277(E) (1987).
- [34] M. Kleman, Phasons and the plastic deformation of quasicrystals, *Eur. Phys. J. B* **31**, 315 (2003).
- [35] M. Kleman, Defects in quasicrystals, revisited I- flips, approximants, phason defects, [arXiv:1303.5563](https://arxiv.org/abs/1303.5563); Defects in quasicrystals, revisited II- perfect and imperfect dislocations, [arXiv:1303.5773](https://arxiv.org/abs/1303.5773).
- [36] L. Radzihovsky and M. Hermele, Fractons from Vector Gauge Theory, *Phys. Rev. Lett.* **124**, 050402 (2020).
- [37] A. Gromov, Chiral Topological Elasticity and Fracton Order, *Phys. Rev. Lett.* **122**, 076403 (2019).
- [38] A. Gromov and P. Surowka, On duality between Cosserat elasticity and fractons, *SciPost Phys.* **8**, 065 (2020).
- [39] D. X. Nguyen, A. Gromov, and S. Moroz, Fracton-elasticity duality of two-dimensional superfluid vortex crystals: defect interactions and quantum melting, *SciPost Phys.* **9**, 076 (2020).
- [40] R. M. Nandkishore and M. Hermele, Fractons, *Annu. Rev. Condens. Matter Phys.* **10**, 295 (2018).
- [41] C. Chamon, Quantum Glassiness in Strongly Correlated Clean Systems: An Example of Topological Overprotection, *Phys. Rev. Lett.* **94**, 040402 (2005).
- [42] J. Haah, Local stabilizer codes in three dimensions without string logical operators, *Phys. Rev. A* **83**, 042330 (2011).

- [43] M. Pretko, Generalized electromagnetism of subdimensional particles, *Phys. Rev. B* **96**, 035119 (2017).
- [44] A. Gromov, A. Lucas, and R. M. Nandkishore, Fracton hydrodynamics, *Phys. Rev. Research* **2**, 033124 (2020).
- [45] J. Feldmeier, P. Sala, G. De Tomasi, F. Pollmann, and M. Knap, Anomalous Diffusion in Dipole- and Higher-Moment-Conserving Systems, *Phys. Rev. Lett.* **125**, 245303 (2020).
- [46] S. Moudgalya, A. Prem, D. A. Huse, and A. Chan, Spectral statistics in constrained many-body quantum chaotic systems, *Phys. Rev. Research* **3**, 023176 (2021).
- [47] L. Radzihovsky, Quantum Smectic Gauge Theory, *Phys. Rev. Lett.* **125**, 267601 (2020).
- [48] A. Kamenev, *Field Theory of Non-Equilibrium Systems* (Cambridge University Press, Cambridge, 2011).
- [49] B. L. van der Waerden, *Spinoranalyse*, *Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl.* **1929**, 100 (1928).
- [50] C. W. Peterson, T. Li, W. Jiang, T. L. Hughes, and G. Bahl, Trapped fractional charges at bulk defects in topological insulators, *Nature (London)* **589**, 376 (2021).
- [51] P. Surówka, Dual gauge theory formulation of planar quasicrystal elasticity and fractons, *Phys. Rev. B* **103**, L201119 (2021).
- [52] D. V. Else, S.-J. Huang, A. Prem, and A. Gromov, Quantum many-body topology of quasicrystals, [arXiv:2103.13393](https://arxiv.org/abs/2103.13393).