

Anomalous electrodynamics and quantum geometry in the Dirac-Harper model for a graphene bilayer

Abigail Timmel  and E. J. Mele ^{*}

Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA



(Received 23 February 2021; accepted 4 August 2021; published 12 August 2021)

Graphene bilayers with layer antisymmetric strains are studied using the Dirac-Harper model for a pair of single-layer Dirac Hamiltonians coupled by a one-dimensional moiré-periodic interlayer tunneling amplitude. This model hosts low-energy, nearly dispersionless bands near charge neutrality that support anomalous polarizations of its charge multipole distributions. These are analyzed introducing a generalized Berry curvature that encodes the field-induced dynamics of multipole fields allowed in a chiral medium with time-reversal symmetry. The formulation identifies a reciprocity relation between responses to layer-symmetric and layer-antisymmetric in-plane electric fields and reveals momentum-space quantum oscillations produced by a spatial pattern of band inversions on the moiré scale.

DOI: [10.1103/PhysRevB.104.075419](https://doi.org/10.1103/PhysRevB.104.075419)

I. INTRODUCTION

The electrodynamic properties of electrons in crystals can include anomalous responses due to the momentum dependence of internal spin and orbital attributes in Bloch bands. There has been interest in interpreting these as geometric phenomena associated with a quantum metric [1,2]. This concept generalizes to driving fields that vary in space. When a smooth spatial variation lowers the symmetry of the system, it can activate responses that would be forbidden for spatially uniform fields. These spatially dispersive responses are sometimes expressed as responses of a system to the field *gradients*. Artificial structures fabricated by stacking atomically thin layers are of great current interest in this regard, and in these systems the field gradient is replaced by a discretized set of driving fields on separate layers [3–5]. In twisted graphene bilayers discretized coupled-layer responses can greatly exceed spatially dispersive responses found in conventional chiral materials [6].

Here we formulate coupled-layer electrodynamic responses in Bloch bands as quantum geometric quantities. We find that charge multipole distributions in the bilayer have dynamics that are naturally represented by a generalization of the Berry curvature to a *class* of curvature forms [7] in a parameter space which combines position and momentum. This approach generalizes the concept of an anomalous charge velocity induced by Berry curvature to a wider class of multipole electrodynamic responses. We illustrate this for the specific case of a graphene bilayer with a one-dimensional moiré periodicity induced by layer-antisymmetric strain.

The paper is organized as follows. In Sec. II for completeness we briefly review the construction of the Dirac-Harper model for our model system, a shear-strained one-dimensional

moiré structure [8]. In Sec. III we describe features of the low-energy spectrum for this model which reveals a set of weakly dispersing modes with charge peaks that counterpropagate in the moiré as a function of the moiré-transverse crystal momentum. In Sec. IV we characterize multipole densities which associate this counterpropagation with a net polarization of a multipole distribution. In Sec. V we formulate a generalized Berry curvature that describes the anomalous response of these multipole densities to the electric field. In Sec. VI we examine the symmetry constraints of both the multipole susceptibilities (Sec. V) and the static multipole polarizations (Sec. IV) which turn out to be distinct phenomena. In Sec. VII we explain the origin of oscillatory behavior found numerically in calculations of the multipole susceptibilities.

II. DIRAC-HARPER MODEL

The system we study is a strained graphene bilayer in which the stacking is modulated in a single crystallographic direction x [Fig. 1(a)] while retaining short-range lattice periodicity in the orthogonal coordinate y . Here for completeness we summarize the formulation of this problem as a spinorial Harper model [8].

The Hamiltonian for this system includes a spatially varying term $v(x)$ which couples the two layers and a set of kinetic terms \mathbf{t} which couple adjacent cells in the same layer. The conserved crystal momentum k_y allows this system to be separated into a family of one-dimensional problems along the x coordinate where the effects of the y -directed couplings are absorbed into a k_y dependence of the kinetic energy terms. The result is a generalization of the scalar Harper equation to a spinorial Dirac-Harper problem [8]:

$$\mathbf{t}(k)\psi_{n+1} + \mathbf{t}^\dagger(k)\psi_{n-1} + \mathbf{t}_0\psi_n + v(x)\psi_n = \epsilon\psi_n. \quad (1)$$

Here, ψ_n are four component fields with amplitudes on two sublattices (a and b) and two layers (1 and 2) with the ordering $\psi^T = (a_1, b_1, a_2, b_2)$. The integers n in Eq. (1) index the

^{*}mele@physics.upenn.edu




FIG. 1. (a) A section of the shear-strained (AC) moiré structure studied in this paper, which is taken to repeat indefinitely in the x and y directions. Typical moiré lengths are around 900 lattice constants in the x direction. (b) Scalar amplitudes of the three mass terms γ_4 , γ_{14} , and γ_{32} normalized by the interlayer coupling constant t' .

primitive cells of the microscopic graphene lattice along the x direction. The $\mathbf{t}(k)$ are block-diagonal 4×4 matrices that can be derived from a nearest-neighbor tight-binding kinetic energy for the individual graphene layers [9]. Defining the Pauli matrices σ_i and τ_i acting on the sublattice index and layer index, respectively, the products $\sigma_x \tau_0 = \gamma_{15}$ and $\sigma_y \tau_0 = \gamma_{25}$ form a basis for \mathbf{t} .

The interlayer coupling $\mathbf{v}(x) = \mathbf{v}(x + L)$ is a smooth spatial variation through three high-symmetry stacking configurations: *AA* where equivalent sublattices in the two layers align vertically, *AB* where a_1 sublattices eclipse b_2 sublattices, and *BA* where b_1 sublattices eclipse a_2 sublattices. The smoothest interpolation between these three configurations is found using a Fourier series in the lowest one-dimensional G vectors,

$$\mathbf{v}(x) = \sum_G \mathbf{v}_G e^{iGx},$$

with matrix-valued coefficients \mathbf{v}_G . The high-symmetry stacking configurations utilize three γ matrices, $\gamma_4 = \sigma_0 \tau_x$, $\gamma_{14} = \sigma_y \tau_y$, and $\gamma_{32} = \sigma_x \tau_x$, so the series can be reexpressed as

$$\mathbf{v}(x) = \sum_\alpha v_\alpha(x) \gamma_\alpha$$

for scalar $v_\alpha(x)$, where $\gamma_\alpha \in \{\gamma_4, \gamma_{14}, \gamma_{32}\}$. The amplitudes $v_\alpha(x)$ are plotted in Fig. 1(b).

The symmetries of this model depend on the orientation of the modulation direction with respect to the underlying graphene lattice, and this crucial feature is encoded in the exact form of the kinetic terms \mathbf{t} . An important limiting case studied here occurs for modulation along the armchair (AC) direction, which is achieved by imposing a layer-antisymmetric shear strain [Fig. 1(a)]. This produces a chiral structure with the twofold rotational symmetries around three orthogonal axes, C_{2x} , C_{2y} , and C_{2z} , but no mirror symmetries. In contrast, modulation along the zigzag (ZZ) direction requires uniaxial strain and the resulting structure respects mirror symmetries. The presence of these mirror symmetries is a nongeneric feature of the Dirac-Harper (DH) model which occurs only for special orientations of the modulation




FIG. 2. (a) Dirac cones produced by the linearized kinetic energy. (b) Dirac cones split by γ_4 to intersect on a critical ring. (c) Gapping of the critical ring by γ_{14} except at two points in the q_y axis. (d) Full moiré spectrum for the AC structure. Inset: Magnification of the low-energy bands within the gap. Each band is twofold degenerate, and there is a total of four *AA*-peaked states and four *AB/BA*-peaked states.

direction. The kinetic terms are expressed in a low-energy effective theory by taking $\psi_{n\pm 1} \approx \psi_n$ in Eq. (1) and linearizing the combination $\mathbf{t}(k) + \mathbf{t}^\dagger(k) + \mathbf{t}_0$ in small momentum q about a Dirac point. For AC, this linearization gives a kinetic energy operator $\hbar v_F (\gamma_{15} q_x - \gamma_{25} q_y)$ where $v_F = \sqrt{3}at/2$, t is the intralayer coupling constant, and a is the graphene lattice constant. To compare, ZZ is a $\pi/2$ rotation of AC which swaps the roles of γ_{15} and γ_{25} .

III. LOW-ENERGY SPECTRA

Figure 2(d) shows the band structure of the AC structure as a function of k_y . Features of the spectrum can be understood in the long-moiré limit directly from the commutation relations of the kinetic terms with the mass terms $\mathbf{v}(x)$ introduced by the interlayer tunneling [8]. In the absence of interlayer coupling, the spectra consist of a pair of layer-degenerate Dirac cones shown in Fig. 2(a). In a semiclassical picture, modification of this spectrum by local mass terms describes behavior of Bloch electrons localized to specific regions within the moiré. The γ_4 mass term describes same-sublattice hopping between the two layers, characteristic of an *AA*-registered region. This mass term commutes with both kinetic terms, splitting two hybridized Dirac cones in energy so that they intersect on a ring in momentum space [Fig. 2(b)] which we refer to as the critical ring. The mass term γ_{14} describes an asymmetry in the coupling between opposite sublattices on the two layers which occurs at a generic position in the superlattice cell and is strongest in the *AB* and *BA* regions. This term anticommutes with γ_{15} , gapping out the critical ring everywhere except at




FIG. 3. (a) Anomalous drift of the charge peaks in AB/BA -peaked bands as a function of $q_r \equiv k_y t' a / \hbar v_F$. The gap closure occurs at $q_r = 1$. (b) Multipole densities ρ_{14} and ρ_5 at $q_r = 0.52$ tagging the peaks with opposite sign.

two nodal points where the critical ring intersects the q_y axis [Fig. 2(c)].

We can use these local four-band features to analyze the full moiré spectrum by noticing that in the long-moiré limit, the slowly varying stacking allows modes to localize in regions of nearly constant stacking. Referring to Fig. 1(b), the AA region is a location where two out of three mass terms have extrema. Indeed, we can readily identify the width of the region supporting zero-energy states in the full moiré spectrum [Fig. 2(d)] as the peak width of the critical ring in AA . This critical ring occurs at $|q| = \hbar v_F / t' a \equiv q_c$ where t' is the interlayer coupling strength. The gap opened around zero energy with two point closures at $k_y = \pm q_c$ arises from γ_{14} changing its sign in this region.

As shown in the inset of Fig. 2(d), a set of nearly dispersionless states appears within this gap. These modes can be understood as produced by sign changes of the interlayer couplings as a function of lateral position x [8]. One set of modes associated with a sign change of the γ_{14} mass confines charge density to the AA region of the moiré. A second set of modes arises from a sign change of the γ_4 mass term giving a k_y -dependent potential that produces a charge density [Fig. 3(a)] with two peaks that counterpropagate as a function of k_y . These features migrate from the AB and BA regions toward the AA region as k_y approaches the gap closure at $k_y/q_c \equiv q_r = 1$. In a semiclassical description, this counterpropagation could be driven by an electric field $\hbar \dot{k}_y = -eE_y$. This counterpropagation transports no charge along x but *can* be associated with transport of higher order multipole distribution.

IV. MULTIPOLE POLARIZATION

We begin by inspecting multipole distributions that are antisymmetric in x because these quantities tag the counterpropagating peaks [Fig. 3(a)] with opposite signs and associate them with net polarization of a charge-neutral multipole field. The lowest nontrivial multipole distribution that

respects C_{2x} , C_{2y} , and C_{2z} symmetries of the shear-strained AC moiré is an octopole. A multipole distribution that is odd in x can therefore be produced by a quantity that has the local rotational symmetry of a quadrupole in the yz plane. Such an octopolar distribution is already explicitly present in the DH model; the mass term $\hat{\gamma}_{14} \sin(Gx)$ is a product of a quadrupolar operator $\hat{\gamma}_{14}$ with an antisymmetric function of x that distinguishes AB and BA registered regions. The multipole distribution $\rho_{14}(x) = \psi^\dagger(x) \gamma_{14} \psi(x)$ manifests this property in an observable by coupling the quadrupolar γ_{14} to an antisymmetric spatial distribution [Fig. 3(b)]. Similarly, a quadrupole density locally measured by the sublattice- and layer-odd operator $\hat{\gamma}_5 = \sigma_z \tau_z$ describes a local layer and sublattice polarization that is also spatially antisymmetric [Fig. 3(b)]. We refer to transverse polarizations intrinsic to these antisymmetric multipole distributions as anomalous polarizations.

V. ELECTRODYNAMIC RESPONSES

The response of an anomalous multipole polarization to an applied electric field can be expressed as a quantum geometric quantity. In a Bloch band with broken time-reversal or inversion symmetry, an anomalous velocity for charge is produced by the Berry curvature $\langle \partial_{\lambda_\mu} u | \partial_{\lambda_\nu} u \rangle - \text{c.c.}$ To extend this to multipole dynamics, we examine the generalized curvature tensor

$$\omega_m = \left\langle \frac{\partial u}{\partial \lambda_\mu} \middle| M \middle| \frac{\partial u}{\partial \lambda_\nu} \right\rangle - \left\langle \frac{\partial u}{\partial \lambda_\nu} \middle| M \middle| \frac{\partial u}{\partial \lambda_\mu} \right\rangle, \quad (2)$$

where $M \equiv m_{\alpha\beta} \otimes \mathbb{I}_N$ and $m_{\alpha\beta}$ is a local multipole operator weighting the sublattice and layer degrees of freedom. For $m_{\alpha\beta} = \mathbb{I}$, we recover the usual Berry curvature responsible for the anomalous Hall conductance. More generally, ω_m can describe the anomalous dynamics of a charge-neutral multipole distribution. A mathematical discussion of this curvature is given in the Supplemental Material [9].

The parameters λ specify a tangent space for a state in the Bloch band which will determine the association of this curvature with a physical response function. To identify appropriate choices for these parameters for the Dirac-Harper model, we observe that the one-dimensional moiré structure is most naturally described using its two extended coordinates: a momentum coordinate in the y direction and a *spatial* coordinate in the x direction. The two dual coordinates k_x and y are insignificant: zone folding reduces the width of the Brillouin zone G_x by a factor of the inverse moiré length $1/L$, and the y coordinate does not appear in the DH equation. For definiteness we choose $k_x = 0$, promoting the model with symmetries that can be only weakly violated at order $1/L$.

We decompose the extended spatial coordinate by introducing three parameters, each of which controls the origin of its respective mass term in the Hamiltonian. These spatial parameters x_α appear in $\mathbf{v}(x)$ as

$$\mathbf{v}(x) = \sum_{\alpha} v_{\alpha}(x + x_{\alpha}) \gamma_{\alpha}.$$

This parametrization allows us to vary the origin for each mass field individually. Each x_α paired with k_y parameterizes a two-dimensional subspace of the projective Hilbert space. Note

that the ground state lies in the one-dimensional sector at $x_\alpha = 0$ where the mass fields are pinned by the moiré structure. Nonetheless the two-dimensional *tangent space* generated by differential changes ∂_{x_α} , ∂_{k_y} will encode an observable property of these states which we denote

$$K_\alpha^{(M)} = \left\langle \frac{\partial u}{\partial k_y} \middle| M \middle| \frac{\partial u}{\partial x_\alpha} \right\rangle - \text{c.c.} \quad (3)$$

Notice that $K_\alpha^{(M)}$ must have the symmetry of an octopole to respect C_2 symmetry. This is satisfied if m has the symmetry of a z -directed dipole, and we use τ_z as a simple and useful example.

The triad of operators ∂_{x_α} , ∂_{k_y} , τ_z can now be identified with two different response functions. The operator ∂_{k_y} couples the band to a perturbing electric field applied in the y direction, and the operator ∂_{x_α} arises from the observable $\hbar \dot{k}_x^{(\alpha)} = [\partial_{x_\alpha}, H]$ which describes an x -directed force on a component of the wave function coupled to the α th mass term. The coupling to individual mass terms will be explained in more detail in the next paragraph, but for now it is adequate to think of this as a force acting on the component of the charge density. To introduce the third member of the triad, τ_z , we can consider either a layer-antisymmetric electric field or a layer-antisymmetric response. For the former, τ_z enters in the driving term $E \tau_z \hat{y}$, and the force is given by

$$f_{y\alpha} = ieE \left(\sum_{m \neq n} \frac{\langle n | [\frac{\partial}{\partial x_\alpha}, H] | m \rangle \langle m | \tau_z | \frac{\partial n}{\partial k_y} \rangle}{\epsilon_m - \epsilon_n} - \text{c.c.} \right), \quad (4)$$

where the first subscript labels the force by an induced lattice effect, to be described later. For the latter, τ_z weights the observable $i\partial_{x_\alpha}$ with opposite sign in each layer:

$$f_{x\alpha} = ieE \left(\sum_{m \neq n} \frac{\langle n | [\tau_z \frac{\partial}{\partial x_\alpha}, H] | m \rangle \langle m | \frac{\partial n}{\partial k_y} \rangle}{\epsilon_m - \epsilon_n} - \text{c.c.} \right). \quad (5)$$

Both of these can be formally expressed as manifestation of the *same* generalized curvature $K_\alpha^{(\tau_z)}$ in Eq. (3), and thus they are equivalent [9]. Numerical calculation for each mass channel α is plotted as a function of q_r in Fig. 4(b).

Equations (3), (4), and (5) are the main results of this work. These results can be regarded as variants of the constitutive relations that describe natural optical activity in chiral media [3,10–12]. Natural optical activity arises from a correlated electric dipole-quadrupole response driven by an optical field, while the multipole susceptibilities obtained here describe an analogous coupling of two different charge multipoles in the static limit. Each mass channel describes a force acting on the component of the charge density confined by the associated mass term. This is demonstrated by examining the differential changes of the charge density produced by shifts of each x_α , shown explicitly in Fig. 4(c). The AA peaks defined by the γ_{14} mass inversion are shifted by x_{14} (i) and the AB/BA peaks defined by γ_4 inversion are shifted with x_4 (ii). Coupling to x_{32} produces almost no effect (iii) because γ_{32} does not confine charge in the moiré cell via a sign change [8]. The combination of all three x_α produces uniform charge shift (iv), so the sum of K_α over mass channels gives the anomalous susceptibility for polarization of a multipole-weighted density (τ_z in this example).




FIG. 4. (a) Reciprocal systems described by (i) Eq. (4) and (ii) Eq. (5). (b) Curvatures $K_\alpha^{(\tau_z)}$ for each of the mass channels. (c) Change of the charge density from (i) x_{14} , primarily affecting the AA charge peak, (ii) x_4 , primarily affecting the AB/BA peaks, (iii) x_{32} , causing little displacement, and (iv) all the mass terms together, inducing a constant shift.

The equivalence of Eqs. (4) and (5) identifies a reciprocity relation stating that a force induced by a multipole-weighted field is equal in magnitude to the multipole-weighted force induced by a uniform field. For our example of a z -directed dipole τ_z , these responses are driven by a layer-antisymmetric field \vec{E}_{\Leftarrow} and a layer symmetric field \vec{E}_{\Rightarrow} , respectively. The layer-antisymmetric field can be thought of the low-frequency limit of a plane wave polarized in the y direction. For the case of \vec{E}_{\Leftarrow} [Eq. (4)], the response summed over mass channels produces a force that acts to depin charge in the negative x direction as shown in Fig. 4(c) (iv). The E_{\Rightarrow} case [Eq. (5)] corresponds to a similar displacement of charge, but weighted in opposite directions in the two layers.

Displacements of various components of the charge density can produce a counterforce acting on the lattice to restore the moiré to an equilibrium stacking arrangement with respect to the charge. The coupling of lattice strain fields to the applied field must respect the C_2 symmetries, so we can deduce the allowed combinations. Focusing our discussion on uniform layer-antisymmetric displacements, the field \vec{E}_{\Leftarrow} (odd in y and z) couples to displacement along y (also odd in y in z), and the field \vec{E}_{\Rightarrow} (odd only in y) couples to displacement along x (odd in x and z). The subscripts in Eqs. (4) and (5) are now clear: f_y (f_x) induces a layer shear along y (x). Intuitively, the y -directed shear transports the full moiré pattern along the x direction [13] to bring the lattice back in phase with the charge, and the x -directed shear counters the layer antisymmetric distortion of the charge along x induced by the initial

electrodynamic response. We can write a general y -directed electric field $E_y(z)$ as a sum of $E_{\rightarrow} + E_{\leftarrow}$, where $E_{\rightarrow} = [E_y(+z) - E_y(-z)]/2$ and $E_{\leftarrow} = [E_y(+z) + E_y(-z)]/2$. Then the induced layer shear acts at a general angle ϕ in the xy plane, which using reciprocity of the curvature is given by $\phi = \tan^{-1}(f_y/f_x) = \tan^{-1}(E_{\rightarrow}/E_{\leftarrow})$. $\phi = 0$ corresponds to opposing layer displacements in the x direction driven by a uniform electric field in the y direction.

VI. SYMMETRY REQUIREMENTS

The response functions described by K_α are symmetric under time reversal \mathcal{T} : since K_α is imaginary and contains only one momentum coordinate, it accumulates two canceling sign changes under \mathcal{T} . This means that the anomalous response described by K_α does not require \mathcal{T} breaking to be present. In contrast to the usual Berry curvature which vanishes everywhere in a Bloch band that respects both time-reversal and inversion symmetry, the generalized susceptibility studied here vanishes everywhere for systems with time-reversal and *mirror* symmetry. We can gain additional insight using $C_{2x}\mathcal{T}$ symmetry, which is a local symmetry in the parameter space (x, k_y) . The symmetry of $m_{\alpha\beta}$ under this operation determines whether the curvature $K^{(M)}$ or its associated multipole density ρ_m must vanish locally in the mixed momentum-position space. For the curvature, one finds [9]

$$\left\langle \frac{\partial u}{\partial k_y} \middle| M \middle| \frac{\partial u}{\partial x_\alpha} \right\rangle = \left\langle \frac{\partial u}{\partial k_y} \middle| M' \middle| \frac{\partial u}{\partial x_\alpha} \right\rangle^*$$

for $M' = (C_{2x}^\dagger M C_{2x})^*$. When $M' = M$, the curvature is equal to its complex conjugate and therefore it vanishes everywhere. Indeed this is the case for the charge susceptibility $M = \mathbb{I}$, but the generalized extensions can incorporate multipoles such as τ_z where $M' = -M$, allowing this response to be nonzero. For the multipole density ρ_m , the same manipulation gives $\rho_m = \rho_m^*$ where m' is the x -projected version of M' . Since ρ_m is real, this quantity vanishes when $m' = -m$. Thus we find that the symmetry of m under $C_{2x}\mathcal{T}$ divides multipole densities into two distinct classes: those that can have nonvanishing density but vanishing curvature and those that have vanishing density but a nonvanishing curvature. The anomalous response measured by Eq. (3) therefore is physically distinct from the construction obtained by antisymmetrically tagging the counterpropagating peaks shown in Fig. 3(a). Dynamical effects occur in quantities with no bulk accumulation of a multipole density, and quantities with bulk accumulation can only display a static polarization.

For unitary multipole operators [9], the generalized Berry curvature is related to the standard Berry curvature by diffeomorphisms on the Hilbert space. This is a consequence of the Darboux theorem which states that any symplectic two-form can be reexpressed in a standard form on an open set via a coordinate change. Thus, we arrive at an alternative interpretation of the generalized Berry curvature as the standard Berry curvature computed on a different sector of the Hilbert space. As an example, we can put the τ_z curvature into standard form by complex conjugation of the second-layer amplitudes of the wave functions, which has the effect of negating the second-layer component of the complex curvature. The new




FIG. 5. (a) Zero energy crossings of the tight-binding spectrum for the $\nu_4(x)$ potential. Tight-binding wave functions plotted against the effective potential for (b) $k_y = 0.08$ and (c) $k_y = 0.086$. As wave functions from consecutive bands, they differ by one peak.

wave functions do not have obvious physical interpretation, but the standard form of the Berry curvature lends computational and conceptual power. Calculations of the generalized Berry curvature can be very difficult in nonstandard form, but after applying the diffeomorphism, a standard Wilson loop can be utilized [9].

VII. QUANTUM OSCILLATIONS

All three of the curvatures shown in Fig. 4(b) oscillate as a function of k_y . This behavior manifests oscillations in the energies of the AB/BA bands, partially shown in Fig. 2(d), which are encoded in the response function via the velocity terms in the Kubo formula. The observed oscillations have a smoothly evolving period which is nearly constant over the width of the band-inverted region $|q_r| < 1$ in momentum space. We can understand this in a minimal model by examining the low-energy behavior of spatially varying γ_4 potential, shown in Fig. 5.

One might be tempted to interpret the spacing of zero energy crossings as a result of zone folding of the critical ring [Fig. 2(b)] into the first moiré Brillouin zone. Such an effect arises from a constant γ_4 potential, which we can inspect numerically using a tight-binding model. This calculation reveals that a non-negligible $\pm q_x$ anisotropy of the critical ring produces two independently evolving series of zero-energy crossings in the band structure that rapidly converge near the edge of the critical ring $q_r = \pm 1$ [9]. Such behavior is quite different from the slowly modulated period of the oscillations we observe in the τ_z curvature. By contrast, a spatially *varying* γ_4 potential produces a nearly uniform pattern of band crossings [Fig. 5(a)] which closely matches the period of the curvature oscillations. This behavior can be understood by squaring the low-energy Dirac equation,

$$\left[i\hbar v_F \gamma_{15} \frac{\partial}{\partial x} + \hbar v_F k_y \gamma_{25} + \nu_4(x) \gamma_4 \right] \psi = \epsilon \psi,$$

to form the effective potential problem

$$\frac{\partial^2}{\partial x^2} \psi = \left(-\frac{\nu_4(x)^2}{\hbar^2 v_F^2} + k_y^2 \right) \psi \equiv V_{\text{eff}} \psi. \quad (6)$$

The spacing of zero crossings in this situation arises from a resonance condition within a basin where V_{eff} is negative. The wave functions produced by the $\nu_4(x)\gamma_4$ potential

in a tight-binding calculation [Fig. 5(b)] oscillate in the region where V_{eff} is negative and decay in regions where it is positive. Wavelengths of the oscillating part of the wave functions match the value of $\sqrt{V_{\text{eff}}}$ averaged over each period. Consecutive zero energy band crossings differ by a single wave function oscillation within the V_{eff} basin, which is indicative that the spacing of the zero-energy crossings is a consequence of wave function resonance within this effective potential.

These observations about the effective potential produced by a pure ν_4 potential carry over to the full DH Hamiltonian. The mass term proportional to $\gamma_{14} \sin(Gx)$ has the effect of opening a gap around the lowest energy band manifold without changing the spacing of these zero energy crossings [9]. Away from the saddle point configuration in the stacking texture (*SP*), γ_{32} decreases to zero and has only a perturbative effect. Therefore, oscillatory behavior in the energies is inherited from the k_y spacing of zero energy crossings intrinsic to the γ_4 part of the potential. Although the full moiré *AB/BA* wave functions decay within the positive effective potential region, the sensitivity of the energies to these special values of k_y suggests that the wave functions retain a small oscillatory part stretching over the positive potential region flanked by the *AB/BA* charge peaks. Interestingly these oscillations would be absent from chiral models for twisted bilayer graphene which exclude this coupling to enforce a chiral symmetry [14]. Here we see that they control a physically measurable response, and measurement of these oscillations can be

used to determine the scale of the breaking of the chiral symmetry.

VIII. CONCLUSION

The curvature forms developed here can also be applied to two-dimensional twisted bilayers. In this case the DH theory generalizes to a two-dimensional theory in a moiré supercell and the forces driving the multipole distributions analogous to Eqs. (4) and (5) are similarly promoted to two-dimensional vectors obtained by integrating over a population in the 2D folded Brillouin zone. The simplest case couples the driving fields to the total charge density recovering the charge pumping phenomena proposed for twisted graphene bilayers [13,15]. The generalized Berry curvature provides a unified formulation that associates this electromechanical response with the more general problem of manipulating multipole distributions using applied fields. The generalized constitutive relations for electric-field-driven multipole responses also provide a geometric formulation of the natural optical activity of twisted graphene bilayers studied experimentally [6] and theoretically [3–5] and to chiral Weyl semimetals [16].

ACKNOWLEDGMENT

This work was supported by the Department of Energy under Grant No. DE-FG02-84ER45118.

-
- [1] R. Resta, The insulating state of matter: A geometrical theory, *Eur. Phys. J. B* **79**, 121 (2011).
 - [2] T. Neupert, C. Chamon, and C. Mudry, Measuring the quantum geometry of Bloch bands with current noise, *Phys. Rev. B* **87**, 245103 (2013).
 - [3] E. Morell, L. Chico, and L. Brey, Twisting Dirac fermions: Circular dichroism in bilayer graphene, *2D Mater.* **4**, 035015 (2017).
 - [4] T. Stauber, T. Low, and G. Gómez-Santos, Chiral Response of Twisted Bilayer Graphene, *Phys. Rev. Lett.* **120**, 046801 (2018).
 - [5] D. Nguyen and D. Son, Electrodynamics of thin sheets of twisted material, [arXiv:2008.02812](https://arxiv.org/abs/2008.02812).
 - [6] C. Kim, A. Sánchez-Castillo, Z. Ziegler, Y. Ogawa, C. Noguez, and J. Park, Chiral atomically thin films, *Nat. Nanotechnol.* **11**, 520 (2016).
 - [7] J. Provost and G. Vallee, Riemannian structure on manifolds of quantum states, *Commun. Math. Phys.* **76**, 289 (1980).
 - [8] A. Timmel and E. J. Mele, Dirac-Harper Theory for One-Dimensional Moiré Superlattices, *Phys. Rev. Lett.* **125**, 166803 (2020).
 - [9] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.104.075419> for the derivation and computational methods.
 - [10] F. Hidalgo, A. Sánchez-Castillo, and C. Noguez, Efficient first-principles method for calculating the circular dichroism of nanostructures, *Phys. Rev. B* **79**, 075438 (2009).
 - [11] C. Noguez, and F. Hidalgo, *Ab initio* electronic circular dichroism of fullerenes, single-walled carbon nanotubes, and ligand-protected metal nanoparticles, *Chirality* **26**, 553 (2014).
 - [12] L. D. Barron, *Molecular Light Scattering and Optical Activity* (Cambridge University Press, Cambridge, 2009).
 - [13] Y. Zhang, Y. Gao, and D. Xiao, Topological charge pumping in twisted bilayer graphene, *Phys. Rev. B* **101**, 041410(R) (2020).
 - [14] G. Tarnopolsky, A. Kruchov, and A. Vishwanath, Origin of Magic Angles in Twisted Bilayer Graphene, *Phys. Rev. Lett.* **122**, 106405 (2019).
 - [15] M. Fujimoto, H. Koschke, and M. Koshino, Topological charge pumping by a sliding moiré pattern, *Phys. Rev. B* **101**, 041112(R) (2020).
 - [16] M. Kargarian, M. Randeria, and N. Trivedi, Theory of Kerr and Faraday rotations and linear dichroism in topological Weyl semimetals, *Sci. Rep.* **5**, 12683 (2015).