


Numerical transfer matrix study of frustrated next-nearest-neighbor Ising models on square lattices

Yi Hu ^{1,*} and Patrick Charbonneau ^{1,2,†}

¹*Department of Chemistry, Duke University, Durham, North Carolina 27708, USA*

²*Department of Physics, Duke University, Durham, North Carolina 27708, USA*

 (Received 7 July 2021; revised 13 September 2021; accepted 15 October 2021; published 29 October 2021)

Ising models with frustrated next-nearest-neighbor interactions present a rich array of modulated phases. These phases, however, assemble and relax slowly, which hinders their computational study. In two dimensions, strong fluctuations further hamper determining their equilibrium phase behavior from theoretical approximations. The exact numerical transfer matrix (TM) method, which bypasses both difficulties, can serve as a benchmark method once its own numerical challenges are surmounted. Building on our recent study [Hu and Charbonneau, *Phys. Rev. B* **103**, 094441 (2021)], in which we evaluated the two-dimensional axial next-nearest-neighbor Ising model with transfer matrices, we here extend the effective usage of the TM method to Ising models with biaxial, diagonal, and third-nearest-neighbor frustration models. The high-accuracy TM numerics help resolve various physical ambiguities about these reference models, thus providing a clearer overview of modulated phase formation in two dimensions.

DOI: [10.1103/PhysRevB.104.144429](https://doi.org/10.1103/PhysRevB.104.144429)

I. INTRODUCTION

A ferromagnetic Ising model frustrated by next-nearest-neighbor antiferromagnetic interactions offers a minimal description of systems with short-range attractive and long-range repulsive (SALR) interactions [1,2]. Because even such a simple model family leads a rich set of modulated morphologies—depending on the strength, length scale, and orientation of the frustration—it has been used to recapitulate the material behavior of systems as diverse as magnetic ordering in metals and superconductors [3,4] and microphase formation in surfactants [5,6].

From a statistical physics standpoint, two-dimensional versions of these models are especially interesting. Strong thermal fluctuations alter the nature of phase transitions and lead to novel equilibrium behaviors, such as floating incommensurability and Kosterlitz-Thouless (KT)-type criticality [7,8]. These models are thus often deemed to be schematic descriptions of experimental systems such as magnetic monolayers, and serve as reference for richer models that more closely capture these solid-state systems [9]. Certain features of these models, however, are challenging to capture in theoretical and numerical studies. As a result, long-standing debates persist, including about the putative existence of the critical incommensurate (IC) phase in the axial and biaxial next-nearest-neighbor Ising (ANNNI and BNNNI, respectively) models [10] as well as the order of the phase transition in the diagonal nearest-neighbor Ising (DNNI) model [11]. Over the last decade, significant advances have been made in surmounting some of the underlying technical difficulties. For

the aforementioned ambiguities in particular, recent studies have confirmed the existence of the IC phase in the ANNNI model [12–14] as well as the order of the transition and the location of the Potts critical point in the DNNI model [15,16]. Yet both qualitative and quantitative uncertainties persist and others emerge in light of these advances (see, for instance, Refs. [17–19]). A benchmark method that provides results that are exact or with well-controlled limitations for target systems would thus be particularly helpful to make complete physical sense of these models.

In this context, the use of numerical transfer matrices (TMs), which provide high-accuracy numerical solutions, appears enticing. In fact, the idea is not new. TMs were used on frustrated Ising models starting in the 1980s [20–22] but then struggled to provide qualitative—let alone quantitative—insight. The challenge is that only strips of finite width L can be solved with TM, hence a careful finite-size scaling analysis must also be part of the thermodynamic extrapolation, in the limit $L \rightarrow \infty$. Given that the algorithmic complexity of TM method grows exponentially with L and that frustrated models exhibit large preasymptotic corrections, limited computational resources then resulted in physical obfuscation. Exponential improvement to computational hardware over the years (following Moore’s Law) coupled with more efficient eigensolvers [23] offer hope that the situation might have since improved. For instance, TMs now provide definitive solutions of even fairly complex (quasi-)one-dimensional continuum-space systems [24–28].

In a recent paper [14], we reported the TM resolution of various long-standing ambiguities about the two-dimensional ANNNI model at a reasonable computational cost, thanks to the combination of various algorithmic optimizations. Because this approach is sufficiently generic to be adapted to related lattice models, this paper revisits a series of frus-

*yi.hu@duke.edu

†patrick.charbonneau@duke.edu

trated lattice models: the ANNNI, DNNI, and BNNNI models as well as the generic third-nearest-neighbor Ising (3NNI) model. We notably resolve long-standing ambiguities surrounding the DNNI model at intermediate frustration and build quantitative phase diagrams for the BNNNI and 3NNI models. The rest of this paper is organized as follows. In Sec. II, we provide a complete description of these models, review their key properties, and discuss some of the remaining phase ambiguities before introducing the TM approach. Section III presents TM results for these models. A brief conclusion follows in Sec. IV.

II. MODELS AND METHODS

In this section, we describe the various next-nearest-neighbor Ising models on a square lattice considered in this paper and highlight some of the existing results and predictions. We also briefly describe the numerical TM method. Additional technical details can be found in Appendixes A and B.

A. ANNNI model

The ANNNI model is a minimal model for lamellar microphase formers. Its Hamiltonian reads

$$\mathcal{H}_{\text{ANNNI}} = -J \sum_{\langle i,j \rangle} s_i s_j + \kappa J \sum_{\langle i,j \rangle_{\text{ANNN}}} s_i s_j - h \sum_i s_i, \quad (1)$$

for spin variables $s_i = \pm 1$, coupling constant $J > 0$, next-nearest-neighbor frustration strength along the axial direction $\kappa > 0$, and an external field h . (For all models considered here, $J = 1$ sets the unit of energy, and we set $h = 0$.) For $\kappa = 0$, this model reduces to the standard ferromagnetic Ising model, and for $\kappa < 1/2$, the order-disorder transition remains part of the Ising universality class. For $\kappa > 1/2$, the antiphase (2) (of period 4, $\uparrow\uparrow\downarrow\downarrow$), which forms the energetic ground state, melts into a critical IC phase at T_{c2} and then becomes fully disordered at T_{c1} . Although the existence of the IC phase has long been debated [10], recent studies provide clear evidence of its persistence, i.e., $T_{c1} > T_{c2}$ for all $\kappa > 1/2$ [12–14].

B. DNNI model

The DNNI model modifies the Ising model by including nonaxial nearest-neighbor interactions. Its Hamiltonian reads

$$\mathcal{H}_{\text{DNNI}} = -J \sum_{\langle i,j \rangle} s_i s_j + \kappa J \sum_{\langle i,j \rangle_{\text{DNN}}} s_i s_j - h \sum_i s_i, \quad (2)$$

using the same parameter notation as for the ANNNI model. (This model is also known as the next-nearest-neighbor or frustrated-Ising model and as the J_1 - J_2 Ising model, but we here denote it DNNI to distinguish it more saliently from the other models considered.) The energetic ground state of the DNNI model is ferromagnetic for $\kappa < 1/2$ and striped with $\langle 1 \rangle$ (of period 2, $\uparrow\downarrow$) for $\kappa > 1/2$. In both cases, melting leads directly to a disordered phase but whether or not the transition remains first order in nature around $\kappa = 1/2$ has long been debated. The putative first-order regime indeed appears to narrow as the quality of numerical studies improves. Predictions for a Potts point dividing the weakly first order

from the continuous transition vary from $\kappa^* \simeq 0.9$ [29,30], 0.67(1) [15,16] to $\simeq 0.54$ [19]. For $\kappa < 1/2$, it has generally been assumed that the transition is part of the Ising universality class, but recent cluster mean-field approximation and effective-field theory studies suggest that a narrow first-order transition regime might then also be present [16,17]. Several simulation studies further suggest a nonzero transition temperature at $\kappa = 1/2$ proper, i.e., $T_c(\kappa = 1/2) > 0$ [18,31,32], in marked contrast to prior works [11,33–36]. Because theoretical approximations behave irregularly around $\kappa = 1/2$ [37], however, a conclusive assessment has thus far remained out of reach.

C. BNNNI and 3NNI models

Including biaxial next-nearest-neighbor interactions, i.e., couplings with Euclidean third-nearest-neighbors, to the DNNI model gives rise to the 3NNI model (also known as the J_1 - J_2 - J_3 Ising model [38]). Its Hamiltonian reads

$$\mathcal{H}_{3\text{NNI}} = -J \sum_{\langle i,j \rangle} s_i s_j + \kappa J \sum_{\langle i,j \rangle_{\text{BNNN}}} s_i s_j + \kappa' J \sum_{\langle i,j \rangle_{\text{DNN}}} s_i s_j - h \sum_i s_i, \quad (3)$$

and setting $\kappa' = 0$ recovers the BNNNI model, which we consider first.

The BNNNI model has a ferromagnetic ground state for $\kappa < 1/2$, and the transition to the high-temperature paramagnetic phase is thought to exhibit Ising universality in that regime. For $\kappa > 1/2$, the model presents two distinct energetic ground states [Fig. 1(d)]: 4×4 checkerboard order, or diagonal stripes of width 2. Early Monte Carlo simulations suggested that melting of these structures proceeds through a first-order transition [39] but later work found the thermodynamics not to be so straightforward. Multiple metastable states indeed develop at intermediate temperatures [40], and a two-step transition involving a critical IC phase at $1/2 < \kappa < \kappa^*$ [22,41–43] has been proposed. It is further unclear whether the Lifshitz point takes place at finite κ^* or whether $\kappa^* \rightarrow \infty$. Even studies suggesting the former offer but a qualitative determination of κ^* [42].

Landau and Binder determined the energetic ground-state structure of the more general 3NNI model with antiferromagnetic Ising interactions [39] [see Fig. 1(d)], which can be mapped onto ferromagnetic Ising interactions by flipping every other spin on the lattice. In short, while the ferromagnetic (F) phase persists for $\kappa + \kappa' < 1/2$, the ground state either follows that of the BNNNI model— 4×4 checkerboard or diagonal stripes of width 2 (both denoted 4×4 , for convenience)—at large κ , or that of the DNNI model— $\langle 1 \rangle$ stripes—at large κ' . At large frustration, the separation line is given by $\kappa' = 2\kappa$ and at intermediate frustration, the $\langle 2 \rangle$ phase is also a ground state. By contrast to the ANNNI model, however, the modulation can here grow along either axial direction (on a square lattice), hence the ground state is eightfold (instead of fourfold) degenerate. The transition is further thought to be first order—as for the eight-state Potts model—instead of continuous [38].

Interestingly, the lattice gas representation of the 3NNI model is also the two-dimensional counterpart to the Widom-




FIG. 1. Schematics of the (a) ANNNI, (b) DNNI, and (c) 3NNI models on a square lattice. Arrows denote spins and lines indicate Ising nearest-neighbor (red), diagonal NN (green), and (bi)axial NNN (blue) interactions. (d) $T = 0$ phase diagram for the 3NNI model.

Wheeler lattice microemulsion model [6] (with h corresponding to the chemical potential). The 3NNI model presents a (2) phase—a lamellar microphase in the language of Ref. [6]—as a result of the competition between DNN and BNNN interactions. Our investigation of the 3NNI model thus concentrates on this particular regime.

D. Numerical TM method

Although these two-dimensional models lack an analytic solution, TMs can numerically solve the exact partition function of a semi-infinite strip of width L , and the results can then be analyzed using finite-size scaling approaches to extrapolate the thermodynamic behavior in the limit $L \rightarrow \infty$.

Generically, TMs encode the interaction between subsequent layers states a and a' as

$$\mathbf{T}_{a,a'} = \exp[-\beta(V_x(a) + V_z(a, a'))], \quad (4)$$

where $\beta = 1/k_B T$ is the inverse temperature (the Boltzmann constant is set to unity, $k_B = 1$) and V_x and V_z are intra- and interlayer interaction energies, respectively. The partition function of the strip of length N is given by $\text{tr}(\mathbf{T}^N)$, and in the limit of $N \rightarrow \infty$, Z is given by the leading eigenvalue of \mathbf{T}^N , λ_0^N . The free energy per spin is thus

$$\beta f = -\frac{\ln \lambda_0}{L}, \quad (5)$$

and the marginal probability of a state a is given by the product of (normalized) left and right eigenvector of λ_0 , $P(a) = \varphi_0^{-1}(a)\varphi_0(a)$ (after normalizing as $\sum_a P(a) = 1$). Given the leading eigenvalue and eigenvector, thermodynamic observables can be obtained exactly, including the internal energy per spin, $\beta u = -\partial(\beta f)/\partial T$, and the specific heat per spin, $c = \partial u/\partial T$.

One of the key advantages of TMs is that they also provide the correlation length ξ , which help identify the location and nature of phase transitions. Given that the conditional probability of finding a subsequent layer state a' after layer a is [28]

$$P(a'|a) = \frac{\mathbf{T}_{a,a'}\varphi_0(a')}{\lambda_0\varphi_0(a)}, \quad (6)$$

the generic conditional probability for a' having a distance N from a is then

$$P^{(N)}(a'|a) = \frac{\mathbf{T}_{a,a'}^N \varphi_0(a')}{\lambda_0^N \varphi_0(a)} = P(a') + \sum_{i=1}^{|\{a\}|-1} \left(\frac{\lambda_i}{\lambda_0}\right)^N \frac{\varphi_i(a)\varphi_i^{-1}(a')\varphi_0(a')}{\varphi_0(a)},$$

$$P^{(N)}(a'|a) - P(a') \sim \exp\left[N \ln\left(\frac{|\lambda_1|}{\lambda_0}\right)\right] = \exp(-N/\xi_1), \quad (7)$$

where $|\{a\}|$ is the total number of states and $\xi_1 = -1/\ln(|\lambda_1|/\lambda_0)$ is the leading correlation length. Subdominant lengths can be analogously defined as $\xi_i = -1/\ln(|\lambda_i|/\lambda_0)$ for $i > 1$.

Beyond that point, each model presents certain peculiarities. (TM prescripts are thus used to distinguish between the ANNNI (A), BNNNI (B), DNNI (D), and 3NNI (3) models as well as to denote the direction of propagation.) For the ANNNI model, because the interaction is anisotropic, the TM can be propagated either perpendicularly (\perp, A TM) [20] or parallel (\parallel, A TM) [21] to the direction of the next-nearest-neighbor interaction. Although the former is markedly smaller $2^L \times 2^L$ (vs $4^L \times 4^L$), in the modulated regime the latter converges much faster to the asymptotic scaling as L increases. For the DNNI model, because interactions reach no further than the first subsequent layer, \perp, D TM can be constructed by modifying \perp, A TM and is thus also of size $2^L \times 2^L$. For the BNNNI and 3NNI models, TM propagation along the diagonal of the square lattice, \diagup TM, has been suggested as preferable at finite L [22] to better capture the ordering in the direction of the antiphase modulation. This choice also results in a TM of size $2^L \times 2^L$. In all cases, the TM size can be significantly reduced by leveraging the model symmetry, as described in Appendix B. (In practice, the TM is not explicitly computed but implicitly represented by a matrix-vector subroutine, as described in Appendix A.)

III. RESULTS AND DISCUSSION

In this section, we present TM results for various frustrated models at finite L as well as the finite-size scaling analysis




FIG. 2. Leading correlation length for the ANNNI model with $\kappa = 0.6$ in (a) \perp TM, $L = 8, 12, \dots, 24$ and (b) \parallel TM, $L = 8, 10, \dots, 16$ (from blue to red lines). The ordering transition position, $T_{c2} = 0.89(1)$ (dashed line) is given as reference. Results from \parallel TM converge more smoothly and cleanly to the thermodynamic limit. Inset: First few leading correlation lengths in (a) \perp TM for $L = 24$ and (b) \parallel TM for $L = 16$. Eigenvalue crossings in the former lead to a complex T dependence of the corresponding ξ_1 .

used to extrapolate the thermodynamic behavior in the limit $L \rightarrow \infty$. We provide results for the ANNNI model that complement our recent analysis of that system [14] and discuss the behavior of the DNNI model, paying particular attention to the physical ambiguities previously reported in the literature. For the more computationally challenging BNNNI model, we propose a phase diagram and discuss various remaining uncertainties. Finally, we examine the $\langle 2 \rangle$ regime of the generic 3NNI model with $\kappa = \kappa'/2$, which are frustration conditions akin to those of a three-dimensional surfactant model.

A. ANNNI model

For the ANNNI model, we first obtain the subleading correlation length using both \perp TM and \parallel TM. These results are mainly intended to serve as a reference for other models. To probe further the existence of a critical IC (or floating) phase, we also compare the domain-wall free energy, using a scheme first proposed for the density matrix renormalization group (DMRG) approach [44].

Comparing the leading correlation length, ξ_1 , obtained from \perp TM and \parallel TM highlights the anisotropic nature of the model (Fig. 2). In particular, as T increases, results from \perp TM decay nonsmoothly due to the crossing of subdominant correlation lengths [Fig. 2(a), inset]. This phenomenon, which

generally accompanies a structural crossover [27], here bespeaks a stepwise change in the modulation period [14]. For $L = 24$, for example, two distinct steps can be identified, both involving $\xi_1 = \xi_2$ (associated with doubly degenerate eigenvalues) crossing the subleading ξ_3 and ξ_4 . These features, however, complicate the evolution of ξ_1 , and hence hinder the local exponent analysis and the determination of the IC phase [14,21,22]. By contrast, results for \parallel TM evolve smoothly with T . The nonmonotonic growth of $\xi_1 = \xi_2$ (associated with complex conjugate eigenvalues) at intermediate T can thus be construed as a signature of the critical IC phase, and its boundaries, $T_{c1} > T_{c2}$, can be identified by analyzing the local exponent [14]. In addition, the subleading correlation lengths ξ_3 and ξ_4 are found to merge at the $\langle 2 \rangle$ -to-IC phase transition temperature, $T_{c2}(L)$, hence providing an additional (and consistent) estimate of T_{c2} to those from Ref. [14].

The stepwise change in the modulation period can also be identified from the domain-wall free energy obtained from comparing two systems under different boundary conditions [45],

$$\tau_L^{(+/-)}(T) = L[f_L^{(+/-)}(T) - f_L^{(+/+)}(T)]\sigma_L, \quad (8)$$

where $f_L^{(+/-)}$ (or $f_L^{(+/+)}$) are the free energy of a system fixing $s_1 = +1$ and $s_L = -1$ (or $+1$), respectively, and $\sigma_L = \pm 1$ in that the ground state $\tau_L^{(+/-)}$ is positive. In the $\langle 2 \rangle$ phase, and for modulations congruent with $L \bmod 4 \equiv 2$, the expected finite-size scaling is [45]

$$\tau_L^{(+/-)}(T) - \tau_L^{(+/-)}(T) \sim L^{-B}, \quad (9)$$

where the exponent is observed to be $B = 2$ in the $\langle 2 \rangle$ phase [45], but preasymptotic corrections grow upon approaching the critical temperature, T_c , at which scaling theory gives $B = 1$ [46].

Following Ref. [44], we first fit the results with $B = 2$ and extrapolate the thermodynamic $\tau^{(+/-)}(T)$ over a broad range of $0.6 < T \lesssim T_{c2}$ (Fig. 3). The result is fully consistent with that former study, predicting $T_{c2} = 0.91(1)$ (vs 0.907 [44]). Alternatively, setting $B = 1$ (the expected critical scaling) gives $T_{c2} = 0.89(1)$. The two estimates thus differ only marginally and are both consistent with recent quantitative estimates for the transition temperature [13,14].

For $T > T_{c2}$, $\tau_L^{(+/-)}(T)$ oscillates around 0, suggesting that the modulation period varies with T , a clear signature of the IC phase. Extrapolating the first peak and valley further suggests that these oscillations coalesce at T_{c2} and thus vanish in the thermodynamic limit $L \rightarrow \infty$. The zigzagging behavior weakens at larger κ yet the smoother oscillations around 0 are still observable at large L for $\kappa \gtrsim 1.5$. Such oscillatory yet vanishing free-energy difference for the $(+/+)$ and $(+/-)$ boundary conditions further supports the floating IC phase scenario. The interfacial results are therefore fully consistent with the length scale analysis of Ref. [14], and are also consistent with the DMRG results [44] obtained for much larger systems ($L \sim 10^2$ vs 10^1 in TMs). However, the interfacial results do not exhibit a clear high temperature boundary for the IC phase, let alone provide a quantitative estimate of T_{c1} .




FIG. 3. Domain-wall free energy $\tau_L^{(+/-)}(T)$ for the ANNNI model with $\kappa = 0.6$ from \perp TM with $L = 14, 18, \dots, 28$ (from blue to red lines). The extrapolation of $\tau^{(+/-)}(T)$ from Eq. (9) with the scaling expected for the antiphase, $B = 2$ (black solid line), vanishes at $T = 0.91(1)$. The extrapolation with the scaling expected for the critical point, $B = 1$ (dashed black line), gives $T_{c2} = 0.89(1)$. Both are numerically consistent with previous estimates [14]. Inset: Extrapolating the positions of local peaks (crosses), valleys (squares), and first zeros (circles) are consistent with T_{c2} (dotted lines are guides for the eye).

B. DNNI model

For the DNNI model, we first compare the correlation length, internal energy and domain wall free energy results with those of the ANNNI model from Ref. [14] and Sec. III A. We then attempt to resolve the phase ambiguities described in Sec. II B.

1. Overview of thermodynamic observables

The evolution of ξ_1 for the DNNI model is smooth and monotonic [see, e.g., Fig. 4(a)]; no eigenvalue crossing or splitting is observed. Unlike the ANNNI model, which presents an algebraic growth, $\xi_1 \sim L^\theta(T)$, over a temperature range [14,21], the DNNI model displays an algebraic scaling only at a single temperature. Also, the anisotropy exponent

is then $\theta = 1$ ($\xi_1/L \approx \text{const}$), as expected for models with isotropic interactions [47]. The crossing point of ξ_1/L further provides an accurate estimate of T_c . Other robust estimators include the crossing point of the internal energy (examined for the ANNNI model for $0 \leq \kappa < 0.5$ [14]) and the peak of the specific heat [Fig. 4(b)]. Because the finite- L transition temperature estimates, $T^*(L)$, change little with L throughout different phase regimes, transition temperatures can often be estimated with up to five significant digits [16]. (Around $\kappa = 1/2$, the situation is more complex, as discussed below.)

We also evaluate the domain-wall free energy using Eq. (8) [Fig. 4(c)]. As expected, $\tau_L^{(+/-)}$ follows the finite-size scaling of Eq. (9) in the ordered striped phase. For $B = 1$, it is extrapolated to vanish at T_c . (The ~ 0.01 deviation from T_c is likely due to preasymptotic corrections to scaling.) Unlike in Fig. 3, here no oscillation is observed for $\tau_L^{(+/-)}(T > T_c)$. This monotonic evolution is robust for various κ both below and above $1/2$. The DNNI model thus clearly lacks an incommensurate phase, or its finite L echo.

2. T_c determination

With an understanding of the behavior for various phase transition estimators in hand, we now quantitatively evaluate T_c . In particular, we wish to determine whether $\lim_{\epsilon \rightarrow 0} T_c(1/2 + \epsilon) = 0$. As shown in Figs. 5(a) and 5(c), a crossing point in the T - u curve is detected for both sides of $\kappa = 1/2$, but is absent right at $\kappa = 1/2$ down to numerical accuracy (in practice, $\sim 10^{-10}$) [Fig. 5(b)]. The crossing thus seemingly takes place at $T = 0$, as does the phase transition, but further evidence is needed.

A systematic comparison reveals that for $\epsilon > 0$ $T_u^*(L)$ barely shifts with L , while for $\epsilon < 0$ significant preasymptotic corrections appear [see Figs. 5(a) and 5(c)]. We thus apply an empirical fitting form to extrapolate T_c ,

$$T^*(L) = T_c + AL^{-t}, \quad (10)$$

with a fitting constant A and empirical exponent t ($t = 1$ for $\kappa > 1/2$). Note that other estimators, such as the location of specific heat and the correlation length peaks [see Fig. 7(c)], $T_c^*(L)$ and $T_\xi^*(L)$, provide consistent estimates [Fig. 5(d)] but




FIG. 4. Finite-size results for various observables of the DNNI model with $\kappa = 0.6$ for $L = 8, 12, \dots, 32$ (from blue to red lines). Recall that $T_c = 0.971(1)$ (dashed vertical lines). (a) The leading correlation length crosses at a single fixed point for all L . Inset: The first few leading correlation lengths for $L = 24$. No correlation length splitting or crossing is observed. (b) Energy per spin u and (inset) specific heat c per spin. Both ξ_1/L and u present a fixed point and c peaks sharply at T_c . (c) Domain-wall free energy for $L = 14, 18, \dots, 28$ (from blue to red lines). The extrapolated $\tau^{(+/-)}$ from Eq. (9) (black solid line) vanishes at T_c (dashed line).




FIG. 5. Phase transition determination for the DNNI model around $\kappa = 1/2$. (a)–(c) Internal energy above the ground state, $u_0 = -(1 + 2|\epsilon|)$, for $\epsilon = \kappa - 1/2 = 0.002, 0, -0.002$ and for $L = 8, 12, \dots, 36$ (from blue to red lines). (d), (e) extrapolation of T_c^* via different estimators indicated in legends (see text for details) for $\epsilon = 0.002$ and 0 , respectively. Dashed lines are fitted using Eq. (10) with empirical exponents $t = 1$ and 0.486 , respectively, and quadratic fits are used for $T_c^*(L)$ and $T_\xi^*(L)$. These different estimators are mutually consistent and for (e) reproduce the results of Ref. [18]. (f) For $\epsilon \rightarrow 0^+$, T_c vanishes logarithmically as in Eq. (11) (red line).

require a quadratic correction to Eq. (10), BL^{-2t} , and are thus less accurate. The estimator $T_u^*(L)$ is therefore uniquely robust in this regime. The underlying theoretical explanation for this observation, however, remains to be developed. Thanks to the exceptionally small finite-size corrections to $T_u^*(L)$ and the high TM accuracy, T_c can be determined down to $\epsilon = 10^{-5}$ [Fig. 5(f)]. (For $\epsilon < 0$, the shift of T_u^* with L makes a comparable extrapolation more haphazard.) Remarkably, for $\epsilon > 0$ the resulting transition temperature scales logarithmically as

$$T_c(\kappa) \approx -\frac{2.16}{\ln(\kappa - 1/2)}. \quad (11)$$

We thus confidently conclude that $T_c(\kappa = 1/2) = 0$.

To better understand why Ref. [18] concluded differently, we replicate their analysis in Fig. 5(e) (with the same $t = 0.486$) and consider $T_\xi^*(L)$ as well. Both extrapolations give $T_c(\kappa = 1/2) = 0.22(1)$, as Ref. [18] found. We thus understand that previous extrapolation attempts have been obfuscated by the complex and significant preasymptotic corrections to various variables around $\kappa = 1/2$. These corrections are further discussed in Sec. III B 3.

3. Order of transition

Another actively debated aspect of the DNNI model is the order of its various ordering phase transitions. In principle,




FIG. 6. Finite-size scaling of the DNNI peak specific heat for various κ : $\delta c^*/\delta L \sim L^{-1}$ for the Ising-type continuous transition ($\kappa = 0.3, 0.45$) and $\sim L^{\alpha/\nu-1}$ with $0 < \alpha/\nu \leq 1$ for the AT-type continuous transition ($\kappa = 0.67, 0.75, 1$). In the first-order regime ($\kappa = 0.48, 0.55, 0.6$), the quantity is expected to approach a constant, but for $1/2 < \kappa < \kappa^*$ the slope of the fitting line instead varies continuously.

this can be determined from the scaling of the peak specific heat:

(1) For an Ising-type continuous phase transition (expected for small κ),

$$c^*(L) \approx A \ln L + c_0, \quad (12)$$

where c_0 denotes the *background* specific heat and A is a fitting constant.

(2) For a first-order transition (expected for $1/2 < \kappa < \kappa^*$ and speculated for $\kappa^\dagger < \kappa < 1/2$),

$$c^*(L) = AL + c_0. \quad (13)$$

For a weakly first-order transition, however, the expected large preasymptotic corrections may lead to an effective scaling that deviates from linear and/or to a delay of the onset of asymptotic regime.

(3) For an Ashkin-Teller (AT)-type phase transition (expected for $\kappa \geq \kappa^*$),

$$c^*(L) = AL^{\alpha/\nu} + c_0, \quad (14)$$

where α is the heat capacity exponent and ν is the correlation length exponent, and $0 < \alpha/\nu \leq 1$. In particular, $\alpha/\nu = 1$ characterizes the Potts critical point.

To eliminate the background correction, we here consider the finite differentiate, which scales as

$$\delta c^*/\delta L = [c^*(L+1) - c^*(L-1)]/2 = AL^b, \quad (15)$$

with $b = \alpha/\nu - 1$ for continuous transitions and with $b = 0$ (a plateau) for first-order transitions. Results for selected κ are reported in Fig. 6. Fitting Eq. (15) gives $\alpha/\nu = -0.01(3)$ and $0.09(10)$ for $\kappa = 0.3$ and 0.45 , respectively, both consistent with an Ising-type transition with $\alpha = 0$. In the (expected) weakly first-order regime $1/2 < \kappa < \kappa^*$, however, the fitting slope decreases with κ . For instance, $\kappa = 0.55$ and 0.6

give $b = 0.32(2)$ and $0.02(1)$, respectively, instead of $b = 0$ throughout. This drift, which was also reported in Monte Carlo simulations of small systems [39], suggests that pronounced finite-size corrections are here at play. The TM approach thus also cannot clearly identify κ^* . Nevertheless, the regime of effective $\alpha/\nu > 1$ deviates sufficiently significantly from the AT scenario ($0 < \alpha/\nu \leq 1$) to marginally favor the weakly first-order over the continuous AT-type transition. By contrast, the drift of α/ν observed at larger κ [e.g., $\kappa = 0.75$ and 1 give $\alpha/\nu = 0.61(2)$ and $0.33(2)$, respectively] is consistent with the AT-type transition with a varying exponent [15,16].

By contrast, the Ising-type (at small κ) and the weakly first-order (speculated for $\kappa^\dagger < \kappa < 1/2$) regimes can be distinguished more straightforwardly from $b = -1$ and 0 , respectively. A clear $b = -1$ scaling persists at least up to $\kappa = 0.45$. For $\kappa = 0.48$, however, $\delta c^*/\delta L$ first decreases with L , and then grows slightly before plateauing. Although preasymptotic features partly muddle the physical picture, this trend clearly deviates from physical expectations for an Ising-type transition. It is instead reminiscent of a weakly first-order transition, and thus support the theoretical speculations of Refs. [16,17] that such a regime should exist for $\kappa^\dagger < \kappa < 1/2$ with $\kappa^\dagger \gtrsim 0.45$. (Reference [18] concluded that a continuous transition takes place for $\kappa = 0.48$, based on the absence of discontinuity in $u(T)$, but did not consider the weakly first-order transition scenario.) Interestingly, a recent computation for a low-connectivity Bethe lattice suggests that such a weakly first-order transition in the vicinity of the multicritical point ($\kappa \lesssim 1/2$ in the DNNI model) is a mean-field feature [48]. This behavior thus clearly differs from the fluctuation-induced discontinuity of the weakly first-order transition for $1/2 < \kappa < \kappa^*$.

This difference might also explain the markedly distinct scaling properties observed on either side of $\kappa = 1/2$ [as noted in Figs. 5(a) and 5(c)]. Slightly above $\kappa = 1/2$, even though preasymptotic corrections prevent a quantitative determination of α/ν (see, e.g., $\kappa = 0.55$ in Fig. 6), the monotonic growth of $c^*(L)$ is qualitatively consistent with a weakly first-order scenario. For example, at $\kappa = 0.502$, $c^*(L)$ grows nearly linearly already for $L \geq 28$ [Fig. 7(a)]. By contrast, slightly below $\kappa = 1/2$ deviations from scaling are confounding even for the sake of qualitative speculations. For instance, for $\kappa = 0.485$, $c^*(L)$ decreases at intermediate L before increasing again. From $\kappa = 0.485$ to 0.499 , this preasymptotic behavior extends to even larger L as $\kappa \rightarrow 1/2$. Moreover, the range of small L growth extends as well, leaving but a purely monotonic growth at $\kappa = 1/2$. The evolution of ξ_1 also hints at a complex finite-size behavior for $\kappa \lesssim 1/2$. For example, for $\kappa = 0.485$, a local peak appears at small L but disappears as L increases [Fig. 7(b)], and then a crossing is recovered around T_c . As κ further approaches $1/2$, this local peak survives for larger systems and is expected to persist for all $L \rightarrow \infty$ at $\kappa = 1/2$ [Fig. 7(c)]. In this limit case, ξ_1 approaches a constant at both low and high T , but a peak persists (the effective exponent $d \ln \xi_1 / d \ln L$ approaches 1), thus suggesting a disorder-disorder transition (albeit possibly shifting to $T = 0$ in the limit $L \rightarrow \infty$).

In summary, while a signature of a first-order transition is observed at $\kappa = 0.48$, a pronounced finite-size dependence




FIG. 7. The DNNI model exhibits pronounced preasymptotic corrections around $\kappa = 1/2$: (a) peak specific heat $c^*(L)$ for around $\kappa = 1/2$ and (b), (c) leading correlation length for $L = 8, 12, \dots, 32$ at $\kappa = 0.485$ and $1/2$ with T_c [dashed line in (b)]. For $\kappa = 0.485$, as L grows the crossing points between subsequent ξ_1/L (asterisk between $L = 28$ and 32) approach T_c , while the local peak vanishes.

of various observables prevents a clear characterization of the range $\kappa^\dagger < \kappa < 1/2$. We nevertheless differentiate sizable preasymptotic corrections that had previously been (incorrectly) associated with a continuous transition [18,35,49]. Particular caution should thus be applied in conducting future studies of this regime.

C. BNNNI model

For the BNNNI model, we mainly analyze the signatures of the phase transition in the antiphase regime ($\kappa > 1/2$) to obtain an overall quantitative phase diagram, which has so far eluded simulation-based approaches.

1. Correlation length scaling

We first consider the T evolution of correlation lengths with \parallel TM and \nearrow TM for $\kappa = 0.6$ (Fig. 8). In both cases, ξ_1 evolves nonmonotonically as a result of multiple eigenvalue crossings. These features are reminiscent of the ANNNI results and markedly differ from their DNNI counterparts, which suggest that a critical phase between T_{c1} and T_{c2} might be present here as well. For \parallel TM, a sharp local peak emerges slightly above the antiphase regime for congruent $L \bmod 4 \equiv 0$, and shifts to lower T as L increases. Quantitatively extrapolating transition temperatures from this observable is, however, not realistic given the limited range of accessible system sizes and the size congruence constraint. By contrast, \nearrow TM presents a much more straightforward trend. As L grows, ξ_1/L (i) seemingly diverges in the commensurate antiphase, (ii) approaches a constant in the putative IC phase, $T_{c2} \leq T \leq T_{c1}$, and (iii) vanishes in the disordered paramagnetic phase [22,47]. From the lowest temperature crossing points of ξ_1/L between systems of two nearby sizes, $T_{\xi,2}^*(L)$, we can extrapolate T_{c2} using an empirical scaling form

$$T_{\xi,2}^*(L) - T_{c2} = A_1 L^{-1} (1 + A_2 L^{-1}), \quad (16)$$




FIG. 8. Leading correlation length for the BNNNI model with $\kappa = 0.6$ in (a) \parallel TM with $L = 8, 12, 16$ (solid lines) and $L = 6, 10, 14$ (dashed lines), and (b) \perp TM with $L = 8, 12, \dots, 32$ (from blue to red lines). Asterisks mark the turning point $T_{\xi_1}^*(L)$, defined in Eq. (17), used to extrapolate T_{c1} . Inset: The first four leading correlation lengths in (a) \parallel TM with $L = 16$ and (b) \perp TM with $L = 24$. Dashed lines denote degenerate correlation length related by complex conjugate subleading eigenvalues. Vertical dashed lines in both panels denote the transition temperature $T_{c2} = 0.283(3)$.

where A_1 and A_2 are fitted constants. Although the scaling of the correction is not known a priori, the resulting extrapolation is nearly linear for all κ between 0.55 and 2 [see Fig. 9, for example], thus lending credence to this scaling form. For $\kappa =$




FIG. 9. Finite-size scaling of characteristic temperatures in \perp TM for the BNNNI model with (a) $\kappa = 0.6$ and (b) $\kappa = 0.8$. The extrapolation of $T_{\xi_2}^*(L)$ (crosses) and $T_{\xi_1}^*(L)$ (squares) with Eq. (16) provide estimates for T_{c2} and T_{c1} , respectively. Empty squares are excluded from the fit. Due to the nonmonotonicity of $T_{\xi_1}^*(L)$, T_{c1} is not estimated in (a).

0.6, in particular, fitting results from $L = 14$ to 30 gives $T_{c2} = 0.283(3)$.

While for the ANNNI model T_{c1} can be determined from the finite-size scaling of the local exponent $Y_L = \delta \ln \xi_1 / \delta \ln L$ via the \parallel TM approach [14], here the situation is not as straightforward. Eigenvalue crossings indeed persist even in the \perp TM analysis [Fig. 8(b)]. As a result, a bumplike peak appears in the plateau regime for $L = 24, 28$ but retreats for $L = 32$. (For $L = 32$, a second peak in ξ_3 appears but does not cross ξ_1 nor ξ_2 .) To understand the physical origin of this effect, recall that the angular argument of this pair of complex conjugate subleading eigenvalues corresponds to the modulation period. A sharp peak thus corresponds to the presence of a real eigenvalue (with wave number $q = \arg(\lambda_1) / (2\pi) = 0$). Because the modulation can propagate along either diagonal directions, this peak indicates that the modulation perpendicular to the TM propagation direction momentarily dominates within the relevant temperature window. Said differently, L is then commensurate with the preferred wave number. The overall scaling trend, however, remains unaffected.

The numerical challenge of determining T_{c1} nevertheless remains. A first option is to consider the scaling of the anisotropy, as for the ANNNI model [14,21]. Although small systems display a smooth evolution [22], those with larger L exhibit complex oscillations. The correlation length scaling in the IC phase regime is thus severely affected by the choice of boundary condition—again possibly resulting from the interference between preferred wave number and L —hence preventing a clear determination of T_{c1} . A second option is to restrict modulation to lie along the TM direction of propagation by examining the leading correlation length associated with a complex eigenvalue, ξ_1' . Given the smooth evolution of ξ_1' with L , it is then reasonable to extrapolate T_{c1} . As a finite- L echo of T_{c1} , we consider the local minimum

$$\left. \frac{\partial \ln \xi_1'}{\partial T} \right|_{T_{\xi_1}^*(L)} = 0, \quad (17)$$

which graphically corresponds to the turning point of $T - \ln(\xi_1'/L)$. As L increases, $T_{\xi_1}^*(L)$ first decreases but the monotonic trend does not persist [e.g., $T_{\xi_1}^*(L = 28) > T_{\xi_1}^*(L = 24)$, see Fig. 9(a)]. As κ increases, both the ξ_1 plateau and its turning onset weaken. Because eigenvalue crossing is absent for the largest L considered and $T_{\xi_1}^*(L)$ evolves smoothly and monotonically, a tentative extrapolation of T_{c1} using Eq. (16) is possible for $\kappa \geq 0.8$ [Fig. 9(b)]. We thus have that $T_{c1} > T_{c2}$, beyond the uncertainty range, up to $\kappa = 1.5$, and the signature of the critical phase is qualitatively visible up to at least $\kappa = 2$. This evidence marginally supports the absence of a Lifshitz point, i.e., $\kappa^* = \infty$. The ANNNI and the BNNNI models both exhibit a similar correlation length plateau, thus suggesting that an IC phase exists in the former as well [14]. We should note that, however, our T_{c1} values are more tentative than those of the ANNNI model [14] because similar finite-size corrections—nonmonotonic trend of $T_{\xi_1}^*(L)$ —may be observed at larger L (although less severely). Hence we cannot exclude as strongly as for the ANNNI model that the BNNNI IC-like phase might disappear in the thermodynamic limit.




FIG. 10. Specific heat for the BNNNI model for $\kappa = 0.6$ in (a) \parallel TM for $L = 8, 12, 16$ and (b) \nearrow TM for $L = 8, 12, \dots, 32$ (from blue to red lines). Vertical dashed lines denote $T_{c2} = 0.283(3)$. The position of the sharp peak in (a) is about 1% higher than that of the local peak of ξ_1 for $L = 12, 16$ [Fig. 8(a)]. Insets: (a) energy per spin for the same systems. (b) The heat capacity peak temperatures, T^* , are extrapolated (using a quadratic fitting form) to merge at $T_{c2} \approx 0.3$ as $1/L \rightarrow 0$.

2. Heat capacity evolution

We next investigate the evolution of the heat capacity. Because \parallel TM and \nearrow TM exhibit different finite-size features, we consider both. For \parallel TM, a sharp peak grows with L , but its temperature is marginally smaller than that of the local peak of ξ_1 . Correspondingly, the internal energy grows stepwise with a decreasing step height as L increases, as observed in Monte Carlo simulations [40] and in the \perp TM solution of the ANNNI model [14]. This behavior is consistent with a Pokrovsky-Talapov type transition, at which the heat capacity divergence is discontinuous (with scaling exponent $\alpha' \rightarrow \infty$) from the antiphase side. For \nearrow TM, the $c(T)$ curve is multiply peaked. The lowest temperature peak is the highest, whereas higher temperature peaks grow and shift to lower T as L increases. These peaks appear to evolve toward T_{c2} [inset of Fig. 10(b)], as does the finite- L -echo of the IC phase in the ANNNI model [14]. The analogy between the two models suggests that for $\kappa > 1/2$ the antiphase of the BNNNI model also undergoes a Pokrovsky-Talapov transition [8] at T_{c2} , followed by critical IC phase [7] that terminates at a KT transition at T_{c1} . The IC phase, if it exists, would then be characterized by an algebraically diverging correlation length and present a stepwise evolution of the modulation in finite systems.




FIG. 11. Phase diagram for the BNNNI model. The TM approach provides phase boundaries for the ferromagnetic (∞)-paramagnetic (squares), degenerate antiphases (AP) to IC (crosses), and paramagnetic-IC (circles) transition. The disorder line (asterisks) subdivides the paramagnetic phase into $q = 0$ and $q > 0$ modulation wave numbers. Configuration snapshots generated by planting [28] with \nearrow TM use blue and yellow points to denote $+1$ and -1 spins, respectively.

3. Phase diagram

Combining the correlation length and heat capacity results offers a consistent phase diagram of the BNNNI model (see Fig. 11). The simple ferromagnetic regime at $\kappa < 1/2$ presents an Ising-type transition at T_c —as identified by ξ_1/L crossings—to the paramagnetic phase. (Quantitative estimates are fully consistent with earlier TM [22] and free fermion approximation [43] results.) An additional disorder line can be identified from the splitting of subleading eigenvalues from a pair of complex conjugates (at high T) into two distinct real numbers (at low T). Being noncritical, these two lines are only marginally affected by finite-size corrections [22]. For $\kappa > 1/2$, two transitions can be identified, $T_{c2} < T_{c1}$, as discussed in Sec. III C 1. Prior estimates for T_{c2} vary dramatically, but our results robustly fall between those of Ref. [39] and those of Ref. [43]. Around $\kappa = 1/2$, the TM approach suggests that the phase boundary has a finite slope on both sides of the multicritical point, thus supporting the free fermion approximation results over those of the renormalization group approach [41]. For T_{c1} , various qualitative proposals have been made [41,42], but to the best of our knowledge no quantitative estimates were reported. Our results, albeit still somewhat imprecise, are consistent with the two-step melting scenario persisting over a wide range of κ and the presence of intermediate critical IC phase.

D. 3NNI model

We finally consider the generic 3NNI model. As shown in Fig. 1(d), the ground-state configuration of this model depends on both κ and κ' . The model is thus expected to present different types of order-disorder transitions, as characterized by the correlation length scaling (Fig. 12). In parameter regimes corresponding to the ferromagnetic, $\langle 1 \rangle$, or 4×4




FIG. 12. Correlation length scaling of the 3NNI model for different frustration parameters. For \parallel TM, solid and dashed lines denote systems with $L = 8, 12, 16$ and $L = 6, 10, 14$ (from blue to red lines). For \nearrow TM, lines denote $L = 8, 12, \dots, 28$ (from blue to red). In (a)–(d), the ground states are ferromagnetic, $\langle 1 \rangle$, 4×4 and $\langle 2 \rangle$, respectively [see Fig. 1(d)]. In (e), (f), the ground state is disordered. Note that results in (e) are truncated at large L and low T , because numerical instability of the eigensolver then gives rise to complex λ_0 .

ground-state configurations, these lengths indeed behave distinctly. Different regimes analogous to those observed in other models can further be identified (Fig. 12).

(1) In the ferromagnetic regime, ξ_1/L curves cross then kink as T increases, as they would for the thermodynamic phase transition T_c and for the disorder line crossover of the BNNNI model (the dotted line in Fig. 11), respectively.

(2) In the $\langle 1 \rangle$ regime, ξ_1/L decays monotonically and presents a single fixed point. The order of phase transition as well as the critical exponent values in the continuous transition regime also vary with the choice of (κ, κ') [38], as in the DNNI model [Fig. 4(a)].

(3) In the 4×4 regime, the ξ_1/L curves from \nearrow TM plateau after the crossing, as in the BNNNI model [Fig. 8(b)].

The $\langle 2 \rangle$ phase of the 3NNI model, however, melts differently from that of the ANNNI model. As shown in [Fig. 12(d)], ξ_1/L curves cross at single point (for congruent $L \bmod 4$), then decay monotonically without exhibiting any shoulder [cf. Fig. 2(b)]. This transition has been identified as being first order [38,39] but the distinction between a first-order and a Pokrovsky-Talapov transition is ambiguous in Monte Carlo simulations [39,40]. These two features here clearly support the former over the latter.

As mentioned in Sec. II C, the three-dimensional version of the 3NNI model was proposed as a minimal model of microemulsions. Frustration parameters were then set to $\kappa = \kappa'/2$ to match the oil-water-surfactant representation [5,6].




FIG. 13. (a) Phase diagram for the 3NNI model for $\kappa = \kappa'/2$. The TM approach provides phase boundaries for the ferromagnetic $\langle \infty \rangle$ -paramagnetic (squares), $\langle 2 \rangle$ antiphases to paramagnetic (crosses), and the disorder line (asterisks) subdivides the paramagnetic phase into $q = 0$ and $q > 0$ modulation wave numbers. Configuration snapshots generated by planting [28] with \parallel TM use blue and yellow dots to denote $+1$ and -1 spins, respectively. (b), (c) The modulation wave number q given by \parallel TM for (b) the 3NNI model with $\kappa = 0.15$, $\kappa' = 0.3$, and (c) ANNNI model with $\kappa = 0.6$ in $L = 8, 12, 16$, from blue to red.

As frustration increases, this parameter choice here results in crossing the $\langle 2 \rangle$ —lamellarlike—regime, and then following the $\langle 1 \rangle$ — 4×4 boundary, at which the system is always disordered. We specifically consider the large frustration regime ($\kappa = \kappa'/2 > 1/2$). For \parallel TM [Fig. 12(e)], ξ_1/L grows monotonically with decreasing T but no crossing is detected. Presumably, ξ_1/L then diverges at a zero-temperature phase transition. For \nearrow TM [Fig. 12(f)], however, the ξ_1/L peak shifts to lower T at large L , a behavior reminiscent of what happens at $\kappa = 1/2$ in the DNNI model [Fig. 7(c)]. Moreover, in the limit $\kappa (= \kappa'/2) \rightarrow \infty$, the model reduces to two penetrating and decoupled DNNI antiferromagnets with $\kappa = 1/2$. The \parallel TM and \nearrow TM approaches for the 3NNI model are then equivalent to the \nearrow TM and \perp TM approaches for the DNNI model, respectively. The model is thus always disordered beyond a zero-temperature phase transition [33].

Figure 13(a) presents a sketch of the κ - T phase diagram for the 3NNI model with $\kappa = \kappa'/2$. The emergence of lamellar microphases at intermediate frustration is characteristic of SALR microphase formers [2,26,50,51]. Specifically, a single first-order transition bounds the $\langle 2 \rangle$ phase for $1/6 < \kappa < 1/2$. The wave number q along the axial direction jumps at the transition [Fig. 13(b)], as expected of a first-order transition scenario. This behavior sharply contrasts with the stepwise decrease of q in the ANNNI model [Fig. 13(c)], which is expected to follow a square-root singularity in the thermodynamic limit [10,44,52]. The disordered phases in these two models are also morphologically different. In the 3NNI model, spin clusters of width 2 ($\uparrow\uparrow$) echo the modulation once it has melted. For the ANNNI model, spins instead form layers of width >2 in the vicinity of $T > T_{c1}$ [14], echoing

the floating IC phase instead. In summary, the first-order scenario for the 3NNI model at $\kappa = \kappa'/2$ is reminiscent of its three-dimensional counterpart, which also exhibit a weakly first-order transition at the melting of the modulated phase [6].

IV. CONCLUSIONS

Using a numerical TM approach, we have resolved various long-standing questions about the phase behavior of a series of two-dimensional frustrated Ising models. For the ANNNI model, the domain-wall free energy results support the existence of the critical IC phase, thus extending our recent analysis [14]. For the DNNI model, the TM results confirm the location of the transition in the limiting case $T_c(\kappa = 1/2) = 0$, and support and distinguish the weakly first-order transition scenario for $\kappa^\dagger < \kappa < 1/2$ and for $1/2 < \kappa < \kappa^*$. For the BNNNI model, a strong signature of the critical IC phase is identified, even though its upper boundary remains imprecise. For the 3NNI model, the lamellar modulated regime has been shown to melt with a single first-order transition, in contrast to that of the ANNNI model. Combining these findings provides a systematic overview of modulated phase formation, and high-accuracy benchmarks for future theoretical and numerical approaches.

The numerical TM method nevertheless still suffers from an insufficiently wide range of system sizes under certain circumstances, such as for determining the BNNNI T_{c1} . Some of these problems will be resolved, in time, as computer architecture improves. Relaxing exactness, such as by using inexact eigensolvers [53] or truncated configuration representations as in the DMRG approach [44,54] might, however, be more time effective. More immediately, the TM approach could certainly be used for other lattice models such as spin-1 and Potts models.

Data relevant to this paper have been archived and can be accessed at the Duke Digital Repository [55].

ACKNOWLEDGMENTS

We acknowledge support from the National Science Foundation Grant No. DMR-1749374 and from the Simons Foundation (No. 454937). The computations were carried out on the Duke Compute Cluster and on Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation Grant No. ACI-1548562.

APPENDIX A: TM DECOMPOSITION

In this Appendix, we detail the transfer matrix decomposition scheme used for two-dimensional frustrated Ising models. Although this approach was first implemented in the 1980s [20–22], earlier reports omitted most technical details. This Appendix and the following aim to fill this gap, and thus facilitate future extensions of these methods.

The general strategy is as follows. Although transfer matrix entries are straightforwardly expressed in Eq. (4), storing the whole matrix \mathbf{T} in memory quickly becomes beyond practical reach as L increases. The solution relies on using iterative eigenvalue algorithms (such as power iteration or Krylov

subspace-based iterations) that only require a matrix-vector multiplication subroutine, with vector \mathbf{v} as input and $\mathbf{w} = \mathbf{T}\mathbf{v}$ as output, and thus avoid explicitly storing \mathbf{T} . Because \mathbf{T} is structured, we further decompose it into a product of sparse matrices, which saves additional memory space as well as computer time. In this Appendix, we first introduce the algorithm in the context of ANNNI model in $d = 2$, and then generalize it to the DNNI and 3NNI (including BNNNI) models.

1. ANNNI model propagated perpendicular to the axial direction

We first consider the \perp TM case for the ANNNI model [Fig. 14(a)]. In this case, we denote the spin layer state $\mathbf{s} = (\pm 1)^L$ as an L dimensional binary vector, which is encoded with an L -bit unsigned integer, a , so $0 \leq a \leq 2^L - 1$ naturally include all possible layer configurations with $a_1, \dots, a_L \in \{0, 1\}$. The physical state \mathbf{s} and machine-expressed state a are related by the simple mapping $a_i = 1 \rightarrow s_i = 1$ and $a_i = 0 \rightarrow s_i = -1$.

The energetic contribution of intralayer interactions of this state is then

$$V_x(a) = -J \sum_{i=1}^L s_i s_{i+1} + \kappa J \sum_{i=1}^L s_i s_{i+2} - h \sum_{i=1}^L s_i. \quad (\text{A1})$$

Technically, this expression can be evaluated using bitwise operations. For convenience, we first define the net number of positive spins as a function of a , such that $\text{netp}(a) = 2 \text{popc}(a) - L$, where popc counts the bits set to 1. One can then write

$$V_x(a) = J \text{netp}(a \wedge \text{rol}(a, 1)) - \kappa J \text{netp}(a \wedge \text{rol}(a, 2)) - h \text{netp}(a), \quad (\text{A2})$$

where rol is rotate-left-shift (ror is similarly rotate-right-shift) and \wedge is bitwise xor. Similarly, the contribution of the energy of the neighboring layer reads

$$V_z(a, a') = -J \sum_{i=1}^L s_i s'_i = J \text{netp}(a \wedge a'). \quad (\text{A3})$$

Formally, the transfer matrix $\hat{\mathbf{T}}$ of size $N_{\text{states}} = 2^L$ (as shorthand for $\perp, \text{A} \hat{\mathbf{T}}$) has entries

$$\hat{\mathbf{T}}_{aa'} = e^{-\beta(V_x(a) + V_z(a, a'))} = \mathbf{T}_{a,a}^x \times \mathbf{T}_{a,a'}^z. \quad (\text{A4})$$

The binary representation of the entry index a, a' naturally gives the spin configuration of the corresponding layer state, as described above. $\hat{\mathbf{T}}$ can further be decomposed into a diagonal matrix \mathbf{T}^x with entries $\mathbf{T}_{a,a}^x = e^{-\beta V_x(a)}$ and a symmetric (and centrosymmetric) matrix \mathbf{T}^z with entries $\mathbf{T}_{a,a'}^z = e^{-\beta V_z(a, a')}$.

The partition function of an N -layer system can be expressed using $\text{tr}(\hat{\mathbf{T}}^N)$, which, however, is not symmetric. To leverage the efficiency of fast numerical eigensolvers for symmetric matrices, we define an alternate symmetric transfer matrix

$$\mathbf{T} = (\mathbf{T}^x)^{1/2} \mathbf{T}^z (\mathbf{T}^x)^{1/2}, \quad (\text{A5})$$




FIG. 14. Schematics of interlayer interaction in (a) $_{\perp, A}$ TM and $_{\perp, D}$ TM, (b) $_{\parallel, A}$ TM and $_{\parallel, 3}$ TM, (c) $_{/, 3}$ TM. Black lines denote spin layers; red, blue, and green lines denote the interlayer nearest-neighbor Ising, (bi)axial NNN, and diagonal interactions, respectively. Note that in (c) the inter-layer nearest-neighbor Ising interaction for (s, s') involves every other spin s_1, s_3, \dots on a layer.

which has the same the eigenvalues as the original matrix, and eigenvectors related by

$$\begin{aligned} \varphi(\mathbf{T}) &= (\mathbf{T}^x)^{-1/2} \varphi(\hat{\mathbf{T}}) \\ \varphi^{-1}(\mathbf{T}) &= \varphi^{-1}(\hat{\mathbf{T}}) (\mathbf{T}^x)^{1/2}, \end{aligned} \quad (\text{A6})$$

where the superscript -1 denotes the left eigenvector. In zero external field, $h = 0$, \mathbf{T} is further centrosymmetric ($\mathbf{T}_{a, a'} = \mathbf{T}_{2^L - a - 1, 2^L - a' - 1}$). As a result of this transformation, we have that (i) all eigenvalues are real, (ii) all eigenvectors are orthogonal, and (iii) every eigenvector is either symmetric or skew symmetric [56] when $h = 0$.

Because the size of \mathbf{T} grows exponentially with L , storing the full matrix in memory becomes first inefficient and then impractical as L increases. However, a subroutine that computes matrix-vector multiplications on the fly can be used to extract the first several leading eigenvalues and eigenvectors. A direct multiplication requires $O(4^L)$ arithmetic operations, but can be reduced by factorizing \mathbf{T}^z into sparse matrices as [57]

$$\mathbf{T}^z = \mathbf{T}^{z, L} \mathbf{T}^{z, L-1} \dots \mathbf{T}^{z, 1}, \quad (\text{A7})$$

where $\mathbf{T}^{z, i}$ has two nonzero entries in each row,

$$\begin{aligned} \mathbf{T}_{a, a'}^{z, i} &= e^{\beta J} \\ \mathbf{T}_{a, a'}^{z, i} &= e^{-\beta J}. \end{aligned} \quad (\text{A8})$$

The off-diagonal indexes $a' = a \wedge \text{rol}(1, i - 1)$ here denote the configurations obtained by flipping the i th spin from a . Note that $\mathbf{T}^{x, i}$ is transformed from $\mathbf{T}^{z, 1}$ by reindexing a to $\text{rol}(a, i - 1)$, formulated by the permutation operation

$$\mathbf{T}^{z, i} = (\mathbf{P}^T)^{i-1} \mathbf{T}^{z, 1} \mathbf{P}^{i-1}, \quad (\text{A9})$$

where \mathbf{P}^i is the permutation matrix with indexes (a, a') to be 1 and 0 otherwise. Inserting Eq. (A9) into Eq. (A7) and knowing $\mathbf{P}^T \mathbf{P} = \mathbf{I}$; $\mathbf{P}^L = \mathbf{I}$ gives

$$\mathbf{T}^z = (\mathbf{P} \mathbf{T}^{z, 1})^L. \quad (\text{A10})$$

In summary, the matrix-vector multiplication can be conducted by a sequence of multiplication with sparse matrices

$$\mathbf{T} \mathbf{v} = (\mathbf{T}^x)^{1/2} (\mathbf{P} \mathbf{T}^{z, 1})^L (\mathbf{T}^x)^{1/2} \mathbf{v}, \quad (\text{A11})$$

with complexity $O(2^L L)$.

In addition, we have that \mathbf{T} is invariant under the permutation of circularly shifting one spin, such that

$$\mathbf{T} = \mathbf{P}^T \mathbf{T} \mathbf{P}, \quad (\text{A12})$$

where $\mathbf{P}^L = \mathbf{I}$. If φ is an eigenvector of \mathbf{T} , then $\mathbf{P}\varphi$ is also an eigenvector of \mathbf{T} associated with same eigenvalue λ , because

$$\mathbf{T}(\mathbf{P}\varphi) = \mathbf{P}\mathbf{T}\varphi = \lambda\mathbf{P}\varphi. \quad (\text{A13})$$

As a result, an eigenvector of nondegenerate eigenvalue is invariant under the permutation of \mathbf{P} , and degenerate eigenvectors associated with degenerate eigenvalue (and their linear combinations) form a cyclic group. This structural property can be used to optimize the extraction of leading eigenvalues, as described in Appendix B.

2. ANNNI model propagated along the axial direction

We now consider $_{\parallel}$ TM for the ANNNI model [Fig. 14(b)]. In this case, we denote three subsequent layers as a, a', a'' where the state of each layer is encoded as an L -bit integer, as above. The energetic contribution of a then includes NN interactions with itself and with a' as well as NNN interaction with a'' :

$$\begin{aligned} V_x(a, a') &= -J \sum_{i=1}^L s_i s_{i+1} - J \sum_{i=1}^L s_i s'_i - h \sum_{i=1}^L s_i \\ V_z(a, a'') &= \kappa J \sum_{i=1}^L s_i s''_i. \end{aligned} \quad (\text{A14})$$

The bitwise operations then reads

$$\begin{aligned} V_x(a, a') &= J \text{netp}(a \wedge \text{rol}(a, 1)) + J \text{netp}(a \wedge a') - h \text{netp}(a) \\ V_z(a, a'') &= -\kappa J \text{netp}(a \wedge a''). \end{aligned} \quad (\text{A15})$$

We define the transfer matrix \mathbf{T} (as shorthand for $_{\parallel, A}$ \mathbf{T}) with entries

$$\begin{aligned} \mathbf{T}_{(a, a'), (a', a'')} &= e^{-\beta(V_x(a, a') + V_z(a, a''))} \\ &= \mathbf{T}_{(a, a'), (a, a')}^x \cdot \mathbf{T}_{(a', a'), (a', a'')}^z. \end{aligned} \quad (\text{A16})$$

The rows and columns of \mathbf{T} are indexed by a combination of two L -bit integers (a, a') and (a', a'') , so $N_{\text{states}} = 2^{2L}$. This construction results in a matrix of size $4^L \times 4^L$. Again we decompose \mathbf{T} into a diagonal matrix \mathbf{T}^x with entries $\mathbf{T}_{(a, a'), (a, a')}^x = e^{-\beta V_x(a, a')}$ and a sparse matrix \mathbf{T}^z with entries $\mathbf{T}_{(a, a'), (a', a'')}^z = e^{-\beta V_z(a, a'')}$ and 0 otherwise. The number of nonzero entries is then 8^L . The partition function of an N -layer system is given by $\text{tr}(\mathbf{T}^N)$. Unlike in Appendix A 1, \mathbf{T}^z is here no longer symmetric. A general eigensolver is then required

to solve the eigenproblem. Although generic eigenvalues can be complex, from the Perron-Frobenius theorem we have that the leading eigenvalue is always a positive real number as are the entries of the leading (left and right) eigenvectors.

Because the value of the nonzero entries in \mathbf{T}^z does not depend on a' , $\|_{\mathbb{A}} \mathbf{T}_{(a,a'),(a',a'')}^z$ can be mapped into $\perp_{\mathbb{A}} \mathbf{T}_{a,a''}^z$ with the interaction strength replaced by κJ [instead of $-J$ in Eq. (A8)]. Given that the complexity for the matrix multiplication with $\perp_{\mathbb{A}} \mathbf{T}^z$ is $O(2^L L)$, and that there are 2^L operations (for different a') in total, the complexity for the matrix-multiplication with $\|_{\mathbb{A}} \mathbf{T}^z$ is $O(4^L L)$. Hence, the matrix-multiplication operation on $\|_{\mathbb{A}} \mathbf{T}$ also has a time complexity of $O(4^L L)$.

3. DNNI model

Because the horizontal and vertical directions of the DNNI model are equivalent, a single transfer matrix can be defined. The contribution of intralayer states,

$$V_x(a) = -J \sum_{i=1}^L s_i s_{i+1} - h \sum_{i=1}^L s_i, \quad (\text{A17})$$

is independent of κ , while the energetic contribution of neighboring layers reads

$$V_z(a, a') = -J \sum_{i=1}^L s_i s'_i + \kappa J \left(\sum_{i=1}^L s_i s'_{i-1} + \sum_{i=1}^L s_i s'_{i+1} \right). \quad (\text{A18})$$

The transfer matrix \mathbf{T} (as shorthand for $\perp_{\mathbb{D}} \mathbf{T}$) can thus be decomposed into intralayer and interlayer interactions as in Eq. (A5). For the DNNI model, the interlayer matrix \mathbf{T}^z is also symmetric (and centrosymmetric when $h = 0$) with entries $\mathbf{T}_{a,a'}^z = e^{-\beta V_z(a,a')}$.

Again, \mathbf{T}^z can be decomposed to reduce the complexity of the matrix-vector multiplication, but we can no longer use Eq. (A10). This scheme drops information about s'_{i-1} after computing the interlayer interaction for s_{i-1} , which, although fine for the ANNNI model, for the DNNI model leaves out the interaction between s_i and its diagonal neighbors, s'_{i-1} and s'_{i+1} [Fig. 14(a)]. To make up for this loss, we introduce an auxiliary spin $t_1 = s'_i$ for spin indexes $i = 1 \dots L$ during the propagation. Because periodic boundary conditions require $s_1 = s_{L+1}$, we introduce an additional auxiliary spin for $r_1 = s'_1$. (In the ANNNI model formulation, s_L does not interact with s'_1 .) The factorization of \mathbf{T}^z is thus

$$\begin{aligned} \mathbf{T}^z &= \mathbf{S}^{-1} \mathbf{T}^{z,L} \mathbf{T}^{z,L-1} \dots \mathbf{T}^{z,1} \mathbf{S} \\ &= \mathbf{S}^{-1} (\mathbf{P} \mathbf{T}^{z,1|r_1}) (\mathbf{P} \mathbf{T}^{z,1})^{L-1} \mathbf{S}. \end{aligned} \quad (\text{A19})$$

The auxiliary matrix \mathbf{S} (and \mathbf{S}^{-1}) maps (recovers) a vector of dimension 2^L to (from) 2^{L+2} , namely,

$$\begin{aligned} (\mathbf{S}x) &(\{s_1, s_2, \dots, s_L, \dots, t_1, r_1\}) \\ &= \begin{cases} x(s_1, \dots, s_L), & t_1 = s_L \text{ and } r_1 = s_1 \\ 0, & \text{otherwise} \end{cases} \end{aligned} \quad (\text{A20})$$

and

$$(\mathbf{S}^{-1}y)(a) = \sum_{t_1, r_1} y(a, t_1, r_1). \quad (\text{A21})$$

The matrix $\mathbf{T}^{z,1}$, which denotes the contribution on the interlayer interaction for one spin, s_1 , has entries

$$\begin{aligned} \mathbf{T}^{z,1} &(\{s_1, s_2, \dots, s_L, t_1, r_1\}, \{s'_1, s'_2, \dots, s'_L, t'_1, r'_1\}) \\ &= \begin{cases} e^{\beta J s_1 s'_1 - \kappa \beta J s_1 (t'_1 + s'_2)}, & s_i = s'_i (i = 2 \dots L) \\ & t_1 = s'_1, r_1 = r'_1 \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \quad (\text{A22})$$

The permutation matrix \mathbf{P} shifts spins $\{s_1, \dots, s_L\}$ to $\{s_2, \dots, s_L, s_1\}$ but does not change auxiliary spins. Note that the term $\mathbf{T}^{z,1|r_1}$ in Eq. (A19) reflects the periodic boundary condition that replaces s'_2 by r_1 .

The overall time complexity of matrix-vector multiplication remains $O(2^L L)$. The size of the temporary vector is quadrupled compared to $\perp_{\mathbb{A}} \mathbf{T}^z$ for the ANNNI model because two auxiliary spins had to be included, but can be halved by tracing r_1 in the code (instead of as a vector index) because it is only invoked for operations on $\mathbf{T}^{z,L}$.

4. BNNNI and 3NNI models

Because the BNNNI model is a special case of the 3NNI model (with $\kappa' = 0$), we only need to consider the latter. One possible transfer matrix construction thus requires but minimal modification from $\|_{\mathbb{A}} \mathbf{T}^z$, namely, including diagonal nearest-neighbor and axial next-nearest-neighbor interactions in the intralayer part $\mathbf{T}_{(a,a'),(a,a')}$,

$$\|_{\mathbb{3}} \mathbf{T}_{(a,a'),(a,a')}^x = \exp\{-\beta[V_x(a, a') + V_{x,B}(a) + V_{x,D}(a, a')]\} \quad (\text{A23})$$

where $V_{x,B}$ and $V_{x,D}$ are missing from the ANNNI model in Eq. (A15):

$$V_{x,B}(a) = -\kappa J \text{netp}(a \wedge \text{rol}(a, 2)), \quad (\text{A24})$$

$$\begin{aligned} V_{x,D}(a, a') &= -\kappa' J [\text{netp}(a \wedge \text{rol}(a', 1)) \\ &\quad + \text{netp}(a \wedge \text{ror}(a', 1))]. \end{aligned} \quad (\text{A25})$$

The structure and complexity of the remaining algorithm then remain unchanged.

However, as stated in the main text (Sec. IID), the checkerboard and diagonal striped phases of the BNNNI and 3NNI models are naturally modulated along the diagonals of a square lattice. To study the correlation length of these modulations and to minimize the finite-size disturbances observed in $\|_{\mathbb{3}} \mathbf{T}$, we thus consider a transfer matrix propagated along the diagonal direction, $\|_{\mathbb{3}} \mathbf{T}$, hence generalizing the approach of Ref. [22]. Note that this arrangement requires L to be even.

As in Eq. (A5), the resulting transfer matrix can be decomposed into intralayer and interlayer contributions. The intralayer matrix can be computed directly, and the inter-layer matrix \mathbf{T}^z can be decomposed similarly as for the DNNI model [Eq. (A19)]

$$\begin{aligned} \mathbf{T}^z &= \mathbf{S}^{-1} \mathbf{T}^{z,L} \mathbf{T}^{z,L-1} \dots \mathbf{T}^{z,2} \mathbf{S} \\ &= \mathbf{S}^{-1} (\mathbf{P} \mathbf{T}^{z,2|t_1, t_2}) (\mathbf{P} \mathbf{T}^{z,2})^{L/2-1} \mathbf{S}. \end{aligned} \quad (\text{A26})$$

The auxiliary matrix \mathbf{S} (and \mathbf{S}^{-1}) then maps (recovers) a vector of dimension 2^L to (from) 2^{L+4} ,

$$(\mathbf{S}x)(\{s_1, s_2, \dots, s_L, \dots, t_1, t_2, r_1, r_2\}) = \begin{cases} x(s_1, \dots, s_L), & (t_1, t_2, r_1, r_2) = (s_{L-1}, s_L, s_1, s_2) \\ 0, & \text{otherwise} \end{cases} \quad (\text{A27})$$

and

$$(\mathbf{S}^{-1}y)(a) = \sum_{t_1, t_2, r_1, r_2} y(a, t_1, t_2, r_1, r_2). \quad (\text{A28})$$

The matrix $\mathbf{T}^{z,2}$, which denotes the contribution on the interlayer interaction for two spins, s_1 and s_2 , [Fig. 14(c)], has entries

$$\mathbf{T}^{z,2}(\{s_1, \dots, s_L, t_1, t_2, r_1, r_2\}, \{s'_1, \dots, s'_L, t'_1, t'_2, r'_1, r'_2\}) = \begin{cases} \exp\{-\beta J[(s'_1(t_2 + s_2) - \kappa(s'_1(t_1 + s_3) + s'_2(t_2 + s_4)) \\ -\kappa'(s'_1 s_1 + s'_2 s_2))]\}, & s_i = s'_i (i = 3 \dots L), (t_1, t_2, r_1, r_2) = (s'_1, s'_2, r'_1, r'_2) \\ 0, & \text{otherwise.} \end{cases} \quad (\text{A29})$$

The permutation matrix \mathbf{P} shifts layer configurations by two spins, i.e., $\{s_1, s_2, \dots, s_L\} \rightarrow \{s_3, \dots, s_L, s_1, s_2\}$. The term $\mathbf{T}^{z,2/r_1, r_2}$ in Eq. (A26) reflects the periodic boundary condition that replaces s'_3, s'_4 by r_1, r_2 .

This arrangement of auxiliary spins can be viewed as a generalization of the approach used for the DNNI model. A spin s_i here involves interactions with $s_{i\pm 2}$, which goes beyond $s_{i\pm 1}$ for the DNNI model. In general, for models with interlayer interactions between s_i and $s'_{i\pm b}$, $2b$ auxiliary spins are needed, among which t spins are associated with an extended vector, then of size 2^{L+b} . The size of this vector controls the space complexity for the matrix-vector multiplication. In addition to the 2^b loops for different choices of r spins, the time complexity is then also larger, $O(2^{L+2b}L)$. For $_{/3}\mathbf{T}$, in particular, the number of operations and the intermediate vector size are $2^4 = 16$ and $2^2 = 4$ times that for $_{\perp A}\mathbf{TM}$, respectively.

APPENDIX B: REDUCING SPACE COMPLEXITY WITH SYMMETRY

In this Appendix, we describe computational schemes used to reduce the size of the transfer matrix, and thus significantly decrease the algorithmic space complexity. The key idea is to identify equivalent states to construct orthogonal bases (or irreducible representations [20], as have been implemented in related models [16,57]). We here adapt this method following the framework of the structured matrix decomposition described in Appendix A. We first derive the general method for structured matrices with certain permutation invariance and then analyze the complexity of the transfer matrix involved in solving the models of interest.

1. General case

The $n \times n$ matrix \mathbf{T} is invariant under permutations $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_g$, such that $\mathbf{P}_i^T \mathbf{T} \mathbf{P}_i = \mathbf{T}$ for $i = 1, \dots, g$ and that these transformations form a symmetry group \mathcal{G} . Matrix indexes are then grouped by these transformations. For example, for a centrosymmetric matrix, under the transformation of $\mathbf{P}^T \mathbf{T} \mathbf{P}$ where $\mathbf{P} = (e_n, e_{n-1}, \dots, e_1)$, row and column indexes are permuted as $0 \rightarrow n-1, 1 \rightarrow n-2$ and so on. The indexes i and $n-i-1$ are then deemed equivalent, and thus there are $n/2$ equivalent sets in total.

We next construct a (nonsquare) matrix with orthogonal columns

$$\mathbf{Q} \equiv \begin{bmatrix} 1/\sqrt{g_1} & 0 & \dots & \dots & 0 \\ 0 & \dots & 1/\sqrt{g_i} & \dots & \dots \\ \dots & \dots & \dots & \dots & 1/\sqrt{g_n} \\ 1/\sqrt{g_1} & \dots & \dots & \dots & \dots \end{bmatrix} \quad (\text{B1})$$

of size $n \times m$, where m is the number of sets of equivalent indexes (not to be confused with the magnetization). Each column in \mathbf{Q} is a column vector corresponding to an equivalent index set of size g_i . The entries in \mathbf{Q}_i are nonzero, and set to $1/\sqrt{g_i}$, if and only if its row index is in the set. Applying the similarity transformation on \mathbf{T} with these bases, we obtain a matrix of size $m \times m$

$$\mathbf{M} = \mathbf{Q}^T \mathbf{T} \mathbf{Q}. \quad (\text{B2})$$

The eigenvalues of \mathbf{M} are also eigenvalues of \mathbf{T} and, specifically, \mathbf{M} and \mathbf{T} have the same leading eigenvalue with eigenvector $\varphi_0(\mathbf{T}) = \mathbf{Q}\psi_0(\mathbf{M})$.

Two-dimensional spin models under periodic boundary conditions are invariant under rotation of one spin (a C_n axis) as well as under counting spins backward (a σ_v reflection), and hence belongs to the C_{nv} point group. In the absence of external field, $h = 0$, the model is also invariant under flipping all spins (a σ_h reflection) and hence it belongs to the D_{nh} group. The dimension of \mathbf{M} is asymptotically reduced by a factor that equals the order of the symmetry group— $2L$ for C_{nv} and $4L$ for D_{nh} (Fig. 15). In this way, the transfer matrix size can be compressed by a factor of $2L$ (or $4L$ if $h = 0$ and only the leading eigenvalue is needed).

2. $_{\perp}\mathbf{TM}$ and $_{/}\mathbf{TM}$

We now adapt this compressed matrix to our transfer matrix calculation. We first consider $_{\perp}\mathbf{TM}$. (The construction of $_{/}\mathbf{TM}$ is very similar to that of $_{\perp}\mathbf{TM}$, as we will see later.) Directly implementing Eq. (A11) to calculate the matrix-vector multiplication

$$\begin{aligned} \mathbf{w} &= \mathbf{M}\mathbf{v} = \mathbf{Q}^T \mathbf{T} \mathbf{Q} \mathbf{v} = \mathbf{Q}^T (\mathbf{T}^x)^{1/2} (\mathbf{P} \mathbf{T}^{z,1})^L (\mathbf{T}^x)^{1/2} \mathbf{Q} \mathbf{v} \\ &= \mathbf{Q}^T \mathbf{w}' \end{aligned} \quad (\text{B3})$$




FIG. 15. The number m of equivalent state sets of different symmetry groups grows with layer length L for (a) \perp TM and (b) \parallel TM. The values obtained follow certain integer series [58]. Insets: m asymptotically scales with N_{states}/L , $N_{\text{states}}/(2L)$ and $N_{\text{states}}/(4L)$ under C_n , C_{nv} , and D_{nh} symmetry, respectively.

gives essentially the same time and space complexity as for calculating $\mathbf{T}\mathbf{v}$. Further optimization is, however, possible. Observing that entries in the intermediate vector \mathbf{w}'_a that belong to the same equivalent states ($a \in b$) are identical, we only need to compute one (among g_b identical entries) for each set of equivalent states in \mathbf{w}' to construct \mathbf{w} , such that

$$\mathbf{w}_b = \sqrt{g_b} \mathbf{w}'_a. \quad (\text{B4})$$

To take advantage of this property, we initialize an array of equivalent states, denoted $[b]$, containing one of the states a in the set as well as the set size g_b . The number of equivalent sets approaches $2^L/(4L)$ for $h = 0$ and $2^L/(2L)$ otherwise. In both cases, $[b]$ has a space complexity of $O(2^L/L)$. This list can be constructed in two ways that each offer a different space/time complexity balance. The first is to set up a temporary array of 2^L bits (thus with a $O(2^L)$ space complexity) and scanning once (thus with a $O(2^L)$ time complexity). The second is to enumerate each of the 2^L states and check all of its equivalent states by bit-wise operations (in a $O(2^L)$ time complexity), and push the state to the array only if it has the smallest index of all equivalent states. The total number of equivalent state $[O(2^L/L)]$ then gives the space complexity.

Equation (B3) can then be evaluated in two parts. First, we compute the interlayer interactions of all states a with L' leftmost spins having the same configuration, a^ℓ . Denoting the remaining $L - L'$ bits to their right as a^r , we hence have $a \equiv a^\ell . a^r$ where “.” is the bit concatenation operation. In practice, we set up an intermediate vector $a^\ell \mathbf{v}$ of size $2^{L-L'}$ such that

$$(({}^0\mathbf{v})^T, ({}^1\mathbf{v})^T, \dots, ({}^{2^L-1}\mathbf{v})^T)^T = \mathbf{Q}\mathbf{v}. \quad (\text{B5})$$

Decoding each $a^\ell \mathbf{v}$ from $[b]$ and \mathbf{v} costs a time $O(L \times m) = O(2^L)$ and needs to run 2^L times. For each $a^\ell \mathbf{v}$, we further compute

$$a^\ell \mathbf{w}'' = a^\ell ((\mathbf{P}\mathbf{T}^{\varepsilon,1})^{L-L'} (\mathbf{T}^x)^{1/2}) \times a^\ell \mathbf{v}. \quad (\text{B6})$$

The time complexity of this step is $O[(L - L')2^{L-L'}]$ and it is run 2^L times (or 2^{L-1} times for $h = 0$, because \mathbf{T} is then centrosymmetric). Second, for every nonequivalent entry in \mathbf{w}' , we also decompose the index $a' = (a^\ell . a^r)$ and increment $\mathbf{w}'_{a'}$ by $\exp[-\beta V_z(a^\ell, a^r)] \times a^\ell \mathbf{w}''_{a^r}$. In summary, this approach gives

$$\mathbf{w}'_{a'} = (\mathbf{T}^x_{a',a'})^{1/2} \sum_{a^\ell} \exp[-\beta V_z(a^\ell, a^r)] \times a^\ell \mathbf{w}''_{a^r}, \quad (\text{B7})$$

and the time complexity for this step is $O(2^L \times m) = O(2^{L+L}/L)$. Comparing the time complexities of Eqs. (B5)–(B7), we choose $L' = 1 + \lfloor \log_2 L \rfloor$ so the total time complexity for matrix-vector multiplication remains $O(2^L L)$ and the space complexity is reduced to $O(2^L/L)$.

The permutation operations that generate equivalent states for \parallel TM slightly differs from \perp TM because the layer then has a zigzag shape. Specifically, the system is invariant after shifting two (instead of one) spins as well as by first shifting one spin and then counting backwards (instead of simply counting backwards). The number of equivalent states m then asymptotically approaches $2^L/L$ (or $2^L/(2L)$ for $h = 0$). Because we consider two spins (s_1, s_2) in every operation in $\mathbf{T}^{\varepsilon,2}$, we choose (an even) $L' = 2(1 + \lfloor \log_2 L/2 \rfloor)$. The time and space complexities remain the same as for \perp TM, but with a larger prefactor.

3. \parallel TM

Similarly to Eq. (B3), for \parallel TM the matrix-vector multiplication is decomposed as

$$\mathbf{w} = \mathbf{M}\mathbf{v} = \mathbf{Q}^T \mathbf{T} \mathbf{Q} \mathbf{v} = \mathbf{Q}^T (\mathbf{T}^x) (\mathbf{P}\mathbf{T}^{\varepsilon,1})^L \mathbf{Q} \mathbf{v} = \mathbf{Q}^T \mathbf{w}'. \quad (\text{B8})$$

Now, however, \mathbf{M} is not symmetric. In addition to permutation invariance, we can also take advantage of the sparsity of \parallel TM (Appendix A 2) to compute $\mathbf{M} \times \mathbf{v}$. The extra space needed is a vector of size 2^L , which is much smaller than the vector size of $m = O(4^L/L)$. The time complexity can also be reduced because \mathbf{w}' has identical entries for equivalent states.

The algorithm is as follows. First, we initialize the array of equivalent states $[b]$, as we did for \perp TM, such that each element is a pair of L -bit integers (a, a') that represents this set. Two extra arrays are stored for later bookkeeping purposes:

- (1) An array of equivalent states for a alone, denoted $[c]$, along with the period of a under cyclic shift. The size of $[c]$

approaches $m' \approx 2^L/(4L)$ for D_{nh} and $m' \approx 2^L/(2L)$ for C_{nv} bases.

(2) An array of indexes $[b']$ for each representing state (a, a') in $[b]$ that records the references in $[b]$ corresponding to the equivalent state with layers swapped, (a', a) , denoted $b'(a', a)$.

The construction of $[b']$ follows the construction of $[b]$. For each (a, a') newly appended to $[b]$, we find $b'(a', a)$. If $b'(a', a) \leq (a, a')$, a binary search finds the index of $b'(a', a)$ in $[b]$. (Each binary search takes on average $O(\ln m) = O(L)$ operations, and hence the overall algorithmic complexity remains unchanged.) In summary, the initialization takes $O(4^L)$ operations, and storing $[b]$ takes space $O(4^L/L)$.

Second, we setup the subroutine for the matrix-vector multiplication of Eq. (B3). Again we denote the indexes of \mathbf{w} and \mathbf{v} as (a, a') and (a', a'') , respectively. For each a' in $[c]$, we construct an intermediate vector ${}^a\mathbf{v}$ such that

$$(({}^{a_1}\mathbf{v})^T, ({}^{a_2}\mathbf{v})^T, \dots, ({}^{a_{m'}}\mathbf{v})^T)^T = \mathbf{Q}\mathbf{v}. \quad (\text{B9})$$

Because each ${}^a\mathbf{v}$ is of size 2^L and m' of these vectors in total, the complexity of this step is $O(2^L m') = O(4^L/L)$.

For each ${}^a\mathbf{v}$, we compute

$${}^a\mathbf{w}'' = (\mathbf{P} \cdot \perp \mathbf{T}^{z,1})^L \times {}^a\mathbf{v}, \quad (\text{B10})$$

with $J' = -\kappa J$ in $\mathbf{T}^{z,1}$. It takes $O(2^L L)$ operations per a , and hence the total complexity of this step is $O(2^L L \times 2^L/L) = O(4^L)$.

The entries in the resulting vector, ${}^a\mathbf{w}''_a$ correspond to those of the intermediate vector \mathbf{w}' with

$$\mathbf{w}'_{b'(a,a')} = \mathbf{T}^x_{b'(a,a'), b'(a,a')} {}^a\mathbf{w}''_a. \quad (\text{B11})$$

Again, we have

$$\mathbf{w}_b = \sqrt{g_b} \mathbf{w}'_a, \quad (\text{B12})$$

and hence Eq. (B11) needs to be evaluated $m = O(4^L/L)$ times.

In summary, the time complexity, $O(4^L)$, is improved by a factor of L over that in Appendix A 2. The extra space needed is $O(m')$ which is marginal given that input and output vectors have sizes $O(m)$.

4. Remarks

Thanks to these compressed TMs, the evaluation of systems with L up to 36 for \perp TM, 32 for $/$ TM and 16 for \parallel TM is accessible within 60 GB memory using the SPECTRALIB package [59]. For these largest systems and an iteration convergence criterion of 10^{-10} , computations take about one day on eight cores for \perp TM and $/$ TM and half a day on a single core for \parallel TM. The time and memory needed for evaluating larger systems exceed currently available computational resources and have thus not been successfully achieved.

Interestingly, when evaluating \mathbf{T} with an iterative eigensolver, convergence slows down markedly around transition temperatures (Fig. 16). (For the DNNI model, the slowdown extends to the entire modulated phase.) This property is related to the underlying physics of these transitions. The condition number, which is defined as the error of the output given a slightly erroneous (or finite-precision) input, is




FIG. 16. Number of iterations N_{iter} needed to solve for the leading eigenvalue of (a) \perp, A TM with $\kappa = 0.6$ and (b) \perp, D TM with $\kappa = 0.6$ for $L = 8, 12, \dots, 24$ (from blue to red lines). Results of the compressed TM (main plots) and the original $2^L \times 2^L$ TM (insets) are compared with the heat capacity for $L = 24$ (light red strip, right axis). While N_{iter} for the original TM scheme exhibits a dramatic numerical slowdown upon approaching the phase transition in both models (vertical dashed lines) following the heat capacity trend, the compressed TM scheme increases by at most a factor of 3.

indeed related to the specific heat, which corresponds to the energy fluctuations, $c = \langle (\Delta u)^2 \rangle / (k_B T^2)$. When c is high, the output of the eigenvectors (density of configurations) is unstable, which corresponds to a large condition number. The slowdown for the compressed TM, however, is much less pronounced. Said differently, the compressed TM is better conditioned, which results in a more numerically stable algorithm and higher-accuracy results. In physical terms, this suggests that part of the c growth is then scaled away through the transformation to a compressed TM.

A potential challenge for this decomposition, however, is that subleading eigenvalues of \mathbf{T} can lie either in $\text{span}(\mathbf{M})$ or in $\text{null}(\mathbf{M})$. In other words, the spectral gap, which gives the leading correlation length, of \mathbf{M} does not necessarily coincide with that of \mathbf{T} . In practice, it is observed that for \parallel, A \mathbf{T} , \perp, D \mathbf{T} , $\parallel, 3$ \mathbf{T} and $/, 3$ \mathbf{T} in $h = 0$, the subleading eigenvectors are all skew symmetric, and are preserved in the compressed TMs with C_{nv} bases. We therefore identify the correlation length from the compressed matrix. For \perp, A \mathbf{T} , however, the subleading eigenvalue is doubly degenerate (as in Ref. [20]), and is observed in the null space of \mathbf{M} obtained from C_{nv} bases. The original transfer matrix is thus used to identify the correlation length in this case. As noted in Ref. [20], this property follows from the symmetry of the bases. For other models a similar argument might also be possible.

- [1] F. Sciortino, S. Mossa, E. Zaccarelli, and P. Tartaglia, *Phys. Rev. Lett.* **93**, 055701 (2004).
- [2] Y. Zhuang and P. Charbonneau, *J. Phys. Chem. B* **120**, 7775 (2016).
- [3] W. Selke and M. E. Fisher, *Phys. Rev. B* **20**, 257 (1979).
- [4] E. Dagotto, *Rev. Mod. Phys.* **85**, 849 (2013).
- [5] B. Widom, *J. Chem. Phys.* **84**, 6943 (1986).
- [6] K. A. Dawson, M. D. Lipkin, and B. Widom, *J. Chem. Phys.* **88**, 5149 (1988).
- [7] J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* **6**, 1181 (1973).
- [8] V. L. Pokrovsky and A. L. Talapov, *Phys. Rev. Lett.* **42**, 65 (1979).
- [9] J. Glasbrenner, I. Mazin, H. O. Jeschke, P. Hirschfeld, R. Fernandes, and R. Valentí, *Nat. Phys.* **11**, 953 (2015).
- [10] W. Selke, *Phys. Rep.* **170**, 213 (1988).
- [11] J. L. Morán-López, F. Aguilera-Granja, and J. M. Sanchez, *Phys. Rev. B* **48**, 3519 (1993).
- [12] T. Shirakura, F. Matsubara, and N. Suzuki, *Phys. Rev. B* **90**, 144410 (2014).
- [13] F. Matsubara, T. Shirakura, and N. Suzuki, *Phys. Rev. B* **95**, 174409 (2017).
- [14] Y. Hu and P. Charbonneau, *Phys. Rev. B* **103**, 094441 (2021).
- [15] S. Jin, A. Sen, and A. W. Sandvik, *Phys. Rev. Lett.* **108**, 045702 (2012).
- [16] S. Jin, A. Sen, W. Guo, and A. W. Sandvik, *Phys. Rev. B* **87**, 144406 (2013).
- [17] A. Bobák, T. Lučivjanský, M. Borovský, and M. Žukovič, *Phys. Rev. E* **91**, 032145 (2015).
- [18] M. Ramazanov, A. Murtazaev, and M. Magomedov, *Solid State Commun.* **233**, 35 (2016).
- [19] H. Li and L.-P. Yang, *Phys. Rev. E* **104**, 024118 (2021).
- [20] W. Pesch and J. Kroemer, *Z. Phys. B: Condens. Matter* **59**, 317 (1985).
- [21] P. D. Beale, P. M. Duxbury, and J. Yeomans, *Phys. Rev. B* **31**, 7166 (1985).
- [22] J. Oitmaa, M. T. Batchelor, and M. N. Barber, *J. Phys. A* **20**, 1507 (1987).
- [23] R. B. Lehoucq, D. C. Sorensen, and C. Yang, *ARPACK Users' Guide: Solution of Large-Scale Eigenvalue Problems with Implicitly Restarted Arnoldi Methods* (SIAM, Philadelphia, PA, 1998).
- [24] M. J. Godfrey and M. A. Moore, *Phys. Rev. E* **91**, 022120 (2015).
- [25] J. F. Robinson, M. J. Godfrey, and M. A. Moore, *Phys. Rev. E* **93**, 032101 (2016).
- [26] Y. Hu and P. Charbonneau, *Soft Matter* **14**, 4101 (2018).
- [27] Y. Hu, L. Fu, and P. Charbonneau, *Mol. Phys.* **116**, 3345 (2018).
- [28] Y. Hu and P. Charbonneau, *Phys. Rev. Res.* **3**, 038001 (2021).
- [29] R. A. dos Anjos, J. R. Viana, and J. R. de Sousa, *Phys. Lett. A* **372**, 1180 (2008).
- [30] A. Kalz, A. Honecker, and M. Moliner, *Phys. Rev. B* **84**, 174407 (2011).
- [31] Y. Boughaleb, M. Nouredine, M. Snina, R. Nassif, and M. Bennai, *Phys. Res. Int.* **2010**, 284231 (2010).
- [32] R. Timmons and K. De'Bell, *Can. J. Phys.* **96**, 912 (2018).
- [33] D. P. Landau, *Phys. Rev. B* **21**, 1285 (1980).
- [34] A. Kalz, A. Honecker, S. Fuchs, and T. Pruschke, *Eur. Phys. J. B* **65**, 533 (2008).
- [35] A. Kalz, A. Honecker, S. Fuchs, and T. Pruschke, in *Journal of Physics: Conference Series* (IOP Publishing, Braunschweig, Germany, 2009), Vol. 145, p. 012051.
- [36] S.-Y. Kim, *Phys. Rev. E* **81**, 031120 (2010).
- [37] J. Oitmaa, *J. Phys. A* **14**, 1159 (1981).
- [38] R. M. Liu, W. Z. Zhuo, S. Dong, X. B. Lu, X. S. Gao, M. H. Qin, and J.-M. Liu, *Phys. Rev. E* **93**, 032114 (2016).
- [39] D. P. Landau and K. Binder, *Phys. Rev. B* **31**, 5946 (1985).
- [40] M. J. Velgakis and J. Oitmaa, *J. Phys. A* **21**, 547 (1988).
- [41] M. Aydin and M. C. Yalabik, *J. Phys. A* **22**, 85 (1989).
- [42] M. Aydin and M. C. Yalabik, *J. Phys. A* **22**, 3981 (1989).
- [43] S. Dasgupta, *J. Phys. A* **24**, 1017 (1991).
- [44] R. Derian, A. Gendiar, and T. Nishino, *J. Phys. Soc. Jpn.* **75**, 114001 (2006).
- [45] H. L. Richards, M. A. Novotny, and P. A. Rikvold, *Phys. Rev. B* **48**, 14584 (1993).
- [46] V. Privman, *Finite Size Scaling and Numerical Simulation of Statistical Systems* (World Scientific, Singapore, 1990).
- [47] M. P. Nightingale, *Physica A* **83**, 561 (1976).
- [48] P. Charbonneau and M. Tarzia, *J. Chem. Phys.* **155**, 024501 (2021).
- [49] J. H. Lee, H. S. Song, J. M. Kim, and S.-Y. Kim, *J. Stat. Mech.: Theory Exp.* (2010) P03020.
- [50] N. G. Almaraz, J. Pękalski, and A. Ciach, *J. Chem. Phys.* **140**, 164708 (2014).
- [51] Y. Zhuang, K. Zhang, and P. Charbonneau, *Phys. Rev. Lett.* **116**, 098301 (2016).
- [52] A. Sato and F. Matsubara, *Phys. Rev. B* **60**, 10316 (1999).
- [53] Z. Wang, Ph.D. thesis, Duke University, 2020.
- [54] T. Nishino, *J. Phys. Soc. Jpn.* **64**, 3598 (1995).
- [55] Duke digital repository, <https://doi.org/10.7924/r4v40tw6p>.
- [56] A. Cantoni and P. Butler, *Linear Algebra Its Appl.* **13**, 275 (1976).
- [57] H. W. J. Blöte and M. P. Nightingale, *Physica A* **112**, 405 (1982).
- [58] For \perp TM: Number of equivalent states under C_n , C_{nv} , and D_{nh} symmetries correspond to OEIS series (<https://oeis.org>) A000031, A000029, and A000011; for \parallel TM: C_n , C_{nv} , and D_{nh} equivalent states correspond to OEIS series A001868, A081720 (fourth column), and A283846.
- [59] Y. Qiu, Spectralib (sparse eigenvalue computation toolkit as a redesigned ARPACK [23]), (2020).