

Lieb-Robinson bounds imply locality of interactionsHenrik Wilming¹ and Albert H. Werner^{2,3}¹*Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland*²*QMATH, Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark*³*NBIA, Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark*

(Received 22 December 2020; revised 22 December 2021; accepted 22 December 2021; published 2 March 2022)

Discrete lattice models are a cornerstone of quantum many-body physics. They arise as effective descriptions of condensed-matter systems and lattice-regularized quantum field theories. Lieb-Robinson bounds imply that if the degrees of freedom at each lattice site only interact locally with each other, correlations can only propagate with a finite group velocity through the lattice, similarly to a light cone in relativistic systems. Here we show that Lieb-Robinson bounds are equivalent to the locality of the interactions: a system with k -body interactions fulfills Lieb-Robinson bounds in exponential form if and only if the underlying interactions decay exponentially in space. In particular, our result already follows from the behavior of two-point correlation functions for single-site observables and generalizes to different decay behaviors as well as fermionic lattice models. As a side result, we thus find that Lieb-Robinson bounds for single-site observables imply Lieb-Robinson bounds for bounded observables with arbitrary support.

DOI: [10.1103/PhysRevB.105.125101](https://doi.org/10.1103/PhysRevB.105.125101)**I. INTRODUCTION**

A crucial feature of the dynamics of a quantum many-body system with local Hamiltonian is the finite group velocity v for the spread of correlations first shown by Lieb and Robinson [1]. If A and B are local observables at locations x and y of a lattice, this finite group velocity implies that two-point correlation functions of the form

$$C_{x,y}(t_1, t_0) = i\langle \Psi [A(t_1), B(t_0)] | \Psi \rangle \quad (1)$$

are exponentially small if (x, t_1) lies outside the effective “light cone” emanating from (y, t_0) with the *Lieb-Robinson velocity* v , see Fig. 2. Such behavior is reminiscent of the strict light cone in relativistic field theories and holds for any lattice Hamiltonian with bounded k -body interactions that decay exponentially in space. It is the basis for a host of important results in condensed-matter and quantum many-body physics. To name just a few, these range from the exponential clustering theorem for ground states of gapped Hamiltonians [2–5], to the quantization of Hall conductance [6,7], Lieb-Schultz-Mattis theorems [2,8], Goldstone’s theorem [9,10], and the area law in one spatial dimension [11]. Besides these theoretical results, the finite group velocity is also crucial to be able to simulate the dynamics of complex quantum systems both on a classical as well as on quantum computers, because it ensures that the dynamics can be well approximated by a local quantum circuit. Many-body systems can now be simulated in experiments, for example, using ultracold atoms in optical lattices, and the Lieb-Robinson cone can be detected experimentally [12–14].

In relativistic field theories, which show a strict light cone, the interactions are usually required to be strictly local. A natural question one may therefore ask is whether a sufficiently quick decay of the two-point functions (1) con-

versely also implies that the dynamics can be generated by a Hamiltonian with local interactions. If true, such a result would, on the one hand, make the connection with relativistic field theories stronger and hence provide valuable insight into the basic properties of quantum many-body systems. On the other hand, it would also have practical consequences for system identification and process tomography, which aims at reconstructing the Hamiltonian of a system from measurements of few observables. This constitutes an important task both from an experimental point of view but also in order to certify the functioning of quantum information processing devices. In fact, it was shown recently that the Hamiltonian may be reconstructed efficiently using local measurements, provided one has the promise that the interactions are indeed local [15–18].

In the following we show such a connection between correlations function and the locality of the Hamiltonian: If the two-point functions $C_{x,y}(t_1, t_0)$ decay at least as quickly as promised by state-of-the-art Lieb-Robinson bounds, then the interactions decay exponentially in space. More generally, our result shows a tight connection between the decay behavior of $C_{x,y}(t_1, t_0)$ for short times and the locality of the interactions (see Fig. 1).

Importantly, to reach our conclusion we only require knowledge about the decay of two-point functions of observables supported on single lattice sites. In practice, these are the relevant quantities in condensed-matter systems as scattering experiments effectively measure two-point correlation functions. However, exponentially decaying interactions in turn imply Lieb-Robinson bounds for all observables. Hence, as a by-product, we thus prove that Lieb-Robinson bounds for on-site observables already imply Lieb-Robinson bounds for bounded observables with arbitrary support without the need for a support-size-dependent correction (as a direct




FIG. 1. Two notions of locality. Left: Only constituents of the many-body system which are close by interact directly with each other (locality of interactions). Right: A local perturbation is not instantaneously detectable everywhere in the system (causal dynamics). Lieb-Robinson bounds show that locality of interactions implies causal dynamics. We prove that Lieb-Robinson bounds conversely imply (quasi-)local interactions.

expansion in terms of a product-operator-basis would require; see Appendix C).

II. SETUP AND LIEB-ROBINSON BOUNDS

Before we state our result, we set up some notation and review the statement of Lieb-Robinson bounds. We consider a lattice \mathbb{Z}^D and associate to each point $x \in \mathbb{Z}^D$ a d -dimensional Hilbert space \mathcal{H}_x . We choose the hypercubic lattice \mathbb{Z}^D purely for simplicity and definiteness—what follows is true for any regular lattice and with slight modifications also on more general graphs. Denote by \mathcal{A}_X the set (algebra) of local observables $\mathcal{A}_X \simeq \mathcal{B}(\mathcal{H}_X)$ supported in a region X . Here, $\mathcal{H}_X = \otimes_{x \in X} \mathcal{H}_x$ denotes the Hilbert space associated to X . If $Y \subseteq X$, then \mathcal{A}_Y is naturally embedded in \mathcal{A}_X by tensoring with identities on the complement of Y in X . We will mostly omit such identities throughout the paper. Conversely, we can consider the reduction map from \mathcal{A}_Λ to \mathcal{A}_X with $X \subseteq \Lambda$, which for any $A \in \mathcal{A}_X$ and $B \in \mathcal{A}_{X^c}$ with $X^c = \Lambda \setminus X$ is given by

$$\Gamma_X^\Lambda[AB] = \text{Tr}[B\mathbb{1}_{X^c}]A, \quad (2)$$

where, as usual, we identify $B \in \mathcal{A}_{X^c}$ with $B \otimes \mathbb{1}_X \in \mathcal{A}_\Lambda$, and $\mathbb{1}_{X^c} := \mathbb{1}_{X^c}/d_{X^c}$ corresponds to the maximally mixed state on X^c . In the case $X = \emptyset$, we simply obtain $\Gamma_\emptyset^\Lambda[B] = \mathbb{1}\text{Tr}[B\mathbb{1}_{X^c}]$.

It is customary to specify the dynamics of many-body systems through the notion of an (interaction) potential Φ . This is a function associating to each finite subset $X \subset \mathbb{Z}^D$ of lattice sites a bounded operator $\Phi(X) = \Phi(X)^\dagger \in \mathcal{A}_X$. For a large but finite subset $\Lambda \subset \mathbb{Z}^D$, we then write

$$H_\Lambda := \sum_{X \subseteq \Lambda} \Phi(X) \quad (3)$$

for the Hamiltonian on Λ induced by the potential. By $\tau_t^\Lambda(\cdot) := e^{iH_\Lambda t} \cdot e^{-iH_\Lambda t}$ we denote the unitary propagator generated by H_Λ , which gives rise to the Heisenberg dynamics of observables in \mathcal{A}_Λ .

The potentials appearing in actual models usually have the property that they only couple at most k spins directly, where k is some finite number independent of the system size. The prototypical examples are spin-spin interactions of the Ising-

or Heisenberg-type or Coulomb interactions for charged particles instead of spins. In both cases we have $k = 2$, but larger values of k appear, for example, as plaquette operators in lattice gauge theories, in the context of topological order, e.g., the toric code, or as effective interactions in perturbation theory [19–22]. We hence call a potential k body if

$$\Phi(X) = 0, \quad \text{if } |X| > k, \quad (4)$$

where $|X|$ denotes the number of lattice sites in X . We emphasize that this condition does not put any restrictions on the range or locality of the interaction. Even with $k = 2$, we, *a priori*, still allow for arbitrarily long-range interactions. Finally, we call a potential *exponentially decaying* if there exist positive constants K and a such that

$$\|\Phi(X)\| \leq K \exp(-a \text{diam}(X)), \quad (5)$$

for any region $X \subset \mathbb{Z}^D$. Here, $\text{diam}(X) = \sup_{x,y \in X} |x - y|$ denotes the diameter of X with $|x - y|$ the (graph-theoretical) distance on the lattice.

Even though a given potential Φ gives rise to a fixed time evolution τ_t^Λ for every Λ , the converse does not hold true: Firstly, the Hamiltonian which generates the time evolution τ_t^Λ is only fixed up to a shift of all the energies. Secondly, a given Hamiltonian can in general be decomposed in many different ways into a potential. The only physically relevant object in the end is the time-evolution operator τ_t^Λ .

Accordingly, the question we are addressing in this paper, namely, whether the interaction is exponentially decaying or not, corresponds to asking whether there exists *some* exponentially decaying potential giving rise to the same time evolution. Our result shows that given *some* k -body potential for which the two-point functions $C_{x,y}(t_1, t_0)$ of the corresponding time evolution show a certain decay behavior in space, then there also exists a k -body potential on Λ generating exactly the same time evolution, which has the same decay behavior in space. Importantly, the following theorem from Ref. [23], which is an improved version of the original bound from Ref. [1], states that any exponentially decaying k -body potential conversely gives rise to suitably bounded two-point functions. We state the theorem here in a form adjusted to our setup. To state the theorem, we introduce $d(X, Y) = \min_{x \in X, y \in Y} |x - y|$ as the distance between two subsets of the lattice.

Theorem 1 (Lieb-Robinson bounds). Given an exponentially decaying k -body potential, there exist constants $\mu, \nu, K > 0$ such that for any $\Lambda, X, Y \subset \mathbb{Z}^D$, we have

$$C_{X,Y}^\Lambda(t) := \max_{A,B} \frac{\|[\tau_t^\Lambda(A), B]\|}{\|A\| \|B\|} \leq 2 \min\{1, g(t)f(X, Y)\}, \quad (6)$$

where the maximization is over operators $A \in \mathcal{A}_X$ and $B \in \mathcal{A}_Y$ supported in X and Y , respectively, and $f(X, Y) = \min\{|X|, |Y|\} K \exp(-\mu d(X, Y))$. The function $g(t)$ is given by

$$g(t) = \begin{cases} \exp(\mu \nu |t|) - 1 & \text{if } d(X, Y) > 0 \\ \exp(\mu \nu |t|) & \text{if } d(X, Y) = 0. \end{cases} \quad (7)$$

Using the fact that the operator norm is invariant under unitary transformations and $[A(t_1), B(t_0)] = \tau_{t_0}^\Lambda([\tau_{t_1-t_0}^\Lambda(A), B])$, Lieb-Robinson bounds imply that the two-point correlation




FIG. 2. Density plot of Lieb-Robinson cone in the form given by Theorem 1. White: Contour of Lieb-Robinson cone as given by the standard exponential form of the Lieb-Robinson bounds. For large times the two agree, but for short times we see a significantly different behavior.

functions in (1) satisfy

$$|\langle \Psi [A(t_1), B(t_0)] | \Psi \rangle| \leq \min\{2, K(e^{\mu v|t_1 - t_0|} - 1)e^{-\mu|x-y|}\}$$

for any state $|\Psi\rangle$ and observables with $\|A\| = \|B\| = 1$. The bound is plotted in Fig. 2 and compared with the conventional bound $\exp[\mu(v|t_1 - t_0| - |x - y|)]$. For most purposes this latter bound is good enough, since it agrees for large times with the one we use here. However, for short times the two bounds differ markedly. In particular, the above version ensures that $C_{X,Y}^\Lambda(0) = 0$ for $d(X, Y) > 0$, and this will be crucial for deriving our result. The functions $g(t)$ and $f(X, Y)$ specify how sharp the Lieb-Robinson cone is localized. If the potential decays slower than exponentially, for example, as a power law, then they have to be replaced by functions that decay slower than exponentially. Indeed, recently a fair amount of literature is devoted to studying Lieb-Robinson bounds for long-range interactions, see, e.g., Refs. [4,24–32]. Furthermore, let us mention that the commutator form of the Lieb-Robinson bound given in (6) can be reformulated in two ways. Firstly, it gives a bound on the detectability of local excitations at distant lattice sites. Secondly, it provides a certificate on how well the evolution of a local observable can be approximated by an evolution under a truncated Hamiltonian. In particular, the first formulation can be interpreted as a bound on the information that can be communicated from one part of the lattice to another part using the given time evolution: it implies that if one tries to send information with nonvanishing capacity using a local encoding of the information at one point of the lattice, then there is a minimal time one has to wait before the information can be decoded at a different part of a lattice.

III. MAIN RESULT

Our main result is a direct converse to the Lieb-Robinson bounds stated in Theorem 1. Importantly, however, it only depends on the behavior of two-point functions for observables located at single lattice sites.

Theorem 2 (Converse to Lieb-Robinson bounds).

Consider a k -body potential and a finite subset $\Lambda \subset \mathbb{Z}^D$. Assume that there exists a function $h(t, x)$, with $h(0, x) = 0$ for $|x| > 0$ and differentiable in t , such that for any two points

$x \neq y \in \Lambda$ we have

$$C_{x,y}^\Lambda(t) := \max_{A,B} \frac{\|[\tau_t^\Lambda(A), B]\|}{\|A\|\|B\|} \leq h(t, |x - y|), \quad (8)$$

where the maximization is over single-site operators situated at x and y : $A \in \mathcal{A}_{\{x\}}, B \in \mathcal{A}_{\{y\}}$. Then there exists a constant m_k only depending on k , and a k -body potential $\hat{\Phi}_\Lambda$ on Λ , giving rise to the same time evolution τ_t^Λ on Λ , with

$$\|\hat{\Phi}_\Lambda(X)\| \leq m_k \partial h(0, \text{diam}(X)) \quad (9)$$

for all $X \subseteq \Lambda$ with $|X| \geq 2$.

Inserting $h(t, x) = g(t)f(x)$ with $f(x) = K \exp(-\mu x)$ and $g(t)$ from the Lieb-Robinson bounds into the theorem, the result shows directly that Lieb-Robinson bounds for two-point functions imply the exponential decay of the interaction potential $\hat{\Phi}_\Lambda$:

$$\|\hat{\Phi}_\Lambda(X)\| \leq m_k \mu v K \exp(-\mu \text{diam}(X)). \quad (10)$$

Moreover, since exponential decay of the interaction implies the Lieb-Robinson bounds in Theorem 1, we find that the decay of two-point functions of single-site observables implies a corresponding decay for arbitrary observables.

More generally, a uniform bound on the decay behavior of two-point functions directly translates into a decay behavior for the potential. For example, if $h(t, x) = g(t)f(x)$ with $f(x)$ being a power law and $g'(0) > 0$, which implies a power-law decay for the two-point functions, then our result also establishes a power-law behavior for the potential.

We may also infer strict locality of the Hamiltonian from the function h . Suppose, for example, that for $|x - y| > R$, $h(t, |x - y|) = t^\alpha f(|x - y|)$ with $\alpha > 1$. Then $\partial_t h[0, \text{diam}(X)] = 0$ if $\text{diam}(X) > R$ and therefore $\hat{\Phi}_\Lambda(X)$ vanishes. The same conclusion holds if h is of the form $h(t, x) = t^{x/R} f(x)$, which can be obtained, for example, from the Lieb-Robinson bound construction based on the commutativity graph of a given Hamiltonian [33]. In general, this shows that strict locality does not manifest itself in the decay of the correlation functions for a fixed time but how their time and space dependence interrelate.

Note that we only obtain bounds for potential terms with $|X| \geq 2$. That is, we do not get bounds on the norm of on-site potentials, such as magnetic fields. This is in accordance with the fact that Lieb-Robinson bounds can also be proven for Hamiltonians with unbounded on-site potentials [34,35]. On the other hand, there are Hamiltonians on bosonic lattices with local but unbounded interactions that allow for signaling at arbitrary speeds [36]. This shows that interactions have to be appropriately bounded to obtain Lieb-Robinson bounds for arbitrary observables, in agreement with our result.

Finally, let us consider the possibility of *strictly causal* behavior in lattice systems: Let us assume that $C_{x,y}^\Lambda(t)$ vanishes identically whenever $vt < |x - y|$ for some v , implying a strict light cone. Then our result directly implies that the interaction terms involving distinct lattice sites vanish identically as well. In other words, the only dynamics on a lattice system, which is strictly causal for all times, is one where the spins do not interact at all.

IV. THE PROOF

A central ingredient in our proof is the reduction map Γ_X^Λ , which we use to define $\hat{\Phi}_\Lambda$ as a certain *canonical form* of a given potential Φ (see also Refs. [37,38]). We say that a potential $\hat{\Phi}_\Lambda$ on Λ is in *canonical form* if

$$\Gamma_X^\Lambda[\hat{\Phi}_\Lambda(Y)] = 0 \quad (11)$$

unless $Y \subseteq X$. Given a Hamiltonian H_Λ on Λ , we can always decompose H_Λ into a potential in canonical form. Moreover, if H_Λ can be written in terms of a k -body potential, then the resulting canonical form will also be k body. These facts are collected in the following Lemma and proven in the Appendixes.

Lemma 3. Consider a k -body potential Φ with associated Hamiltonians H_Λ . Then for every finite subset $\Lambda \subseteq \mathbb{Z}^D$, the potential defined by

$$\hat{\Phi}_\Lambda(Z) = \sum_{X \subseteq Z} (-1)^{|Z|-|X|} \Gamma_X^\Lambda[H_\Lambda] \quad (12)$$

is a k -body potential in canonical form on Λ giving rise to the Hamiltonian H_Λ .

Given this lemma, we are now ready to give the essential steps of the proof of Theorem 2. To lighten the notation, we now fix Λ and omit it as a sub- or superscript from the potential, the reduction map, the propagator, and the quantity $C_{x,y}(t)$. Let us consider two points $x \neq y \in \Lambda$ and operators $A \in \mathcal{A}_{\{x\}}$ and $B \in \mathcal{A}_{\{y\}}$ with $\|A\| = \|B\| = 1$. Since $[A, B] = 0$, we then have

$$\lim_{\delta t \rightarrow 0} \frac{\|[\tau_{\delta t}(A), B]\|}{\delta t} = \|[[H_\Lambda, A], B]\|. \quad (13)$$

On the other hand, we can make use of the assumption on $C_{x,y}(t)$ to find

$$\begin{aligned} \lim_{\delta t \rightarrow 0} \frac{\|[\tau_{\delta t}(A), B]\|}{\delta t} &\leq \lim_{\delta t \rightarrow 0} \frac{C_{x,y}(\delta t)}{\delta t} \\ &\leq \lim_{\delta t \rightarrow 0} \frac{h(\delta t, |x-y|)}{\delta t} \\ &= \partial_t h(0, |x-y|), \end{aligned} \quad (14)$$

where we used $h(0, x) = 0$ in the last step. We thus finally get

$$\|[[H_\Lambda, A], B]\| \leq \partial_t h(0, |x-y|). \quad (15)$$

In the following let us write $\varepsilon_{x,y} := \partial_t h(0, |x-y|)$ and $x^c := \Lambda \setminus \{x\}$. Since (15) holds for all $B \in \mathcal{A}_{\{y\}}$ with $\|B\| = 1$, a Lemma from Ref. [39], whose proof we provide in the Supplemental Material (Lemma 4), implies

$$\|(1 - \Gamma_{y^c})[H_\Lambda], A\| \leq \varepsilon_{x,y}. \quad (16)$$

Here we used $\Gamma_{y^c}[H_\Lambda A] = \Gamma_{y^c}[H_\Lambda]A$ and $\Gamma_{y^c}[AH_\Lambda] = A\Gamma_{y^c}[H_\Lambda]$, which holds because A is supported within y^c . Since (16) again holds for all normalized $A \in \mathcal{A}_{\{x\}}$, we can use the same argument again to obtain

$$\|(1 - \Gamma_{x^c})[(1 - \Gamma_{y^c})[H_\Lambda]]\| \leq \varepsilon_{x,y}. \quad (17)$$

Now let $\hat{\Phi}$ be the potential in canonical form representing the Hamiltonian H_Λ . We then have $\Gamma_X[H_\Lambda] = H_X$ and $\hat{\Phi}(\emptyset) \propto 1$ (and the identity operator only appears in this term). Thus,

using $\Gamma_{x^c} \circ \Gamma_{y^c} = \Gamma_{x^c \cap y^c}$,

$$\begin{aligned} (1 - \Gamma_{x^c})[(1 - \Gamma_{y^c})[H_\Lambda]] &= H_\Lambda - H_{y^c} - H_{x^c} + H_{y^c \cap x^c} \\ &= \sum_{Z \ni x,y} \hat{\Phi}(Z). \end{aligned} \quad (18)$$

Therefore Eq. (17) says that for any pair of lattice sites $x \neq y$ we have

$$\left\| \sum_{Z' \ni x,y} \hat{\Phi}(Z') \right\| \leq \varepsilon_{x,y}. \quad (19)$$

We now fix a set Z and take two points $x, y \in Z$ such that $|x-y| = \text{diam}(Z)$ and make use of the fact that operators can only become smaller in norm under reductions to subsystems, i.e., for any $C = C^\dagger \in \mathcal{A}_\Lambda$ and $X \subseteq \Lambda$ we have

$$\|\Gamma_X[C]\| \leq \|\Gamma_X[\mathbb{1}]\| \|C\| = \|C\|, \quad (20)$$

where we used that Γ_X is a completely positive, unital map. This property allows us to restrict the bound in (19) to potential terms supported only within Z :

$$\left\| \sum_{\substack{Z' \ni x,y \\ Z' \subseteq Z}} \hat{\Phi}(Z') \right\| \leq \varepsilon_{x,y}. \quad (21)$$

Unfortunately, this bound does not yield a bound for each of the terms in the sum by $\varepsilon_{x,y}$ individually. This is because we cannot assume that the potential terms are positive operators due to our demand that the potential is in canonical form. We therefore now make use of the fact that $|Z| \leq k$ and use the inverse triangle inequality to obtain

$$\|\hat{\Phi}(Z)\| \leq \varepsilon_{x,y} + \left\| \sum_{\substack{Z' \ni x,y \\ Z' \subset Z}} \hat{\Phi}(Z') \right\|. \quad (22)$$

Since the sum on the right-hand side is over strict subsets of Z , each of the potential terms on the right-hand side couples at most $|Z| - 1$ spins. Clearly, the above equation gives $\|\hat{\Phi}(Z)\| \leq \varepsilon_{x,y}$ for $|Z| = 2$, since the only possible set is $Z = \{x, y\}$. Hence consider $|Z| = 3$. Then the norm on the right-hand side only contains one set Z' , which has cardinality $|Z'| = 2$. Thus $\|\hat{\Phi}_\Lambda(Z)\| \leq 2\varepsilon_{x,y}$. This reasoning allows us to recursively bound the potential terms $\Phi(Z)$ with $|Z| = m$ in terms of the those with $|Z| = m - 1$. Since we are assuming the interaction potential to be k body, the recursion stops after a constant number of steps. Hence there is a finite number m_k only depending on k such that

$$\|\hat{\Phi}_\Lambda(Z)\| \leq m_k \varepsilon_{x,y} = m_k \partial_t h(0, \text{diam}(Z)), \quad (23)$$

where we inserted the definition of $\varepsilon_{x,y}$ and used that $\text{diam}(Z) = |x-y|$. This finishes the proof of Theorem 2.

V. FERMIONIC LATTICE SYSTEMS

So far we only considered spin systems. In the Supplemental Material we generalize our result to fermionic lattice systems. The only difference to the case of spins is that we need to allow the observables A, B in the formulation of the

(assumed) bound in Theorem 2 to act not only on the respective lattice points but also on potentially present fermionic auxiliary systems. Such an auxiliary system would make no difference in the case of spins. However, due to the anticommutation relations of fermions and the parity superselection rule, they are important to consider in the case of a fermionic lattice system.

VI. DISCUSSION

In this work we have shown that whenever the two-point functions $C_{x,y}(t_1, t_0)$ show a certain decay behavior in space, then there also exists a potential generating exactly the same time evolution, which has the same decay behavior in space. In particular, this shows that Lieb-Robinson bounds in exponential form are equivalent to the exponential decay of interactions. Our result implies that whenever a system has *some* long-range interactions, then this can be diagnosed by measuring a two-point function in a suitable initial state. Furthermore, considering the reformulation of Lieb-Robinson bounds in terms of the detectability of local excitations, we see that there is a direct correspondence between the possibility of sending information through the lattice with finite capacity and the locality of the interactions. Let us conclude by discussing some open problems and avenues for further research.

First, we derived our results in the setting of continuous-time, unitary dynamics. One can ask whether similar results also hold for nonunitary dynamics or when one has discrete-time unitary dynamics. Suppose, for example, that one only has promises for the time evolution in discrete time steps δt , i.e., for the quantities $C_{A,B}^\Lambda(j \delta t)$ with j an integer. Then the dynamics is given by $\tau_{\delta t}(A) = UAU^*$ (omitting Λ for clarity), and U is a quasilocal unitary implementing the time evolution for a time step δt . Compared to large distances on the lattice, this time step is arbitrarily small. Going to very large distances, we could therefore imagine that on large scales the dynamics is then generated by a quasilocal Hamiltonian. Interestingly, according to Refs. [40–42], this is not the case; there exist *strictly local* unitary dynamics in discrete time which cannot be generated by quasilocal Hamiltonians, despite the fact that they can always be generated by local, finite depth quantum circuits [43]. In the Appendixes we present a very simple example of this behavior [44]. We thus conclude that it is indeed necessary to have promises for arbitrary short times. Whether our results generalize to continuous-time, dissipative, Markovian time evolution is an open problem. Let us remind the reader that Lieb-Robinson bounds also hold for dissipative systems, whose time evolution is generated by quasilocal Lindbladians [3,45–48]. Here, however, we would have to show that any long-ranged term in a Lindbladian leads to a violation of the corresponding Lieb-Robinson bound. It is not clear whether this is true for purely dissipative evolution. Certainly the techniques used in this work would have to be extended to prove such a result. If a counterexample indeed exists, it would show that there exists a close connection between unitarity of time evolution and locality of the dynamics in the sense of information propagation, which would be remarkable. We leave the investigation of this interesting point for future work.

Secondly, our results only need a promise over correlation functions of single-site observables for very short times and in this sense are rather friendly to experimental settings. However, to derive our results, we require a promise on the behavior of such correlation functions for arbitrary states $|\Psi\rangle$. In practice, such information is usually not available, since we do not prepare arbitrary states in real experiments. On the other hand, we are usually also not interested in the dynamics for arbitrary states but only in some subset of states, for example, from the low-energy sector. It is an interesting open problem to find out whether having a bound of the form

$$\langle \Psi | [\tau_t^\Lambda(A), B] | \Psi \rangle \leq \|A\| \|B\| g(t) \exp(-\mu|x-y|) \quad (24)$$

for all states $|\Psi\rangle$ in some experimentally relevant subset of states implies that the dynamics for this subset of states can be represented by an exponentially decaying interaction.

Thirdly, we have shown our results for spin models and fermionic lattice models. It would be interesting to investigate whether similar results as ours can be sensibly formulated and proven for the case of bosonic systems (restricting to bounded operators as observables). In this setting it is still an open problem to formulate and prove general Lieb-Robinson bounds for interacting bosonic systems (see, however, Refs. [34,35,49]). Indeed, without additional assumptions they cannot be proven due to the results in Ref. [36]. The assumption of reasonably bounded correlation functions naturally rules out interactions that lead to violations of Lieb-Robinson bounds. Methods similar to those we used in this work may thus help in identifying the general form of bosonic interactions for which Lieb-Robinson bounds can be proven.

ACKNOWLEDGMENTS

We would like to thank Terry Farrelly and Zoltán Zimborás for interesting comments and discussions. H.W. acknowledges support through the National Centre of Competence in Research, *Quantum Science and Technology* (QSIT). A.H.W. thanks the VILLUM FONDEN for its support with a Villum Young Investigator Grant (Grant No. 25452) and its support via the QMATH Centre of Excellence (Grant No. 10059).

APPENDIX A: PROOF OF LEMMA 3

In this section we provide the proof of Lemma 3, i.e., the statement that the potential defined by

$$\hat{\Phi}_\Lambda(Z) = \sum_{X \subseteq Z} (-1)^{|Z|-|X|} \Gamma_X^\Lambda[H_\Lambda] \quad (A1)$$

is a k -body potential in canonical form on Λ giving rise to the Hamiltonian H_Λ . This form of the potential is motivated by the *inclusion-exclusion* principle (see, e.g., Ref. [50]).

We need to show the following facts: (i) the potential gives rise to the same Hamiltonian H_Λ , (ii) the potential is in canonical form, and (iii) the potential is k body. Before we come to the details of this, we collect some simple facts about sums over subsets. The crucial fact that we will need is an elementary form of the inclusion-exclusion principle. It states

that for any finite set Z , we have

$$\sum_{X \subseteq Z} (-1)^{|X|} = \delta_{\emptyset}(Z) = \begin{cases} 0 & \text{if } Z \neq \emptyset \\ 1 & \text{if } Z = \emptyset. \end{cases} \quad (\text{A2})$$

This equality is very simple to prove: Clearly it is true for $Z = \emptyset$. Therefore consider any Z with $Z \neq \emptyset$ and let $z \in Z$ be an arbitrary point. Then

$$\begin{aligned} \sum_{X \subseteq Z} (-1)^{|X|} &= \sum_{X \subseteq Z, z \in X} (-1)^{|X|} + \sum_{X \subseteq Z, z \notin X} (-1)^{|X|} \\ &= \sum_{X \subseteq Z \setminus \{z\}} (-1)^{|X \cup \{z\}|} + \sum_{X \subseteq Z \setminus \{z\}} (-1)^{|X|} \\ &= \sum_{X \subseteq Z \setminus \{z\}} (-1 + 1)(-1)^{|X|} = 0. \end{aligned} \quad (\text{A3})$$

We will also use that for any function of sets f , we can write

$$\sum_{Z: X \subseteq Z \subseteq \Lambda} f(Z) = \sum_{Z \subseteq \Lambda \setminus X} f(Z \cup X). \quad (\text{A4})$$

In particular, we have

$$\sum_{Z: X \subseteq Z \subseteq \Lambda} (-1)^{|Z|} = \sum_{Z \subseteq \Lambda \setminus X} (-1)^{|Z| + |X|} = (-1)^{|X|} \delta_{\emptyset}(\Lambda \setminus X). \quad (\text{A5})$$

Finally, we require the identity,

$$\begin{aligned} \sum_{Z \subseteq \Lambda} \sum_{X \subseteq Z} f(X) g(X, Z) &= \sum_{X \subseteq \Lambda} \sum_{Z: X \subseteq Z \subseteq \Lambda} f(X) g(X, Z) \\ &= \sum_{X \subseteq \Lambda} f(X) \sum_{Z: X \subseteq Z \subseteq \Lambda} g(X, Z), \end{aligned} \quad (\text{A6})$$

which holds for arbitrary functions of sets f and g . With these ingredients in place, we now start with the proof of the Lemma.

We first show that the potential sums up to give the Hamiltonian,

$$\begin{aligned} \sum_{Z \subseteq \Lambda} \hat{\Phi}_{\Lambda}(Z) &= \sum_{Z \subseteq \Lambda} \sum_{X \subseteq Z} (-1)^{|Z| - |X|} \Gamma_X^{\Lambda} [H_{\Lambda}] \\ &= \sum_{X \subseteq \Lambda} \Gamma_X^{\Lambda} [H_{\Lambda}] \underbrace{\sum_{Z: X \subseteq Z \subseteq \Lambda} (-1)^{|Z| - |X|}}_{\delta_{\emptyset}(\Lambda \setminus X)} \\ &= \Gamma_{\Lambda}^{\Lambda} [H_{\Lambda}] = H_{\Lambda}, \end{aligned} \quad (\text{A7})$$

where we made use of (A2) and $\Gamma_{\Lambda}^{\Lambda} = 1$. To show that the potential is in canonical form, we consider $Y \subset \Lambda$ and $Z \subset \Lambda$ such that $Z \cap Y^c \neq \emptyset$ (i.e., $Z \setminus (Z \cap Y) \neq \emptyset$). We then find, using $\Gamma_Y^{\Lambda} \circ \Gamma_X^{\Lambda} = \Gamma_{X \cap Y}^{\Lambda}$,

$$\Gamma_Y^{\Lambda} [\hat{\Phi}(Z)] = \sum_{X \subseteq Z} (-1)^{|Z| - |X|} \Gamma_{X \cap Y}^{\Lambda} [H_{\Lambda}]. \quad (\text{A8})$$

Let us decompose Z into $Z \cap Y$ and $Z \setminus (Z \cap Y) \neq \emptyset$. For any function of sets f , a sum over subsets of Z can be decomposed as

$$\sum_{X \subseteq Z} f(X) = \sum_{X' \subseteq Z \setminus (Z \cap Y)} \sum_{X'' \subseteq Z \cap Y} f(X' \cup X''). \quad (\text{A9})$$

We thereby find

$$\begin{aligned} \Gamma_Y^{\Lambda} [\hat{\Phi}_{\Lambda}(Z)] &= \underbrace{\sum_{X' \subseteq Z \setminus (Z \cap Y)} (-1)^{|Z| - |X'|}}_{\delta_{\emptyset}(Z \setminus (Z \cap Y)) (-1)^{|Z|} = 0} \\ &\times \sum_{X'' \subseteq Z \cap Y} (-1)^{-|X''|} \Gamma_{X''}^{\Lambda} [H_{\Lambda}] = 0. \end{aligned} \quad (\text{A10})$$

Finally, let us show that $\hat{\Phi}_{\Lambda}$ is k body. To see this, consider Z with $|Z| > k$. We have to show that $\hat{\Phi}_{\Lambda}(Z) = 0$. First we use the original k -body potential Φ to write

$$\begin{aligned} \hat{\Phi}_{\Lambda}(Z) &= \sum_{X \subseteq Z} (-1)^{|Z| - |X|} \Gamma_X^{\Lambda} \left[\sum_{Y \subseteq \Lambda, |Y| \leq k} \Phi(Y) \right] \\ &= \sum_{Y \subseteq \Lambda, |Y| \leq k} \sum_{X \subseteq Z} (-1)^{|Z| - |X|} \Gamma_X^{\Lambda} [\Phi(Y)]. \end{aligned} \quad (\text{A11})$$

We now show that for each Y the corresponding term is zero. To do this we again split the sum over X into a sum over $Z \cap Y$ and $Z \setminus (Z \cap Y)$ to get

$$\begin{aligned} \sum_{X \subseteq Z} (-1)^{|Z| - |X|} \Gamma_X^{\Lambda} [\Phi(Y)] &= \sum_{X' \subseteq Z \setminus (Z \cap Y)} (-1)^{|Z| - |X'|} \\ &\times \sum_{X'' \subseteq Z \cap Y} (-1)^{|X''|} \Gamma_{X' \cup X''}^{\Lambda} [\Phi(Y)]. \end{aligned} \quad (\text{A12})$$

However, $\Phi(Y) \in \mathcal{A}_Y$, and the regions X' do not overlap Y by definition. Accordingly, we have $\Gamma_{X' \cup X''}^{\Lambda} [\Phi(Y)] = \Gamma_{X''}^{\Lambda} [\Phi(Y)]$. We thus find

$$\begin{aligned} \sum_{X \subseteq Z} (-1)^{|Z| - |X|} \Gamma_X^{\Lambda} [\Phi(Y)] &= \underbrace{\sum_{X' \subseteq Z \setminus (Z \cap Y)} (-1)^{|Z| - |X'|}}_{\delta_{\emptyset}(Z \setminus (Z \cap Y)) (-1)^{|Z|}} \\ &\times \sum_{X'' \subseteq Z \cap Y} (-1)^{|X''|} \Gamma_{X''}^{\Lambda} [\Phi(Y)] \\ &= \delta_{\emptyset}(Z \setminus (Z \cap Y)) (-1)^{|Z|} \\ &\times \sum_{X'' \subseteq Z \cap Y} (-1)^{|X''|} \Gamma_{X''}^{\Lambda} [\Phi(Y)]. \end{aligned} \quad (\text{A13})$$

Now, since $\Phi(Y)$ is zero for $|Y| > k$ the expression vanishes in this case. On the other hand, if $|Y| \leq k$, we can make use of our assumption that $|Z| > k$. This implies that $Z \setminus (Z \cap Y) \neq \emptyset$ and hence the δ function vanishes. This finishes the proof.

APPENDIX B: REDUCTION MAP AND COMMUTATOR

For completeness, we provide in this section a proof of a well-known result about the quality of local restrictions of an observable, given that it almost commutes with all observables on the complement of this restriction [39].

Lemma 4. Let $A \in \mathcal{A}_{\Lambda}$ and assume that $\|[A, B]\| \leq \varepsilon \|B\|$ for all $B \in \mathcal{A}_Y$ for some subset $Y \subset \Lambda$. Then

$$\|(1 - \Gamma_{Y^c})[A]\| \leq \varepsilon. \quad (\text{B1})$$

Proof. The reduction map $\Gamma_{Y^c}^\Lambda$ onto the complement of Y can be implemented as a twirl over the unitary group on Y , i.e.,

$$\Gamma_{Y^c}^\Lambda[A] = \int dU_Y (\mathbb{1}_{Y^c} \otimes U) A (\mathbb{1}_{Y^c} \otimes U^*), \quad (\text{B2})$$

where the integral runs over the normalized Haar measure on Y . Accordingly, the norm difference of A and its reduction to Y^c can be upper bounded by

$$\begin{aligned} \|(1 - \Gamma_{Y^c})[A]\| &= \left\| A - \int dU_Y (\mathbb{1}_{Y^c} \otimes U) A (\mathbb{1}_{Y^c} \otimes U^*) \right\| \\ &\leq \int dU_Y \| [A, (\mathbb{1}_{Y^c} \otimes U)] (\mathbb{1}_{Y^c} \otimes U^*) \| \leq \varepsilon, \end{aligned} \quad (\text{B3})$$

where in the last step we used the unitary invariance of the operator norm, our assumption on A , and the normalization of the Haar measure.

APPENDIX C: FROM LIEB-ROBINSON BOUNDS OF SINGLE-SITE OBSERVABLES TO GENERAL LIEB-ROBINSON BOUNDS

We have emphasized in the main text that our result implies that Lieb-Robinson bounds for single-site observables imply Lieb-Robinson bounds for observables with arbitrary support. Indeed, this follows immediately by combining our main result with Theorem 1. One may be tempted to think that this follows directly from 1, since any operator A supported on region X may be expressed as

$$A = \sum_{i_1, \dots, i_{|X|}} c_{i_1 \dots i_{|X|}} S_{i_1} \otimes \dots \otimes S_{i_{|X|}}, \quad (\text{C1})$$

where the S_i with $i = 1, \dots, d^2$ provide an operator basis for the space of operators acting on a single site of the lattice. Now using the Leibniz rule for the commutator and the triangle inequality, a bound for single-site observables implies a bound for more general observables. However, it is important to realize that this procedure yields an additional prefactor $d^{2|X|}$ on the right-hand side of the Lieb-Robinson bounds. For large regions X , for example, half the system, this prefactor diverges exponentially quickly for large systems, making the resulting bound useless. Our result, on the other hand, shows that a Lieb-Robinson bound for single-site observables implies a corresponding bound for arbitrary observables without such a prefactor.

APPENDIX D: FERMIONIC SYSTEMS

In the main text we discussed the setting of a lattice of spins. Here we explain how our results transfer to fermionic lattice systems. While the basic idea and proof strategy are exactly identical, we need to be a bit more careful due to the parity superselection rule for fermions. To do this, let us first briefly recapitulate the formalism of fermionic lattice systems (see, e.g., Ref. [38] for an introduction to the mathematical formalism of fermionic systems). For recent Lieb-Robinson bounds in the context of fermionic lattice systems, see Ref. [51], which also includes a detailed discussion

of the formalism of fermionic lattice systems from which we took inspiration. However, our discussion of the reduction map is quite different and might be of independent interest.

1. The basic setup of fermionic lattice systems

To every point of our lattice \mathbb{Z}^D , we associate an index set $I_x = \{1, \dots, d\}$. For every index $j \in I_x$ we define annihilation and creation operators $f_{x,j}, f_{x,j}^\dagger$ fulfilling the canonical anti-commutation relations:

$$\{f_{x,j}, f_{y,k}^\dagger\} = \delta_{x,y} \delta_{j,k} \mathbb{1}, \quad \{f_{x,j}, f_{y,k}\} = \{f_{x,j}^\dagger, f_{y,k}^\dagger\} = 0. \quad (\text{D1})$$

The algebra \mathcal{A}_X of local operators associated to a finite region X is then generated by arbitrary monomials of the $f_{x,j}, f_{y,k}^\dagger$ with $x, y \in X$. Every \mathcal{A}_X is isomorphic to the matrix algebra $M_{2^{|X|}}$ of $2^{|X|} \times 2^{|X|}$ matrices. A special role is played by the operator,

$$P_X = \prod_{x \in X} \prod_{j \in I_x} (-1)^{n_{x,j}}, \quad (\text{D2})$$

where $n_{x,j} := f_{x,j}^\dagger f_{x,j}$ is the *number operator* associated to the *mode* (x, j) . Due to the canonical anticommutation relations, the ordering of the operators in the definition of P_X does not matter. P_X is called the *parity operator* associated to the region X and fulfills $P_X^2 = \mathbb{1}$ and $P_X^\dagger = P_X$. Every $A \in \mathcal{A}_X$ can be decomposed into an *even* and *odd* part as

$$A^+ = \frac{A + P_X A P_X}{2}, \quad A^- = \frac{A - P_X A P_X}{2}. \quad (\text{D3})$$

Then $[A^+, P_X] = 0$ and $\{A^-, P_X\} = 0$. We write

$$\begin{aligned} \mathcal{A}_X &= \mathcal{A}_X^+ + \mathcal{A}_X^-, \\ \mathcal{A}_X^+ &:= \{A^+ | A \in \mathcal{A}_X\}, \quad \mathcal{A}_X^- := \{A^- | A \in \mathcal{A}_X\}. \end{aligned} \quad (\text{D4})$$

Due to the parity superselection rule of fermionic systems, all *physical observables* are even self-adjoint operators, and hence part of some \mathcal{A}_X^+ . We note that \mathcal{A}_X^+ is an algebra containing the identity, but \mathcal{A}_X^- is not an algebra, since $a, b \in \mathcal{A}_X^-$ implies $ab \in \mathcal{A}_X^+$. Alternatively, we can characterize \mathcal{A}_X^+ as being generated by monomials of an even number of the $f_{x,j}, f_{y,k}^\dagger$ with $x, y \in X$, while \mathcal{A}_X^- is generated by monomials of an odd number of the creation and annihilation operators. Due to the parity superselection rule, the algebra of physical operators \mathcal{A}_X^+ is *not* isomorphic to a full matrix algebra but isomorphic to a direct sum,

$$\begin{aligned} \mathcal{A}_X^+ &\simeq M_{2^{d|X|-1}} \oplus M_{2^{d|X|-1}} \\ &\simeq M_{2^{d|X|-1}} \otimes |0\rangle\langle 0| + M_{2^{d|X|-1}} \otimes |1\rangle\langle 1|, \end{aligned} \quad (\text{D5})$$

where the two direct summands correspond to the $+1$ and -1 eigenspaces of P_X , whose projectors we denote by P_X^\pm and which, in the above decomposition, are given by $P_X^+ = \mathbb{1} \otimes |0\rangle\langle 0|$, $P_X^- = \mathbb{1} \otimes |1\rangle\langle 1|$. Alternatively, we can write them as $P_X^\pm = (\mathbb{1} \pm P_X)/2$. Any operator $A \in \mathcal{A}_X$ can be decomposed as

$$A = (P_X^+ + P_X^-) A (P_X^+ + P_X^-) = \underbrace{A_{++} + A_{--}}_{A^+} + \underbrace{A_{+-} + A_{-+}}_{A^-} \quad (\text{D6})$$

with $A_{++} = P_X^+ A P_X^+$, $A_{+-} = P_X^+ A P_X^-$ and so forth. For any algebra \mathcal{A}_X there is a unique *even tracial state* ω^{Tr} , i.e., a

positive and normalized linear function, which vanishes on \mathcal{A}_X^- and fulfills

$$\omega^{\text{Tr}}[AB] = \omega^{\text{Tr}}[A]\omega^{\text{Tr}}[B], \quad (\text{D7})$$

for $A \in \mathcal{A}_{X_1}, B \in \mathcal{A}_{X_2}$ with $X_1 \cap X_2 = \emptyset$, and for any two $A, B \in \mathcal{A}_X$ we have $\omega^{\text{Tr}}[AB] = \omega^{\text{Tr}}[BA]$.

2. The reduction map

We also need to define a reduction map as in the case of spin systems. Here an additional difficulty arises in the setting of fermionic systems. To understand this issue, we should remember that in any given physical situation we should be allowed to add auxiliary systems to our description which do not take part in the time evolution of the system of interest. For spin systems, such "innocent bystanders" do not make any difference. However, for fermionic systems the situation is different due to the parity superselection rule and the canonical anticommutation relations. Anticipating this issue, let us therefore consider, next to the lattice of fermions, also another system S of $|S|$ fermions, described by associated annihilation and creation operators f_s^\dagger, f_s for $s = 1, \dots, S$ with $\{f_{x,j}^\dagger, f_s\} = 0$ and $\{f_s, f_{x,j}\} = 0$ for any $x \in \mathbb{Z}^D$. For $X \subseteq \Lambda \subset \mathbb{Z}^D$, we then introduce a reduction map

$$\Gamma_X^{\Lambda S} : \mathcal{A}_{\Lambda S} \rightarrow \mathcal{A}_{\Lambda S} \quad (\text{D8})$$

that maps physical operators on Λ to physical operators on X , $\Gamma_X^{\Lambda S} : \mathcal{A}_\Lambda^+ \rightarrow \mathcal{A}_X^+$, and fulfills for any $A \in \mathcal{A}_{\Lambda S}^+$ (with $X^c = \Lambda \setminus X$),

$$[\Gamma_X^{\Lambda S}[A], B] = 0 \quad \forall B \in \mathcal{A}_{X^c S}^+. \quad (\text{D9})$$

Importantly, we will see later that the map $\Gamma_X^{\Lambda S} : \mathcal{A}_\Lambda^+ \rightarrow \mathcal{A}_X^+$ does not depend on S as long as $|S| > 1$.

Let us now show how the reduction map is constructed. As shown in the proof of Lemma 4, in the spin case the reduction map can be written as

$$\Gamma_X^{\Lambda} [A] = \int dU_{X^c} (\mathbb{1}_X \otimes U) A (\mathbb{1}_X \otimes U^\dagger), \quad (\text{Spins}) \quad (\text{D10})$$

where dU_{X^c} denotes the normalized Haar measure on X^c . For the fermionic case, the construction we need is more involved. First note that we can decompose

$$\begin{aligned} \mathcal{A}_{X^c S}^+ &\simeq (\mathcal{A}_{X^c}^+ \otimes \mathcal{A}_S^+) \oplus (\mathcal{A}_{X^c}^- \otimes \mathcal{A}_S^-) \\ &\simeq M_{2^{d|X^c|+S-1}} \otimes |0\rangle\langle 0| + M_{2^{d|X^c|+S-1}} \otimes |1\rangle\langle 1|, \end{aligned} \quad (\text{D11})$$

where the last decomposition is with respect to the parity operator $P_{X^c S}$. The group $\mathcal{U}_{X^c S}^+$ of *even unitaries* on $X^c S$ thus corresponds to $U(2^{d|X^c|+S-1}) \oplus U(2^{d|X^c|+S-1})$. We now define the reduction maps as

$$\Gamma_X^{\Lambda S} [A] = \int dU U A U^\dagger, \quad (\text{D12})$$

where dU is the normalized Haar measure over $\mathcal{U}_{X^c S}^+$. Since this group corresponds to two copies of the group $U(2^{d|X^c|+S-1})$, one for each parity sector on $X^c S$, we can

rewrite this map as

$$\begin{aligned} \Gamma_X^{\Lambda S} [A] &= \int dU_+ \int dU_- (U_+ P_{X^c S}^+ + U_- P_{X^c S}^-) \\ &\quad \times A (U_+ P_{X^c S}^+ + U_- P_{X^c S}^-)^\dagger \\ &= \int dU_+ U_+ P_{X^c S}^+ A P_{X^c S}^+ U_+^\dagger \\ &\quad + \int dU_- U_- P_{X^c S}^- A P_{X^c S}^- U_-^\dagger \\ &\quad + \left[\int dU_+ U_+ P_{X^c S}^+ A P_{X^c S}^- \int dU_- U_-^\dagger + \text{H.c.} \right] \\ &= \int dU_+ U_+ P_{X^c S}^+ A P_{X^c S}^+ U_+^\dagger \\ &\quad + \int dU_- U_- P_{X^c S}^- A P_{X^c S}^- U_-^\dagger, \end{aligned} \quad (\text{D13})$$

where dU_\pm are normalized Haar measures over $U(2^{d|X^c|+S-1})$ and we used that $\int dU_\pm U_\pm = 0$. For the Haar measure we have

$$\begin{aligned} \int dU_+ U_+ P_{X^c S}^+ A P_{X^c S}^+ U_+^\dagger &= \frac{\text{Tr}_{X^c S} [A P_{X^c S}^+]}{2^{d|X^c|+|S|-1}} P_{X^c S}^+ \\ &= 2\omega_{X^c S}^{\text{Tr}} [A P_{X^c S}^+ P_{X^c S}^+], \end{aligned} \quad (\text{D14})$$

where $\omega_{X^c S}^{\text{Tr}}$ denotes the *partial* tracial state or maximally mixed state on $X^c S$. On operators of the form $A_X B_{X^c S}$, it acts as

$$\omega_{X^c S}^{\text{Tr}} [A_X B_{X^c S}] = A_X \omega^{\text{Tr}} [B_{X^c S}]. \quad (\text{D15})$$

Note that when an operator is supported in the support of the tracial state, we can omit the subscript. We summarize this result in the following lemma.

Lemma 5. The reduction map is a unital, completely positive map which can be written as

$$\Gamma_X^{\Lambda S} [A] = 2(\omega_{X^c S}^{\text{Tr}} [A P_{X^c S}^+] P_{X^c S}^+ + \omega_{X^c S}^{\text{Tr}} [A P_{X^c S}^-] P_{X^c S}^-). \quad (\text{D16})$$

It fulfills the following:

- (i) If $A, C \in \mathcal{A}_X^+$ and $B \in \mathcal{A}_{X^c S}^+$, we have $\Gamma_X^{\Lambda S} [ABC] = A \Gamma_X^{\Lambda S} [B] C$.
- (ii) If $B \in \mathcal{A}_{X^c S}^-$, then $\Gamma_X^{\Lambda S} [B] = 0$.
- (iii) If $|S| \geq 1$, the restriction to \mathcal{A}_Λ maps to \mathcal{A}_X^+ , $\Gamma_X^{\Lambda S} : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_X^+$, and the result does not depend on S .
- (iv) For $X \subseteq \Lambda, Y \subseteq \Lambda$ and when restricted to \mathcal{A}_Λ^+ , we have

$$\Gamma_X^{\Lambda S} \circ \Gamma_Y^{\Lambda S} = \Gamma_Y^{\Lambda S} \circ \Gamma_X^{\Lambda S} = \Gamma_{X \cap Y}^{\Lambda S}. \quad (\text{D17})$$

Proof. 1. Is obvious. 2. Follows from the fact that $\omega_{X^c S}^{\text{Tr}}$ is an even state and hence vanishes on odd operators. For 3. we make the following observations. First, if $A \in \mathcal{A}_\Lambda$ is in $\mathcal{A}_X \otimes \mathcal{A}_{X^c}^-$, then the reduction vanishes by 1. and 2. Therefore consider operators of the form $A_X A_{X^c}$ with $A_{X^c} \in \mathcal{A}_{X^c}^+$. By 1. we can forget about the A_X part and in the following consider $A \in \mathcal{A}_{X^c}^+$. We use that for $|S| \geq 1$, we have

$$\begin{aligned} P_{X^c S}^+ &= \frac{\mathbb{1} + P_{X^c S}}{2} = \frac{1}{2} [(P_{X^c}^+ + P_{X^c}^-)(P_S^+ + P_S^-) \\ &\quad + (P_{X^c}^+ - P_{X^c}^-)(P_S^+ - P_S^-)] = P_{X^c}^+ P_S^+ + P_{X^c}^- P_S^-, \end{aligned} \quad (\text{D18})$$

$$\begin{aligned}
P_{X^c S}^- &= \frac{\mathbb{1} - P_{X^c S}}{2} = \frac{1}{2}[(P_{X^c}^+ + P_{X^c}^-)(P_S^+ + P_S^-) \\
&\quad - (P_{X^c}^+ - P_{X^c}^-)(P_S^+ - P_S^-)] = P_{X^c}^+ P_S^- + P_{X^c}^- P_S^+. \quad (D19)
\end{aligned}$$

Since $P_{X^c}^+ P_{X^c S}^\pm = P_{X^c}^+ P_S^\pm$ and $P_{X^c}^- P_{X^c S}^\pm = P_{X^c}^- P_S^\mp$, we have

$$AP_{X^c S}^\pm = A(P_{X^c}^+ + P_{X^c}^-)P_{X^c S}^\pm = A(P_{X^c}^+ P_S^\pm + P_{X^c}^- P_S^\mp). \quad (D20)$$

Using that $\omega_{X^c S}^{\text{Tr}}$ is a product state and $\omega_S^{\text{Tr}}[P_S^\pm] = 1/2$, we find

$$\begin{aligned}
\omega_{X^c S}^{\text{Tr}}[AP_{X^c S}^+] &= \omega_{X^c}^{\text{Tr}}[AP_{X^c}^+] \omega_S^{\text{Tr}}[P_S^+] + \omega_{X^c}^{\text{Tr}}[AP_{X^c}^-] \omega_S^{\text{Tr}}[P_S^-] \\
&= \frac{1}{2} \omega_{X^c}^{\text{Tr}}[A(P_{X^c}^+ + P_{X^c}^-)] = \frac{1}{2} \omega_{X^c}^{\text{Tr}}[A] \quad (D21)
\end{aligned}$$

and

$$\begin{aligned}
\omega_{X^c S}^{\text{Tr}}[AP_{X^c S}^-] &= \omega_{X^c}^{\text{Tr}}[AP_{X^c}^+] \omega_S^{\text{Tr}}[P_S^-] + \omega_{X^c}^{\text{Tr}}[AP_{X^c}^-] \omega_S^{\text{Tr}}[P_S^+] \\
&= \frac{1}{2} \omega_{X^c}^{\text{Tr}}[A(P_{X^c}^+ + P_{X^c}^-)] = \frac{1}{2} \omega_{X^c}^{\text{Tr}}[A]. \quad (D22)
\end{aligned}$$

Consequently,

$$\Gamma_X^{\Lambda S}[A] = \omega_{X^c}^{\text{Tr}}[A](P_{X^c S}^+ + P_{X^c S}^-) = \omega_{X^c}^{\text{Tr}}[A] \mathbb{1} \in \mathcal{A}_X^+. \quad (D23)$$

Finally, we need to show 4. We consider any operators $A \in \mathcal{A}_{(X \cup Y)^c}$, $B \in \mathcal{A}_{X \setminus Y}$, $C \in \mathcal{A}_{Y \setminus X}$, $D \in \mathcal{A}_{X \cap Y}$ such that $ABCD \in \mathcal{A}_\Lambda^+$. Then

$$\Gamma_{X \cap Y}^{\Lambda S}[ABCD] = \Gamma_{X \cap Y}^{\Lambda S}[ABC]D = \omega^{\text{Tr}}[ABC]D. \quad (D24)$$

Similarly, we have

$$\begin{aligned}
\Gamma_X^{\Lambda S} \circ \Gamma_Y^{\Lambda S}[ABCD] &= \Gamma_X^{\Lambda S}[\Gamma_Y^{\Lambda S}[AB]CD] \\
&= \Gamma_X^{\Lambda S}[C]D \omega^{\text{Tr}}[AB] \\
&= D \omega^{\text{Tr}}[C] \omega^{\text{Tr}}[AB] = \omega^{\text{Tr}}[ABC]D, \quad (D25)
\end{aligned}$$

where we used that ω^{Tr} is a product state. For the reversed order,

$$\begin{aligned}
\Gamma_Y^{\Lambda S} \circ \Gamma_X^{\Lambda S}[ABCD] &= \pm \Gamma_Y^{\Lambda S}[B \Gamma_X^{\Lambda S}[AC]D] \\
&= \pm \Gamma_Y^{\Lambda S}[B]D \omega^{\text{Tr}}[AC] = \pm D \omega^{\text{Tr}}[ACB], \quad (D26)
\end{aligned}$$

where the negative sign only appears if both A and B are odd, in which case the expression vanishes. We thus conclude that all three expressions vanish if any of the A, B, C are odd and otherwise give the same result. Hence they always give the same result.

We emphasize that $\Gamma_X^{\Lambda S}$ does not map all observables to observables supported in X . For example, the parity operator $P_{X^c S}$ is mapped to itself. This is easy to see from the definition via the average over even unitaries but can also be seen from the above form of the map using $P_{X^c S} = P_{X^c S}^+ - P_{X^c S}^-$:

$$\begin{aligned}
\Gamma_X^{\Lambda S}[P_{X^c S}^+ - P_{X^c S}^-] &= 2(\omega_{X^c S}^{\text{Tr}}[P_{X^c S}^+] P_{X^c S}^+ - \omega_{X^c S}^{\text{Tr}}[P_{X^c S}^-] P_{X^c S}^-) \\
&= 2(\frac{1}{2} P_{X^c S}^+ - \frac{1}{2} P_{X^c S}^-) = P_{S^c X}. \quad (D27)
\end{aligned}$$

We close the discussion of the reduction map for fermionic lattice systems by proving the analog of Lemma 4.

Lemma 6. Suppose $|S| \geq 1$ and let $A \in \mathcal{A}_\Lambda^+$ be such that

$$\| [A, B] \| \leq \varepsilon \| B \| \quad \forall B \in \mathcal{A}_{X^c S}^+. \quad (D28)$$

Then $\Gamma_X^{\Lambda S}[A] \in \mathcal{A}_X^+$ and

$$\| A - \Gamma_X^{\Lambda S}[A] \| \leq \varepsilon. \quad (D29)$$

Proof. That $\Gamma_X^{\Lambda S}[A] \in \mathcal{A}_X^+$ for $|S| \geq 1$ was shown in the previous lemma. The second claim follows as in the proof of Lemma 4:

$$\begin{aligned}
\| A - \Gamma_X^{\Lambda S}[A] \| &\leq \int_{\mathcal{U}_{X^c}^+} dU \| AUU^\dagger - UAU^\dagger \| \\
&= \int_{\mathcal{U}_{X^c}^+} dU \| [A, U] \| \leq \varepsilon, \quad (D30)
\end{aligned}$$

where we used the triangle inequality and normalization of the Haar measure in the first step and unitary invariance of the norm in the second step.

As shown in Lemma 5, the map $\Gamma_X^{\Lambda S} : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_X^+$ does not in fact depend on S . We can hence define

$$\Gamma_X^\Lambda : \mathcal{A}_\Lambda \rightarrow \mathcal{A}_X^+, \quad A \mapsto \Gamma_X^{\Lambda S}[A], \quad \text{for some } |S| \geq 1, \quad (D31)$$

and this map fulfills all the relevant properties that we needed in the case of spins.

3. Dynamics and the fermionic result

Similarly as for spin systems, we describe dynamics by a potential which now associates *even* operators $\Phi(X) = \Phi(X)^\dagger \in \mathcal{A}_X^+$ to finite regions X of the lattice. Therefore the Hamiltonian H_Λ is even and the induced time evolution τ_t^Λ maps \mathcal{A}_Λ^\pm to \mathcal{A}_Λ^\pm as required. Analogously to the spin setting, we call such a potential k body if $\Phi(X) = 0$ for $|X| > k$. As for spin systems, exponentially decaying k -body potentials lead to Lieb-Robinson bounds also in the case of fermions (see, e.g., Ref. [51] for a recent account). Similarly as for the reduction map, we explicitly take into account the possibility of fermionic auxiliary systems S_1, S_2 that are left invariant under the time evolution. This allows us to recover our result also in the case of fermionic lattice systems:

Theorem 7 (Fermionic converse to Lieb-Robinson bounds). Consider a fermionic lattice system with k -body potential and assume that there exists a function $h(t, x)$ with $h(0, x) = 0$, with $h(0, x) = 0$ for $|x| > 0$ and differentiable in t , such that for any finite subset $\Lambda \subset \mathbb{Z}^D$, any two points $x \neq y \in \mathbb{Z}^D$ and any two fermionic systems S_1, S_2 we have

$$C_{x,y}^\Lambda(t) := \max_{A,B} \frac{\| [\tau_t^\Lambda(A), B] \|}{\| A \| \| B \|} \leq h(t, |x - y|), \quad (D32)$$

where the maximization is over physical operators on $\{x\} \cup S_1$ and $\{y\} \cup S_2$: $A \in \mathcal{A}_{\{x\} \cup S_1}^+$, $B \in \mathcal{A}_{\{y\} \cup S_2}^+$. Then there exists a constant m_k only depending on k , and a k -body potential $\hat{\Phi}_\Lambda$ on Λ , giving rise to the same time evolution τ_t^Λ , with

$$\| \hat{\Phi}_\Lambda(X) \| \leq m_k \partial_t h(0, \text{diam}(X)) \quad (D33)$$

for all $X \subseteq \Lambda$ with $|X| \geq 2$.

Proof. The proof is essentially identical to the spin case using the fermionic reduction map defined in (D31).

APPENDIX E: STRICTLY LOCAL, DISCRETE-TIME, UNITARY DYNAMICS THAT CANNOT BE GENERATED LOCALLY IN CONTINUOUS TIME

In this section we briefly describe an example of discrete-time, unitary dynamics that is strictly local but cannot be generated in continuous time with a (quasi-)local Hamiltonian, namely, the shift. We discuss the example in first-quantized language but remark that it can be second-quantized to yield a fermionic lattice model. Indeed, any local second-quantized Hamiltonian would necessarily have to reproduce the first-quantized Hamiltonian in the single-particle sector, which as we argue here has to violate Lieb-Robinson bounds in continuous time. The model describes a particle hopping on a one-dimensional line, which we can view as \mathbb{Z} , with Hilbert space given by $l^2(\mathbb{Z})$. An orthonormal basis is given by associating one basis vector $|x\rangle$ with each lattice site $x \in \mathbb{Z}$, which is interpreted as the particle being located at the site x . The unitary dynamics that we consider simply amounts to shifting the particle by one site:

$$U|x\rangle = |x+1\rangle. \quad (\text{E1})$$

In other words, $U = \exp(iP)$, where P is the lattice momentum, which generates translations. It is clear that this discrete-time dynamics is strictly local. However, the matrix elements of the lattice momentum in position space fulfill (for $x \neq y$)

$$|\langle y|P|x\rangle| = \frac{1}{|y-x|}. \quad (\text{E2})$$

Therefore, if we choose $A = |x\rangle\langle x|$, $B = |y\rangle\langle y|$ for $x \neq y$, we find that for small times t ,

$$\| [e^{iPt} A e^{-iPt}, B] \| = \frac{t}{|x-y|} + \mathcal{O}(t^2). \quad (\text{E3})$$

Thus while the dynamics is strictly local for discrete times, it becomes long-ranged for intermediate times.

We remark that this is not a complete argument since the choice of P as the logarithm of $\exp(iP)$ is not unique and there could in principle exist a local choice. However, it follows from the index theory for strictly local unitary evolutions that no such choice exists, see [52]. Also see [53] for examples of quantum cellular automata that cannot be generated from a local time-independent evolution.

-
- [1] E. H. Lieb and D. W. Robinson, The finite group velocity of quantum spin systems, *Commun. Math. Phys.* **28**, 251 (1972).
- [2] M. B. Hastings, Lieb-Schultz-Mattis in higher dimensions, *Phys. Rev. B* **69**, 104431(2004).
- [3] M. B. Hastings, Locality in Quantum and Markov Dynamics on Lattices and Networks, *Phys. Rev. Lett.* **93**, 140402 (2004).
- [4] M. B. Hastings and T. Koma, Spectral gap and exponential decay of correlations, *Commun. Math. Phys.* **265**, 781 (2006).
- [5] B. Nachtergaele and R. Sims, Lieb-Robinson bounds and the exponential clustering theorem, *Commun. Math. Phys.* **265**, 119 (2006).
- [6] M. B. Hastings and S. Michalakis, Quantization of Hall conductance for interacting electrons on a torus, *Commun. Math. Phys.* **334**, 433 (2015).
- [7] S. Bachmann, W. D. Roeck, and M. Fraas, The adiabatic theorem in a quantum many-body setting, in *Analytic Trends in Mathematical Physics*, edited by H. Abdul-Rahman, R. Sims, and A. Young, Contemporary Mathematics Vol. 741 (American Mathematical Society, 2020), pp. 1-21.
- [8] B. Nachtergaele and R. Sims, A multi-dimensional Lieb-Schultz-Mattis theorem, *Commun. Math. Phys.* **276**, 437 (2007).
- [9] W. F. Wreszinski, Goldstone's theorem for quantum spin systems of finite range, *J. Math. Phys.* **17**, 109 (1976).
- [10] W. F. Wreszinski, Charges and symmetries in quantum theories without locality, *Fortschr. Phys.* **35**, 379 (1987).
- [11] M. B. Hastings, An area law for one-dimensional quantum systems, *J. Stat. Mech.: Theory Exp.* (2007) P08024.
- [12] M. Cheneau, P. Barmettler, D. Poletti, M. Endres, P. Schauß, T. Fukuhara, C. Gross, I. Bloch, C. Kollath, and S. Kuhr, Light-cone-like spreading of correlations in a quantum many-body system, *Nature (London)* **481**, 484 (2012).
- [13] P. Jurcevic, B. P. Lanyon, P. Hauke, C. Hempel, P. Zoller, R. Blatt, and C. F. Roos, Quasiparticle engineering and entanglement propagation in a quantum many-body system, *Nature (London)* **511**, 202 (2014).
- [14] P. Richerme, Z.-X. Gong, A. Lee, C. Senko, J. Smith, M. Foss-Feig, S. Michalakis, A. V. Gorshkov, and C. Monroe, Non-local propagation of correlations in quantum systems with long-range interactions, *Nature (London)* **511**, 198 (2014).
- [15] E. Bairey, I. Arad, and N. H. Lindner, Learning a Local Hamiltonian from Local Measurements, *Phys. Rev. Lett.* **122**, 020504 (2019).
- [16] Z. Li, L. Zou, and T. H. Hsieh, Hamiltonian Tomography via Quantum Quench, *Phys. Rev. Lett.* **124**, 160502 (2020).
- [17] E. Bairey, C. Guo, D. Poletti, N. H. Lindner, and I. Arad, Learning the dynamics of open quantum systems from their steady states, *New J. Phys.* **22**, 032001 (2020).
- [18] A. Anshu, S. Arunachalam, T. Kuwahara, and M. Soleimanifar, *Sample-efficient learning of quantum many-body systems*, in *IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)* (IEEE, Piscataway, NJ, 2020), pp. 685–691.
- [19] A. Kitaev, Fault-tolerant quantum computation by anyons, *Ann. Phys.* **303**, 2 (2003).
- [20] J. Kempe, A. Kitaev, and O. Regev, The complexity of the local Hamiltonian problem, in *FSTTCS 2004: Foundations of Software Technology and Theoretical Computer Science* (Springer, Berlin, Heidelberg, 2004), pp. 372–383.
- [21] A. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys.* **321**, 2 (2006).
- [22] S. P. Jordan and E. Farhi, Perturbative gadgets at arbitrary orders, *Phys. Rev. A* **77**, 062329 (2008).
- [23] B. Nachtergaele and R. Sims, Locality estimates for quantum spin systems, in *New Trends in Mathematical Physics, Selected Contributions of the XVth International Congress on Mathematical Physics*, edited by S. Sidoravicius (Springer Verlag, Berlin, 2009), pp. 591–614.

- [24] J. Eisert, M. van den Worm, S. R. Manmana, and M. Kastner, Breakdown of Quasilocality in Long-Range Quantum Lattice Models, *Phys. Rev. Lett.* **111**, 260401 (2013).
- [25] P. Hauke and L. Tagliacozzo, Spread of Correlations in Long-Range Interacting Quantum Systems, *Phys. Rev. Lett.* **111**, 207202 (2013).
- [26] Z.-X. Gong, M. Foss-Feig, S. Michalakis, and A. V. Gorshkov, Persistence of Locality in Systems with Power-Law Interactions, *Phys. Rev. Lett.* **113**, 030602 (2014).
- [27] M. Foss-Feig, Z.-X. Gong, C. W. Clark, and A. V. Gorshkov, Nearly Linear Light Cones in Long-Range Interacting Quantum Systems, *Phys. Rev. Lett.* **114**, 157201 (2015).
- [28] T. Matsuta, T. Koma, and S. Nakamura, Improving the Lieb–Robinson bound for long-range interactions, *Ann. Henri Poincaré* **18**, 519 (2017).
- [29] T. Kuwahara and K. Saito, Strictly Linear Light Cones in Long-Range Interacting Systems of Arbitrary Dimensions, *Phys. Rev. X* **10**, 031010 (2020).
- [30] C.-F. Chen and A. Lucas, Finite Speed of Quantum Scrambling with Long Range Interactions, *Phys. Rev. Lett.* **123**, 250605 (2019).
- [31] D. V. Else, F. Machado, C. Nayak, and N. Y. Yao, Improved Lieb-Robinson bound for many-body Hamiltonians with power-law interactions, *Phys. Rev. A* **101**, 022333 (2020).
- [32] M. C. Tran, C.-F. Chen, A. Ehrenberg, A. Y. Guo, A. Deshpande, Y. Hong, Z.-X. Gong, A. V. Gorshkov, and A. Lucas, Hierarchy of Linear Light Cones with Long-Range Interactions, *Phys. Rev. X* **10**, 031009 (2020).
- [33] Z. Wang and K. R. A. Hazzard, Tightening the Lieb-Robinson bound in locally interacting systems, *PRX Quantum* **1**, 010303 (2020).
- [34] B. Nachtergaele, H. Raz, B. Schlein, and R. Sims, Lieb-Robinson bounds for harmonic and anharmonic lattice systems, *Commun. Math. Phys.* **286**, 1073 (2009).
- [35] B. Nachtergaele and R. Sims, On the dynamics of lattice systems with unbounded on-site terms in the Hamiltonian, [arXiv:1410.8174](https://arxiv.org/abs/1410.8174).
- [36] J. Eisert and D. Gross, Supersonic Quantum Communication, *Phys. Rev. Lett.* **102**, 240501 (2009).
- [37] H. Araki, Conditional expectations relative to a product state and the corresponding standard potentials, *Commun. Math. Phys.* **246**, 113 (2004).
- [38] O. Bratteli and D. W. Robinson, Models in quantum statistical mechanics, in *Operator Algebras and Quantum Statistical Mechanics, Equilibrium States* (Springer-Verlag, Berlin, Heidelberg, 1981), Vol. 2.
- [39] S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims, Automorphic equivalence within gapped phases of quantum lattice systems, *Commun. Math. Phys.* **309**, 835 (2012).
- [40] T. Farrelly, A review of quantum cellular automata, *Quantum* **4**, 368 (2000).
- [41] Z. Zimborás, T. C. Farrelly, S. Farkas, and L. Masanes, Does causal dynamics imply local interactions? [arXiv:2006.10707](https://arxiv.org/abs/2006.10707).
- [42] D. Ranard, M. Walter, and F. Witteveen, A converse to Lieb-Robinson bounds in one dimension using index theory, [arXiv:2012.00741](https://arxiv.org/abs/2012.00741).
- [43] P. Arrighi, V. Nesme, and R. Werner, Unitarity plus causality implies localizability, *J. Comput. Syst. Sci.* **77**, 372 (2011).
- [44] We thank Terry Farrelly and Zoltán Zimborás for reminding us of this example.
- [45] D. Poulin, Lieb-Robinson Bound and Locality for General Markovian Quantum Dynamics, *Phys. Rev. Lett.* **104**, 190401 (2010).
- [46] B. Nachtergaele, A. Vershynina, and V. A. Zagrebnov, Lieb-Robinson bounds and existence of the thermodynamic limit for a class of irreversible quantum dynamics, *AMS Cont. Math.* **552**, 161 (2011).
- [47] T. Barthel and M. Kliesch, Quasilocality and Efficient Simulation of Markovian Quantum Dynamics, *Phys. Rev. Lett.* **108**, 230504 (2012).
- [48] M. Kliesch, C. Gogolin, and J. Eisert, Lieb-Robinson bounds and the simulation of time evolution of local observables in lattice systems, in *Many-Electron Approaches in Physics, Chemistry and Mathematics* (Springer International Publishing, Cham, Switzerland, 2014), pp. 301–318.
- [49] B. Nachtergaele, B. Schlein, R. Sims, S. Starr, and V. Zagrebnov, On the existence of the dynamics for anharmonic quantum oscillator systems, *Rev. Math. Phys.* **22**, 207 (2010).
- [50] I. M. Gessel and R. P. Stanley, Algebraic enumeration, in *Handbook of Combinatorics* (MIT Press, Cambridge, MA, 1996), Vol. 2, pp. 1021–1063.
- [51] B. Nachtergaele, R. Sims, and A. Young, Lieb-Robinson bounds, the spectral flow, and stability of the spectral gap for lattice fermion systems, in *Mathematical Problems in Quantum Physics*, edited by F. Bonetto, D. Borthwick, E. Harrell, and M. Loss, Contemporary Mathematics Vol. 717 (American Mathematical Society, Providence, RI, 2018), pp. 93–115.
- [52] D. Gross, V. Nesme, H. Vogts, and R. F. Werner, Index theory of one dimensional quantum walks and cellular automata, *Commun. Math. Phys.* **310**, 419 (2012).
- [53] Z. Zimborás, T. Farrelly, S. Farkas, and L. Masanes, Does causal dynamics imply local interactions? [arXiv:2006.10707](https://arxiv.org/abs/2006.10707).