





# Thermally enhanced Majorana-mediated spin transport in the Kitaev model

Hirokazu Taguchi , Yuta Murakami , and Akihisa Koga 

*Department of Physics, Tokyo Institute of Technology, Meguro, Tokyo 152-8551, Japan*

 (Received 29 January 2022; revised 15 March 2022; accepted 16 March 2022; published 28 March 2022)

We study how stable the Majorana-mediated spin transport in a quantum spin Kitaev model is against thermal fluctuations. Using the time-dependent thermal pure quantum state method, we examine finite-temperature spin dynamics in the Kitaev model. The model exhibits two characteristic temperatures,  $T_L$  and  $T_H$ , which correspond to energy scales of the local flux and the itinerant Majorana fermion, respectively. At low temperatures ( $T \ll T_L$ ), an almost flux-free state is realized, and the spin excitation propagates in a similar way to that for the ground state. Namely, after the magnetic pulse is introduced at one of the edges, the itinerant Majorana fermions propagate the spin excitations even through the quantum spin liquid state region, and oscillations in the spin moment appear in the other edge with a tiny magnetic field. When  $T \sim T_L$ , larger oscillations in the spin moments are induced in the other edge, compared with the results at the ground state. At higher temperatures, excited  $Z_2$  fluxes disturb the coherent motion of the itinerant Majorana fermions, which suppresses the spin propagation. Our results demonstrate a crucial role of thermal fluctuations in the Majorana-mediated spin transport.

DOI: [10.1103/PhysRevB.105.125137](https://doi.org/10.1103/PhysRevB.105.125137)

## I. INTRODUCTION

Recently, spin transport has been attracting much interest. One example is the spin current induced by a polarized electric current in the ferromagnetic metals [1–8]. Another example is the spin current in the magnetic insulators, where magnons carry spins without the electric current [9–12]. In both cases, the spin current flows in materials with magnetic orders. On the other hand, it has been revealed that the spin transport is also realized in quantum spin liquids (QSLs) [13–15], where no magnetic order is realized due to strong quantum fluctuations [16–22]. One of the typical examples is provided by an antiferromagnetic  $S = 1/2$  Heisenberg chain. The anisotropic negative spin Seebeck effect in a candidate material,  $\text{Sr}_2\text{CuO}_3$ , indicates spin current mediated by spinons [23], which are magnetic elementary excitations in this system.

Another interesting playground for QSLs is given by the Kitaev model [24], which is composed of direction-dependent Ising interaction between  $S = 1/2$  spins on the honeycomb lattice. In the model, quantum spins are fractionalized into itinerant Majorana fermions and local fluxes due to quantum many-body effects. The itinerant Majorana fermions have been observed as a half-quantized plateau in thermal quantum Hall experiments [25,26] in the candidate  $\alpha\text{-RuCl}_3$  [27]. Furthermore, it has been reported that the itinerant Majorana fermions play a crucial role for spin transport without spin oscillations [13–15]. It is known that Majorana and flux excitations have distinct energy scales, which leads to interesting thermodynamic properties such as double peaks in the specific heat and a plateau in the entropy [28–32]. Therefore it is highly desired to clarify how stable such Majorana-related phenomena are against thermal fluctuations. This should be important to realize spintronics devices with Majorana fermions.

To answer this question, we deal with the Kitaev model with edges and consider the spin transport at finite temperatures. By means of the time-dependent thermal pure quantum (TPQ) state method [33–35], we examine the dynamics of the system after the magnetic pulse is introduced at one of the edges. Then, we discuss how thermal fluctuations affect the Majorana-mediated spin transport.

The paper is organized as follows. In Sec. II, we introduce the Kitaev model on the honeycomb lattice and explain the time-dependent TPQ method. In Sec. III, we discuss how stable the spin propagation in the Kitaev model is against thermal fluctuations. A summary is given in the last section.

## II. MODEL AND METHOD

We consider the Kitaev model on a two-dimensional honeycomb lattice, which is given by the following Hamiltonian:

$$H_K = -J \sum_{\langle i,j \rangle_x} S_i^x S_j^x - J \sum_{\langle i,j \rangle_y} S_i^y S_j^y - J \sum_{\langle i,j \rangle_z} S_i^z S_j^z, \quad (1)$$

where  $\langle i, j \rangle_\mu$  indicates the nearest-neighbor pair on the  $\mu$  ( $= x, y, z$ ) bonds. The  $x, y$ , and  $z$  bonds are shown as green, red, and blue lines in Fig. 1.  $S_i^\mu$  is the  $\mu$  component of an  $S = 1/2$  spin operator at the  $i$ th site, and  $J$  is the exchange coupling between the nearest-neighbor spins.

An important feature is that the Kitaev model has local conserved quantities. The operator  $W_p$  is defined as

$$W_p = 2^6 \cdot S_{p_1}^x S_{p_2}^y S_{p_3}^z S_{p_4}^x S_{p_5}^y S_{p_6}^z, \quad (2)$$

where  $p_i$  ( $i = 1, 2, \dots, 6$ ) is the site in the plaquette  $p$  [see Fig. 1(b)]. Since  $[H_K, W_p] = 0$ ,  $[W_p, W_q] = 0$  ( $p \neq q$ ), and  $W_p^2 = 1$ , the operator  $W_p$  is a  $Z_2$  local conserved quantity. Each eigenstate of the Kitaev model is classified by the Hilbert space specified by a set  $\{w_p\}$ , where  $w_p$  ( $= \pm 1$ ) is the eigenvalue of  $W_p$ . Since the spin operator changes the sign of the




FIG. 1. (a) Kitaev model with armchair edges. Green, red, and blue lines indicate  $x$ ,  $y$ , and  $z$  bonds, respectively. The bond length is  $a/\sqrt{3}$ . The static magnetic field  $h_R$  is applied in the right (R) region, and no magnetic field is applied in the middle (M) region. A time-dependent pulsed magnetic field is introduced in the left (L) region. (b) Plaquette  $p$  with sites marked  $p_1, p_2, \dots, p_6$  shown for the operator  $W_p$ .

corresponding eigenvalues  $w_p$  for a certain state, the existence of the local conserved quantity guarantees the absence of local magnetic moments  $\langle S_i^\mu \rangle$  and long-range spin-spin correlations  $\langle C_{ij}^\mu \rangle$  in the Kitaev model, where  $C_{ij}^\mu = S_i^\mu S_j^\mu$ . The ground state is realized in the space with  $w_p = 1$  for each plaquette, which can be regarded as the flux-free space [24]. Besides the flux degrees of freedom, the other remains: itinerant Majorana fermions. It is known that gapless dispersion with the velocity  $v [= (\sqrt{3}/4)aJ]$  appears in the itinerant Majorana excitation in the flux-free space, where  $a$  is a lattice constant. It is also known that finite energy is needed to create adjacent fluxes in the system [24]. The energy scales of the itinerant Majorana and the flux excitations are distinct from each other. In the following, we discuss how the energy difference affects the spin transport at finite temperatures.

To study the spin transport in the Kitaev model, we treat the system with armchair edges, as shown in Fig. 1(a). We note that qualitatively the same behavior discussed below can be observed also for a system with zigzag edges. The system is composed of L, M, and R regions, where the distinct magnetic fields are applied in the  $z$  direction. In the L region on the left edge, a time-dependent pulsed magnetic field  $h_L(t)$  is introduced around  $t = 0$ . No magnetic field is applied to the M region, while the static magnetic field  $h_R$  is applied to the R region. The model Hamiltonian is given as

$$H(t) = H_0 + H_1(t), \quad (3)$$

$$H_0 = H_K - h_R \sum_{i \in R} S_i^z, \quad (4)$$

$$H_1(t) = -h_L(t) \sum_{i \in L} S_i^z. \quad (5)$$

We note that in the regions under the finite magnetic field, the local operator  $W_p$  is no longer a conserved quantity. For example, the local operator on the plaquette composed of sites 5, 12, 13, 14, 7, and 6 shown in Fig. 1(a) does not commute

with  $-h_R S_7^z$ . Therefore, in general, this leads to the finite magnetizations in the R region.

In this paper, we examine the real-time dynamics in the model at finite temperatures after the magnetic pulse is introduced in the L region. The expectation value at time  $t$  for an operator  $\hat{O}$  is given as

$$\langle \hat{O}(t) \rangle = \frac{1}{Z_0} \text{Tr}[\hat{O}(t) e^{-\beta H_0}], \quad (6)$$

where  $\beta = 1/T$ ,  $T$  is the temperature,  $Z_0 (= \text{Tr}[e^{-\beta H_0}])$  is the partition function, and  $\hat{O}(t) = U^\dagger(t) \hat{O} U(t)$  with the time-evolution operator  $U(t)$ . At zero temperature ( $T = 0$ ), the localized  $Z_2$  fluxes freeze into the flux-free state, and the Majorana mean-field approach should work to evaluate the expectation values [13, 15, 36–38]. On the other hand, at finite temperatures, treating both excitations with distinct energy scales should be hard with the above method. Thus we use the TPQ state method [33, 34], where local quantities are efficiently evaluated without the trace calculations [39–45]. An important point is that this numerical method takes several energy scales into account on an equal footing and thereby has been successfully used in several systems such as the Heisenberg model on frustrated lattices [33–35, 46–49] and the Kitaev models [30, 50–55].

In the TPQ method, the expectation value (6) is described by means of the TPQ state as

$$\langle \hat{O}(t) \rangle = \langle \Psi_T | \hat{O}(t) | \Psi_T \rangle = \langle \Psi_T(t) | \hat{O} | \Psi_T(t) \rangle, \quad (7)$$

where  $|\Psi_T\rangle$  is the TPQ state at the temperature  $T$  and  $|\Psi_T(t)\rangle = U(t) |\Psi_T\rangle$ . The time evolution of the physical quantities can be evaluated by the time evolution of the TPQ state [35].

Here, we briefly explain the TPQ method. A TPQ state at  $T \rightarrow \infty$  is simply given by a random vector,

$$|\Psi_0\rangle = \sum c_i |i\rangle, \quad (8)$$

where  $\{c_i\}$  is a set of random complex numbers satisfying  $\sum_i |c_i|^2 = 1$  and  $|i\rangle$  is an arbitrary Hilbert basis. By multiplying a certain TPQ state by the Hamiltonian, the TPQ states at lower temperatures are constructed. The  $k$ th TPQ state is represented as

$$|\Psi_k\rangle = \frac{(L - H_0) |\Psi_{k-1}\rangle}{\|(L - H_0) |\Psi_{k-1}\rangle\|}, \quad (9)$$

where  $L$  is a constant value, which is larger than the maximum eigenvalue of the Hamiltonian  $H_0$ . The corresponding temperature is given by

$$T_k = \frac{L - E_k}{2k}, \quad (10)$$

where  $E_k (= \langle \Psi_k | H_0 | \Psi_k \rangle)$  is the internal energy. Thermodynamic quantities such as the entropy and specific heat can be obtained from the internal energy and temperature.

We repeat this procedure until  $T_k = T$  and obtain the TPQ state  $|\Psi_T\rangle$ . Then, we calculate the time evolution of the TPQ state  $|\Psi_T(t)\rangle$  in terms of the Lanczos time-evolution methods [56–61]. We can efficiently obtain the expectation value  $\langle \hat{O}(t) \rangle$ . When we discuss the real-time dynamics by applying the pulsed magnetic field, it is useful to consider a change in

the quantities as

$$\Delta O(t) = \langle \hat{O}(t) \rangle - \langle \hat{O} \rangle_0, \quad (11)$$

where  $\langle \cdots \rangle_0$  is the expectation value for the static Hamiltonian  $H_0$ .

When the TPQ method is applied to the finite cluster, the obtained results are sensitive to its size and/or shape. This is due to, at least, two effects. One of them is that low-energy properties in the thermodynamic limit cannot be described correctly in terms of finite clusters. Therefore the large system size dependence of the physical quantities appears at low temperatures although the TPQ method reproduces the correct results at higher temperatures. The other is the random dependence in the initial TPQ state. This should become negligible, by taking a statistical average of the results for independent TPQ states. Nevertheless, we sometimes meet with difficulty in evaluating time-dependent quantities, since each TPQ state is not an eigenstate of the Hamiltonian. Namely, ill oscillations appear in the physical quantities with respect to time even without time-dependent perturbations, unless the quantities are conserved ones. Although this oscillation should be neglected in the statistical average, the sample dependence is somewhat large even at high temperatures. To avoid this problem, we construct two time-dependent TPQ states from the common TPQ state as  $|\Psi_T(t)\rangle$  and  $|\Psi_T^0(t)\rangle = U_0(t)|\Psi_T\rangle$ , where  $U_0(t)$  is the time-evolution operator for the system described by  $H_0$ . Then, we calculate  $\langle \hat{O}(t) \rangle_0 = \langle \Psi_T^0(t) | \hat{O} | \Psi_T^0(t) \rangle$  instead of  $\langle \hat{O} \rangle_0$  and evaluate the change in the quantities (11), where unphysical oscillations should be canceled. This allows us to obtain  $\Delta O(t)$  efficiently and to discuss correctly how the external field affects the Kitaev system at finite temperatures. We have confirmed that, in a 16-site cluster, our TPQ results are in good agreement with the results obtained by the finite-temperature exact diagonalization (not shown). Even in a large cluster ( $N = 28$ ), the TPQ results at the lowest temperature  $T/J = 0.01$  agree well with the results at zero temperature obtained by the Lanczos method, which will be shown later. Therefore we believe that the time evolution of physical quantities can be examined quantitatively by means of the TPQ method. In this paper, we mainly consider the Kitaev model with  $N = 28$ , where  $N$  is the total number of sites. The model is schematically shown in Fig. 1(a). The static magnetic field in the R region is set to be  $h_R = 0.01J$ , which is smaller than the critical values  $h_c$  [36,37,62]. Before discussing the time evolution, we first demonstrate equilibrium quantities of the Kitaev model. Applying the TPQ method to the Hamiltonian  $H_0$  with  $h_R/J = 0.01$  on the 28-site cluster with armchair edges, we obtain the entropy  $S$ , specific heat  $C$ , internal energy  $E$ , and expectation value  $\langle W_p \rangle$ . The results are shown in Fig. 2. We clearly find double peaks in the specific heat at  $T_L/J \sim 0.019$  and  $T_H/J \sim 0.36$  and shoulder behavior in the entropy around  $T/J \sim 0.1$ . It is also found that  $E$  is largely changed around  $T_H$ , while  $\langle W_p \rangle$  is changed around  $T_L$ . These results are consistent with the fact that  $T_L$  and  $T_H$  correspond to typical energy scales of the local fluxes and the itinerant Majorana fermions, respectively. The residual entropy originates from the existence of the edge states in the cluster. As mentioned above, low-temperature properties are sensitive to the cluster since low-energy excitations depend on the size and/or shape. Nevertheless, the spin fractionalization inherent in the Kitaev




FIG. 2. (a) Specific heat  $C$  (blue line) and entropy  $S$  (orange line), and (b) internal energy  $E$  (blue line) and expectation values of local conserved quantities  $\langle W \rangle_p$  (orange line) as a function of  $T/J$  in the Kitaev system ( $N = 28$ ) with the armchair edges described by  $H_0$ .

model can be captured even in the 28-site system with edges. This allows us to discuss how thermal fluctuations affect the spin propagation in the Kitaev model qualitatively.

### III. RESULTS

Now, we study the real-time dynamics of the Kitaev system at finite temperatures after the Gaussian magnetic pulse is introduced in the L region. The form of the Gaussian pulse is given as

$$h_L(t) = \frac{A}{\sqrt{2\pi}\sigma} \exp\left[-\frac{t^2}{2\sigma^2}\right], \quad (12)$$

where  $A$  and  $\sigma$  are the strength and width of the pulse. Here, we set  $\sigma = 2/J$  and  $A = 1$ . Taking the average over more than 100 independent TPQ states, we calculate the time evolution of local physical quantities. It is known that the spin transport through the Kitaev QSL region is mediated by the Majorana fermions [13,15]. To avoid discussion of the reflection around the right edge, we define the arrival time of oscillations triggered by the magnetic pulse at  $x$  as  $t^* = x/v$ , where  $x$  is the coordinate of the  $i$ th site or the midpoint of the bond [see Fig. 1(a)].

Figure 3 shows the change in the local magnetizations in the L, M, and R regions at  $T/J = 0.01$ . We note that the cluster treated here has translational and mirror symmetries in the perpendicular direction to the  $x$  axis, and four sites with a certain  $x$  coordinate are topologically equivalent. In the L




FIG. 3. Real-time evolution of the changes in the local magnetizations  $\Delta S_i^z$  in the system with  $h_R/J = 0.01$  after the introduction of the pulsed magnetic field with  $A = 1$  and  $\sigma = 2/J$ . Solid and dashed lines represent the results at the temperatures  $T/J = 0.01$  and  $T/J = 0$ , respectively. For clarity,  $\Delta S_i^z(t)$  for sites 11 and 14 are plotted on a scale multiplied by 10.

region (site 8), no magnetic field is applied at  $t \rightarrow -\infty$ , and thus no magnetic moment appears. We find that the pulsed magnetic field induces the magnetic moment  $\Delta S_8^z(t)$  at the same time as the pulse is introduced. On the other hand, no magnetic moments are induced in the M region (site 11). This is consistent with the fact that the existence of local conserved quantities guarantees the absence of the magnetic moments even after the magnetic pulse is introduced. In the R region, the tiny static magnetic field  $h_R$  is applied, and the magnetic moment appears with  $\langle S_{14}^z \rangle \sim 0.092$  at  $t = -\infty$ . We find that the spin oscillation is induced at site 14 around  $t \sim 3/J$ . This means that the wave packet triggered by the magnetic pulse in the L region reaches the R region through the M region without spin oscillations. The peculiar spin transport is mediated by itinerant Majorana fermions [13]. We also apply exact diagonalization to this system and calculate the spin oscillation at zero temperature. The obtained results are shown as the dashed lines in Fig. 3. We find that the spin oscillation for the ground state is slightly different from that at  $T/J = 0.01$ . This should imply that few excited fluxes influence the motion of the itinerant Majorana fermions.

To clarify how the Majorana-mediated spin transport is modified at finite temperatures, we first focus on the time evolution of the nearest-neighbor spin-spin correlations  $C_{ij}^\mu(t)$  on the  $\mu$  bond. This quantity is proportional to the bond energy, and thereby the oscillation indicates the energy flow for the Majorana-mediated transport [14]. Figure 4 shows the real-time evolution of the change in the spin-spin correlations in the M region. Note that in the M region the spin moments  $\langle S_i^z(t) \rangle = 0$  (not shown). We find that at  $T = 0$ , the oscillations of  $\Delta C_{3,10}^z(t)$  and  $\Delta C_{11,18}^z(t)$  start at  $t \sim -2.5J$  and  $t \sim -1.5J$ , respectively (see insets of Fig. 4). This difference in time means that the energy injected by the magnetic pulse in the L region in turn flows through the M region, which is in contrast to there being no oscillations in the magnetic moments (see Fig. 3). For both bonds, the oscillation is little changed at  $T/J < 0.01$ . On the other hand, at  $T/J \gtrsim 0.01$ , it




FIG. 4. (a) and (b) Real-time evolution of the change in the spin-spin correlation in the M region at several temperatures. The results at zero temperature are obtained by the exact diagonalization. Dashed vertical lines represent the time  $t^*$  for the corresponding bonds. The insets of (a) and (b) are corresponding magnified graphs at  $-3 < tJ < 1$  with the same magnification.

is rapidly changed, and its intensity monotonically decreases with increasing temperatures. This behavior seems a general feature in the correlated systems, where the propagation smears due to thermal fluctuations.

By contrast, different behavior appears in the change in the magnetic moment in the R region. The results with sev-




FIG. 5. Real-time evolution of the change in the local magnetization  $\Delta S_{14}^z(t)$  in the R region. The dashed vertical line represents the time  $t_{14}^*$ .




FIG. 6. The orange circles represent  $\Delta S_{14}^z(t)$  at  $t = t_{14}^*$ , and the blue line represents the magnetic susceptibility of the  $N = 28$  cluster (see the inset). The orange arrow represents the result in the ground state obtained by the exact diagonalization. The three dashed vertical lines indicate  $T_L/J$ ,  $T_\chi/J$ , and  $T_H/J$ .

eral temperatures are shown in Fig. 5. At low temperatures ( $T/J \lesssim 0.01$ ), shoulder behavior in  $\Delta S_{14}^z$  appears around  $t = t^*$ . Beyond  $T = T_L$ , shoulder behavior smears, and the time evolution becomes monotonic. The increase in  $|\Delta S_{14}^z(t)|$  implies that spin oscillations triggered by the magnetic pulse are enhanced by thermal fluctuations. With further increasing temperatures, its magnitude decreases and almost vanishes when  $T \gtrsim T_H$ . This should originate from thermal fluctuations for both  $Z_2$  fluxes and itinerant Majorana fermions.

Now, we discuss the oscillation of the local moment in the R region in more detail, regarding  $|\Delta S_{14}^z(t)|$  at  $t = t_{14}^*$  as its representative magnitude. Figure 6 shows the temperature dependence of the quantity (orange circles). It is found that, at zero temperature, Majorana-mediated spin transport appears with  $|\Delta S_{14}^z(t_{14}^*)| \sim 0.0022$ . This value changes little when  $T/J \lesssim 0.01$ . With increasing temperatures,  $|\Delta S_{14}^z(t_{14}^*)|$  increases and takes a maximum around  $T/J \sim 0.03$ . Further increase in the temperature decreases the value monotonically due to thermal fluctuations. This nonmonotonic behavior reminds us of the magnetic susceptibility of the bulk system [63–68].

Then, we consider a 28-site Kitaev cluster with the periodic boundary conditions along the  $x$  and  $y$  directions (see the inset of Fig. 6). Applying the TPQ method to this cluster with the tiny uniform magnetic field  $h$  ( $= 0.01J$ ) in the  $z$  direction, we obtain the static susceptibility  $\chi = \sum_i^N S_i^z / (hN)$ . We show the results with the solid line in Fig. 6. One can see that the susceptibility behaves nonmonotonically against the temperature and shows a broad peak around  $T = T_\chi$  ( $\sim 0.027J$ ). This nonmonotonic behavior originates from two competing effects. One is the thermal fluctuation, which tends to suppress

the susceptibility. The other is the existence of the finite gap of the  $Z_2$  fluxes. Due to the spin fractionalization, one spin is represented by the flux and itinerant Majorana fermion.

The flux becomes thermally activated at finite temperatures, in particular, around  $T_L$ , which makes the spin sensitive against the external field. Our results show that the temperature dependences of the magnetic oscillations induced by the magnetic pulse and the susceptibility are essentially the same. This implies that, also for the spin transport, the competition between the thermal fluctuations and the thermal activation of  $Z_2$  fluxes plays a role similar to the case of the susceptibility, leading to a nonmonotonic temperature dependence.

#### IV. CONCLUSION

We have investigated how stable the Majorana-mediated spin transport in a quantum spin Kitaev model is against thermal fluctuations. The finite-temperature spin dynamics has been examined by applying the time-dependent thermal pure quantum state method to the Kitaev model. At low temperatures ( $T \ll T_L$ ), an almost flux-free state is realized, and the spin excitation propagates similarly to that for the ground state. When  $T \sim T_L$ , larger oscillations in the spin moments are observed, comparing with the results at the ground state. At high temperature ( $T \sim T_H$ ), both itinerant Majorana fermions and localized  $Z_2$  fluxes strongly fluctuate thermally, which suppresses the spin oscillations. We have also confirmed that such a phenomenon appears even in the Kitaev cluster with zigzag edges. Therefore our results demonstrate a crucial role of thermal fluctuations in the Majorana-mediated spin transport in the Kitaev model.

We have found the enhancement of the spin oscillation around  $T \sim T_\chi$  by considering the ferromagnetic Kitaev model of a finite cluster. In future, it is important to elucidate the case of the antiferromagnetic Kitaev model. In addition, since the thermally activated fluxes also exist in Kitaev clusters with different numbers of sites, thermally enhanced spin transport should not be sensitive to the system size, but it is also important to clarify how robust this nonmonotonic behavior is against changes in the size and whether it survives in more realistic setups with the Heisenberg terms and/or disorders.

#### ACKNOWLEDGMENTS

Parts of the numerical calculations were performed on the supercomputing systems at ISSP, The University of Tokyo. This work was supported by Grants-in-Aid for Scientific Research from JSPS [KAKENHI Grants No. JP19H05821, No. JP18K04678, and No. JP17K05536 (A.K.) and No. JP20H05265, No. JP20K14412, and No. JP21H05017 (Y.M.)] and by JST CREST Grant No. JPMJCR1901 (Y.M.).

- [1] J. C. Slonczewski, *Phys. Rev. B* **39**, 6995 (1989).
- [2] L. Berger, *Phys. Rev. B* **54**, 9353 (1996).
- [3] J. Slonczewski, *J. Magn. Magn. Mater.* **159**, L1 (1996).
- [4] R. D. R. Bhat and J. E. Sipe, *Phys. Rev. Lett.* **85**, 5432 (2000).
- [5] M. Tsoi, A. G. M. Jansen, J. Bass, W.-C. Chiang, V. Tsoi, and P. Wyder, *Nature (London)* **406**, 46 (2000).

- [6] J. König, M. C. Bønsager, and A. H. MacDonald, *Phys. Rev. Lett.* **87**, 187202 (2001).
- [7] I. Žutić, J. Fabian, and S. Das Sarma, *Rev. Mod. Phys.* **76**, 323 (2004).
- [8] N. Ogawa, R. Yoshimi, K. Yasuda, A. Tsukazaki, M. Kawasaki, and Y. Tokura, *Nat. Commun.* **7**, 12246 (2016).

- [9] D. C. Tsui, R. E. Dietz, and L. R. Walker, *Phys. Rev. Lett.* **27**, 1729 (1971).
- [10] J. S. Moodera, L. R. Kinder, T. M. Wong, and R. Meservey, *Phys. Rev. Lett.* **74**, 3273 (1995).
- [11] K. Kajiwara, Y. nd Harii, S. Takahashi, J. Ohe, K. Uchida, M. Mizuguchi, H. Umezawa, H. Kawai, K. Ando, K. Takanashi, S. Maekawa, and E. Saitoh, *Nature (London)* **464**, 262 (2010).
- [12] L. J. Cornelissen, J. Liu, R. A. Duine, J. B. Youssef, and B. J. van Wees, *Nat. Phys.* **11**, 1022 (2015).
- [13] T. Minakawa, Y. Murakami, A. Koga, and J. Nasu, *Phys. Rev. Lett.* **125**, 047204 (2020).
- [14] A. Koga, T. Minakawa, Y. Murakami, and J. Nasu, *J. Phys. Soc. Jpn.* **89**, 033701 (2020).
- [15] H. Taguchi, Y. Murakami, A. Koga, and J. Nasu, *Phys. Rev. B* **104**, 125139 (2021).
- [16] P. Anderson, *Mater. Res. Bull.* **8**, 153 (1973).
- [17] N. Read and B. Chakraborty, *Phys. Rev. B* **40**, 7133 (1989).
- [18] X. G. Wen, *Phys. Rev. B* **44**, 2664 (1991).
- [19] L. Balents, *Nature (London)* **464**, 199 (2010).
- [20] C.-Z. Chen, Q.-f. Sun, F. Wang, and X. C. Xie, *Phys. Rev. B* **88**, 041405(R) (2013).
- [21] L. Savary and L. Balents, *Rep. Prog. Phys.* **80**, 016502 (2017).
- [22] Y. Zhou, K. Kanoda, and T.-K. Ng, *Rev. Mod. Phys.* **89**, 025003 (2017).
- [23] D. Hirobe, M. Sato, T. Kawamata, Y. Shiomi, K. i. Uchida, R. Iguchi, Y. Koike, S. Maekawa, and E. Saitoh, *Nat. Phys.* **13**, 30 (2017).
- [24] A. Kitaev, *Ann. Phys. (Amsterdam)* **321**, 2 (2006).
- [25] J. Nasu, J. Yoshitake, and Y. Motome, *Phys. Rev. Lett.* **119**, 127204 (2017).
- [26] Y. Kasahara, T. Ohnishi, Y. Mizukami, O. Tanaka, S. Ma, K. Sugii, N. Kurita, H. Tanaka, J. Nasu, Y. Motome, T. Shibauchi, and Y. Matsuda, *Nature (London)* **559**, 227 (2018).
- [27] K. W. Plumb, J. P. Clancy, L. J. Sandilands, V. V. Shankar, Y. F. Hu, K. S. Burch, H.-Y. Kee, and Y.-J. Kim, *Phys. Rev. B* **90**, 041112(R) (2014).
- [28] J. Nasu, M. Udagawa, and Y. Motome, *Phys. Rev. Lett.* **113**, 197205 (2014).
- [29] J. Nasu, M. Udagawa, and Y. Motome, *Phys. Rev. B* **92**, 115122 (2015).
- [30] A. Koga, H. Tomishige, and J. Nasu, *J. Phys. Soc. Jpn.* **87**, 063703 (2018).
- [31] Q. Luo, S. Hu, and H.-Y. Kee, *Phys. Rev. Research* **3**, 033048 (2021).
- [32] S. S. Jahromi, H. Yarloo, and R. Orús, *Phys. Rev. Research* **3**, 033205 (2021).
- [33] S. Sugiura and A. Shimizu, *Phys. Rev. Lett.* **108**, 240401 (2012).
- [34] S. Sugiura and A. Shimizu, *Phys. Rev. Lett.* **111**, 010401 (2013).
- [35] H. Endo, C. Hotta, and A. Shimizu, *Phys. Rev. Lett.* **121**, 220601 (2018).
- [36] J. Nasu, Y. Kato, Y. Kamiya, and Y. Motome, *Phys. Rev. B* **98**, 060416(R) (2018).
- [37] S. Liang, M.-H. Jiang, W. Chen, J.-X. Li, and Q.-H. Wang, *Phys. Rev. B* **98**, 054433 (2018).
- [38] J. Nasu and Y. Motome, *Phys. Rev. Research* **1**, 033007 (2019).
- [39] A. Wietek, P. Corboz, S. Wessel, B. Normand, F. Mila, and A. Honecker, *Phys. Rev. Research* **1**, 033038 (2019).
- [40] C. Hickey, C. Berke, P. P. Stavropoulos, H.-Y. Kee, and S. Trebst, *Phys. Rev. Research* **2**, 023361 (2020).
- [41] A. Wietek, Y.-Y. He, S. R. White, A. Georges, and E. M. Stoudenmire, *Phys. Rev. X* **11**, 031007 (2021).
- [42] H. Shackleton, A. Wietek, A. Georges, and S. Sachdev, *Phys. Rev. Lett.* **126**, 136602 (2021).
- [43] C. Hickey, M. Gohlke, C. Berke, and S. Trebst, *Phys. Rev. B* **103**, 064417 (2021).
- [44] G. Sala, M. B. Stone, B. K. Rai, A. F. May, P. Laurell, V. O. Garlea, N. P. Butch, M. D. Lumsden, G. Ehlers, G. Pokharel, A. Podlesnyak, D. Mandrus, D. S. Parker, S. Okamoto, G. B. Halász, and A. D. Christianson, *Nat. Commun.* **12**, 171 (2021).
- [45] H. Suzuki, H. Liu, J. Bertinshaw, K. Ueda, H. Kim, S. Laha, D. Weber, Z. Yang, L. Wang, H. Takahashi, K. Fürsich, M. Minola, B. V. Lotsch, B. J. Kim, H. Yavaş, M. Daghofer, J. Chaloupka, G. Khaliullin, H. Gretarsson, and B. Keimer, *Nat. Commun.* **12**, 4512 (2021).
- [46] Y. Yamaji, T. Suzuki, T. Yamada, S.-i. Suga, N. Kawashima, and M. Imada, *Phys. Rev. B* **93**, 174425 (2016).
- [47] T. Suzuki and Y. Yamaji, *J. Phys. Soc. Jpn.* **88**, 115001 (2019).
- [48] R. Schäfer, I. Hagymási, R. Moessner, and D. J. Luitz, *Phys. Rev. B* **102**, 054408 (2020).
- [49] T. Shimokawa, *Phys. Rev. B* **103**, 134419 (2021).
- [50] H. Tomishige, J. Nasu, and A. Koga, *Phys. Rev. B* **97**, 094403 (2018).
- [51] A. Koga, S. Nakauchi, and J. Nasu, *Phys. Rev. B* **97**, 094427 (2018).
- [52] J. Oitmaa, A. Koga, and R. R. P. Singh, *Phys. Rev. B* **98**, 214404 (2018).
- [53] A. Koga and J. Nasu, *Phys. Rev. B* **100**, 100404(R) (2019).
- [54] C. Hickey and S. Trebst, *Nat. Commun.* **10**, 530 (2019).
- [55] K. Morita and T. Tohyama, *Phys. Rev. Research* **2**, 013205 (2020).
- [56] T. J. Park and J. C. Light, *J. Chem. Phys.* **85**, 5870 (1986).
- [57] Y. Saad, *SIAM J. Numer. Anal.* **29**, 209 (1992).
- [58] V. Druskin and L. Knizhnerman, *Numer. Linear Algebra Appl.* **2**, 205 (1995).
- [59] M. Hochbruck and C. Lubich, *SIAM J. Numer. Anal.* **34**, 1911 (1997).
- [60] M. Hochbruck, C. Lubich, and H. Selhofer, *SIAM J. Sci. Comput.* **19**, 1552 (1998).
- [61] M. Hochbruck and C. Lubich, *BIT Numer. Math.* **39**, 620 (1999).
- [62] K. Ido and T. Misawa, *Phys. Rev. B* **101**, 045121 (2020).
- [63] Y. Singh and P. Gegenwart, *Phys. Rev. B* **82**, 064412 (2010).
- [64] Y. Kubota, H. Tanaka, T. Ono, Y. Narumi, and K. Kindo, *Phys. Rev. B* **91**, 094422 (2015).
- [65] J. A. Sears, M. Songvilay, K. W. Plumb, J. P. Clancy, Y. Qiu, Y. Zhao, D. Parshall, and Y.-J. Kim, *Phys. Rev. B* **91**, 144420 (2015).
- [66] F. Freund, S. C. Williams, R. D. Johnson, R. Coldea, P. Gegenwart, and A. Jesche, *Sci. Rep.* **6**, 35362 (2016).
- [67] J. Yoshitake, J. Nasu, and Y. Motome, *Phys. Rev. Lett.* **117**, 157203 (2016).
- [68] J. Yoshitake, J. Nasu, and Y. Motome, *Phys. Rev. B* **96**, 064433 (2017).