

Anomaly inflow for subsystem symmetries

Fiona J. Burnell,^{1,2} Trithip Devakul,^{3,4} Pranay Gorantla⁴,,⁴ Ho Tat Lam,^{3,4} and Shu-Heng Shao^{2,5}

¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

²*School of Natural Sciences, Institute for Advanced Study, Princeton, New Jersey 08540, USA*

³*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

⁴*Physics Department, Princeton University, Princeton, New Jersey 08544, USA*

⁵*C. N. Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook, New York 11794, USA*



(Received 10 May 2022; revised 22 July 2022; accepted 27 July 2022; published 8 August 2022)

We study 't Hooft anomalies and the related anomaly inflow for subsystem global symmetries. These symmetries and anomalies arise in a number of exotic systems, including models with fracton order such as the X-cube model. As is the case for ordinary global symmetries, anomalies for subsystem symmetries can be canceled by anomaly inflow from a bulk theory in one higher dimension; the corresponding bulk is therefore a nontrivial subsystem symmetry protected topological (SSPT) phase. We demonstrate these phenomena in several examples with continuous and discrete subsystem global symmetries, as well as time-reversal symmetry. For each example we describe the boundary anomaly, and present classical continuum actions for the corresponding bulk SSPT phases, which describe the response of background gauge fields associated with the subsystem symmetries. Interestingly, we show that the anomaly does not uniquely specify the bulk SSPT phase. In general, the latter may also depend on how the symmetry and the associated foliation structure on the boundary are extended into the bulk.

DOI: [10.1103/PhysRevB.106.085113](https://doi.org/10.1103/PhysRevB.106.085113)

I. INTRODUCTION

Global symmetry is one of the central tools in analyzing strongly coupled quantum systems. In recent years, a new kind of global symmetry, known as the *subsystem global symmetry*, has featured prominently in many exotic lattice systems, including the gapless model of Ref. [1] and many gapped fracton models [2,3].¹ (See Refs. [5,6] for reviews on fractons.) In this paper, we will discuss anomaly inflow [7] and symmetry-protected topological (SPT) phases for subsystem global symmetries [8–13]. We will be working under the framework developed in Refs. [14–22] for these exotic field theories with subsystem global symmetries.

Unlike for ordinary global symmetry, the generator of a subsystem global symmetry acts only on a subspace $\mathcal{S}$ of the whole spatial manifold.² Different choices of the subspace generally give rise to independent conserved charges. On a lattice, the number of independent conserved charges therefore grow subextensively with the number of sites. In the low-energy limit, this leads to an infinite number of charges, which underlies many of the peculiarities in these exotic models.

These include the surprising UV/IR mixing in some of the physical observables [15–27].

It is useful to compare the subsystem global symmetry with another generalized symmetry, the higher-form global symmetry [28]. For both kinds of global symmetries, the conserved charges are supported on closed, higher-codimensional manifolds $\mathcal{S}$ in space. But the charges, especially in the continuum limit, are different in many ways for these two kinds of symmetries. The charge $Q(\mathcal{S})$ of a higher-form symmetry depends on $\mathcal{S}$ topologically, i.e., $Q(\mathcal{S}) = Q(\mathcal{S}')$ if $\mathcal{S}$ and $\mathcal{S}'$ are homologous to each other. Relatedly, there is no restriction on the choice of the manifold $\mathcal{S}$ of a given codimension. However, the charge of a subsystem global symmetry depends not only on the topology of $\mathcal{S}$, but possibly also on the shape and the location of $\mathcal{S}$. Furthermore, the charge might only be allowed to be on certain $\mathcal{S}$ but not all manifolds of a given codimension. (For example, $\mathcal{S}$ may be restricted to be straight lines along certain preferred directions, rather than be a general curve.) See Refs. [14,29] for related discussions.

Just as for ordinary global symmetry, one can attempt to gauge a subsystem global symmetry by coupling to dynamical gauge fields. This is, however, not always possible. The obstruction to gauging a global symmetry is known as the 't Hooft anomaly.

A. Review of anomaly inflow

The 't Hooft anomaly of a quantum system $\mathcal{T}$ in d space-time dimensions, with either ordinary or subsystem global symmetry, can be diagnosed as follows. We couple the system

¹Subsystem global symmetries have also appeared in some earlier references such as Ref. [4].

²Similar to the term higher-form global symmetry, the adjective “global” does not mean that the charges act globally on the whole space. Rather, it is used to distinguish the case of interest from that of the gauge symmetry.

$\mathcal{T}$ to background gauge fields A and denote the partition function by $Z_{\mathcal{T}}[A]$. For an ordinary global symmetry, A are one-form gauge fields. For a subsystem global symmetry, they are tensor gauge fields, as we describe in detail in the main text. When an 't Hooft anomaly is present, under a background gauge transformation $A \rightarrow A^g$, the partition function is not invariant but transforms with an anomalous phase:

$$Z_{\mathcal{T}}[A^g] = e^{i \int_{M^{(d)}} \theta(g, A)} Z_{\mathcal{T}}[A], \quad (1.1)$$

where $M^{(d)}$ is the spacetime d -dimensional manifold. We can always *change* the anomalous phase $\theta(g, A)$ by adding d -dimensional local counterterms of the background gauge fields A . However, the 't Hooft anomaly is characterized by the fact that no choice of d -dimensional local counterterms can remove the anomalous phase.

Another powerful way to describe the anomaly is using a classical field theory in one dimension higher. This classical field theory is the continuum description of the SPT phase. Let the partition function for this classical field theory of the background gauge fields A in $d + 1$ spacetime dimensions be

$$\exp \left[i \int_{N^{(d+1)}} \omega(A) \right]. \quad (1.2)$$

If $N^{(d+1)}$ has no boundary, then this partition function is gauge invariant. When $N^{(d+1)}$ has a boundary, then there can be a boundary term under the background gauge transformation. Let $N^{(d+1)}$ be a $d + 1$ -dimensional manifold whose boundary is $M^{(d)}$, i.e., $\partial N^{(d+1)} = M^{(d)}$. While a genuine anomaly of $\mathcal{T}$ cannot be canceled by a d -dimensional local counterterm, it can generally be canceled by the anomalous gauge transformation of a classical field theory in $d + 1$ spacetime dimensions:

$$\begin{aligned} & \exp \left[i \int_{N^{(d+1)}} \omega(A^g) \right] \\ &= \exp \left[-i \int_{M^{(d)}} \theta(g, A) \right] \exp \left[i \int_{N^{(d+1)}} \omega(A) \right]. \end{aligned} \quad (1.3)$$

In other words, the original d -dimensional system $\mathcal{T}$ coupled to a $d + 1$ -dimensional bulk classical field theory

$$Z_{\mathcal{T}}[A] \exp \left[i \int_{N^{(d+1)}} \omega(A) \right] \quad (1.4)$$

is invariant under the background gauge transformation.

We emphasize that there is nothing inconsistent with the original system $\mathcal{T}$ in d spacetime dimensions with an anomalous (subsystem) global symmetry. Such a system can be defined without the need of a bulk in one dimension higher. We simply cannot gauge the global symmetry in d spacetime dimensions.³

B. Anomaly inflow for subsystem symmetries

In this paper, we will discuss several examples of anomaly inflow for subsystem global symmetries and the corresponding subsystem symmetry-protected topological phases (SSPT)

³Another consequence of the 't Hooft anomaly of a global symmetry is that it cannot be realized in an on-site manner in a Hamiltonian lattice model with tensor product factorized Hilbert space.

in one dimension higher.⁴ Our examples include discrete and continuous subsystem symmetries, and for each one of them we will also discuss an analogous system with an anomalous ordinary global symmetry in the Appendices.

The simplest example of an 't Hooft anomaly in a model with continuous subsystem symmetry is the $U(1) \times U(1)$ anomaly of the 2+1 dimensions (2+1d) continuum field theory of Ref. [15], which had been first introduced in Ref. [1]:

$$\mathcal{L} = \frac{\mu_0}{2} (\partial_\tau \phi)^2 + \frac{1}{2\mu} (\partial_x \partial_y \phi)^2, \quad \phi \sim \phi + 2\pi. \quad (1.5)$$

(See also Refs. [21,22,24,30–35] for related discussions on this theory.) This anomaly, both in the continuum and on the lattice, was discussed in Refs. [21,22]. Here we will further present its SSPT in 3+1d, which can be described as a Euclidean Lagrangian of the classical bulk tensor gauge fields (A_τ, A_{xy}, A_z) , $(\tilde{A}_\tau, \tilde{A}_{xy}, \tilde{A}_z)$:

$$\begin{aligned} & \frac{i}{2\pi} [\tilde{A}_{xy} (\partial_\tau A_z - \partial_z A_\tau) + \tilde{A}_z (\partial_\tau A_{xy} \\ & - \partial_x \partial_y A_\tau) - \tilde{A}_\tau (\partial_z A_{xy} - \partial_x \partial_y A_z)]. \end{aligned} \quad (1.6)$$

We will discuss this 3+1d SSPT, and the associated tensor gauge fields, in more detail in Sec. II A 2. Interestingly, this anomaly can be viewed as a higher-rank analog of a mixed anomaly between the momentum $U(1)$ and the winding $U(1)$ symmetry in the ordinary 1+1d compact boson.

In this work, we also analyze a number of systems with anomalies in their subsystem symmetries and SSPTs that have not been previously discussed in the literature. A noteworthy example is the 3+1d $\mathbb{Z}_N$ X-cube model [36], one of the simplest gapped fracton models. The X-cube model has two sets of $\mathbb{Z}_N$ subsystem global symmetries, supported on strips and lines, respectively.⁵ On the lattice, these symmetries are simply the logical operators that map between different ground state sectors. In the continuum field theory, they become the Wilson operators of the underlying tensor gauge fields [17,40]. We show that these two subsystem symmetries have a mixed 't Hooft anomaly, which we describe explicitly using the field theory developed in Refs. [17,40] (see also Refs. [23,26,41]). An immediate consequence of this anomaly is that the two $\mathbb{Z}_N$ subsystem symmetry operators do not commute with each other, leading to the subextensive ground state degeneracy [17]. Moreover, we identify a 4+1d SSPT that cancels the anomaly of these subsystem global symmetries in the X-cube model.

An analogy can be drawn between the X-cube model and the 2+1d toric code [42]. The toric code has two $\mathbb{Z}_N$ stringlike logical operators which act within the space of ground states. In the IR, they flow to the two generators of an emergent $\mathbb{Z}_N \times \mathbb{Z}_N$ one-form global symmetry of the 2+1d $\mathbb{Z}_N$ gauge theory, the continuum description of the toric code. The non-trivial commutation relation between the two $\mathbb{Z}_N$ operators

⁴For the rest of this paper, we will use SSPT and the classical field theory interchangeably.

⁵Many gapped fracton models arise as the gauge theory of a subsystem symmetry [36–39]. Here we discuss the subsystem global symmetry of the X-cube model, not the gauge symmetry.




FIG. 1. Two different extensions of the 1+1d boundary (at $y = 0$) foliation into the 2+1d bulk. The time direction is not displayed in the figures.

can be interpreted as a mixed 't Hooft anomaly between the two $\mathbb{Z}_N$ one-form global symmetries [28,43,44]. (See also Ref. [45] for a parallel discussion from the condensed matter viewpoint.) This anomaly can be canceled by a 3+1d SPT [28,44,46,47], which is the low-energy limit of a Walker-Wang model [48].⁶ Our 4+1d SSPT for the X -cube model is analogous to this ordinary 3+1d SPT.

In all of our examples, the subsystem symmetries are associated with a foliation structure in space. The foliation is typically specified by leaves defined by setting one of the spatial coordinates to be a constant. The importance of the choice of the foliation in models with subsystem symmetries has been emphasized in Refs. [20,23,26,38,41,49–55].

Given a choice of the foliation on the boundary system with subsystem symmetry anomalies, there is typically more than one way to extend the foliation structure into the bulk. Each extension of the foliation is associated with a distinct bulk SSPT that can be used to cancel the same boundary anomaly. We demonstrate this phenomenon in a 1+1d system with a $\mathbb{Z}_N$ subsystem symmetry. The two different bulk foliation structures are shown in Fig. 1.

C. Organization

This paper is organized as follows. In the main text, we will discuss various systems with subsystem global symmetries. We will analyze their 't Hooft anomalies and the corresponding SSPTs in one higher dimension. In parallel, in Appendix, we will review analogous systems with ordinary global symmetries, 't Hooft anomalies, and the corresponding SPTs.

Section II A discusses the anomaly of the $U(1) \times U(1)$ subsystem symmetry and the corresponding 3+1d SSPT of the 2+1d ϕ theory of Ref. [15]. This is to be compared with the mixed anomaly between the momentum $U(1)$ and the winding $U(1)$ symmetry in the ordinary 1+1d compact boson, which we review in Appendix A 1. In Sec. II B, we then discuss a $U(1)$ subsystem anomaly in a chiral version

of the scalar field theory in Ref. [18]. This anomaly is analogous to that of an ordinary 1+1d chiral boson, discussed in Appendix A 1 b.

Section III A discusses the anomaly and the SSPT of the two $\mathbb{Z}_N$ subsystem global symmetries of the X -cube field theory. The discussion is parallel to that of the $\mathbb{Z}_N \times \mathbb{Z}_N$ one-form global symmetry in the ordinary 2+1d $\mathbb{Z}_N$ gauge theory, the low-energy limit of the toric code. We will review this one-form symmetry anomaly in Appendix A 2 a.

In Sec. III B, we turn to a 1+1d system with a $\mathbb{Z}_N$ subsystem symmetry. Its 't Hooft anomaly can be canceled by two distinct 2+1d SSPTs with different foliation structures.

Finally, in Sec. IV we consider the 2+1d $U(1)$ tensor gauge theory of Ref. [15] with a θ -angle. At $\theta = \pi$, there is a mixed anomaly between a $U(1)$ subsystem symmetry and the time-reversal symmetry. This is analogous to the mixed anomaly between the $U(1)$ one-form symmetry and the time-reversal symmetry in the ordinary 1+1d $U(1)$ gauge theory [56,57], which we review in Appendix A 3.

Note added: In a recent paper [58], the author also studies the 3+1d chiral ϕ theory with some overlapping results.

II. ANOMALIES OF $U(1)$ SUBSYSTEM SYMMETRIES

A. 2+1d $U(1) \times U(1)$ subsystem symmetry

It is well-known that the 1+1d compact boson conformal field theory (CFT), which describes the gapless phase of the 1+1d XY model, has a $U(1) \times U(1)$ mixed 't Hooft anomaly. A mixed 't Hooft anomaly between two global symmetries means that gauging one symmetry breaks the other, and vice versa. The mixed anomaly of the 1+1d compact boson can be canceled by a 2+1d $U(1) \times U(1)$ SPT, whose classical action is given by the mixed Chern-Simons term (see Appendix A 1 a for a review). Here, we demonstrate an analogous anomaly for a $U(1) \times U(1)$ subsystem symmetry in the 2+1d ϕ theory of Ref. [15]. This anomaly has previously been discussed in Refs. [21,22]. Below, we review the nature of the anomaly, and present the 3+1d SSPT that cancels it.

1. 2+1d ϕ theory

The 2+1d ϕ theory has a Euclidean Lagrangian

$$\mathcal{L}_{2+1}^{(\phi)} = \frac{\mu_0}{2} (\partial_\tau \phi)^2 + \frac{1}{2\mu} (\partial_x \partial_y \phi)^2. \quad (2.1)$$

The field ϕ is subject to the identification:

$$\phi(\tau, x, y) \sim \phi(\tau, x, y) + 2\pi n^x(x) + 2\pi n^y(y), \quad n^i(x^i) \in \mathbb{Z}. \quad (2.2)$$

Because of this, there exist nontrivial winding configurations of ϕ and they are summed over in the path integral (see Ref. [15] for details).⁷ This continuum Lagrangian has been

⁷An example of winding configurations of ϕ on a torus of size ℓ_x, ℓ_y is

$$\phi = 2\pi \left[\frac{x}{\ell_x} \Theta(y - y_0) + \frac{y}{\ell_y} \Theta(x - x_0) - \frac{xy}{\ell_x \ell_y} \right], \quad (2.3)$$

where $\Theta(x)$ is the Heaviside step function.

⁶More specifically, this is a Walker-Wang model whose input braided tensor category is modular. In this case, the low-energy limit is invertible and has no bulk topological order.

studied extensively in the literature [21,22,24,30–35]. It was motivated by the 2+1d XY-plaquette lattice model of rotors (which are related to the compact scalar field by $e^{i\phi}$) introduced in Ref. [1].

The 2+1d ϕ theory has a $U(1)$ momentum subsystem symmetry that shifts⁸

$$\phi(\tau, x, y) \rightarrow \phi(\tau, x, y) + \alpha_x(x) + \alpha_y(y). \quad (2.4)$$

The symmetry is generated by the current

$$J_\tau = i\mu_0 \partial_\tau \phi, \quad J^{xy} = \frac{i}{\mu} \partial_x \partial_y \phi, \quad \partial_\tau J_\tau = \partial_x \partial_y J^{xy}. \quad (2.5)$$

The theory also has a $U(1)$ winding subsystem symmetry generated by the current

$$\tilde{J}_\tau = \frac{1}{2\pi} \partial_x \partial_y \phi, \quad \tilde{J}^{xy} = \frac{1}{2\pi} \partial_\tau \phi, \quad \partial_\tau \tilde{J}_\tau = \partial_x \partial_y \tilde{J}^{xy}. \quad (2.6)$$

The winding subsystem symmetry does not act on the fundamental field ϕ , but there are (discontinuous) winding configurations, such as Eq. (2.3), carrying nontrivial charge under this symmetry. This action can be seen explicitly in a dual version of the model, where the winding symmetry shifts the field dual to ϕ in a way similar to Eq. (2.4). We refer the readers to Ref. [15] for details.

The momentum and winding $U(1)$ symmetries can be coupled to background tensor gauge fields (A_τ, A_{xy}) and $(\tilde{A}_\tau, \tilde{A}_{xy})$. The Lagrangian after coupling becomes

$$\begin{aligned} \mathcal{L}_{2+1}^{(\phi)}[A_\tau, A_{xy}; \tilde{A}_\tau, \tilde{A}_{xy}] \\ = \frac{\mu_0}{2} (\partial_\tau \phi - A_\tau)^2 + \frac{1}{2\mu} (\partial_x \partial_y \phi - A_{xy})^2 \\ + \frac{i}{2\pi} \tilde{A}_\tau (\partial_x \partial_y \phi - A_{xy}) + \frac{i}{2\pi} \tilde{A}_{xy} (\partial_\tau \phi - A_\tau). \end{aligned} \quad (2.7)$$

It is not invariant under the two $U(1)$ gauge transformations

$$\begin{aligned} A_\tau &\sim A_\tau + \partial_\tau \alpha, & A_{xy} &\sim A_{xy} + \partial_x \partial_y \alpha, & \phi &\sim \phi + \alpha, \\ \tilde{A}_\tau &\sim \tilde{A}_\tau + \partial_\tau \tilde{\alpha}, & \tilde{A}_{xy} &\sim \tilde{A}_{xy} + \partial_x \partial_y \tilde{\alpha}. \end{aligned} \quad (2.8)$$

Rather, it is shifted by

$$\mathcal{L}_{2+1}^{(\phi)} \rightarrow \mathcal{L}_{2+1}^{(\phi)} + \frac{i}{2\pi} \tilde{\alpha} (\partial_\tau A_{xy} - \partial_x \partial_y A_\tau). \quad (2.9)$$

As discussed in the Introduction, we are always free to add 2+1d counterterms involving just the background gauge fields (A_τ, A_{xy}) , $(\tilde{A}_\tau, \tilde{A}_{xy})$ to the Lagrangian (2.7). However, there is no way to completely remove the anomalous gauge transformation (2.9) by adding these 2+1d local counterterms. For example, one may try to remove the anomaly using the

⁸Here, by momentum, we mean the conjugate momentum of the field ϕ in the target space as opposed to the momentum in coordinate space. Indeed, the temporal current J_τ of the momentum symmetry is the conjugate momentum of ϕ .




FIG. 2. The foliation on the 2+1d boundary (at $z=0$) is extended to the 3+1d bulk. The time direction is not displayed in the figure.

following counterterm:

$$\frac{i}{2\pi} (\tilde{A}_\tau A_{xy} + \tilde{A}_{xy} A_\tau). \quad (2.10)$$

This does not help because the Lagrangian now transforms as

$$\mathcal{L}_{2+1}^{(\phi)} \rightarrow \mathcal{L}_{2+1}^{(\phi)} - \frac{i}{2\pi} \alpha (\partial_\tau \tilde{A}_{xy} - \partial_x \partial_y \tilde{A}_\tau). \quad (2.11)$$

It is easy to see that other quadratic terms in the gauge fields cannot remove the gauge noninvariance. Furthermore, higher order terms in the gauge fields have dimensionful coefficients, so they too cannot remove the gauge noninvariance. We conclude that there is a mixed 't Hooft anomaly between the $U(1)$ momentum and winding subsystem symmetries.

We emphasize that this mixed 't Hooft anomaly is absent in the original 2+1d XY-plaquette lattice model defined in Ref. [1], since the winding subsystem symmetry is only emergent in the low-energy limit. However, it is possible to realize both momentum and winding subsystem symmetry, as well as their mixed 't Hooft anomaly, exactly on a 3D lattice, where the third direction corresponds to discrete time [21]. Because of the mixed 't Hooft anomaly, the long-distance theory of the latter lattice model is always gapless and is described by the 2+1d ϕ theory in the continuum [22].

2. 3+1d SSPT

The mixed anomaly (2.9) can be canceled by coupling the theory to a 3+1d SSPT. Denote the radial bulk coordinate by z . The 3+1d geometry will be taken to be $S^1_\tau \times \mathbb{R}_{z \geq 0} \times \Sigma$, with the 2+1d ϕ theory living on the boundary $S^1_\tau \times \Sigma$ at $z=0$. Here Σ is a 2-manifold with a foliation structure. The leaves of the foliation on Σ are specified by either the constant x or constant y conditions.⁹ We extend the foliation structure of Σ into the bulk, but we do not introduce additional leaves specified by constant z (see Fig. 2). For this reason, the bulk SSPT will be called a 2-foliated SSPT.

⁹For example, we can take Σ to be a rectangular torus. More generally, Σ can be a twisted torus with a choice of the X and Y cycles that wrap finitely many times. See Ref. [20] for a related discussion.

The 3+1d SSPT is protected by a $U(1) \times U(1)$ subsystem symmetry whose conserved charges are supported on the constant x leaves and the constant y leaves at a fixed time. The subsystem symmetry can be coupled to background gauge fields in the bulk with gauge transformations

$$\begin{aligned} A_\tau &\sim A_\tau + \partial_\tau \alpha, & A_{xy} &\sim A_{xy} + \partial_x \partial_y \alpha, & A_z &\sim A_z + \partial_z \alpha, \\ \tilde{A}_\tau &\sim \tilde{A}_\tau + \partial_\tau \tilde{\alpha}, & \tilde{A}_{xy} &\sim \tilde{A}_{xy} + \partial_x \partial_y \tilde{\alpha}, & \tilde{A}_z &\sim \tilde{A}_z + \partial_z \tilde{\alpha}. \end{aligned} \quad (2.12)$$

The components $A_z, \tilde{A}_z$ are the analogs of the radial components of the bulk gauge fields in ordinary anomaly inflow.

The 3+1d SSPT is described by the classical Euclidean Lagrangian of the background gauge fields:

$$\begin{aligned} \mathcal{L}_{3+1}[A_\tau, A_{xy}, A_z; \tilde{A}_\tau, \tilde{A}_{xy}, \tilde{A}_z] \\ = \frac{i}{2\pi} [\tilde{A}_{xy}(\partial_\tau A_z - \partial_z A_\tau) + \tilde{A}_z(\partial_\tau A_{xy} - \partial_x \partial_y A_\tau) \\ - \tilde{A}_\tau(\partial_z A_{xy} - \partial_x \partial_y A_z)]. \end{aligned} \quad (2.13)$$

Under gauge transformations, the Lagrangian is shifted by

$$\begin{aligned} \mathcal{L}_{3+1} \rightarrow \mathcal{L}_{3+1} + \frac{i}{2\pi} \partial_x [\partial_y \tilde{\alpha}(\partial_\tau A_z - \partial_z A_\tau)] \\ - \frac{i}{2\pi} \partial_y [\tilde{\alpha} \partial_x (\partial_\tau A_z - \partial_z A_\tau)] \\ + \frac{i}{2\pi} \partial_z [\tilde{\alpha}(\partial_\tau A_{xy} - \partial_x \partial_y A_\tau)] \\ - \frac{i}{2\pi} \partial_\tau [\tilde{\alpha}(\partial_z A_{xy} - \partial_x \partial_y A_z)]. \end{aligned} \quad (2.14)$$

The bulk action is invariant up to a boundary term at $z = 0$:

$$\begin{aligned} S_{3+1} &= \int d\tau dx dy dz \mathcal{L}_{3+1} \\ &\rightarrow S_{3+1} - \frac{i}{2\pi} \int_{z=0} d\tau dx dy \tilde{\alpha}(\partial_\tau A_{xy} - \partial_x \partial_y A_\tau), \end{aligned} \quad (2.15)$$

which cancels the anomaly (2.9) of the boundary ϕ theory.

We can also place this SSPT on a 3+1d geometry with only a boundary at $x = 0$ or $y = 0$. We will not discuss the anomaly inflow for those boundaries here.

If we set $\tilde{A} = A$ in Eq. (2.13), we will find that the Lagrangian is a total derivative, i.e., there is no Chern-Simons-like terms for (A_τ, A_{xy}, A_z) . This implies that the diagonal $U(1)$ subsystem symmetry of the 2+1d ϕ theory is anomaly free.

B. 3+1d $U(1)$ subsystem symmetry

The compact boson CFT in 1+1d has a chiral counterpart, whose anomaly is well-known (see Appendix A 1 b for a review). Here, we show that an analogous theory of chiral bosons exists in 3+1d, with $U(1)$ subsystem symmetry acting along lines. It can be viewed as a chiral version of the 3+1d ϕ theory of Ref. [18] (see also Ref. [59]).¹⁰ We show that, like

¹⁰The 2+1d ϕ theory (2.1) does not have a chiral counterpart, since the naive chiral Lagrangian is a total derivative.

its 1+1d cousin, this theory has an 't Hooft anomaly, which can be canceled by a 4+1d bulk SSPT.

1. 3+1d chiral ϕ theory

The 3+1d (nonchiral) ϕ theory of Ref. [18] has the Euclidean Lagrangian

$$\mathcal{L} = \frac{\mu_0}{2} (\partial_\tau \phi)^2 + \frac{1}{2\mu} (\partial_x \partial_y \partial_z \phi)^2. \quad (2.16)$$

The field ϕ is subject to the identification:

$$\begin{aligned} \phi(\tau, x, y, z) \sim \phi(\tau, x, y, z) + 2\pi n^{xy}(x, y) + 2\pi n^{yz}(y, z) \\ + 2\pi n^{zx}(z, x), \quad n^{ij}(x^i, x^j) \in \mathbb{Z}. \end{aligned} \quad (2.17)$$

Because of this, there exist nontrivial winding configurations for ϕ (see Ref. [18] for more details).¹¹

Here we consider a chiral version of this theory with the Euclidean Lagrangian

$$\mathcal{L}_{3+1}^{(\phi)} = \frac{iN}{4\pi} \partial_\tau \phi \partial_x \partial_y \partial_z \phi, \quad (2.19)$$

where $N \in \mathbb{Z}$. The field ϕ obeys the same identification (2.17) and has the same winding configuration, such as Eq. (2.18), as in the nonchiral theory. The theory has a gauge symmetry

$$\begin{aligned} \phi(\tau, x, y, z) \sim \phi(\tau, x, y, z) + g_{xy}(\tau, x, y) \\ + g_{yz}(\tau, y, z) + g_{zx}(\tau, z, x). \end{aligned} \quad (2.20)$$

The theory also has a momentum subsystem global symmetry that shifts

$$\phi(\tau, x, y, z) \rightarrow \phi(\tau, x, y, z) + f(x, y, z), \quad (2.21)$$

where $f(x, y, z)$ obeys the same global conditions as ϕ does. The symmetry is generated by the current

$$J_\tau = -\frac{N}{2\pi} \partial_x \partial_y \partial_z \phi, \quad \partial_z J_\tau = 0. \quad (2.22)$$

We can couple the current J_τ to a $U(1)$ background tensor gauge field (A_τ, A_{xyz}) . This modifies the Lagrangian according to

$$\begin{aligned} \mathcal{L}_{3+1}^{(\phi)}[A_\tau, A_{xyz}] \\ = \frac{iN}{4\pi} \partial_\tau \phi \partial_x \partial_y \partial_z \phi - \frac{iN}{2\pi} A_\tau \partial_x \partial_y \partial_z \phi + \frac{iN}{4\pi} A_\tau A_{xyz}. \end{aligned} \quad (2.23)$$

Note that since the current only has a J_τ component, the background gauge fields A_{xyz} does not couple to any current. Here we include in the Lagrangian a classical counterterm

¹¹An example of winding configurations of ϕ on a torus of size ℓ_x, ℓ_y, ℓ_z is

$$\begin{aligned} \phi = 2\pi \left[\frac{xyz}{\ell_x \ell_y \ell_z} - \frac{yz}{\ell_y \ell_z} \Theta(x - x_0) - \frac{xz}{\ell_x \ell_z} \Theta(y - y_0) \right. \\ - \frac{xy}{\ell_x \ell_y} \Theta(z - z_0) + \frac{x}{\ell_x} \Theta(y - y_0) \Theta(z - z_0) \\ \left. + \frac{y}{\ell_y} \Theta(x - x_0) \Theta(z - z_0) + \frac{z}{\ell_z} \Theta(x - x_0) \Theta(y - y_0) \right]. \end{aligned} \quad (2.18)$$

$\frac{iN}{4\pi} A_\tau A_{xyz}$ for later convenience. This does not affect the 't Hooft anomaly.

The Lagrangian (2.23) is not invariant under the $U(1)$ gauge transformation

$$\varphi \sim \varphi + \alpha, \quad (A_\tau, A_{xyz}) \sim (A_\tau + \partial_\tau \alpha, A_{xyz} + \partial_x \partial_y \partial_z \alpha). \quad (2.24)$$

Rather, it transforms as

$$\mathcal{L}_{3+1}^{(\varphi)}[A_\tau, A_{xyz}] \rightarrow \mathcal{L}_{3+1}^{(\varphi)}[A_\tau, A_{xyz}] - \frac{iN}{4\pi} \alpha (\partial_\tau A_{xyz} - \partial_x \partial_y \partial_z A_\tau). \quad (2.25)$$

It is straightforward to see that no 3+1d local counterterms of (A_τ, A_{xyz}) can be added to cancel the anomalous gauge transformation (2.25). This signals an 't Hooft anomaly of the $U(1)$ subsystem symmetry.

2. 4+1d SSPT

Just as the anomaly of a 1+1d chiral boson can be canceled by a 2+1d $U(1)$ SPT described by a classical Chern-Simons action, we now show that the anomaly (2.25) can be canceled by coupling the theory to a 4+1d SSPT described by

$$\begin{aligned} & \frac{iN}{4\pi} [\partial_\tau (\alpha \partial_w A_{xyz}) - \partial_w (\alpha \partial_\tau A_{xyz})] + \frac{iN}{4\pi} [\partial_x (\partial_y \partial_z \alpha \partial_\tau A_w) - \partial_y (\partial_z \alpha \partial_\tau \partial_x A_w) + \partial_z (\alpha \partial_\tau \partial_x \partial_y A_w) - \partial_\tau (\alpha \partial_x \partial_y \partial_z A_w)] \\ & - \frac{iN}{4\pi} [\partial_x (\partial_y \partial_z \alpha \partial_w A_\tau) - \partial_y (\partial_z \alpha \partial_w \partial_x A_\tau) + \partial_z (\alpha \partial_w \partial_x \partial_y A_\tau) - \partial_w (\alpha \partial_x \partial_y \partial_z A_\tau)]. \end{aligned} \quad (2.28)$$

The bulk action is invariant up to a boundary term at $w = 0$:

$$S_{4+1}[A_\tau, A_{xyz}, A_w] \rightarrow S_{4+1}[A_\tau, A_{xyz}, A_w] + \frac{iN}{4\pi} \int_{w=0} d\tau dx dy dz \alpha (\partial_\tau A_{xyz} - \partial_x \partial_y \partial_z A_\tau). \quad (2.29)$$

which cancels the anomaly (2.25) of the boundary chiral φ theory.

III. ANOMALIES OF $\mathbb{Z}_N$ SUBSYSTEM SYMMETRIES

A. 3+1d X-cube field theory

We now consider the continuum field theory [17,40] that describes the low-energy physics of the $\mathbb{Z}_N$ X-cube model [36]. The low-energy field theory has two kinds of $\mathbb{Z}_N$ subsystem global symmetries, one supported along strips, and the other supported along lines [17]. The corresponding subsystem global symmetry operators of the continuum field theory descend from the logical operators of the lattice X-cube model.

We will show that there is a mixed 't Hooft anomaly between the two $\mathbb{Z}_N$ subsystem symmetries. One manifestation of the anomaly is that the two subsystem symmetry operators fail to commute when the strips and the lines intersect in space. The states in the Hilbert space have to transform under representations of this nontrivial algebra. In particular, the subextensive ground state degeneracy of the X-cube field theory can be viewed as a direct consequence of this mixed anomaly [17].

This mixed anomaly is analogous to that between the two $\mathbb{Z}_N$ one-form symmetries of the 2+1d $\mathbb{Z}_N$ gauge theory. Both

a classical Chern-Simons-like action. Denote the radial bulk coordinate by $w \geq 0$. We will place our chiral φ theory at the $w = 0$ boundary.

The 4+1d SSPT is protected by a $U(1)$ subsystem symmetry whose conserved charges are supported on either wx , wy , or wz planes. As in our previous example (see Fig. 2), these bulk planes can be viewed as a minimal extension of the boundary foliation to the bulk, with no additional leaves added parallel to the boundary. The subsystem symmetry can be coupled to background gauge fields

$$(A_\tau, A_{xyz}, A_w) \sim (A_\tau + \partial_\tau \alpha, A_{xyz} + \partial_x \partial_y \partial_z \alpha, A_w + \partial_w \alpha). \quad (2.26)$$

The 4+1d SSPT is described by the classical Euclidean Lagrangian

$$\begin{aligned} \mathcal{L}_{4+1}[A_\tau, A_{xyz}, A_w] = & \frac{iN}{4\pi} [A_\tau (\partial_w A_{xyz} - \partial_x \partial_y \partial_z A_w) \\ & - A_w (\partial_\tau A_{xyz} - \partial_x \partial_y \partial_z A_\tau) \\ & + A_{xyz} (\partial_\tau A_w - \partial_w A_\tau)]. \end{aligned} \quad (2.27)$$

Under the gauge transformation, the Lagrangian is shifted by

the symmetry operators and the charged objects of the one-form symmetries are the Wilson lines of the 2+1d $\mathbb{Z}_N$ gauge theory. The one-form symmetry operators in the low-energy field theory descend from the stringlike logical operators of the microscopic toric code [42]. Another manifestation of this anomaly is the nontrivial braiding between the electric and the magnetic Wilson lines. We will review this mixed anomaly in the ordinary 2+1d $\mathbb{Z}_N$ gauge theory in Appendix A 2 a.

Below we will encounter gauge fields transforming in tensor representations of the spatial S_4 rotation symmetry. These can be described with spatial indices (i, j, k) , which we will always take to be cyclically ordered and, in particular, nonequal, i.e., $(i, j, k) = (x, y, z)$, (y, z, x) , or (z, x, y) . We will use $[ij]$ to denote antisymmetrization over the indices i and j , and (ij) to indicate symmetrization thereof. The tensors we will encounter include the symmetric off-diagonal tensor $T^{ij} = T^{ji}$ with three component T^{xy}, T^{yz}, T^{zx} ; the partially antisymmetric tensor $T^{[ij]k}$, with three components $T^{[xy]z}, T^{[yz]x}, T^{[zx]y}$ and a common gauge symmetry $T^{[ij]k} \sim T^{[ij]k} + c$; the partially symmetric tensor $T^{i(jk)}$ with three components that obey $T^{x(yz)} + T^{y(zx)} + T^{z(xy)} = 0$; and $T_{[ij]k}$ with three components that obey $T_{[xy]z} + T_{[yz]x} + T_{[zx]y} = 0$. Note that we distinguish upper and lower indices. Fields with indices $T^{[ij]k}$ and $T^{k(ij)}$ are related by $T^{k(ij)} \equiv T^{[kij]} - T^{[jki]}$. For more details on this notation and its connection to representations of spatial S_4 rotation group, see Refs. [16,17,21].

The $\mathbb{Z}_N$ X -cube model can be described by a low-energy continuum field theory using two sets of $U(1)$ tensor gauge fields

$$(A_\tau, A_{ij}) \sim (A_\tau + \partial_\tau \alpha, A_{ij} + \partial_i \partial_j \alpha),$$

$$(\hat{A}_\tau^{(ijk)}, \hat{A}^{ij}) \sim (\hat{A}_\tau^{(ijk)} + \partial_\tau \hat{\alpha}^{i(jk)}, \hat{A}^{ij} + \partial_k \hat{\alpha}^{k(ij)}). \quad (3.1)$$

The Euclidean Lagrangian of the low-energy continuum field theory is [17,40]

$$\mathcal{L}_{3+1}^{XC} = \frac{iN}{2\pi} [A_{ij}(\partial_\tau \hat{A}^{ij} - \partial_k \hat{A}_\tau^{k(ij)}) + A_\tau \partial_i \partial_j \hat{A}^{ij}]. \quad (3.2)$$

The simplest gauge-invariant operator that can be constructed from A is

$$W(x, y, z) = \exp \left[i \oint d\tau A_\tau \right], \quad (3.3)$$

which is the worldline of an immobile fracton. In addition, we have

$$W(z_1, z_2, C) = \exp \left[i \int_{z_1}^{z_2} dz \oint_C (\partial_z A_\tau d\tau + A_{zx} dx + A_{yz} dy) \right], \quad (3.4)$$

where C is a curve in (τ, x, y) representing the world strip of a dipole of fractons separated in the z direction, which are mobile in the (x, y) plane.

The gauge-invariant operators constructed from $\hat{A}$ have the form

$$\hat{W}^z(x, y, \hat{C}) = \exp \left[i \oint_{\hat{C}} (\hat{A}_\tau^{z(xy)} d\tau + \hat{A}^{xy} dz) \right], \quad (3.5)$$

where $\hat{C}$ is a curve in the (τ, z) plane, which describes the world-line of a z -lineon, which can move only along the z direction. The analogous operators $\hat{W}^x(y, z, \hat{C})$ and $\hat{W}^y(z, x, \hat{C})$ represent the motion of x and y lineons.

1. Global symmetries and anomaly

We now discuss the two $\mathbb{Z}_N$ subsystem global symmetries of the continuum field theory for the X -cube model.

2. $\mathbb{Z}_N$ subsystem global symmetries

The theory has a $\mathbb{Z}_N$ subsystem global symmetry generated by the $\hat{W}$ operators in Eq. (3.5), with $\hat{C}$ chosen to be a straight line at a fixed time. Due to the commutation relations implied by Eq. (3.2), such an operator shifts (A_τ, A_{ij}) by a flat $\mathbb{Z}_N$ tensor gauge field. The corresponding gauge-invariant charged operators are the W operators.

We can couple the symmetry to the following background tensor gauge fields¹²

$$(C_{\tau ij}, C_{[ij]k}) \sim (C_{\tau ij} + \partial_\tau \lambda_{ij} - \partial_i \partial_j \lambda_\tau, C_{[ij]k} - \partial_i \lambda_{jk} + \partial_j \lambda_{ik}). \quad (3.6)$$

The gauge parameters $(\lambda_\tau, \lambda_{ij})$ have their own gauge symmetry

$$(\lambda_\tau, \lambda_{ij}) \sim (\lambda_\tau + \partial_\tau \gamma, \lambda_{ij} + \partial_i \partial_j \gamma). \quad (3.7)$$

These background gauge fields are $U(1)$ gauge fields. We can restrict them to $\mathbb{Z}_N$ gauge fields by coupling them to the dynamical fields $\hat{\phi}^{[ij]k}$:

$$\mathcal{L}_C = \frac{iN}{2\pi} \hat{\phi}^{[ij]k} (\partial_\tau C_{[ij]k} + \partial_i C_{\tau jk} - \partial_j C_{\tau ik}). \quad (3.8)$$

The theory has another $\mathbb{Z}_N$ subsystem global symmetry generated by the W operator (3.4) with C chosen to be at a fixed time. This operator shifts $(\hat{A}_\tau^{k(ij)}, \hat{A}^{ij})$ by a flat $\mathbb{Z}_N$ tensor gauge field. The corresponding gauge-invariant charged operators are the $\hat{W}$ operators.

We can couple this second symmetry to the following background tensor gauge fields¹³

$$(\hat{C}_\tau^{ij}, \hat{C}) \sim (\hat{C}_\tau^{ij} + \partial_\tau \hat{\lambda}^{ij} - \partial_k \hat{\lambda}_\tau^{k(ij)}, \hat{C} + \partial_i \partial_j \hat{\lambda}^{ij}), \quad (3.9)$$

where the gauge parameters $(\hat{\lambda}_\tau^{k(ij)}, \hat{\lambda}^{ij})$ have their own gauge symmetry

$$(\hat{\lambda}_\tau^{k(ij)}, \hat{\lambda}^{ij}) \sim (\hat{\lambda}_\tau^{k(ij)} + \partial_\tau \hat{\gamma}^{k(ij)}, \hat{\lambda}^{ij} + \partial_k \hat{\gamma}^{k(ij)}). \quad (3.10)$$

Again, these background gauge fields are $U(1)$ gauge fields. To restrict them to $\mathbb{Z}_N$ gauge fields, we couple them to a dynamical scalar ϕ :

$$\mathcal{L}_\phi = \frac{iN}{2\pi} \phi (\partial_\tau \hat{C} - \partial_i \partial_j \hat{C}_\tau^{ij}). \quad (3.11)$$

3. Mixed 't Hooft anomaly

Having introduced the background gauge fields for the two $\mathbb{Z}_N$ subsystem symmetries, we now discuss how they are coupled to the X -cube model at low energy. This coupling is described by the Lagrangian

$$\mathcal{L}_{3+1}^{XC}[C, \hat{C}] = \mathcal{L}_C + \mathcal{L}_\phi + \frac{iN}{2\pi} [A_{ij}(\partial_\tau \hat{A}^{ij} - \partial_k \hat{A}_\tau^{k(ij)} - \hat{C}_\tau^{ij}) + A_\tau(\partial_i \partial_j \hat{A}^{ij} - \hat{C}) + \hat{A}^{ij} C_{\tau ij} + \hat{A}_\tau^{[ij]k} C_{[ij]k}]. \quad (3.12)$$

To ensure that this Lagrangian is invariant under the dynamical gauge symmetry of the X -cube model, we require that under the gauge transformations (3.1), the dynamical scalar fields transform according to

$$\phi \sim \phi - \alpha, \quad \hat{\phi}^{[ij]k} \sim \hat{\phi}^{[ij]k} + \hat{\alpha}^{[ij]k}, \quad (3.13)$$

where $\hat{\alpha}^{[ij]k}$ obeys $\hat{\alpha}^{k(ij)} = \hat{\alpha}^{[ki]j} - \hat{\alpha}^{[jk]i}$ and has a gauge symmetry $\hat{\alpha}^{[ij]k} \sim \hat{\alpha}^{[ij]k} + c$. Additionally, under the gauge transformations (3.6) and (3.9) of the background gauge fields, the dynamical fields of the original X -cube model transform according to

$$(A_\tau, A_{ij}) \sim (A_\tau + \lambda_\tau, A_{ij} + \lambda_{ij}),$$

$$(\hat{A}_\tau^{k(ij)}, \hat{A}^{ij}) \sim (\hat{A}_\tau^{k(ij)} + \hat{\lambda}_\tau^{k(ij)}, \hat{A}^{ij} + \hat{\lambda}^{ij}). \quad (3.14)$$

¹²The $U(1)$ and the $\mathbb{Z}_N$ tensor gauge theories of $(C_{\tau ij}, C_{[ij]k})$ were studied in Ref. [18].

¹³The $U(1)$ and the $\mathbb{Z}_N$ tensor gauge theories of $(\hat{C}_\tau^{ij}, \hat{C})$ were studied in Ref. [18].

Thus, we see that under these background gauge transformations, the Lagrangian is not invariant; rather, it is shifted by

$$\frac{iN}{2\pi}[-\lambda_{ij}\hat{C}_\tau^{ij} - \lambda_\tau\hat{C} + \hat{\lambda}^{ij}(C_{\tau ij} + \partial_\tau\lambda_{ij} - \partial_i\partial_j\lambda_\tau) + \hat{\lambda}_\tau^{[ij]k}(C_{[ij]k} - \partial_i\lambda_{jk} + \partial_j\lambda_{ik})]. \quad (3.15)$$

This signals a mixed 't Hooft anomaly between the two $\mathbb{Z}_N$ subsystem global symmetries.

4. 4+1d SSPT

The mixed 't Hooft anomaly described by Eq. (3.15) can be canceled by coupling the system to a 4+1d SSPT protected by two $\mathbb{Z}_N$ subsystem symmetries, as we now describe. We denote the radial bulk coordinate by $w \geq 0$, and place our X -cube field theory at the $w = 0$ boundary.

Our bulk theory has two sets of background tensor gauge fields, which couple to the two subsystem $\mathbb{Z}_N$ symmetries. The first set is comprised of the fields

$$(C_{\tau ij}, C_{[\tau w]}, C_{[ij]k}, C_{wij}) \sim (C_{\tau ij} + \partial_\tau\lambda_{ij} - \partial_i\partial_j\lambda_\tau, C_{[\tau w]} + \partial_\tau\lambda_w - \partial_w\lambda_\tau, C_{[ij]k} - \partial_i\lambda_{jk} + \partial_j\lambda_{ik}, C_{wij} + \partial_w\lambda_{ij} - \partial_i\partial_j\lambda_w). \quad (3.16)$$

As in our boundary theory, the gauge parameters $(\lambda_\tau, \lambda_{ij}, \lambda_w)$ have their own gauge symmetry

$$(\lambda_\tau, \lambda_{ij}, \lambda_w) \sim (\lambda_\tau + \partial_\tau\gamma, \lambda_{ij} + \partial_i\partial_j\gamma, \lambda_w + \partial_w\gamma). \quad (3.17)$$

Since these are $U(1)$ gauge fields, we couple them to a dynamical gauge field

$$(\hat{B}_\tau^{k(ij)}, \hat{B}^{ij}, \hat{B}_w^{k(ij)}) \sim (\hat{B}_\tau^{k(ij)} + \partial_\tau\hat{\beta}^{k(ij)}, \hat{B}^{ij} + \partial_k\hat{\beta}^{k(ij)}, \hat{B}_w^{k(ij)} + \partial_w\hat{\beta}^{k(ij)}), \quad (3.18)$$

with the Euclidean Lagrangian

$$\mathcal{L}_{4+1}^{(C)} = \frac{iN}{2\pi}[\hat{B}_\tau^{[ij]k}(\partial_w C_{[ij]k} + \partial_i C_{wj} - \partial_j C_{wi}) - \hat{B}_w^{[ij]k}(\partial_\tau C_{[ij]k} + \partial_i C_{\tau j} - \partial_j C_{\tau i}) - \hat{B}^{ij}(\partial_i\partial_j C_{[\tau w]} + \partial_\tau C_{wij} - \partial_w C_{\tau ij})]. \quad (3.19)$$

The second $\mathbb{Z}_N$ symmetry couples to the following tensor gauge fields:

$$(\hat{C}_\tau^{ij}, \hat{C}_{[\tau w]}^{k(ij)}, \hat{C}, \hat{C}_w^{ij}) \sim (\hat{C}_\tau^{ij} + \partial_\tau\hat{\lambda}^{ij} - \partial_k\hat{\lambda}_\tau^{k(ij)}, \hat{C}_{[\tau w]}^{k(ij)} + \partial_\tau\hat{\lambda}_w^{k(ij)} - \partial_w\hat{\lambda}_\tau^{k(ij)}, \hat{C} + \partial_i\partial_j\hat{\lambda}^{ij}, \hat{C}_w^{ij} + \partial_w\hat{\lambda}^{ij} - \partial_k\hat{\lambda}_w^{k(ij)}), \quad (3.20)$$

where the gauge parameters $(\hat{\lambda}_\tau^{ij}, \hat{\lambda}^{ij}, \hat{\lambda}_w^{k(ij)})$ have their own gauge symmetry

$$(\hat{\lambda}_\tau^{ij}, \hat{\lambda}^{ij}, \hat{\lambda}_w^{k(ij)}) \sim (\hat{\lambda}_\tau^{ij} + \partial_\tau\hat{\gamma}^{k(ij)}, \hat{\lambda}^{ij} + \partial_k\hat{\gamma}^{k(ij)}, \hat{\lambda}_w^{k(ij)} + \partial_w\hat{\gamma}^{k(ij)}). \quad (3.21)$$

To restrict the symmetry to $\mathbb{Z}_N$, we introduce the dynamical gauge fields

$$(B_\tau, B_{ij}, B_w) \sim (B_\tau + \partial_\tau\beta, B_{ij} + \partial_i\partial_j\beta, B_w + \partial_w\beta), \quad (3.22)$$

which couple to $\hat{C}$ via the Euclidean Lagrangian

$$\mathcal{L}_{4+1}^{(\hat{C})} = \frac{iN}{2\pi}[B_\tau(\partial_w\hat{C} - \partial_i\partial_j\hat{C}_w^{ij}) - B_w(\partial_\tau\hat{C} - \partial_i\partial_j\hat{C}_\tau^{ij}) - B_{ij}(\partial_k\hat{C}_{[\tau w]}^{k(ij)} + \partial_\tau\hat{C}_w^{ij} - \partial_w\hat{C}_\tau^{ij})]. \quad (3.23)$$

Our 4+1d SSPT is described by coupling these two subsystem-symmetric theories, via the classical Euclidean Lagrangian

$$\mathcal{L}_{4+1d} = \mathcal{L}_{4+1}^{(C)} + \mathcal{L}_{4+1}^{(\hat{C})} + \frac{iN}{2\pi}(C_{\tau ij}\hat{C}_w^{ij} - C_{wij}\hat{C}_\tau^{ij} + C_{[\tau w]}\hat{C} - C_{[ij]k}\hat{C}_{[\tau w]}^{[ij]k}). \quad (3.24)$$

For the coupling between the C and $\hat{C}$ gauge fields to be gauge-invariant, the dynamical gauge fields B and $\hat{B}$ also have to transform under the background gauge symmetries of $(\lambda_\tau, \lambda_{ij}, \lambda_w)$ and $(\hat{\lambda}_\tau^{k(ij)}, \hat{\lambda}^{ij}, \hat{\lambda}_w^{k(ij)})$ as follows:

$$(B_\tau, B_{ij}, B_w) \sim (B_\tau - \lambda_\tau, B_{ij} - \lambda_{ij}, B_w - \lambda_w), \\ (\hat{B}_\tau^{k(ij)}, \hat{B}^{ij}, \hat{B}_w^{k(ij)}) \sim (\hat{B}_\tau^{k(ij)} + \hat{\lambda}_\tau^{k(ij)}, \hat{B}^{ij} + \hat{\lambda}^{ij}, \hat{B}_w^{k(ij)} + \hat{\lambda}_w^{k(ij)}). \quad (3.25)$$

It is straightforward to check that the resulting action is invariant under the background gauge transformations up to a boundary term at $w = 0$, which cancels the anomaly (3.15) of the boundary X -cube field theory (3.2).

B. 1+1d system with a $\mathbb{Z}_N$ subsystem symmetry

We now consider a 1+1d system with a $\mathbb{Z}_N$ subsystem symmetry. As we show in Sec. III B 5, this system is related to the 1+1d chiral boson by gauging a $\mathbb{Z}$ subsystem symmetry.

Consider the Euclidean action

$$S = \frac{iN}{4\pi} \oint dx \left[\int_{\tau_*}^{\tau_* + \ell_\tau} d\tau \partial_\tau \phi(\tau, x) \partial_x \phi(\tau, x) + 2\pi \partial_x \phi(\tau_*, x) w_\tau(x) \right]. \quad (3.26)$$

The field ϕ is subject to the identification

$$\phi(\tau, x) \sim \phi(\tau, x) + 2\pi m(x), \quad m(x) \in \mathbb{Z}. \quad (3.27)$$

It can wind in the Euclidean time and the spatial directions:

$$\phi(\tau + \ell_\tau, x) = \phi(\tau, x) + 2\pi w_\tau(x), \quad w_\tau(x) \in \mathbb{Z}, \\ \phi(\tau, x + \ell_x) = \phi(\tau, x) + 2\pi w_x, \quad w_x \in \mathbb{Z}. \quad (3.28)$$

Here $w_\tau(x)$ is a single-valued, integer function. Indeed, under the identification (3.27), the action is shifted by

$$S \rightarrow S + 2\pi iN \oint dx w_\tau(x) \partial_x m(x), \quad (3.29)$$

which is an integer multiple of $2\pi i$. Therefore the partition function is invariant under this subsystem symmetry.

The action (3.26) is very similar to that of an ordinary chiral boson (A12), but they differ in the second term. We now explain the importance of this term. Because of the position-dependent winding of ϕ in the τ direction, the first term in the action (3.26) is not well-defined on its own. This is precisely fixed by adding the second correction term. With the correction term, the full action (3.26) is independent of the choice of the reference time τ_* , and it is invariant modulo $2\pi i$ under the identification (3.27) [see Eq. (3.29)]. This correction term is similar to the correction term in the action of the quantum mechanics of N degenerate ground states reviewed in Appendix A 2 b.

In addition to the subsystem symmetry, the theory (3.26) has a gauge symmetry

$$\phi(\tau, x) \sim \phi(\tau, x) + g(\tau), \quad (3.30)$$

where $g(\tau)$ can wind in time, $g(\tau + \ell_\tau) - g(\tau) \in 2\pi\mathbb{Z}$.

1. Global symmetry and anomaly

The action (3.26) is invariant under the $\mathbb{Z}_N$ subsystem global symmetry:

$$\phi(\tau, x) \rightarrow \phi(\tau, x) + \frac{2\pi m(x)}{N}, \quad m(x) \in \mathbb{Z}. \quad (3.31)$$

Note that the symmetry with constant $m(x) = m$ is not a global symmetry, but part of the gauge symmetry (3.30). The subsystem global symmetry is generated by the gauge invariant operator

$$U(x_1, x_2) = \exp[i\phi(x_2) - i\phi(x_1)], \quad (3.32)$$

with $x_1 < x_2$. It shifts $\phi(\tau, x) \rightarrow \phi(\tau, x) + \frac{2\pi}{N}$ only within an interval, $x \in (x_1, x_2)$.

Using the commutation relation,

$$[\phi(x_1), \partial_x \phi(x_2)] = \frac{2\pi i}{N} \delta(x_1 - x_2), \quad (3.33)$$

we find that the symmetry operators obey a nontrivial commutation relation

$$\begin{aligned} & U(x_1, x_2)U(x_3, x_4) \\ &= \begin{cases} e^{-2\pi i/N} U(x_3, x_4)U(x_1, x_2) & x_1 < x_3 < x_2 < x_4, \\ e^{2\pi i/N} U(x_3, x_4)U(x_1, x_2) & x_3 < x_1 < x_4 < x_2, \\ U(x_3, x_4)U(x_1, x_2) & \text{otherwise.} \end{cases} \end{aligned} \quad (3.34)$$

This signals an 't Hooft anomaly of the $\mathbb{Z}_N$ subsystem symmetry. In other words, the $\mathbb{Z}_N$ subsystem symmetry acts projectively on the Hilbert space.

The 't Hooft anomaly can also be detected by coupling the system to the background gauge field A_τ for the $\mathbb{Z}_N$ subsystem

symmetry. The action after coupling becomes

$$S = \frac{iN}{4\pi} \oint d\tau dx (\partial_\tau \phi \partial_x \phi - 2A_\tau \partial_x \phi). \quad (3.35)$$

Here we omit the correction term, and restrict the holonomies of the $U(1)$ background gauge fields A_τ to be $\mathbb{Z}_N$ -valued. The background gauge symmetry is

$$\phi \sim \phi + \alpha, \quad A_\tau \sim A_\tau + \partial_\tau \alpha. \quad (3.36)$$

It shifts the action by

$$S \rightarrow S - \frac{iN}{4\pi} \oint d\tau dx (\partial_\tau \alpha \partial_x \alpha + 2A_\tau \partial_x \alpha), \quad (3.37)$$

which signals an 't Hooft anomaly of the $\mathbb{Z}_N$ subsystem symmetry.

2. 2+1d SSPT

We now construct the 2+1d SSPTs that cancel the anomaly via anomaly inflow. We will show that there are two such SSPTs, and they differ in their global symmetry and foliation. Denote the radial bulk coordinate by $y \geq 0$. We place our 1+1d system on the $y = 0$ boundary.

3. 1-foliated SSPT

First, we consider a 2+1d SSPT protected by a 1-foliated $\mathbb{Z}_N$ subsystem global symmetry. The subsystem symmetry is generated by distinct symmetry line operators, that extend in the y direction, at different x . This is illustrated on the left in Fig. 1. Hence, we refer to it as a 1-foliated symmetry, and the corresponding SSPT as 1-foliated SSPT.

We can couple the subsystem symmetry to a background gauge field (A_τ, A_y) . The SSPT that we are interested in is described by the Euclidean Lagrangian

$$\mathcal{L}_{2+1} = \frac{iN}{2\pi} \Phi (\partial_\tau A_y - \partial_y A_\tau) - \frac{iN}{2\pi} A_\tau \partial_x A_y, \quad (3.38)$$

where Φ is a dynamical field of mass dimension +1 that constrains the classical $U(1)$ gauge field (A_τ, A_y) to a $\mathbb{Z}_N$ gauge field. The background gauge transformation,

$$\Phi \sim \Phi + \partial_x \alpha, \quad (A_\tau, A_y) \sim (A_\tau + \partial_\tau \alpha, A_y + \partial_y \alpha), \quad (3.39)$$

shifts the Lagrangian by total derivatives

$$\begin{aligned} & - \frac{iN}{4\pi} [\partial_x (\partial_\tau \alpha \partial_y \alpha + 2A_y \partial_\tau \alpha) + \partial_y (\partial_\tau \alpha \partial_x \alpha + 2A_\tau \partial_x \alpha) \\ & - \partial_\tau (\partial_x \alpha \partial_y \alpha + 2A_y \partial_x \alpha)]. \end{aligned} \quad (3.40)$$

It leads to an anomaly inflow to the $y = 0$ boundary, which cancels the anomaly (3.37) of the boundary 1+1d system.

4. 2-foliated SSPT

Second, we show that the anomaly can also be canceled by a 2+1d SSPT protected by a 2-foliated $\mathbb{Z}_N$ subsystem symmetry. The subsystem symmetry is generated by symmetry line operators that extend in either the x or the y direction. This is illustrated on the right in Fig. 1. These symmetry lines are all distinct operators except that the products of symmetry lines that extend in the x and y directions are the same.

We can couple the subsystem symmetry to a $\mathbb{Z}_N$ background tensor gauge field (A_τ, A_{xy}) . The SSPT that we are interested in is described by the classical Lagrangian

$$\mathcal{L}_{2+1} = \frac{iN}{2\pi} \Phi^{xy} (\partial_\tau A_{xy} - \partial_x \partial_y A_\tau) - \frac{iN}{2\pi} A_\tau A_{xy}, \quad (3.41)$$

where Φ^{xy} is a dimensionless dynamical field that constrains the classical $U(1)$ gauge field (A_τ, A_{xy}) to a $\mathbb{Z}_N$ gauge field. The gauge symmetry is

$$A_\tau \sim A_\tau + \partial_\tau \alpha, \quad A_{xy} \sim A_{xy} + \partial_x \partial_y \alpha, \quad \Phi^{xy} \sim \Phi^{xy} - \alpha. \quad (3.42)$$

The Lagrangian is invariant under the gauge symmetry up to total derivatives

$$-\frac{iN}{4\pi} [\partial_x (\partial_\tau \alpha \partial_y \alpha - 2\alpha \partial_y A_\tau) + \partial_y (\partial_\tau \alpha \partial_x \alpha + 2A_\tau \partial_x \alpha) - \partial_\tau (\partial_x \alpha \partial_y \alpha - 2\alpha A_{xy})]. \quad (3.43)$$

These terms signal an anomaly inflow to the $y = 0$ boundary, which cancels the anomaly (3.37) of the boundary 1+1d system.

5. Relation to the ordinary chiral boson

We now show that starting from the 1+1d ordinary chiral boson theory (reviewed in Appendix A 1 b), we can gauge a $\mathbb{Z}$ subsystem symmetry to obtain the 1+1d system (3.26). We will see that the correction term in Eq. (3.26) arises naturally from this gauging.

In the absence of the velocity term $(\partial_x \varphi)^2$, the ordinary chiral boson theory (A12) of φ has a $\mathbb{Z}$ subsystem symmetry that acts as

$$\varphi(\tau, x) \rightarrow \varphi(\tau, x) + 2\pi w_\tau(x), \quad w_\tau(x) \in \mathbb{Z}. \quad (3.44)$$

Note that the conjugate momentum of φ is $\pi = \frac{N}{2\pi} \partial_x \varphi$. Gauging the subsystem symmetry amounts to inserting the symmetry operators

$$\begin{aligned} & \exp \left[2\pi i \oint dx w_\tau(x) \pi(\tau_*, x) \right] \\ &= \exp \left[iN \oint dx w_\tau(x) \partial_x \varphi(\tau_*, x) \right], \end{aligned} \quad (3.45)$$

at a given time τ_* , and then summing over all possible $w_\tau(x)$.¹⁴ Inserting these symmetry operators is equivalent to modifying the action to

$$\begin{aligned} S &= \frac{iN}{4\pi} \oint dx \left[\int_{\tau_*}^{\tau_* + \ell_\tau} d\tau \partial_\tau \varphi(\tau, x) \partial_x \varphi(\tau, x) \right. \\ &\quad \left. + 4\pi w_\tau(x) \partial_x \varphi(\tau_*, x) \right], \end{aligned} \quad (3.46)$$

where we sum over the winding configurations (A13) of φ in the path integral.

Next, we define a new field ϕ as

$$\phi(\tau, x) = \varphi(\tau, x) + 2\pi w_\tau(x) \Theta(\tau - \tau_*), \quad (3.47)$$

¹⁴Note that the symmetry operator with constant $w_\tau(x) = n$ is a trivial operator since it shifts $\varphi \rightarrow \varphi + 2\pi n$.

and rewrite the path integral using this field. In contrast to the old field φ , the new field ϕ can have nontrivial, x -dependent winding in the Euclidean time direction. The action then becomes (3.26), and in the path integral, we now sum over the winding configurations (3.28) of ϕ .

IV. ANOMALIES OF TIME-REVERSAL AND SUBSYSTEM SYMMETRIES

For our final example, we consider a time-reversal invariant $U(1)$ tensor gauge theory with a θ -angle $\theta = \pi$. We show that this system exhibits a mixed 't Hooft anomaly between time-reversal symmetry and a subsystem symmetry. This anomaly is analogous to the mixed anomaly between the electric one-form symmetry and the time-reversal symmetry in the ordinary 1+1d $U(1)$ gauge theory at $\theta = \pi$ [56,57]. See Appendix A 3 for a review.

Importantly, the tensor gauge fields (A_τ, A_{xy}) below are dynamical gauge fields, rather than background gauge fields for a response action. The subsystem symmetry in question is analogous to a one-form symmetry, whose charged objects are extended Wilson strips rather than points.

A. 2+1d $U(1)$ tensor gauge theory

Consider a 2+1d $U(1)$ tensor gauge theory of a symmetric tensor gauge field

$$(A_\tau, A_{xy}) \sim (A_\tau + \partial_\tau \alpha, A_{xy} + \partial_x \partial_y \alpha), \quad (4.1)$$

with a Euclidean Lagrangian [15]

$$\mathcal{L}_{2+1}^{(A)} = \frac{1}{g_e^2} E_{xy}^2 + \frac{i\theta}{2\pi} E_{xy}. \quad (4.2)$$

Here $E_{xy} = \partial_\tau A_{xy} - \partial_x \partial_y A_\tau$ is the gauge-invariant electric field. This continuum Lagrangian was studied in the context of higher order topological insulator in Ref. [32].

The θ -parameter has a 2π periodicity, i.e., $\theta \sim \theta + 2\pi$, since the electric flux is quantized as follows:

$$\oint d\tau dx dy E_{xy} \in 2\pi \mathbb{Z}. \quad (4.3)$$

To see this quantization explicitly, observe that the gauge field configuration

$$A_\tau = 0, \quad A_{xy} = \frac{2\pi\tau}{\ell_\tau} \left[\frac{1}{\ell_x} \delta(y - y_0) + \frac{1}{\ell_y} \delta(x - x_0) - \frac{1}{\ell_x \ell_y} \right] \quad (4.4)$$

carries nontrivial electric flux. This is a valid configuration since $A_{xy}(\tau + \ell_\tau, x, y) - A_{xy}(\tau, x, y) = \partial_x \partial_y g$ with the transition function g given by Eq. (2.3). See Ref. [15] for a more detailed discussion of the fluxes in this $U(1)$ tensor gauge theory.

Because θ is periodic, the theory at $\theta = 0, \pi$ has a time-reversal symmetry

$$(\tau, x, y) \rightarrow (-\tau, x, y), \quad (A_\tau, A_{xy}) \rightarrow (-A_\tau, A_{xy}). \quad (4.5)$$

If the gauge field is classical, the θ -term at $\theta = \pi$ can be viewed as a response action for a 2+1d SSPT with gapless corner modes protected by time-reversal and $U(1)$ subsystem

symmetry [32]. Here, we instead consider the situation where the tensor gauge field is dynamical.

In this case, at all values of θ , the theory has a $U(1)$ electric tensor symmetry that shifts (A_τ, A_{xy}) by a flat tensor gauge field. The symmetry is generated by a current

$$J_\tau^{xy} = \frac{2i}{g_e^2} E_{xy} - \frac{\theta}{2\pi},$$

$$\partial_\tau J_\tau^{xy} = 0, \quad \partial_x \partial_y J_\tau^{xy} = 0. \quad (4.6)$$

The operators charged under this symmetry are the following Wilson strips:

$$W_x(y_1, y_2) = \exp \left(\int_{y_1}^{y_2} dy \oint dx A_{xy} \right),$$

$$W_y(x_1, x_2) = \exp \left(\int_{x_1}^{x_2} dx \oint dy A_{xy} \right). \quad (4.7)$$

We will show below that this electric tensor symmetry has a mixed 't Hooft anomaly with the time-reversal symmetry at $\theta = \pi$.

We can couple the electric tensor symmetry to a $U(1)$ background tensor gauge field B_τ^{xy} . The gauge transformation acts as

$$A_\tau \sim A_\tau + \lambda_\tau, \quad A_{xy} \sim A_{xy} + \lambda_{xy},$$

$$B_\tau^{xy} \sim B_\tau^{xy} + \partial_\tau \lambda_{xy} - \partial_x \partial_y \lambda_\tau. \quad (4.8)$$

The gauge parameters $(\lambda_\tau, \lambda_{xy})$ are tensor gauge fields themselves with gauge symmetry

$$(\lambda_\tau, \lambda_{xy}) \sim (\lambda_\tau + \partial_\tau \gamma, \lambda_{xy} + \partial_x \partial_y \gamma). \quad (4.9)$$

The Lagrangian that couples to the background gauge field is

$$\mathcal{L}_{2+1}^{(A)}[B_\tau^{xy}] = \frac{1}{g_e^2} (E_{xy} - B_\tau^{xy})^2 + \frac{i\theta}{2\pi} (E_{xy} - B_\tau^{xy}). \quad (4.10)$$

$$(B_\tau^{xy}, B_{[\tau z]}, B_z^{xy}) \sim (B_\tau^{xy} + \partial_\tau \lambda_{xy} - \partial_x \partial_y \lambda_\tau, B_{[\tau z]} + \partial_\tau \lambda_z - \partial_z \lambda_\tau, B_z^{xy} + \partial_z \lambda_{xy} - \partial_x \partial_y \lambda_z). \quad (4.14)$$

The gauge parameters $(\lambda_\tau, \lambda_{xy}, \lambda_z)$ are gauge fields themselves with gauge transformations

$$(\lambda_\tau, \lambda_{xy}, \lambda_z) \sim (\lambda_\tau + \partial_\tau \gamma, \lambda_{xy} + \partial_x \partial_y \gamma, \lambda_z + \partial_z \gamma). \quad (4.15)$$

The 3+1d SSPT can be described by the classical Euclidean Lagrangian

$$\mathcal{L}_{3+1} = \frac{i}{2} (\partial_\tau B_z^{xy} - \partial_z B_\tau^{xy} + \partial_x \partial_y B_{[\tau z]}). \quad (4.16)$$

On a closed manifold, the theory is invariant under the time-reversal symmetry transformation

$$(\tau, x, y) \rightarrow (-\tau, x, y),$$

$$(B_\tau^{xy}, B_{[\tau z]}, B_z^{xy}) \rightarrow (-B_\tau^{xy}, -B_{[\tau z]}, B_z^{xy}). \quad (4.17)$$

Here we have used the quantization of the following fluxes:

$$\oint d\tau dx dy dz (\partial_\tau B_z^{xy} - \partial_z B_\tau^{xy} + \partial_x \partial_y B_{[\tau z]}) \in 2\pi \mathbb{Z}. \quad (4.18)$$

We now examine the action of time-reversal symmetry at $\theta = 0, \pi$. Time-reversal symmetry acts on the background gauge field according to $B_\tau^{xy} \rightarrow -B_\tau^{xy}$, whereas $E_{xy} \rightarrow -E_{xy}$. At $\theta = 0$, this leaves both Eqs. (4.2) and (4.10) invariant; hence, the total theory is time-reversal symmetric. However, at $\theta = \pi$, under time-reversal symmetry the Lagrangian transforms as

$$\mathcal{L}_{2+1}^{(A)}[B_\tau^{xy}] \rightarrow \mathcal{L}_{2+1}^{(A)}[B_\tau^{xy}] + iB_\tau^{xy}. \quad (4.11)$$

Here we have dropped the term iE_{xy} since its integral is always $2\pi i\mathbb{Z}$. In contrast, the integral of iB_τ^{xy} is generally not $2\pi i\mathbb{Z}$. One may try to fix the anomalous time-reversal transformation (4.11) by adding the following local counterterm

$$\frac{i}{2} B_\tau^{xy}. \quad (4.12)$$

However, this will break the background gauge symmetry of B_τ^{xy} . Hence, our theory exhibits a mixed 't Hooft anomaly between the time-reversal and the $U(1)$ electric tensor symmetry at $\theta = \pi$.

B. 3+1d SSPT

We can restore the time-reversal symmetry at $\theta = \pi$ by coupling the system to a 3+1d SSPT. Denote the radial bulk coordinate by $z \geq 0$. We will place our theory at the $z = 0$ boundary.

The 3+1d SSPT is protected by time-reversal symmetry and a $U(1)$ tensor symmetry generated by the currents $(J_\tau^{xy}, J_{[\tau z]}, J_z^{xy})$ that obey

$$\partial_\tau J_\tau^{xy} + \partial_z J_z^{xy} = 0,$$

$$\partial_\tau J_{[\tau z]} + \partial_x \partial_y J_z^{xy} = 0,$$

$$\partial_z J_{[\tau z]} = \partial_x \partial_y J_\tau^{xy}. \quad (4.13)$$

We can couple the SSPT to a background tensor gauge field

For example, a gauge field configuration that carries nontrivial flux is

$$B_\tau^{xy} = \frac{2\pi z}{\ell_\tau \ell_z} \left[\frac{1}{\ell_x} \delta(y - y_0) + \frac{1}{\ell_y} \delta(x - x_0) - \frac{1}{\ell_x \ell_y} \right],$$

$$B_{[\tau z]} = 0, \quad B_z^{xy} = 0. \quad (4.19)$$

This is a valid configuration since $B_\tau^{xy}(z + \ell_z) - B_\tau^{xy}(z) = \partial_\tau \lambda_{xy}$ with λ_{xy} the configuration in Eq. (4.4). On a manifold with boundary at $z = 0$, the SSPT is time-reversal invariant up to a boundary term

$$S_{3+1} \rightarrow -S_{3+1} = S_{3+1d} - i \int_{z=0} d\tau dx dy B_\tau^{xy}, \quad (4.20)$$

which cancels the anomalous time-reversal transformation (4.11) of the 2+1d $U(1)$ tensor gauge theory on the boundary.

V. DISCUSSION AND OUTLOOK

In this work, we have discussed a number of qualitatively different examples of 't Hooft anomalies that arise in the

context of subsystem symmetry, and shown how these anomalies can be canceled by a suitably chosen bulk theory in one higher dimension via the anomaly inflow mechanism. Our examples illustrate that the diversity of possible 't Hooft anomalies in systems with subsystem global symmetry parallels that known to exist for ordinary global symmetry; hence, a comparably rich classification of SSPT phases presumably exists. Specifically, we have demonstrated that anomalies occur in systems with discrete and continuous subsystem symmetry, whose charged operators can be pointlike or extended objects.

The subsystem symmetries examined in this work are all naturally associated with a certain foliation structure in space. More generally, there are lattice models exhibiting fractal subsystem symmetries, i.e., the symmetry operators on fractal geometric objects. These include the fracton topological order of Ref. [3] and also the fractal SPT of Ref. [60]. It is a challenging open question to develop a continuum framework to understand these fractal subsystem symmetries.

Even for the examples discussed here, where the anomalous theory has a natural foliation structure, there can be more than one possible extension of the boundary foliation structure into the bulk. For example, in Sec. III B, we encountered an example of a $\mathbb{Z}_N$ subsystem symmetry anomaly in 1+1d that can be canceled by two distinct 2+1d SSPTs with different bulk foliation structures. Thus the correspondence between anomalies and SPTs appears to be more subtle than for ordinary global symmetries, where anomalies can be classified by SPTs in one dimension higher. To achieve a classification of subsystem anomalies via SSPTs in one dimension higher, one needs to first understand the possible extensions of the boundary foliation structure into the bulk.¹⁵

The perspective adopted in this paper is to start from a boundary system with a given anomaly and then identify a bulk theory to cancel this anomaly. Alternatively, one can start with a bulk SSPT, and analyze the anomaly inflow into the boundary. Since our SSPT does not have continuous spatial rotation symmetry, the anomaly inflow depends sensitively on the choice of the boundary. For example, the anomalies might be different for the boundaries along different directions in space. This possibility also makes the correspondence between the boundary anomaly and the bulk SSPT more intricate.

We leave a systematic investigation of these questions for future studies.

¹⁵The foliation structure of a theory with subsystem symmetries is somewhat analogous to the tangential structure (e.g., the spin structure) in the discussion of ordinary global symmetries. For example, various physical observables, such as the ground state degeneracy, depend not only on the geometry, but also on the choice of a foliation in systems with subsystem symmetries [49,50]. This is analogous to the dependence on the choice of the spin structure in a fermionic theory. In the ordinary correspondence between the boundary anomaly and the bulk SPT, one assumes the tangential structure is *stable* in the sense that it can be defined in all spacetime dimensions. It would be interesting to understand the corresponding mathematical structure for foliated manifolds. We thank Kantaro Ohmori for discussions on this point.

ACKNOWLEDGMENTS

We thank N. Seiberg for suggesting this topic to us and for initial collaborations in this project. We also thank Z. Komargodski and K. Ohmori for discussions. F.J.B. is grateful for the financial support of the Carnegie corporation of New York, and of the National Science foundation NSF Grant No. DMR 1928166. P.G. was supported by the Physics Department of Princeton University. H.T.L. was supported in part by a Centennial Fellowship and a Charlotte Elizabeth Procter Fellowship from Princeton University, a Croucher fellowship from the Croucher Foundation, the Packard Foundation, the Physics Department of Princeton University and the Center for Theoretical Physics at MIT. S.-H.S. was supported by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651440, NS). The authors of this paper were ordered alphabetically.

APPENDIX: ANOMALIES IN ORDINARY QUANTUM FIELD THEORIES

In this Appendix, we will review several well-known examples of systems with 't Hooft anomalies of ordinary global symmetries, and the associated symmetry protected topological (SPT) phases that cancel the anomalies via anomaly inflow.

1. Anomalies of $U(1)$ symmetries

a. $U(1) \times U(1)$ symmetry: 1+1d compact boson

Consider the 1+1d compact boson theory described by the Euclidean Lagrangian

$$\mathcal{L}_{1+1}^{(\phi)} = \frac{\beta}{2} [(\partial_\tau \phi)^2 + (\partial_x \phi)^2], \quad (\text{A1})$$

where ϕ is a compact scalar, i.e., $\phi \sim \phi + 2\pi$. In the path integral, we sum over winding configurations of ϕ . On a torus of size ℓ_τ, ℓ_x , they obey the boundary conditions

$$\begin{aligned} \phi(\tau + \ell_\tau, x) &= \phi(\tau, x) + 2\pi w_\tau, & w_\tau &\in \mathbb{Z}, \\ \phi(\tau, x + \ell_x) &= \phi(\tau, x) + 2\pi w_x, & w_x &\in \mathbb{Z}. \end{aligned} \quad (\text{A2})$$

The Lagrangian (A1) is invariant under the $U(1)$ momentum global symmetry that shifts

$$\phi(\tau, x) \rightarrow \phi(\tau, x) + c, \quad (\text{A3})$$

where c is a constant. The Noether current for this symmetry are

$$\begin{aligned} J_\tau &= i\beta \partial_\tau \phi, & J_x &= i\beta \partial_x \phi, \\ \partial_\tau J_\tau + \partial_x J_x &= 0. \end{aligned} \quad (\text{A4})$$

There is also a $U(1)$ winding symmetry with Noether current

$$\begin{aligned} \tilde{J}_\tau &= \frac{1}{2\pi} \partial_x \phi, & \tilde{J}_x &= -\frac{1}{2\pi} \partial_\tau \phi, \\ \partial_\tau \tilde{J}_\tau + \partial_x \tilde{J}_x &= 0. \end{aligned} \quad (\text{A5})$$

A field configuration that carries a nontrivial winding charge is $\phi = 2\pi x/\ell_x$, which satisfies the boundary condition (A2).

We can couple the theory to the background gauge fields $A = (A_\tau, A_x)$ and $\tilde{A} = (\tilde{A}_\tau, \tilde{A}_x)$ of the $U(1)$ momentum and

the $U(1)$ winding symmetry. The Lagrangian after coupling is

$$\mathcal{L}_{1+1}^{(\phi)}[A, \tilde{A}] = \frac{\beta}{2} [(\partial_\tau \phi - A_\tau)^2 + (\partial_x \phi - A_x)^2] + \frac{i}{2\pi} [\tilde{A}_\tau (\partial_x \phi - A_x) - \tilde{A}_x (\partial_\tau \phi - A_\tau)]. \quad (\text{A6})$$

It is not invariant under the gauge symmetry:

$$\phi \sim \phi + \alpha, \quad A \sim A + d\alpha, \quad \tilde{A} \sim \tilde{A} + d\tilde{\alpha}. \quad (\text{A7})$$

Instead, it is shifted by

$$\mathcal{L}_{1+1}^{(\phi)}[A, \tilde{A}] \rightarrow \mathcal{L}_{1+1}^{(\phi)}[A, \tilde{A}] + \frac{i}{2\pi} \tilde{\alpha} (\partial_\tau A_x - \partial_x A_\tau). \quad (\text{A8})$$

This signals an 't Hooft anomaly of the $U(1) \times U(1)$ global symmetry.

Consider a 2+1d SPT protected by a $U(1) \times U(1)$ global symmetry described by the classical Euclidean Lagrangian¹⁶

$$\begin{aligned} \mathcal{L}_{2+1}[A, \tilde{A}] &= \frac{i}{2\pi} \tilde{A} dA \\ &= \frac{i}{2\pi} [\tilde{A}_\tau (\partial_x A_y - \partial_y A_x) - \tilde{A}_x (\partial_\tau A_y - \partial_y A_\tau) \\ &\quad + \tilde{A}_y (\partial_\tau A_x - \partial_x A_\tau)], \end{aligned} \quad (\text{A9})$$

where $A = (A_\tau, A_x, A_y)$ and $\tilde{A} = (\tilde{A}_\tau, \tilde{A}_x, \tilde{A}_y)$ are background $U(1) \times U(1)$ gauge fields with gauge symmetry,

$$A \sim A + d\alpha, \quad \tilde{A} \sim \tilde{A} + d\tilde{\alpha}. \quad (\text{A10})$$

The Lagrangian (A9) is invariant under the gauge symmetry up to a total derivative:

$$\begin{aligned} \frac{i}{2\pi} d(\tilde{\alpha} dA) &= \frac{i}{2\pi} \{ \partial_\tau [\tilde{\alpha} (\partial_x A_y - \partial_y A_x)] \\ &\quad - \partial_x [\tilde{\alpha} (\partial_\tau A_y - \partial_y A_\tau)] \\ &\quad + \partial_y [\tilde{\alpha} (\partial_\tau A_x - \partial_x A_\tau)] \}. \end{aligned} \quad (\text{A11})$$

On a manifold with boundary at $y = 0$, the anomaly inflow cancels the anomaly (A8).

b. $U(1)$ symmetry: 1+1d compact chiral boson

Consider the 1+1d compact chiral boson described by the Euclidean action

$$S_{1+1}^{(\varphi)} = \frac{iN}{4\pi} \oint d\tau dx \partial_\tau \varphi \partial_x \varphi, \quad (\text{A12})$$

where $N \in \mathbb{Z}$ and φ is a compact scalar, i.e., $\varphi \sim \varphi + 2\pi$. More generally, one can add a velocity term $(\partial_x \varphi)^2$ to the Lagrangian, but we will consider the special case when this term is absent.

On a torus of size ℓ_τ, ℓ_x , we sum over the winding configurations that obey the boundary conditions¹⁷

$$\begin{aligned} \varphi(\tau + \ell_\tau, x) &= \varphi(\tau, x), \\ \varphi(\tau, x + \ell_x) &= \varphi(\tau, x) + 2\pi w_x, \quad w_x \in \mathbb{Z}, \end{aligned} \quad (\text{A13})$$

in the path integral. The theory has a gauge symmetry

$$\varphi(\tau, x) \sim \varphi(\tau, x) + g(\tau), \quad (\text{A14})$$

where $g(\tau + \ell_\tau) = g(\tau)$.

The theory has a $LU(1)/U(1)$ momentum global symmetry that shifts [61,62]

$$\varphi(\tau, x) \rightarrow \varphi(\tau, x) + f(x), \quad (\text{A15})$$

where $f(x)$ is a map from S^1 to $U(1)$. In particular, $f(x)$ can wind with $f(x + \ell_x) = f(x) + 2\pi \mathbb{Z}$. The zero mode of $f(x)$ is not a global symmetry but a gauge symmetry included in Eq. (A14). Because of this, the global symmetry has a $U(1)$ quotient. The symmetry is generated by the current

$$J_\tau = -\frac{N}{2\pi} \partial_x \varphi, \quad \partial_\tau J_\tau = 0. \quad (\text{A16})$$

Although a constant shift of φ is a gauge symmetry, the zero-mode charge $Q = \oint dx J_\tau$ can be nontrivial since the winding configuration $\varphi = 2\pi x / \ell_x$ carries a nontrivial charge $Q = -N$.

We can couple the current J_τ to a $U(1)$ background gauge field $A = (A_\tau, A_x)$ [63]

$$S_{1+1}^{(\varphi)}[A_\tau, A_x] = \frac{iN}{4\pi} \oint d\tau dx (\partial_\tau \varphi \partial_x \varphi - 2A_\tau \partial_x \varphi + A_\tau A_x). \quad (\text{A17})$$

Note that since the current only has the J_τ component, the background gauge field A_x is not coupled to any current. Here we include in the Lagrangian a classical counterterm $\frac{iN}{4\pi} A_\tau A_x$ for later convenience. This does not affect the 't Hooft anomaly. The Lagrangian is not invariant under the gauge symmetry,

$$\varphi \sim \varphi + \alpha, \quad A \sim A + d\alpha. \quad (\text{A18})$$

Rather, it is shifted by

$$S_{1+1}^{(\varphi)}[A_\tau, A_x] \rightarrow S_{1+1}^{(\varphi)}[A_\tau, A_x] - \frac{iN}{4\pi} \oint d\tau dx \alpha (\partial_\tau A_x - \partial_x A_\tau). \quad (\text{A19})$$

It signals an 't Hooft anomaly.

This anomaly can be canceled by coupling the system to a 2+1d $U(1)$ SPT described by the classical Euclidean

¹⁶In this Appendix, we will often expand the differential forms in components to compare with the analogous expressions for the subsystem symmetries in the main texts. We will often ignore the volume form, such as $d\tau \wedge dx \wedge dy$, in these expressions.

¹⁷More generally, in the path integral, we can include configurations that wind in the time direction. But using the gauge symmetry (A14), we can always shift a winding configuration $\varphi(\tau, x)$ by a $g(\tau)$ with opposite winding such that in the path integral we can restrict to configurations that do not wind in the time direction.

Lagrangian

$$\begin{aligned}\mathcal{L}_{2+1}[A_\tau, A_x, A_y] &= -\frac{iN}{4\pi}AdA \\ &= -\frac{iN}{4\pi}[A_\tau(\partial_x A_y - \partial_y A_x) + A_x(\partial_y A_\tau \\ &\quad - \partial_\tau A_y) + A_y(\partial_\tau A_x - \partial_x A_\tau)].\end{aligned}\quad (\text{A20})$$

The Lagrangian is invariant under the gauge symmetry up to a total derivative

$$\begin{aligned}-\frac{iN}{4\pi}d(\alpha dA) &= -\frac{iN}{4\pi}\{\partial_\tau[\alpha(\partial_x A_y - \partial_y A_x)] \\ &\quad - \partial_x[\alpha(\partial_\tau A_y - \partial_y A_\tau)] \\ &\quad + \partial_y[\alpha(\partial_\tau A_x - \partial_x A_\tau)]\}.\end{aligned}\quad (\text{A21})$$

On a manifold with a $y = 0$ boundary, the anomaly inflow cancels the anomaly (A19).

2. Anomalies of $\mathbb{Z}_N$ symmetries

a. $\mathbb{Z}_N \times \mathbb{Z}_N$ one-form symmetry: 2+1d $\mathbb{Z}_N$ gauge theory

The Euclidean Lagrangian of the 2+1d $\mathbb{Z}_N$ gauge theory is [64–66]

$$\mathcal{L}_{2+1} = \frac{iN}{2\pi}\tilde{A}dA, \quad (\text{A22})$$

where $A = (A_\tau, A_x, A_y)$ and $\tilde{A} = (\tilde{A}_\tau, \tilde{A}_x, \tilde{A}_y)$ are $U(1)$ gauge fields with gauge symmetry

$$A \sim A + d\alpha, \quad \tilde{A} \sim \tilde{A} + d\tilde{\alpha}. \quad (\text{A23})$$

Their equations of motion constrain each other to be $\mathbb{Z}_N$ gauge fields. This theory describes the low energy physics of the $\mathbb{Z}_N$ toric code.

The theory (A22) has an electric $\mathbb{Z}_N$ one-form global symmetry, and a magnetic $\mathbb{Z}_N$ one-form global symmetry. The electric one-form symmetry is generated by the Wilson loops of $\tilde{A}$, $\tilde{W}(\tilde{C}) = \exp(i\oint_{\tilde{C}} \tilde{A})$, and the charged operators are the Wilson loops of A , $W(C) = \exp(i\oint_C A)$. However, the magnetic one-form symmetry is generated by the Wilson loops of A , and the charged operators are the Wilson loops of $\tilde{A}$.

These operators satisfy

$$\begin{aligned}W(C)^N &= \tilde{W}(\tilde{C})^N = 1, \\ W(C)^n \tilde{W}(\tilde{C})^m &= e^{\frac{2\pi i}{N} n m I(C, \tilde{C})} \tilde{W}(\tilde{C})^m W(C)^n,\end{aligned}\quad (\text{A24})$$

where C and $\tilde{C}$ are restricted to curves in space, and $I(C, \tilde{C})$ is their intersection number. The fact that the two symmetry operators are charged under each other signals a mixed 't Hooft anomaly between the two $\mathbb{Z}_N$ one-form symmetries [28,43–45]. One consequence of this anomaly is that, on a torus, the $\mathbb{Z}_N$ gauge theory has N^2 degenerate ground states that live in the minimal representation of the operator algebra (A24).

The Wilson lines W and $\tilde{W}$ are the worldlines of the electric and magnetic anyons in the microscopic toric code. The non-trivial braiding between the electric and the magnetic anyons becomes the 't Hooft anomaly of the emergent $\mathbb{Z}_N \times \mathbb{Z}_N$ one-form global symmetry.

The anomaly can also be detected by coupling the theory (A22) to background two-form gauge fields B and $\tilde{B}$ of the

$\mathbb{Z}_N \times \mathbb{Z}_N$ one-form global symmetry:

$$\mathcal{L}_{2+1}[B, \tilde{B}] = \frac{iN}{2\pi}[\tilde{A}(dA - B) - A\tilde{B}] + \frac{iN}{2\pi}\tilde{\phi}dB + \frac{iN}{2\pi}\phi d\tilde{B}, \quad (\text{A25})$$

where B and $\tilde{B}$ are classical $U(1)$ two-form gauge fields, and $\tilde{\phi}$ and ϕ are dynamical compact scalar fields that constrain B and $\tilde{B}$ to be $\mathbb{Z}_N$ two-form gauge fields. The dynamical gauge symmetry (A23) acts on ϕ and $\tilde{\phi}$ as

$$\phi \sim \phi - \alpha, \quad \tilde{\phi} \sim \tilde{\phi} - \tilde{\alpha}. \quad (\text{A26})$$

The background one-form gauge symmetry acts as

$$\begin{aligned}A &\sim A + \lambda, \quad B \sim B + d\lambda, \\ \tilde{A} &\sim \tilde{A} + \tilde{\lambda}, \quad \tilde{B} \sim \tilde{B} + d\tilde{\lambda},\end{aligned}\quad (\text{A27})$$

where the one-form gauge parameters λ and $\tilde{\lambda}$ have their own gauge symmetries

$$\lambda \sim \lambda + d\gamma, \quad \tilde{\lambda} \sim \tilde{\lambda} + d\tilde{\gamma}. \quad (\text{A28})$$

The Lagrangian (A25) is not invariant under the gauge symmetry (A27):

$$\mathcal{L}_{2+1}[B, \tilde{B}] \rightarrow \mathcal{L}_{2+1}[B, \tilde{B}] - \frac{iN}{2\pi}[\tilde{\lambda}B + \lambda(\tilde{B} + d\tilde{\lambda})]. \quad (\text{A29})$$

This signals an 't Hooft anomaly of the $\mathbb{Z}_N \times \mathbb{Z}_N$ one-form symmetry.

Consider a 3+1d SPT protected by the $\mathbb{Z}_N \times \mathbb{Z}_N$ one-form symmetry described by the Lagrangian [44,47]

$$\mathcal{L}_{3+1}[B, \tilde{B}] = \frac{iN}{2\pi}\tilde{C}dB + \frac{iN}{2\pi}Cd\tilde{B} - \frac{iN}{2\pi}B\tilde{B}, \quad (\text{A30})$$

where B and $\tilde{B}$ are $U(1)$ two-form background gauge fields, and C and $\tilde{C}$ are dynamical one-form gauge fields that constrain $\tilde{B}$ and B to be $\mathbb{Z}_N$ two-form gauge fields. The gauge symmetry is

$$\begin{aligned}C &\sim C + d\chi + \lambda, \quad B \sim B + d\lambda, \\ \tilde{C} &\sim \tilde{C} + d\tilde{\chi} + \tilde{\lambda}, \quad \tilde{B} \sim \tilde{B} + d\tilde{\lambda},\end{aligned}\quad (\text{A31})$$

where the background one-form gauge parameters λ and $\tilde{\lambda}$ have their own gauge symmetries

$$\lambda \sim \lambda + d\gamma, \quad \tilde{\lambda} \sim \tilde{\lambda} + d\tilde{\gamma}. \quad (\text{A32})$$

The 3+1d SPT (A30) is the low-energy limit of a Walker-Wang model [48].

The Lagrangian (A30) is invariant under the background gauge symmetry (A31) up to total derivatives

$$\mathcal{L}_{3+1}[B, \tilde{B}] \rightarrow \mathcal{L}_{3+1}[B, \tilde{B}] - \frac{iN}{2\pi}d[\tilde{\lambda}B + \lambda(\tilde{B} + d\tilde{\lambda})]. \quad (\text{A33})$$

On a manifold with boundary at $z = 0$, the anomaly inflow cancels the anomaly (A29).

b. $\mathbb{Z}_N \times \mathbb{Z}_N$ symmetry: Quantum mechanics of N degenerate ground states

Consider the quantum mechanics of N degenerate ground states. When the Euclidean time is noncompact, the theory is

described by the Euclidean action

$$S_{0+1} = \frac{iN}{2\pi} \int d\tau p(\tau) \dot{q}(\tau), \quad (\text{A34})$$

where $N \in \mathbb{Z}$ and $p(\tau), q(\tau)$ are both circle-valued fields,

$$p(\tau) \sim p(\tau) + 2\pi, \quad q(\tau) \sim q(\tau) + 2\pi. \quad (\text{A35})$$

If the time is compact with a periodicity ℓ_τ , then the fields $p(\tau), q(\tau)$ can wind around the Euclidean time circle

$$\begin{aligned} p(\tau + \ell_\tau) &= p(\tau) + 2\pi n_p, & n_p &\in \mathbb{Z}, \\ q(\tau + \ell_\tau) &= q(\tau) + 2\pi n_q, & n_q &\in \mathbb{Z}, \end{aligned} \quad (\text{A36})$$

and all values of n_p, n_q are summed over in the path integral. As a result, the action (A34) is no longer well-defined unless it is supplemented with a correction term [21,67–69]:

$$S_{0+1} = \frac{iN}{2\pi} \int_{\tau_*}^{\tau_* + \ell_\tau} d\tau p(\tau) \dot{q}(\tau) - iN n_p q(\tau_*). \quad (\text{A37})$$

It is straightforward to see that the action (A37) is independent of the reference Euclidean time τ_* , and is invariant (modulo $2\pi i\mathbb{Z}$) under the identifications (A35).

The action (A37) has a $\mathbb{Z}_N \times \mathbb{Z}_N$ global symmetry that shifts

$$\begin{aligned} p(\tau) &\rightarrow p(\tau) + \frac{2\pi m_p}{N}, & m_p &\in \mathbb{Z}, \\ q(\tau) &\rightarrow q(\tau) + \frac{2\pi m_q}{N}, & m_q &\in \mathbb{Z}. \end{aligned} \quad (\text{A38})$$

It is generated by $U = e^{iq}$ and $V = e^{ip}$, which satisfy

$$U^N = V^N = 1, \quad UV = e^{\frac{2\pi i}{N}} VU. \quad (\text{A39})$$

Quantizing the action leads to a Hilbert space of N degenerated ground states, which are in the minimal representation of this algebra. The nontrivial commutation relations in Eq. (A39) mean that the Hilbert space transforms projectively under the $\mathbb{Z}_N \times \mathbb{Z}_N$ symmetry. This implies an 't Hooft anomaly of the symmetry.

The anomaly can also be detected by coupling the system to $\mathbb{Z}_N \times \mathbb{Z}_N$ background gauge fields:

$$S_{0+1}[A_\tau, \tilde{A}_\tau] = \frac{iN}{2\pi} \oint d\tau [p(\dot{q} - A_\tau) + \tilde{A}_\tau q]. \quad (\text{A40})$$

Here, we omit the correction terms, and restrict the holonomies of the $U(1)$ background gauge fields A_τ and $\tilde{A}_\tau$

to be $\mathbb{Z}_N$ -valued.¹⁸ The background gauge symmetry acts as

$$\begin{aligned} q &\sim q + \alpha, & A_\tau &\sim A_\tau + \dot{\alpha}, \\ p &\sim p + \tilde{\alpha}, & \tilde{A}_\tau &\sim \tilde{A}_\tau + \dot{\tilde{\alpha}}. \end{aligned} \quad (\text{A43})$$

The action (A40) is not invariant under the gauge symmetry (A43),

$$S_{0+1}[A_\tau, \tilde{A}_\tau] \rightarrow S_{0+1}[A_\tau, \tilde{A}_\tau] + \frac{iN}{2\pi} \oint d\tau (\alpha \tilde{A}_\tau - \tilde{\alpha} A_\tau - \tilde{\alpha} \dot{\alpha}), \quad (\text{A44})$$

which signals an 't Hooft anomaly of the $\mathbb{Z}_N \times \mathbb{Z}_N$ global symmetry.

Consider a 1+1d SPT protected by a $\mathbb{Z}_N \times \mathbb{Z}_N$ global symmetry. The SPT is described by the Euclidean Lagrangian

$$\begin{aligned} \mathcal{L}_{1+1}[A, \tilde{A}] &= \frac{iN}{2\pi} [\tilde{\Phi} dA + \Phi d\tilde{A} - A\tilde{A}] \\ &= \frac{iN}{2\pi} [\tilde{\Phi}(\partial_\tau A_x - \partial_x A_\tau) + \Phi(\partial_\tau \tilde{A}_x - \partial_x \tilde{A}_\tau) \\ &\quad - (A_\tau \tilde{A}_x - A_x \tilde{A}_\tau)], \end{aligned} \quad (\text{A45})$$

where $\tilde{\Phi}$ and Φ are dynamical compact scalar fields that constrain the $U(1)$ background gauge fields $A = (A_\tau, A_x)$ and $\tilde{A} = (\tilde{A}_\tau, \tilde{A}_x)$ to be $\mathbb{Z}_N$ -valued. The gauge symmetry acts as

$$\begin{aligned} A &\sim A + d\alpha, & \Phi &\sim \Phi - \alpha, \\ \tilde{A} &\sim \tilde{A} + d\tilde{\alpha}, & \tilde{\Phi} &\sim \tilde{\Phi} + \tilde{\alpha}. \end{aligned} \quad (\text{A46})$$

The Lagrangian (A45) is invariant under the gauge symmetry up to total derivatives:

$$\begin{aligned} & - \frac{iN}{2\pi} d(\alpha \tilde{A} - \tilde{\alpha} A - \tilde{\alpha} d\alpha) \\ &= - \frac{iN}{2\pi} [\partial_\tau (\alpha \tilde{A}_x - \tilde{\alpha} A_x - \tilde{\alpha} \partial_x \alpha) - \partial_x (\alpha \tilde{A}_\tau \\ &\quad - \tilde{\alpha} A_\tau - \tilde{\alpha} \partial_\tau \alpha)]. \end{aligned} \quad (\text{A47})$$

On a manifold with boundary at $x = 0$, the anomaly inflow cancels the anomaly (A44).

3. Time-reversal and $U(1)$ one-form symmetry: 1+1d $U(1)$ gauge theory

Consider the 1+1d $U(1)$ Maxwell gauge theory with a θ -term,

$$\mathcal{L}_{1+1}^{\text{Max}} = \frac{1}{g^2} E_x^2 + \frac{i\theta}{2\pi} E_x, \quad (\text{A48})$$

¹⁸We can impose these restrictions by adding to the action (A40) the following terms:

$$\frac{i}{2\pi} \oint d\tau [\tilde{\chi}(\dot{\psi} - NA_\tau) + \chi(\dot{\tilde{\psi}} - N\tilde{A}_\tau)], \quad (\text{A41})$$

where $\chi, \tilde{\chi}$ are real-valued Lagrange multipliers, and $\psi, \tilde{\psi}$ are circle-valued. The latter fields transform under the background gauge symmetry (A43) as

$$\psi \sim \psi + N\alpha, \quad \tilde{\psi} \sim \tilde{\psi} + N\tilde{\alpha}. \quad (\text{A42})$$

Since these terms do not affect the anomaly, we omit them in Eq. (A40).

where $E_x = \partial_\tau A_x - \partial_x A_\tau$ is the electric field. The one-form gauge field $A = (A_\tau, A_x)$ has the ordinary gauge symmetry:

$$A \sim A + d\alpha. \quad (\text{A49})$$

The θ -parameter has 2π periodicity because of flux quantization $\oint d\tau dx E_x \in 2\pi\mathbb{Z}$.

The theory (A48) has a $U(1)$ electric one-form symmetry with Noether current

$$J_{[\tau x]} = \frac{2i}{g^2} E_x - \frac{\theta}{2\pi},$$

$$\partial_\tau J_{[\tau x]} = 0, \quad \partial_x J_{[\tau x]} = 0. \quad (\text{A50})$$

At $\theta = 0, \pi$, the theory also has a time-reversal symmetry:

$$(\tau, x) \rightarrow (-\tau, x), \quad (A_\tau, A_x) \rightarrow (-A_\tau, A_x). \quad (\text{A51})$$

We can couple Eq. (A48) to the background two-form gauge field $B = B_{[\tau x]}$ of the electric one-form symmetry:

$$\mathcal{L}_{1+1}^{\text{Max}}[B] = \frac{1}{g^2} (E_x - B_{[\tau x]})^2 + \frac{i\theta}{2\pi} (E_x - B_{[\tau x]}). \quad (\text{A52})$$

The one-form background gauge symmetry is

$$A \sim A + \lambda, \quad B \sim B + d\lambda, \quad (\text{A53})$$

where the one-form gauge parameter $\lambda = (\lambda_\tau, \lambda_x)$ itself has a gauge symmetry,

$$\lambda \sim \lambda + d\gamma. \quad (\text{A54})$$

With a nontrivial $B_{[\tau x]}$, the time-reversal symmetry is unbroken at $\theta = 0$ but it is broken at $\theta = \pi$ with the anomalous

transformation

$$\mathcal{L}_{1+1}^{\text{Max}}[B] \rightarrow \mathcal{L}_{1+1}^{\text{Max}}[B] + iB_{[\tau x]}. \quad (\text{A55})$$

This signals a mixed 't Hooft anomaly between the electric one-form symmetry and the time reversal symmetry at $\theta = \pi$ [56,57].

We can restore the time-reversal symmetry at $\theta = \pi$ by coupling the system to a 2+1d SPT protected by the time-reversal and the $U(1)$ one-form symmetry. For simplicity, we will assume the bulk manifold is orientable and refer the readers to Ref. [57] for a more general discussion.

In the bulk, the $U(1)$ one-form symmetry is coupled to a two-form background gauge field $B = (B_{[\tau x]}, B_{[\tau y]}, B_{[xy]})$ with gauge symmetry

$$B \sim B + d\lambda, \quad (\text{A56})$$

where the one-form gauge parameter $\lambda = (\lambda_\tau, \lambda_x, \lambda_y)$ itself has a gauge symmetry

$$\lambda \sim \lambda + d\gamma. \quad (\text{A57})$$

The 2+1d SPT is described by the Euclidean Lagrangian

$$\mathcal{L}_{2+1}[B] = -\frac{i}{2} dB = -\frac{i}{2} (\partial_\tau B_{[xy]} - \partial_x B_{[\tau y]} + \partial_y B_{[\tau x]}). \quad (\text{A58})$$

On a closed orientable manifold, the SPT is time-reversal invariant because of flux quantization $\oint dB \in 2\pi\mathbb{Z}$. On a manifold with boundary at $y = 0$, it is time-reversal invariant up to a boundary term:

$$S_{2+1}[B] \rightarrow -S_{2+1}[B] = S_{2+1}[B] - i \int_{y=0} d\tau dx B_{[\tau x]}, \quad (\text{A59})$$

which cancels the anomalous time-reversal transformation (A55) of the 1+1d $U(1)$ Maxwell theory on the boundary.

-
- [1] A. Paramekanti, L. Balents, and M. P. A. Fisher, Ring exchange, the exciton Bose liquid, and Bosonization in two dimensions, *Phys. Rev. B* **66**, 054526 (2002).
 - [2] C. Chamon, Quantum Glassiness in Strongly Correlated Clean Systems: An Example of Topological Overprotection, *Phys. Rev. Lett.* **94**, 040402 (2005).
 - [3] J. Haah, Local stabilizer codes in three dimensions without string logical operators, *Phys. Rev. A* **83**, 042330 (2011).
 - [4] V. J. Emery, E. Fradkin, S. A. Kivelson, and T. C. Lubensky, Quantum Theory of the Smectic Metal State in Stripe Phases, *Phys. Rev. Lett.* **85**, 2160 (2000).
 - [5] R. M. Nandkishore and M. Hermele, Fractons, *Annu. Rev. Condens. Matter Phys.* **10**, 295 (2019).
 - [6] M. Pretko, X. Chen, and Y. You, Fracton phases of matter, *Int. J. Mod. Phys. A* **35**, 2030003 (2020).
 - [7] C. G. Callan Jr. and J. A. Harvey, Anomalies and fermion zero modes on strings and domain walls, *Nucl. Phys. B* **250**, 427 (1985).
 - [8] R. Raussendorf, C. Okay, D.-S. Wang, D. T. Stephen, and H. P. Nautrup, Computationally Universal Phase of Quantum Matter, *Phys. Rev. Lett.* **122**, 090501 (2019).

- [9] Y. You, T. Devakul, F. J. Burnell, and S. L. Sondhi, Subsystem symmetry protected topological order, *Phys. Rev. B* **98**, 035112 (2018).
- [10] T. Devakul, D. J. Williamson, and Y. You, Classification of subsystem symmetry-protected topological phases, *Phys. Rev. B* **98**, 235121 (2018).
- [11] D. T. Stephen, H. P. Nautrup, J. Bermejo-Vega, J. Eisert, and R. Raussendorf, Subsystem symmetries, quantum cellular automata, and computational phases of quantum matter, *Quantum* **3**, 142 (2019).
- [12] Y. You, T. Devakul, F. Burnell, and S. Sondhi, Symmetric fracton matter: Twisted and enriched, *Ann. Phys.* **416**, 168140 (2020).
- [13] T. Devakul, W. Shirley, and J. Wang, Strong planar subsystem symmetry-protected topological phases and their dual fracton orders, *Phys. Rev. Research* **2**, 012059(R) (2020).
- [14] N. Seiberg, Field theories with a vector global symmetry, *SciPost Phys.* **8**, 050 (2020).
- [15] N. Seiberg and S.-H. Shao, Exotic symmetries, duality, and fractons in 2+1-dimensional quantum field theory, *SciPost Phys.* **10**, 027 (2021).

- [16] N. Seiberg and S.-H. Shao, Exotic $U(1)$ symmetries, duality, and fractons in 3+1-dimensional quantum field theory, *SciPost Phys.* **9**, 046 (2020).
- [17] N. Seiberg and S.-H. Shao, Exotic $\mathbb{Z}_N$ symmetries, duality, and fractons in 3+1-dimensional quantum field theory, *SciPost Phys.* **10**, 003 (2021).
- [18] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, More exotic field theories in 3+1 dimensions, *SciPost Phys.* **9**, 073 (2020).
- [19] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, FCC lattice, checkerboards, fractons, and quantum field theory, *Phys. Rev. B* **103**, 205116 (2021).
- [20] T. Rudelius, N. Seiberg, and S.-H. Shao, Fractons with twisted boundary conditions and their symmetries, *Phys. Rev. B* **103**, 195113 (2021).
- [21] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, A modified villain formulation of fractons and other exotic theories, *J. Math. Phys.* **62**, 102301 (2021).
- [22] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, Low-energy limit of some exotic lattice theories and UV/IR mixing, *Phys. Rev. B* **104**, 235116 (2021).
- [23] K. Slagle, Foliated Quantum Field Theory of Fracton Order, *Phys. Rev. Lett.* **126**, 101603 (2021).
- [24] Y. You, J. Bibo, T. L. Hughes, and F. Pollmann, Fractonic critical point proximate to a higher-order topological insulator: How does UV blend with IR?, [arXiv:2101.01724](https://arxiv.org/abs/2101.01724).
- [25] Z. Zhou, X.-F. Zhang, F. Pollmann, and Y. You, Fractal Quantum Phase Transitions: Critical Phenomena Beyond Renormalization, [arXiv:2105.05851](https://arxiv.org/abs/2105.05851).
- [26] P.-S. Hsin and K. Slagle, Comments on foliated gauge theories and dualities in 3+1d, *SciPost Phys.* **11**, 032 (2021).
- [27] Y. You and R. Moessner, Fractonic plaquette-dimer liquid beyond renormalization, [arXiv:2106.07664](https://arxiv.org/abs/2106.07664).
- [28] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, Generalized Global Symmetries, *J. High Energy Phys.* **02** (2015) 172.
- [29] M. Qi, L. Radzihovsky, and M. Hermele, Fracton phases via exotic higher-form symmetry-breaking, *Ann. Phys.* **424**, 168360 (2021).
- [30] T. Tay and O. I. Motrunich, Possible realization of the exciton Bose liquid phase in a hard-core boson model with ring-only exchange interactions, *Phys. Rev. B* **83**, 205107 (2011).
- [31] Y. You, Z. Bi, and M. Pretko, Emergent fractons and algebraic quantum liquid from plaquette melting transitions, *Phys. Rev. Research* **2**, 013162 (2020).
- [32] Y. You, F. J. Burnell, and T. L. Hughes, Multipolar topological field theories: Bridging higher-order topological insulators and fractons, *Phys. Rev. B* **103**, 245128 (2021).
- [33] A. Gromov, A. Lucas, and R. M. Nandkishore, Fracton hydrodynamics, *Phys. Rev. Research* **2**, 033124 (2020).
- [34] A. Karch and A. Raz, Reduced conformal symmetry, *J. High Energy Phys.* **04** (2021) 182.
- [35] E. Lake, Renormalization group and stability in the exciton Bose liquid, *Phys. Rev. B* **105**, 075115 (2022).
- [36] S. Vijay, J. Haah, and L. Fu, Fracton topological order, generalized lattice gauge theory and duality, *Phys. Rev. B* **94**, 235157 (2016).
- [37] D. J. Williamson, Fractal symmetries: Ungauging the cubic code, *Phys. Rev. B* **94**, 155128 (2016).
- [38] W. Shirley, K. Slagle, and X. Chen, Foliated fracton order from gauging subsystem symmetries, *SciPost Phys.* **6**, 041 (2019).
- [39] A. Gromov, Towards Classification of Fracton Phases: The Multipole Algebra, *Phys. Rev. X* **9**, 031035 (2019).
- [40] K. Slagle and Y. B. Kim, Quantum field theory of X -cube fracton topological order and robust degeneracy from geometry, *Phys. Rev. B* **96**, 195139 (2017).
- [41] K. Slagle, D. Aasen, and D. Williamson, Foliated field theory and string-membrane-net condensation picture of fracton order, *SciPost Phys.* **6**, 043 (2019).
- [42] A. Yu. Kitaev, Fault tolerant quantum computation by anyons, *Ann. Phys.* **303**, 2 (2003).
- [43] J. Gomis, Z. Komargodski, and N. Seiberg, Phases of adjoint QCD₃ and dualities, *SciPost Phys.* **5**, 007 (2018).
- [44] P.-S. Hsin, H. T. Lam, and N. Seiberg, Comments on one-form global symmetries and their gauging in 3D and 4D, *SciPost Phys.* **6**, 039 (2019).
- [45] X.-G. Wen, Emergent anomalous higher symmetries from topological order and from dynamical electromagnetic field in condensed matter systems, *Phys. Rev. B* **99**, 205139 (2019).
- [46] A. Kapustin and R. Thorngren, Higher symmetry and gapped phases of gauge theories, in *Algebra, Geometry, and Physics in the 21st Century*, edited by D. Auroux, L. Katzarkov, T. Pantev, Y. Soibelman, and Y. Tschinkel, Progress in Mathematics, Vol. 324 (Springer, Cham, Birkhäuser, 2017), pp. 177–202.
- [47] R. Thorngren and C. von Keyserlingk, Higher SPT's and a generalization of anomaly in-flow, [arXiv:1511.02929](https://arxiv.org/abs/1511.02929).
- [48] K. Walker and Z. Wang, (3+1)-TQFTs and topological insulators, *Front. Phys.* **7**, 150 (2012).
- [49] K. Slagle and Y. B. Kim, X -cube model on generic lattices: Fracton phases and geometric order, *Phys. Rev. B* **97**, 165106 (2018).
- [50] W. Shirley, K. Slagle, Z. Wang, and X. Chen, Fracton Models on General Three-Dimensional Manifolds, *Phys. Rev. X* **8**, 031051 (2018).
- [51] W. Shirley, K. Slagle, and X. Chen, Fractional excitations in foliated fracton phases, *Ann. Phys.* **410**, 167922 (2019).
- [52] W. Shirley, K. Slagle, and X. Chen, Foliated fracton order in the checkerboard model, *Phys. Rev. B* **99**, 115123 (2019).
- [53] A. Gromov, Chiral Topological Elasticity and Fracton Order, *Phys. Rev. Lett.* **122**, 076403 (2019).
- [54] K. Slagle, A. Prem, and M. Pretko, Symmetric tensor gauge theories on curved spaces, *Ann. Phys.* **410**, 167910 (2019).
- [55] W. Shirley, K. Slagle, and X. Chen, Twisted foliated fracton phases, *Phys. Rev. B* **102**, 115103 (2020).
- [56] D. Gaiotto, A. Kapustin, Z. Komargodski, and N. Seiberg, Theta, time reversal, and temperature, *J. High Energy Phys.* **05** (2017) 091.
- [57] Z. Komargodski, A. Sharon, R. Thorngren, and X. Zhou, Comments on Abelian Higgs models and persistent order, *SciPost Phys.* **6**, 003 (2019).
- [58] S. Yamaguchi, Gapless edge modes in (4+1)-dimensional topologically massive tensor gauge theory and anomaly inflow for subsystem symmetry, *Prog. Theor. Exp. Phys.* **2022**, 033B08 (2022).
- [59] S. Yamaguchi, Supersymmetric quantum field theory with exotic symmetry in 3+1 dimensions and fermionic fracton phases, *Prog. Theor. Exp. Phys.* **2021**, 063B04 (2021).
- [60] T. Devakul, Y. You, F. J. Burnell, and S. L. Sondhi, Fractal symmetric phases of matter, *SciPost Phys.* **6**, 007 (2019).
- [61] G. W. Moore and N. Seiberg, Taming the conformal zoo, *Phys. Lett. B* **220**, 422 (1989).

- [62] S. Elitzur, G. W. Moore, A. Schwimmer, and N. Seiberg, Remarks on the canonical quantization of the Chern-Simons-Witten theory, *Nucl. Phys. B* **326**, 108 (1989).
- [63] J. Sonnenschein, Chiral bosons, *Nucl. Phys. B* **309**, 752 (1988).
- [64] J. M. Maldacena, G. W. Moore, and N. Seiberg, D-brane charges in five-brane backgrounds, *J. High Energy Phys.* **10** (2001) 005.
- [65] T. Banks and N. Seiberg, Symmetries and strings in field theory and gravity, *Phys. Rev. D* **83**, 084019 (2011).
- [66] A. Kapustin and N. Seiberg, Coupling a QFT to a TQFT and duality, *J. High Energy Phys.* **04** (2014) 001.
- [67] M. Bauer, G. Girardi, R. Stora, and F. Thuillier, A Class of topological actions, *J. High Energy Phys.* **08** (2005) 027.
- [68] C. Córdova, D. S. Freed, H. T. Lam, and N. Seiberg, Anomalies in the space of coupling constants and their dynamical applications I, *SciPost Phys.* **8**, 001 (2020).
- [69] C. Córdova, D. S. Freed, H. T. Lam, and N. Seiberg, Anomalies in the space of coupling constants and their dynamical applications II, *SciPost Phys.* **8**, 002 (2020).