# Wire-width and electron-density dependence of the crossover in the peak of the static structure factor from $2k_F \rightarrow 4k_F$ in one-dimensional paramagnetic electron gases

Ankush Girdhar<sup>1</sup>,<sup>1</sup> Vinod Ashokan<sup>1</sup>,<sup>\*</sup> Rajesh O. Sharma<sup>1</sup>,<sup>2</sup> N. D. Drummond<sup>1</sup>,<sup>3</sup> and K. N. Pathak<sup>4</sup>

<sup>1</sup>Department of Physics, Dr. B. R. Ambedkar National Institute of Technology, Jalandhar, Punjab 144027, India

<sup>2</sup>Department of Physics, Indian Institute of Science, Bengaluru 560012, India

<sup>3</sup>Department of Physics, Lancaster University, Lancaster LA1 4YB, United Kingdom

<sup>4</sup>Centre for Advanced Study in Physics, Panjab University, Chandigarh 160014, India

(Received 29 November 2022; revised 17 February 2023; accepted 6 March 2023; published 15 March 2023)

We use the variational quantum Monte Carlo (VMC) method to study the wire-width (*b*) and electron-density ( $r_s$ ) dependences of the ground-state properties of quasi-one-dimensional paramagnetic electron fluids. The onset of a quasi-Wigner crystal phase is known to depend on electron density and the crossover occurs in the low density regime. We study the effect of wire width on the crossover of the dominant peak in the static structure factor from  $k = 2k_F$  to  $k = 4k_F$ . It is found that, for a fixed electron density, in the charge structure factor the crossover from the dominant peak occurring at  $2k_F$  to  $4k_F$  occurs as the wire width decreases. Our study suggests that the crossover is due to the interplay of both  $r_s$  and  $b < r_s$ . The finite wire-width correlation effect is reflected in the peak height of the charge and spin structure factors. We fit the dominant peaks of the charge and spin structure factors assuming fit functions based on our finite wire-width theory and clues from bosonization, resulting in a good fit of the VMC data. The pronounced peaks in the charge and spin structure factors at  $4k_F$  and  $2k_F$ , respectively, indicate the complete decoupling of the charge and spin degrees of freedom. Furthermore, the wire-width dependence of the electron correlation energy and the Tomonaga-Luttinger parameter  $K_{\rho}$  is found to be significant.

DOI: 10.1103/PhysRevB.107.115414

## I. INTRODUCTION

Many-body correlation effects in one-dimensional (1D) electron systems give rise to fascinating properties with diverse technological applications [1,2]. Advances in fabrication techniques have made the realization of extremely thin wires feasible, although only a few aspects of theoretical predictions have yet been experimentally tested. This is due to the experimental difficulties of realizing ideal and controllable 1D systems. There are several features of 1D systems, such as spin-charge separation [3], power-law behavior of correlation functions, and other physical quantities such as conductance [4], charge fractionalization [5], Wigner crystallization [6–9], and so on, which make them unique and interesting. The physics of 1D electron systems cannot be explained using Landau's Fermi liquid theory due to the fact that singleparticle excitation energies and their inverse lifetimes are of the same order of magnitude.

The exactly solvable model by Tomonaga and Luttinger [10–12] describes the low-energy spectrum of a 1D homogeneous electron gas (HEG) assuming short-range interactions

and a linear dispersion relation. The assumption of short-range interactions is quite successfully used in studies of the properties of various 1D structures such as quasi-one-dimensional conductors where screening between adjacent chains leads to effective short-range interactions within each chain [13]. However, the electrons in systems which do not have effective screening, such as isolated metallic carbon nanotubes [8], interact via the true long-range Coulomb potential V(x) = $e^2/|x|$ ]. For the purpose of theoretical studies of 1D electron systems, there are several choices of confinement model available such as hard wall [14,15], harmonic [16,17], and Coulombic [18]. In all these cases, the effective electronelectron interaction is known to have a 1/|x| long-range tail in the lowest subband of the transverse motion. In this study, we consider electrons interacting through an effective longrange Coulomb interaction using a harmonic confinement model.

One-dimensional interacting electron systems were studied using several techniques such as density functional theory (DFT) [18], exact diagonalization [19], the random phase approximation (RPA) [16,20–24], density matrix renormalization group (DMRG) [25,26], quantum Monte Carlo (QMC) [27–31], lattice regularized diffusion Monte Carlo (LRDMC) [32,33], and bosonization [34]. This last was used by Schulz in an interesting paper [34] studying a 1D electron gas with long-range interactions assuming a linearized kinetic energy dispersion. The major finding of the paper was an extremely slowly decaying  $4k_{\rm F}$  component in the charge-charge correlations, signifying the presence of a 1D Wigner crystal state

<sup>\*</sup>ashokanv@nitj.ac.in

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

even for the weakest long-range Coulomb interaction. However, it is not clear how the finite width of a wire and the coupling strength will affect the Wigner crystallization. In fact, one of the purposes of this paper is to find possible answers to this question. At the same time, in the spin correlation function, a finite  $2k_F$  peak was obtained whereas the  $4k_F$  peak was absent.

The experimental observation of a Wigner crystal is a tricky task since it requires extreme conditions such as low temperatures and low densities [9]. In addition, we show in this paper that it also requires small wire widths. Examining such a fragile crystal requires the use of noninvasive imaging probes. In the case of 1D infinite systems, true long-range order is forbidden due to thermal and quantum fluctuations. However, in finite systems, physicists studied 1D Wigner crystals and produced crystalline correlations due to quasi-long-range order. Introducing long-range Coulomb interactions results in a  $4k_{\rm F}$  oscillation in charge correlations whose decay rate is slower than any power law [34]. The experimental realization of long-range interactions was made possible in a recent work by Shapir et al. [8], where a few electrons (<10) were housed in a carbon nanotube. A second nanotube was used as a scanning probe to measure the spatial charge distribution of electrons in the first nanotube. The measured charge densities were found to exhibit the features of a quasi-Wigner crystal. There is no critical density associated with the onset of the 1D Wigner crystal.

Our aim in this work is to study the effects of confinement and electron correlations on the ground-state properties of a paramagnetic harmonic wire. Unlike our previous work in Ref. [31] we investigate the emergence of quasi-Wigner crystal phases at low density and we examine crossover in the position of the dominant peak in the static structure factor for various wire widths at low density. We employ a variational quantum Monte Carlo (VMC) technique to calculate the ground-state properties of the system under consideration. The paper is structured as follows. The model for confinement of electrons in one dimension is described in Sec. II. The energies for the paramagnetic harmonic wire are presented in Sec. III A. The dependence of the  $2k_F \rightarrow 4k_F$  crossover on wire width is discussed in detail in Sec. III B. In Sec. III C, we present momentum density (MD) data around  $k \sim k_{\rm F}$ as a function of wire width. Our conclusions are drawn in Sec. IV. Hartree atomic units ( $\hbar = |e| = m_e = 4\pi\epsilon_0 = 1$ ) are used throughout this paper.

### **II. MODEL AND CALCULATION DETAILS**

The form of the Hamiltonian used for simulating a 1D electron fluid is

$$\hat{H} = -\frac{1}{2} \sum_{i=1}^{N} \frac{\partial^2}{\partial x_i^2} + \sum_{i < j} \tilde{V}(x_{ij}) + \frac{N}{2} V_{\text{Mad}}, \qquad (1)$$

where  $\tilde{V}(x_{ij})$  and  $V_{\text{Mad}}$  describe the Ewald interaction and Madelung energy, respectively. The Ewald-like interaction for a harmonic wire in a periodic cell of length *L* is

[29,35]

$$\tilde{V}(x_{ij}) = \sum_{m=-\infty}^{\infty} \left[ \frac{\sqrt{\pi}}{2b} e^{(x_{ij}-mL)^2/(4b)^2} \operatorname{erfc}\left(\frac{|x_{ij}-mL|}{2b}\right) - \frac{1}{|x_{ij}-mL|} \operatorname{erf}\left(\frac{|x_{ij}-mL|}{2b}\right) \right] + \frac{2}{L} \sum_{n=1}^{\infty} E_1[(bGn)^2] \cos(Gnx_{ij}), \quad (2)$$

where  $G = 2\pi/L$ , *b* is the wire width and  $E_1$  is the exponential integral function.

We considered a 1D HEG in a simulation cell of length  $L = 2Nr_s$  subject to periodic boundary conditions, with N being the total number of electrons in the periodic cell. We keep  $N^{\uparrow} = N^{\downarrow}$  odd to avoid shell degeneracy effects. A Slater-Jastrow-backflow wave function was used in the VMC calculations [36,37]. The orbitals in the Slater determinant were chosen to be plane waves with wave vectors up to  $k_F = \pi/(4r_s)$  for the paramagnetic systems. The VMC method as implemented in the CASINO code [38] was used to calculate ground-state expectation values. More simulation details can be found in our previous works [29–31].

In a 1D electron gas with an electron-electron interaction potential that is appropriate for a finite wire of width b, opposite-spin electrons have a small but nonzero contact probability density. By contrast, in an infinitely thin 1D electron gas, the contact probability density is zero for both samespin and opposite-spin electrons. QMC calculations for 1D electron gases with finite-width interactions are challenging because the width introduces features on the small lengthscale b into the wave function near coalescence points. Typically bwill be much smaller than other lengthscales in the problem, such as the density parameter  $r_s$ .

One issue that affects diffusion quantum Monte Carlo calculations of finite-width wires is ergodicity; to explore all of configuration space, opposite-spin electrons have to be able to pass one another. However the "quasinodes" at opposite-spin coalescence points lead to a very low (but nonzero) probability of electrons passing each other, resulting in very long correlation times. This is much less of an issue in VMC, where there is no need to work in the limit of small time steps so that electron moves past coalescence points are permitted. A second issue is simply the challenge of creating and optimizing a trial wave function that contains features on the small length scale b as well as the large lengthscale L of the simulation cell.

#### **III. RESULTS AND DISCUSSION**

#### A. Energies

The ground-state energy per electron for the harmonic wire is calculated for  $r_s = 0.5$ , 1, 2, and 5 and for b = 0.1, 0.3, 0.5, 0.7, 1, 2 a.u. We calculate the energies for N = 30, 50, 74, 98 electrons and then extrapolate them to obtain the thermodynamic limit. The ground-state energy is found to scale with system size as  $E(N) = E_{\infty} + BN^{-2}$  [29] where *B* and  $E_{\infty}$  are fitting parameters. We also calculate the correlation energy



FIG. 1. Correlation energy per particle ( $E_c$ ) as a function of wire width *b* for  $r_s = 0.5$ , 1, and 2 (top to bottom).

per electron using the ground-state energy. These energies are tabulated in Table I of the Appendix A. The wire-width dependence of correlation energy per particle for different values of density parameter  $r_s$  is drawn in Fig. 1.

# **B.** Structure and correlation function

We calculate the VMC charge and spin static structure factors (SSFs), which are defined as

$$S_{\rho\rho}(k) = S_{\uparrow\uparrow} + S_{\uparrow\downarrow}, \qquad (3)$$

$$S_{\sigma\sigma}(k) = S_{\uparrow\uparrow} - S_{\uparrow\downarrow}, \qquad (4)$$

respectively, for a paramagnetic quantum wire. Here  $S_{\uparrow\uparrow}$  and  $S_{\uparrow\downarrow}$  are parallel-spin and antiparallel-spin SSFs. The SSF is a crucial property of electron fluids that is used to look into the structural properties of the electron system in reciprocal space. The charge SSF, also known as the total SSF, describes correlations between electron pairs irrespective of their spins. The spin PCF describes the difference between pair correlations for parallel- and antiparallel-spin electron pairs. To study the charge-charge and spin-spin correlation functions in real space, we define alternative quantities, namely, charge and spin pair correlation functions (PCFs), as

$$g_{\rho\rho} = \frac{g_{\uparrow\uparrow} + g_{\uparrow\downarrow}}{2},\tag{5}$$

$$g_{\sigma\sigma} = \frac{g_{\uparrow\uparrow} - g_{\uparrow\downarrow}}{2},\tag{6}$$

respectively. Here  $g_{\uparrow\uparrow}$  and  $g_{\uparrow\downarrow}$  are parallel-spin and antiparallel-spin PCFs, respectively. A peak at  $k = 4k_F$  in the charge structure factor corresponds to slowly decaying oscillations of period  $2r_s$ , which is the average interparticle spacing. This is a signature of crossover to a ordered phase known as a Wigner crystal. For the case of ferromagnetic infinitely thin wires, the singularity at  $k = 4k_F$  is absent for densities  $r_s < 15$  a.u., but a small feature at  $4k_F$  starts to develop at  $r_s \approx 15$  [29]. The effect of singularity at  $4k_F$  was observed recently [39] for the case of electron-electron biwires, where the parallel wire effectively provides an extra spin degree of freedom to electrons. In the case of a paramagnetic quasiquantum wire, it was verified in previous works [27,28] that the dominant  $4k_{\rm F}$  peak height grows sublinearly with system size *N* at low density, which is a signature of a quasi-Wigner crystal. The adjective "quasi" has been used to emphasize the absence of true long-range order due to quantum fluctuations. Further, it was found that, for any fixed wire width, this crossover occurs as one decreases the electron density (i.e., low density with large values of  $r_{\rm s}$ ). In this work, we observed a similar correlation effect by decreasing the wire

width at fixed coupling parameter  $r_s$ .

The charge structure factor  $S_{\rho\rho}(k)$  is plotted for several values of *b* and  $r_s$  for N = 98 electrons in Fig. 2. In Fig. 2(a),  $S_{\rho\rho}(k)$  shows a peak at  $k = 2k_F$  and no peak at  $4k_F$  is present. This leads to oscillations of period  $4r_s$  in charge-charge PCF, as can be seen in Fig. 2(d). It is evident from Fig. 2(b) that, as *b* decreases for  $r_s = 2$ , the  $4k_F$  peak grows and the  $2k_F$  peak is suppressed, indicating the crossover  $(2k_F \rightarrow 4k_F)$  to an ordered phase. This crossover leads to a change in the period of the oscillations in the charge-charge PCF from  $4r_s$  to  $2r_s$ , as apparent in Fig. 2(e). Further, it can be noted from Fig. 2(c) that the  $2k_F$  peak is absent, but decreasing *b* leads to an increase in the height of the  $4k_F$  peak. This is also reflected in a more structured behavior of  $g_{\rho\rho}$ , with oscillation period  $2r_s$  in Fig. 2(f).

The spin structure factor  $S_{\sigma\sigma}(k)$  is plotted for several values of b and  $r_s$  for N = 98 electrons in Fig. 3.  $S_{\sigma\sigma}(k)$  is found to show a peak at  $k = 2k_{\rm F}$ , which grows as b decreases for all the densities considered [see Figs. 3(a) to 3(c)]. This is followed by the spin-spin PCF in Figs. 3(d) to 3(f), where the amplitude of oscillations of period  $4r_s$  is enhanced as b decreases. It can also be noted from Figs. 3(a) to 3(c) that the peak height increases with increasing  $r_s$  for a given value of b. In the present work, we observed complete decoupling of the charge and spin degrees of freedom in the SSFs. For example, in  $r_s = 2$ , there is a crossover from  $2k_F \rightarrow 4k_F$  behavior, with complete disappearance of  $2k_{\rm F}$  peak at b = 0.5, in the charge SSF [Fig. 2(b)], while for the same values of band  $r_s$  the spin SSF shows a peak at  $k = 2k_F$  [see Fig. 3(c)]. The spin and charge excitations in the SSFs manifest the complete separation of spin and charge of electrons due to strong Coulomb correlation effects induced by the small transverse confinement width.

The crossover observed is between (i) a state in which opposite spin electrons can easily slide past each other (large  $b/r_s$ ; in this case oscillations in both charge and spin PCFs have period  $4r_s$  because correlation is only important for same-spin electrons and negligible for opposite-spin pairs) and (ii) a state in which opposite spin electrons cannot slide past each other (small  $b/r_s$ ; in this case the system is like a floating antiferromagnetic crystal lattice; the spin PCF has period  $4r_s$ , whereas the charge PCF has period  $2r_s$ ). The "fluid-to-crystal" crossover can be visualized in terms of the behavior of interlocking lattices of spin-up electrons and spin-down electrons, with the crossover occurring at the point at which the lattices start being able to slide past each other.

On the basis of our finite wire-width theory [31], the SSF is a function of  $b/r_s$ . In view of this, and the clues from bosonization theory [25,34], we propose that the dependence of the peak heights  $S_{\rho\rho}(4k_F, b)$  and  $S_{\sigma\sigma}(2k_F, b)$  on wire width



FIG. 2. (a)–(c) Charge structure factor  $S_{\rho\rho}(k)$  plotted against  $k/k_{\rm F}$  and (d)–(f) charge-charge pair correlation function plotted against  $r/r_{\rm s}$  for  $r_{\rm s} = 0.5$ , 2, and 5 for various *b* values with N = 98 electrons. The insets in (b) and (c) show the peak at  $k = 4k_{\rm F}$  in greater detail.

can be represented as

$$S_{\rho\rho}(4k_{\rm F},b) = a_0 \frac{L r_{\rm s}}{b} \exp\left(-4c\sqrt{\ln\frac{L r_{\rm s}}{b}}\right) + a_1, \quad (7)$$

and

$$S_{\sigma\sigma}(2k_{\rm F},b) = -a_2\left(\sqrt{\ln\frac{L\,r_{\rm s}}{b}} + \frac{1}{c}\right)\exp\left(-c\sqrt{\ln\frac{L\,r_{\rm s}}{b}}\right) + a_3,$$
(8)

where  $a_0$ ,  $a_1$ ,  $a_2$ ,  $a_3$ , and c are fitting parameters and  $L = 2Nr_s$ .

Finite wire-width effects are studied in Fig. 4. The  $4k_F$  and  $2k_F$  peak heights in the charge SSF are plotted against the wire width in Fig. 4 (top) for  $r_s = 0.5$ , 1, and 2. The dependence

of the peak heights on b is found to be significant for  $r_s = 1$ and 2, and for these coupling parameters both the  $4k_{\rm F}$  and  $2k_{\rm F}$ peaks are present. On the other hand, for  $r_s = 0.5$ , there is no  $4k_{\rm F}$  peak present for any of the wire widths considered and even the  $2k_{\rm F}$  peak height depends only weakly on b (at least until  $b \ll r_s$ ). The  $4k_F$  peak heights were fitted to Eq. (7), which was deduced on the basis of bosonization and which shows a good representation of the numerically calculated data. The wire-width dependence of the  $2k_{\rm F}$  peak height in the spin SSF is plotted in Fig. 4 (bottom) for  $r_s = 0.5$ , 1, and 2. It is observed that the peak height in the spin SSF depends on both  $r_s$  and b. It is also observed that, for a fixed value of electron coupling, the amplitude of the peak height at  $k = 2k_{\rm F}$  in the spin SSF strongly depends on the confinement width. Further, the  $2k_{\rm F}$  peak heights are well fitted by Eq. (8).



FIG. 3. (a)–(c) Spin structure factor  $S_{\sigma\sigma}(k)$  plotted against  $k/k_{\rm F}$  and (d)–(f) spin-spin pair correlation function plotted against  $r/r_{\rm s}$  for  $r_{\rm s} = 0.5$ , 1, and 2 for various b values with N = 98 electrons. The insets (a), (b), and (c) show the peak at  $k = 2k_{\rm F}$  in greater detail.

It should be noted from Fig. 4 that the SSF of the finitewidth paramagnetic wire is not expected to converge to that of a ferromagnetic infinitely thin wire in the limit  $b \rightarrow 0$ because the latter system has only parallel spin correlations whereas the former system has both parallel and antiparallel spin correlations. Further, the quasi-long-range order in ferromagnetic and paramagnetic electron gases is represented by dominant peaks in the charge structure factor  $S_{\rho\rho}(k)$  at  $k = 2k_F$  and  $k = 4k_F$ , respectively, when b is small. The dominant peaks in  $S_{\rho\rho}(k)$  for these two cases correspond to the same  $2r_s$  oscillation period in the charge-charge PCF. A nonmonotonic shift of the dominant peak from  $4k_F$  to  $2k_F$ for  $b \rightarrow 0$  in a paramagnetic wire is not therefore expected. This behavior is consistent with Eq. (7), the model formula obtained from the bosonization prediction [25,34]. It is known that the Lieb-Mattis theorem is not applicable for infinitely thin wires, and in fact, the ferromagnetic and paramagnetic states are energetically degenerate in this limit [29]. However, physical observables which do not commute with the Hamiltonian, such as the SSF, need not be same for infinitely thin paramagnetic and ferromagnetic wires.

### C. Momentum density and TL parameters

TL theory suggests that the 1D MD has a peculiar power-law behavior, which is continuous at  $k_F$  with a singular first-order derivative and takes the form  $n(k) = n(k_F) + A[sgn(k - k_F)]|k - k_F|^{\alpha}$  close to  $k = k_F$ , where  $n(k_F)$ , A, and  $\alpha$  are constants [11,40]. The method used to estimate the inter-



FIG. 4. (Top) Charge SSF peak height at  $k = 4k_F$  as a function of wire width *b* for  $r_s = 0.5$ , 1, and 2 for N = 98 electrons. The peak heights at  $k = 4k_F$  are fitted by Eq. (7) and the fits are shown by solid lines, whereas the  $2k_F$  peak heights are joined by dashed lines. (Bottom) Height of spin SSF plotted against *b* for  $r_s = 0.5$ , 1, and 2 for N = 98 electrons and fitted by Eq. (8).

action exponent  $\alpha$  is discussed in great detail in our previous works [29–31].

In Fig. 5, the MD is plotted for N = 98 electrons at several values of *b* and  $r_s$ . The effect of wire width on the behavior of the MD is clearly visible in the figure. It can be seen that as *b* decreases or  $r_s$  increases, the power-law behavior becomes more appreciable. It can also be noted that the value of n(k = 0) decreases from 1 and the value of n(k) increases for  $k > k_F$  as *b* decreases or  $r_s$  increases.

We also compare our QMC MD data with bosonization results [25,34], finding a good fit of the jump in the MD at  $k = k_F$  at finite size by the asymptotic scaling function predicted by bosonization theory

$$\Delta n(k_{\rm F}) \sim L e^{-c(\ln L)^{3/2}}.$$
(9)

Figure 6 shows that Eq. (9) provides quite a good numerical representation of the system-size scaling of the jump in the MD at  $k = k_{\rm F}$ .

The TL parameter  $K_{\rho}$  for repulsive interactions ( $K_{\rho} < 1$ ) and exponent  $\alpha$  are plotted against *b* for  $r_s = 0.5$ , 1, and 2 in



FIG. 5. MD against  $k/k_{\rm F}$  for different values of wire width b and  $r_{\rm s}$  for N = 98 electrons.

Fig. 7. It was observed that, as *b* decreases or  $r_s$  increases,  $K_\rho$  decreases and the interacting electrons become more strongly correlated. Hence it is found that  $K_\rho$  significantly depends on the confinement width *b* and  $r_s$ .

# **IV. CONCLUSION**

We studied the wire-width dependence of the ground-state properties of quasi-1D quantum wires for several electron densities. We analyzed the effects of adjusting the wire width on the charge structure factor, where a crossover from a peak at  $2k_F$  to a peak at  $4k_F$  was observed as *b* decreases. The crossover indicates the onset of a quasi-Wigner crystalline state in the 1D electron fluid. The PCF was also found to



FIG. 6. Jump in the MD at  $k = k_F$  at finite size, plotted against system size and fitted by  $\Delta n(k_F) = aLe^{-c(\ln L)^{3/2}}$ . The fitted parameter values are a = 1.03(2) a.u. and c = 0.731(2) for  $r_s = 1$  and b = 1; and a = 0.93(5) a.u. and c = 0.777(7) for  $r_s = 2$  and b = 2.



FIG. 7. TL parameter  $K_{\rho}$  against wire width *b* for different values of the density parameter  $r_s$  as indicated in the plot. The TL parameter is obtained by fitting MD data to appropriate fitting functions to obtain the exponent  $\alpha$  and  $K_{\rho}$ . The exponent  $\alpha$  is plotted against *b* in the inset.

show a more structured behavior as *b* decreases, which again suggested a crossover to a quasi-Wigner crystal phase. We found the crossover does not appear for all values of  $r_s$  and *b*, a result at variance with the bosonization findings of Schulz in Ref. [34]. We also fitted the dominant peaks of the charge and spin structure factors assuming fitting functions on the basis of our finite wire-width theory and clues from bosonization,

giving a good fit of the VMC data. The crossover occurred only for those density and wire-width values which brought strong correlations in the system.

The correlation energy as a function of *b* and  $r_s$  was represented by a formula which fits our VMC data perfectly. The MD n(k) was found to show a power-law behavior. We found that n(k) at k = 0 reduces from 1, whereas n(k) increases beyond  $k_F$  as *b* decreases or  $r_s$  increases. We find the TL liquid exponent  $\alpha$  by fitting the MD data to an appropriate fitting function near  $k_F$ . The TL parameter  $K_{\rho}$  was found to depend significantly on the wire width at a constant  $r_s$ .

# ACKNOWLEDGMENTS

A.G. would like to thank MHRD for financial support. K.N.P. acknowledges the National Academy of Sciences of India, Prayagraj for partial financial assistance. R.O.S. acknowledges SERB for providing NPDF Grant No. PDF/2021/000546. V.A. acknowledges financial support from DST-SERB Grant No. EEQ/2019/000528 and the National PARAM Supercomputing Facility at C-DAC Pune for computational resources.

#### APPENDIX

The thermodynamic limit of the VMC ground-state energy per electron  $E_{\infty}$  and the correlation energy per electron  $E_{c}$ are calculated by extrapolation from calculations performed at several different values of system size N at different values of b and  $r_{s}$ ; these are tabulated in Table I.

TABLE I. VMC energies in a.u. per electron [E(N)] for N = 30, 50, 74, and 98 for a paramagnetic harmonic wire.  $E_{\infty}$  and  $E_c$  denote the ground-state energy per electron extrapolated to the thermodynamic limit and the correlation energy per electron, respectively.

$(r_{\rm s},b)$	<i>E</i> (30)	<i>E</i> (50)	<i>E</i> (74)	E(98)	$E_\infty$	$E_{ m c}$
(0.5, 0.1)	-0.937394(8)	-0.935599(7)	-0.935042(5)	-0.934791(5)	-0.93455(2)	-0.14477(2)
(0.5, 0.3)	-0.336230(3)	-0.334478(2)	-0.333931(2)	-0.333725(2)	-0.333473(7)	-0.032178(7)
(0.5, 0.5)	-0.127807(2)	-0.126094(1)	-0.1255490(9)	-0.1253469(9)	-0.125100(9)	-0.014176(9)
(0.5, 0.7)	-0.0145572(9)	-0.0128579(7)	-0.0123206(6)	-0.0121214(5)	-0.011877(8)	-0.008008(8)
(0.5, 1.0)	0.0853158(5)	0.0869805(4)	0.0875067(3)	0.0877014(3)	0.087942(8)	-0.004262(8)
(0.5, 2.0)	0.225663(2)	0.2272236(1)	0.22772541(9)	0.22791182(8)	0.228151(5)	-0.001176(5)
(1.0, 0.1)	-0.899453(5)	-0.898910(4)	-0.898695(3)	-0.898505(3)	-0.89847(6)	-0.23295(6)
(1.0, 0.3)	-0.478440(2)	-0.477901(2)	-0.477725(2)	-0.477671(1)	-0.477589(4)	-0.074459(4)
(1.0, 0.5)	-0.328023(1)	-0.327485(1)	-0.327323(1)	-0.3272540(9)	-0.327179(4)	-0.036280(4)
(1.0, 0.7)	-0.245460(1)	-0.2449252(8)	-0.2447628(7)	-0.2446985(6)	-0.244622(2)	-0.021498(2)
(1.0, 1.0)	-0.1710782(7)	-0.1705563(5)	-0.1703899(4)	-0.1703292(4)	-0.170254(3)	-0.011984(3)
(1.0, 2.0)	-0.0610819(3)	-0.0605826(2)	-0.0604240(2)	-0.0603641(1)	-0.060292(3)	-0.003585(3)
(2.0, 0.1)	-0.687882(6)	-0.687592(7)	-0.687429(6)	-0.687309(2)	-0.68726(3)	-0.24309(3)
(2.0, 0.3)	-0.428893(1)	-0.428709(1)	-0.4286485(9)	-0.4285848(9)	-0.42858(2)	-0.11954(2)
(2.0, 0.5)	-0.321088(1)	-0.3209185(8)	-0.3208670(7)	-0.3208426(6)	-0.320819(2)	-0.072522(2)
(2.0, 0.7)	-0.2583233(8)	-0.2581581(7)	-0.2581048(6)	-0.2580852(5)	-0.258061(1)	-0.048212(1)
(2.0, 1.0)	-0.2006410(7)	-0.2004732(5)	-0.2004218(5)	-0.2004024(4)	-0.2003781(3)	-0.0292266(3)
(2.0, 2.0)	-0.1146782(3)	-0.1145145(2)	-0.1144626(2)	-0.1144422(2)	-0.114419(1)	-0.009582(1)
(5.0, 0.1)	-0.393883(2)	-0.393808(3)	-0.393718(2)	-0.3935256(8)	-0.39352(6)	-0.16401(6)
(5.0, 0.3)	-0.284488(3)	-0.283437(9)	-0.284425(1)	-0.2843393(4)	-0.28433(5)	-0.10955(5)
(5.0, 0.5)	-0.234260(4)	-0.234304(6)	-0.234296(8)	-0.2342142(4)	-0.23420(3)	-0.08464(3)
(5.0, 0.7)	-0.201875(3)	-0.2019512(7)	-0.2019229(7)	-0.2019149(2)	-0.201911(9)	-0.068787(9)
(5.0, 1.0)	-0.168710(3)	-0.1687108(6)	-0.1686853(6)	-0.1686792(2)	-0.168670(4)	-0.052681(4)
(5.0, 2.0)	-0.1102775(8)	-0.1102383(7)	-0.1102273(5)	-0.1102212(1)	-0.1102150(4)	-0.0260165(4)

- G. F. Giuliani and G. Vignale, *Quantum Theory of the Electron Liquid* (Cambridge University Press, Cambridge, England, 2005).
- [2] T. Giamarchi, *Quantum Physics in One Dimension* (Clarendon, Oxford, 2004).
- [3] O. M. Auslaender, H. Steinberg, A. Yakoby, Y. Tserkovnyak, B. I. Halperin, K. W. Baldwin, L. N. Pfeiffer, and K. W. West, Science 308, 88 (2005).
- [4] M. Bockrath, D. H. Cobden, J. Lu, A. G. Rinzler, R. E. Smalley, L. Balents, and P. L. McEuen, Nature (London) 397, 598 (1999).
- [5] H. Steinberg, G. Barak, A. Yacoby, L. N. Pfeiffer, K. W. West, B. I. Halperin, and K. Le Hur, Nat. Phys. 4, 116 (2008).
- [6] V. V. Deshpande and M. Bockrath, Nat. Phys. 4, 314 (2008).
- [7] S. Pecker, F. Kuemmeth, A. Secchi, M. Rontani, D. C. Ralph, P. L. McEuen, and S. Ilani, Nat. Phys. 9, 576 (2013).
- [8] I. Shapir, A. Hamo, S. Pecker, C. Moca, Ó. Legeza, G. Zarand, and S. Ilani, Science 364, 870 (2019).
- [9] N. T. Ziani, F. Cavaliere, K. G. Becerra, and M. Sassetti, Crystals 11, 20 (2020).
- [10] S. Tomonaga, Prog. Theor. Phys. 5, 544 (1950).
- [11] J. M. Luttinger, J. Math. Phys. 4, 1154 (1963).
- [12] F. D. M. Haldane, Phys. Rev. Lett. 47, 1840 (1981).
- [13] H. J. Schulz, J. Phys. C 16, 6769 (1983).
- [14] A. Gold and A. Ghazali, Phys. Rev. B 41, 7626 (1990).
- [15] B. Y.-K. Hu and S. Das Sarma, Phys. Rev. B 48, 5469 (1993).
- [16] W. I. Friesen and B. Bergersen, J. Phys. C 13, 6627 (1980).
- [17] G. Y. Hu and R. F. O'Connell, Phys. Rev. B 42, 1290 (1990).
- [18] Y. Sun and G. Kirczenow, Phys. Rev. B 47, 4413 (1993).
- [19] D. Poilblanc, S. Yunoki, S. Maekawa, and E. Dagotto, Phys. Rev. B 56, R1645 (1997).
- [20] R. Bala, R. K. Moudgil, S. Srivastava, and K. N. Pathak, Eur. Phys. J. B 87, 5 (2014).

- [21] V. Ashokan, R. Bala, K. Morawetz, and K. N. Pathak, Eur. Phys. J. B 91, 29 (2018).
- [22] K. Morawetz, V. Ashokan, R. Bala, and K. N. Pathak, Phys. Rev. B 97, 155147 (2018).
- [23] V. Ashokan, R. Bala, K. Morawetz, and K. N. Pathak, Phys. Rev. B 101, 075130 (2020).
- [24] L. Calmels and A. Gold, Europhys. Lett. 39, 539 (1997).
- [25] G. Fano, F. Ortolani, A. Parola, and L. Ziosi, Phys. Rev. B 60, 15654 (1999).
- [26] Z.-H. Li, J. Phys.: Condens. Matter **31**, 255601 (2019).
- [27] A. Malatesta and G. Senatore, J. Phys. IV France 10, Pr5-341 (2000).
- [28] M. Casula and G. Senatore, ChemPhysChem 6, 1902 (2005).
- [29] R. M. Lee and N. D. Drummond, Phys. Rev. B 83, 245114 (2011).
- [30] V. Ashokan, N. D. Drummond, and K. N. Pathak, Phys. Rev. B 98, 125139 (2018).
- [31] A. Girdhar, V. Ashokan, N. D. Drummond, K. Morawetz, and K. N. Pathak, Phys. Rev. B 105, 115140 (2022).
- [32] M. Casula, S. Sorella, and G. Senatore, Phys. Rev. B **74**, 245427 (2006).
- [33] L. Shulenburger, M. Casula, G. Senatore, and R. M. Martin, Phys. Rev. B 78, 165303 (2008).
- [34] H. J. Schulz, Phys. Rev. Lett. 71, 1864 (1993).
- [35] V. R. Saunders, C. Freyria-Fava, R. Dovesi, and C. Roetti, Comput. Phys. Commun. 84, 156 (1994).
- [36] N. D. Drummond, M. D. Towler, and R. J. Needs, Phys. Rev. B 70, 235119 (2004).
- [37] P. López Ríos, A. Ma, N. D. Drummond, M. D. Towler, and R. J. Needs, Phys. Rev. E 74, 066701 (2006).
- [38] R. J. Needs, M. D. Towler, N. D. Drummond, P. López Ríos, and J. R. Trail, J. Chem. Phys. 152, 154106 (2020).
- [39] R. O. Sharma, N. D. Drummond, V. Ashokan, K. N. Pathak, and K. Morawetz, Phys. Rev. B 104, 035149 (2021).
- [40] D. C. Mattis and E. H. Lieb, J. Math. Phys. 6, 304 (1965).