

Gapped lineon and fracton models on graphs

Pranay Gorantla¹, Ho Tat Lam², Nathan Seiberg³, and Shu-Heng Shao⁴

¹Physics Department, Princeton University, Princeton, New Jersey 08544, USA

²Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

³School of Natural Sciences, Institute for Advanced Study, Princeton, New Jersey 08540, USA

⁴C. N. Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook, New York 11794, USA



(Received 26 November 2022; revised 15 February 2023; accepted 16 February 2023; published 9 March 2023)

We introduce a \mathbb{Z}_N stabilizer code that can be defined on any spatial lattice of the form $\Gamma \times C_{L_z}$, where Γ is a general graph. We also present the low-energy limit of this stabilizer code as a Euclidean lattice action, which we refer to as the anisotropic \mathbb{Z}_N Laplacian model. It is gapped, robust (i.e., stable under small deformations), and has lineons. Its ground-state degeneracy (GSD) is expressed in terms of a “mod N -reduction” of the Jacobian group of the graph Γ . In the special case when space is an $L \times L \times L_z$ cubic lattice, the logarithm of the GSD depends on L in an erratic way and grows no faster than $O(L)$. We also discuss another gapped model, the \mathbb{Z}_N Laplacian model, which can be defined on any graph. It has fractons and a similarly strange GSD.

DOI: [10.1103/PhysRevB.107.125121](https://doi.org/10.1103/PhysRevB.107.125121)

I. INTRODUCTION

In recent years, there has been a rapid development in the study of exotic lattice models in condensed matter systems. Some models, known as fractons [1–3] exhibit a variety of surprising features. These include a robust ground-state degeneracy (GSD) that grows sub-extensively in the system size [4], as well as particle excitations with restricted mobility. Many of these unusual properties can be understood as following from their exotic global symmetries. These symmetries are also the underlying reasons why these lattice models defy a conventional continuum limit. See [5–10] for reviews on these novel topological phases of matter and their exotic global symmetries.

Most of these exotic lattice models are defined on a cubic lattice, or lattices with additional structure such as foliation [11–21]. It is then natural to ask if there are exotic models that can be defined on a general lattice graph. Recently, two such lattice models, the Laplacian ϕ -theory and the $U(1)$ Laplacian gauge theory, were proposed in [22,23] using the discrete Laplacian operator Δ_L . (See also [24] for a model along this line.) The former has a large GSD being the number of spanning trees of the spatial graph, which is a common measure of complexity, but it does not have fractons. The latter has defects representing immobile fracton particles, but it has no large GSD.

The \mathbb{Z}_N version of the $U(1)$ Laplacian gauge theory has a large GSD and fractons, but the GSD is not robust against perturbations by local operators. This motivates us to consider a certain anisotropic generalization, which we call the *anisotropic \mathbb{Z}_N Laplacian model*.¹ It has the following salient

features, some of which are reminiscent of the celebrated Haah’s code [2]:

(1) It can be placed on a spatial lattice of the form $\Gamma \times C_{L_z}$, where Γ is a general graph and C_{L_z} is a cycle graph on L_z vertices, or a 1D periodic chain with L_z sites. See Fig. 1.

(2) The GSD is robust² and is given by

$$\text{GSD} = |\text{Jac}(\Gamma, N)|^2, \quad (1.1)$$

where $\text{Jac}(\Gamma, N)$ is a “mod N -reduction” of the Jacobian group $\text{Jac}(\Gamma)$ of Γ .

(3) It has lineons that can only move in the z direction if Γ is an infinite two-dimensional square lattice.

(4) In the special case when the spatial lattice is a $L_x \times L_y \times L_z$ cubic lattice and when $N = p$ is prime, we have

$$\begin{aligned} \log_p \text{GSD} &= 2 \dim_{\mathbb{Z}_p} \frac{\mathbb{Z}_p[X, Y]}{(Y(X-1)^2 + X(Y-1)^2, X^{L_x} - 1, Y^{L_y} - 1)}. \end{aligned} \quad (1.2)$$

The definition and the explicit evaluation of this formula are discussed in Appendix C. It depends on the number-theoretic properties of L_x, L_y . Interestingly, there exists a sequence of L_x, L_y going to infinity such that the $\log_p \text{GSD} \sim O(L_x, L_y)$, but there is also a sequence such that $\log_p \text{GSD}$ stays at order 1 if $p > 2$. See Fig. 2.

We present this model both in terms of the low-energy limit of a stabilizer code in the Hamiltonian formalism, and in terms

¹The relation between the \mathbb{Z}_N Laplacian model and its anisotropic uplift is analogous to that between the 2+1D \mathbb{Z}_N Ising plaquette model [25] and the 3+1D anisotropic lineon model in [13,26].

²On a general graph, there is no notion of locality, and therefore we cannot discuss local operators and match them between the UV and the IR theories. Consequently, the discussion of robustness is ambiguous. This is not the case on regular lattices where the usual discussion of local operators and robustness applies. In that case, the anisotropic model is robust as we will show in Sec. III B 4.




FIG. 1. The spatial lattice $\Gamma \times C_{L_z}$: the black lines correspond to the edges of the graph Γ , and the green lines represent the z -links between two copies of Γ . Each site of the lattice is labeled as (i, z) , where i denotes a vertex of the graph Γ and z denotes a vertex of C_{L_z} .

of a Euclidean lattice model using an integer BF action [27]. We compare the four Laplacian lattice models in Table I.³

Following [22,23,25–35], we focus on the exotic global symmetries of this model. The symmetries of the models on Γ (the first three models in Table I) are not subsystem global symmetries. The symmetry operators are supported on most (or all) of the sites of Γ , rather than on a small subset of them. The precise subset depends delicately on the details of Γ . Yet there are many such symmetries. In this sense these symmetries are generalizations of the dipole symmetries [36,37] on cubic lattices, which are also supported on the entire lattice. The difference is that the dipole symmetries have simple dependence on the coordinates, while here the dependence on the coordinates is more complicated.

This is not the case in the anisotropic \mathbb{Z}_N Laplacian model (the fourth model in Table I). Here the symmetries act at fixed z and in that sense they are subsystem symmetries. (See [38,39] for other anisotropic fractal models with symmetries that act at fixed z .) In fact, as we will discuss below, at low energy these symmetries are independent of z and are similar to one-form global symmetries.⁴

The rest of the paper is organized as follows. In Sec. II we introduce some necessary graph theory background, including

³In the table we assume the θ angle of the $U(1)$ Laplacian theory is not π , otherwise the GSD is 2.

⁴More generally, we can classify symmetries by the difference operators that annihilate the transformation parameters α . For example, on a regular lattice, an ordinary symmetry has $\Delta_x \alpha = \Delta_y \alpha = \Delta_z \alpha = 0$, a dipole symmetry has $\Delta_x \Delta_x \alpha = \Delta_x \Delta_y \alpha = \Delta_y \Delta_y \alpha = 0$, etc. (And, of course, α can carry more indices for the various fields or directions in spacetime.) More interesting examples arise in theories associated with Haah’s code [2], where the difference equations are




FIG. 2. The logarithm of the ground-state degeneracy \log_N GSD of the 3+1D anisotropic \mathbb{Z}_N Laplacian model of lineons on a cubic lattice with $L \times L \times L_z$ sites (i.e., $\Gamma = C_L \times C_L$) for $N = 2$ (red) and $N = 3$ (blue), and $1 \leq L \leq 100$. While \log_N GSD grows steadily as $\sim 4L$ for $N = 2$, it behaves erratically for $N = 3$. In fact, for $N = 3$, there are infinitely many L for which it is just 2, and also infinitely many L (powers of 3) for which it is $\sim 4L$. Here the GSD for both $N = 2, 3$ was calculated using techniques from commutative algebra explained in Appendix C 1. In Sec. IV B, we give a simpler derivation of the GSD for $N = 2$ by relating the 3+1D anisotropic \mathbb{Z}_2 Laplacian model to the 3+1D anisotropic \mathbb{Z}_2 lineon model [13,26] on a tilted lattice.

the discrete Laplace operator and the Jacobian group of a graph. In Sec. III we introduce the stabilizer code and the Euclidean integer BF action for the anisotropic \mathbb{Z}_N Laplacian model. We derive the general expression for the GSD (1.1) and discuss the restricted mobility of the lineon defects from timelike symmetries.⁵ Section IV considers the special case when the spatial lattice is a cubic lattice and when N is prime. The GSD reduces to (1.2) and we discuss its asymptotic behaviors. Appendix A discusses \mathbb{Z}_N -valued discrete harmonic functions on a general graph. Appendixes B and C contain the detailed computation of the GSD and the mobility restrictions for the anisotropic \mathbb{Z}_N Laplacian model on a cubic lattice with N a prime number. In Appendix D, we study the \mathbb{Z}_N Laplacian model of fractons, which is not robust.

II. GRAPH THEORY PRIMER

In this section, we review some well-known facts about a finite graph, and \mathbb{Z}_N -valued functions on the graph. A good reference on this subject is [40]. See also [22] for more discussion on these topics in related lattice models.

Let Γ be a simple, undirected, connected graph on N vertices.⁶ Here *simple* means there is at most one edge between

$$(\Delta_x + \Delta_y + \Delta_z)\alpha = 0 \text{ and } [\Delta_x \Delta_y + \Delta_y \Delta_z + \Delta_z \Delta_x + 2(\Delta_x + \Delta_y + \Delta_z)]\alpha = 0.$$

⁵See [35] for a definition of spacelike and timelike global symmetries and their applications to the ground-state degeneracy and restricted mobility constraints.

⁶Note that N in \mathbb{Z}_N is different from N , the number of vertices of Γ .

TABLE I. The comparison of four exotic lattice models that can be defined on a general graph Γ . The model is robust if it has no relevant local operator. See Sec. III B 4 for more details on what we mean by robustness.

Model	Spatial lattice	GSD	Defects	Robust
Laplacian ϕ theory [22,23]	Γ	$ \text{Jac}(\Gamma) $	None	No
$U(1)$ Laplacian gauge theory [22,23]	Γ	1	Fracton	Yes
\mathbb{Z}_N Laplacian model Appendix D	Γ	$ \text{Jac}(\Gamma, N) $	Fracton	No
Anisotropic \mathbb{Z}_N Laplacian model Secs. III and IV	$\Gamma \times C_{L_c}$	$ \text{Jac}(\Gamma, N) ^2$	Lineon	Yes

any two vertices and no self-loop on any vertex, *undirected* means the edges do not have any orientation, and *connected* means there is a path between any two vertices of the graph. We use i to denote a vertex (or site), and $\langle i, j \rangle$ to denote an edge (or link) of the graph. We write $\langle i, j \rangle \in \Gamma$ if there is an edge between vertices i and j in Γ .

Let d_i be the *degree* of vertex i , i.e., the number of edges incident on i . The *Laplacian matrix* L of Γ is an $N \times N$ symmetric matrix defined as follows: $L_{ii} = d_i$ for every vertex i , $L_{ij} = -1$ if there is an edge $\langle i, j \rangle$ between vertices i and j , and $L_{ij} = 0$ otherwise.

A. Discrete Laplacian operator Δ_L and its Smith decomposition

Consider a \mathbb{Z}_N -valued function $f(i)$ on the vertices of the graph. We define the *discrete Laplacian operator* Δ_L as

$$\begin{aligned} \Delta_L f(i) &:= \sum_j L_{ij} f(j) = d_i f(i) - \sum_{j:\langle i,j \rangle \in \Gamma} f(j) \\ &= \sum_{j:\langle i,j \rangle \in \Gamma} [f(i) - f(j)], \end{aligned} \tag{2.1}$$

where the equalities are modulo N . This is one of the most natural and universal difference operators that can be defined on any such graph Γ .

We are interested in the following two questions:

(1) What are all the \mathbb{Z}_N -valued functions $h(i)$ that satisfy the *discrete Laplacian equation*

$$\Delta_L h(i) = 0 \pmod N? \tag{2.2}$$

They are known as the \mathbb{Z}_N -valued *discrete harmonic functions*, and we denote the set of such functions as $\mathcal{H}(\Gamma, \mathbb{Z}_N)$.⁷

(2) We define an equivalence class of \mathbb{Z}_N -valued functions by saying that two functions $g(i)$ and $\tilde{g}(i)$ belong to the same class if there is a \mathbb{Z}_N -valued function $f(i)$ such that

$$\tilde{g}(i) - g(i) = \Delta_L f(i) \pmod N. \tag{2.3}$$

In this case, we write $\tilde{g}(i) \sim g(i)$. What are all the distinct equivalence classes under the equivalence relation “ \sim ”?

Interestingly, both questions can be answered using the *Smith decomposition* [42] of the Laplacian matrix L [43]. The *Smith normal form* of L is given by three matrices R, P , and Q , such that

$$R = PLQ, \quad \text{or} \quad R_{ab} = \sum_{i,j} P_{ai} L_{ij} Q_{jb}, \tag{2.4}$$

where $P, Q \in GL_N(\mathbb{Z})$, and $R = \text{diag}(r_1, \dots, r_N)$. Here r_a 's are nonnegative integers, known as the *invariant factors* of L , such that r_a divides r_{a+1} for $a = 1, \dots, N - 1$. While R is uniquely determined by L , the matrices P and Q are not. For a connected graph Γ , we have $r_a > 0$ for $a = 1, \dots, N - 1$, and $r_N = 0$.

We state the answers to the two questions here (see Appendix A for details):

(1) Any \mathbb{Z}_N -valued discrete harmonic function takes the form

$$h(i) = \sum_{a=1}^N \frac{N Q_{ia} p_a}{\text{gcd}(N, r_a)} \pmod N, \tag{2.5}$$

where $p_a = 0, \dots, \text{gcd}(N, r_a) - 1$ for $a = 1, \dots, N$. In other words, $\mathcal{H}(\Gamma, \mathbb{Z}_N)$ is isomorphic to the finite Abelian group $\prod_{a=1}^N \mathbb{Z}_{\text{gcd}(N, r_a)}$. Here the group operation is simply the sum. It is well defined because if $h_1, h_2 \in \mathcal{H}(\Gamma, \mathbb{Z}_N)$, then $h_1 + h_2 \in \mathcal{H}(\Gamma, \mathbb{Z}_N)$.

(2) Any equivalence class is uniquely represented by the \mathbb{Z}_N -valued function

$$g(i) = \sum_{a=1}^N p_a (Q^{-1})_{ai} \pmod N, \tag{2.6}$$

where $p_a = 0, \dots, \text{gcd}(N, r_a) - 1$ for $a = 1, \dots, N$. In other words, the set of all equivalence classes is isomorphic to the finite Abelian group $\prod_{a=1}^N \mathbb{Z}_{\text{gcd}(N, r_a)}$. Here the group operation is simply the sum which is well defined because if $\tilde{g}_1 \sim g_1$ and $\tilde{g}_2 \sim g_2$, then $\tilde{g}_1 + \tilde{g}_2 \sim g_1 + g_2$.

It follows that the number of \mathbb{Z}_N -valued discrete harmonic functions and the number of equivalence classes are both $\prod_{a=1}^N \text{gcd}(N, r_a)$, which is the order of the finite Abelian group $\prod_{a=1}^N \mathbb{Z}_{\text{gcd}(N, r_a)}$. We will have more to say about this group below.

B. Jacobian group of a graph

The finite Abelian group encountered above is intimately related to the *Jacobian group* $\text{Jac}(\Gamma)$, which is a natural finite Abelian group associated with a general graph Γ .⁸ In terms of the invariant factors of the Laplacian matrix L , we have the

⁸It has several different names in the graph theory literature, including the *sandpile group* [44], the *group of components* [45], or the *critical group* [46] of Γ , and it is related to the *group of bicycles* [41] of Γ .

⁷It is also known as the *group of balanced vertex weightings* [41].

following isomorphism

$$\text{Jac}(\Gamma) \cong \prod_{a=1}^{N-1} \mathbb{Z}_{r_a}. \quad (2.7)$$

The order of $\text{Jac}(\Gamma)$ is the most fundamental and well-studied notion of *complexity* in graph theory. What we have here is a “mod N -reduction” of the Jacobian group:

$$\text{Jac}(\Gamma, N) \cong \prod_{a=1}^N \mathbb{Z}_{\text{gcd}(N, r_a)}. \quad (2.8)$$

As we will see below, the group $\text{Jac}(\Gamma, N)$ plays an crucial role in the \mathbb{Z}_N Laplacian models.

III. ANISOTROPIC \mathbb{Z}_N LAPLACIAN MODEL ON A GRAPH

In this section, we study a robust gapped lineon model on a spatial lattice of the form $\Gamma \times C_{L_z}$, where Γ is a simple, connected, undirected graph, and L_z is the number of sites in the z direction (see Fig. 1). We refer to it as the anisotropic \mathbb{Z}_N Laplacian model because it is the anisotropic extension along the z direction of the \mathbb{Z}_N Laplacian model analyzed in Appendix D.

A. Hamiltonian for the stabilizer code

In the Hamiltonian formulation of the anisotropic \mathbb{Z}_N Laplacian model, there are a \mathbb{Z}_N variable $U(i, z)$ and its conjugate variable $V(i, z)$, i.e., $U(i, z)V(i, z) = e^{2\pi i/N} V(i, z)U(i, z)$, on every site of $\Gamma \times C_{L_z}$. There are also a \mathbb{Z}_N variable $U_z(i, z + \frac{1}{2})$ and its conjugate variable $V_z(i, z + \frac{1}{2})$, i.e., $U_z(i, z + \frac{1}{2})V_z(i, z + \frac{1}{2}) = e^{2\pi i/N} V_z(i, z + \frac{1}{2})U_z(i, z + \frac{1}{2})$, on every z -link of $\Gamma \times C_{L_z}$.

The Hamiltonian is

$$H = -\gamma_1 \sum_{i,z} G(i, z) - \gamma_2 \sum_{i,z} F\left(i, z + \frac{1}{2}\right) + \text{H.c.}, \quad (3.1)$$

where

$$\begin{aligned} G(i, z) &= V_z\left(i, z + \frac{1}{2}\right)^\dagger V_z\left(i, z - \frac{1}{2}\right) \\ &\quad \times \prod_{j:(i,j) \in \Gamma} V(i, z)V(j, z)^\dagger, \\ F\left(i, z + \frac{1}{2}\right) &= U(i, z + 1)^\dagger U(i, z) \\ &\quad \times \prod_{j:(i,j) \in \Gamma} U_z\left(i, z + \frac{1}{2}\right)U_z\left(j, z + \frac{1}{2}\right)^\dagger. \end{aligned} \quad (3.2)$$

The two kinds of terms are shown in Fig. 3. Since all the terms in this Hamiltonian commute with each other, it is a stabilizer code.

The model enjoys the duality transformation

$$\begin{aligned} U(i, z) &\rightarrow V_z\left(i, z + \frac{1}{2}\right), & U_z\left(i, z + \frac{1}{2}\right) &\rightarrow V(i, z), \\ V(i, z) &\rightarrow U_z\left(i, z + \frac{1}{2}\right)^\dagger, & V_z\left(i, z + \frac{1}{2}\right) &\rightarrow U(i, z)^\dagger. \end{aligned} \quad (3.3)$$




FIG. 3. The two kinds of stabilizer terms in the Hamiltonian (3.1).

It exchanges the two kind of terms in the Hamiltonian and therefore it maps the model with (γ_1, γ_2) to the model with (γ_2, γ_1) . As a result, for $\gamma_1 = \gamma_2$, this model is self-dual.

The ground states satisfy $G(i, z) = 1$ and $F(i, z + \frac{1}{2}) = 1$ for all i, z . The excited states are violations of $G = 1$ or $F = 1$, which we call electric and magnetic excitations respectively. These excitations are mobile along the z direction so they are at least z -lineons. Their mobility constraints along the graph Γ are more complicated. We postpone that discussion to Sec. III B 3.

We could also take $\gamma_1, \gamma_2 \rightarrow \infty$, in which case, the Hilbert space consists of only the ground states, and the Hamiltonian is trivial. The Euclidean presentation of this model in this limit will be discussed later in Sec. III B.

We are particularly interested in those operators that commute with the Hamiltonian (3.1) and act nontrivially on its ground states. They are the global symmetry operators of the model in the low-energy limit, and they are also known as the logical operators of the stabilizer code. We choose a basis of these symmetry operators as follows: the electric symmetry operators are⁹

$$\begin{aligned} \tilde{W}_z(a) &= \prod_{i,z} V(i, z)^{(Q^{-1})_{ai}}, & a = 1, \dots, N, \\ \tilde{W}\left(a; z + \frac{1}{2}\right) &= \prod_i V_z\left(i, z + \frac{1}{2}\right)^{\frac{N}{\text{gcd}(N, r_a)} Q_{ia}}, & (3.4) \\ &z = 0, \dots, L_z - 1, \end{aligned}$$

and the magnetic symmetry operators are

$$\begin{aligned} W_z(a) &= \prod_{i,z} U_z\left(i, z + \frac{1}{2}\right)^{(Q^{-1})_{ai}}, & a = 1, \dots, N, \\ W(a; z) &= \prod_i U(i, z)^{\frac{N}{\text{gcd}(N, r_a)} Q_{ia}}, & z = 0, \dots, L_z - 1, \end{aligned} \quad (3.5)$$

⁹In fact, the operator $\prod_z V(i, z)$, which is local in Γ and extends in the z direction, also commutes with the Hamiltonian. Here we choose to work with the basis of $\tilde{W}_z(a)$ because the latter has a simpler commutation relation with $W(a; z)$. Similarly, $\prod_z U_z(i, z + \frac{1}{2})$ also commutes with the Hamiltonian and is local in Γ , but we choose to work in the basis of $W_z(a)$ for the same reason.

where Q and r_a are defined in (2.4). These operators generate a $\text{Jac}(\Gamma, N)^2$ electric symmetry and a $\text{Jac}(\Gamma, N)^2$ magnetic symmetry.

For each a , the four operators in (3.4) and (3.5) are all $\mathbb{Z}_{\text{gcd}(N, r_a)}$ operators. Clearly, $W(a; z)^{\text{gcd}(N, r_a)} = \tilde{W}(a; z + \frac{1}{2})^{\text{gcd}(N, r_a)} = 1$. [In fact, when acting on the ground states, the operators $\tilde{W}(a; z + \frac{1}{2})$ and $W(a; z)$ are independent of z .] Moreover, the operator $W_z(a)$ satisfies $W_z(a)^{r_a} = 1$ when acting on the ground states,¹⁰ which when combined with the obvious relation $W_z(a)^N = 1$ gives the relation $W_z(a)^{\text{gcd}(N, r_a)} = 1$. The same conclusion holds for $\tilde{W}_z(a)$ as well.

The basis of symmetry operators defined in (3.4) and (3.5) is chosen such that they satisfy the following commutation relations:

$$W(a; z)\tilde{W}_z(b) = \exp\left[\frac{2\pi i\delta_{ab}}{\text{gcd}(N, r_a)}\right]\tilde{W}_z(b)W(a; z),$$

$$a, b = 1, \dots, N, \quad (3.6)$$

and similarly for the other pair. So for each $a = 1, \dots, N$, there are two independent copies of $\mathbb{Z}_{\text{gcd}(N, r_a)}$ Heisenberg algebras, leading to a ground-state degeneracy of¹¹

$$\text{GSD} = \prod_{a=1}^N \text{gcd}(N, r_a)^2 = |\text{Jac}(\Gamma, N)|^2. \quad (3.7)$$

B. Euclidean presentation

We now discuss the Euclidean presentation of the anisotropic \mathbb{Z}_N Laplacian model. We place the theory on a Euclidean spacetime lattice $C_{L_\tau} \times \Gamma \times C_{L_z}$, where $\Gamma \times C_{L_z}$ is the spatial slice. We use (τ, i, z) to label a site in the spacetime lattice, where i denotes a vertex of the graph Γ .

We use the integer BF formulation of [27]. The integer BF action of the anisotropic \mathbb{Z}_N Laplacian model is

$$S = \frac{2\pi i}{N} \sum_{\tau, i, z} \left(-\tilde{m}_\tau \left(\tau, i, z + \frac{1}{2} \right) \left[\Delta_z m \left(\tau, i, z + \frac{1}{2} \right) - \Delta_L m_z \left(\tau, i, z + \frac{1}{2} \right) \right] + \tilde{m}_z \left(\tau + \frac{1}{2}, i, z \right) \times \left[\Delta_\tau m \left(\tau + \frac{1}{2}, i, z \right) - \Delta_L m_\tau \left(\tau + \frac{1}{2}, i, z \right) \right] + \tilde{m} \left(\tau + \frac{1}{2}, i, z + \frac{1}{2} \right) \left[\Delta_\tau m_z \left(\tau + \frac{1}{2}, i, z + \frac{1}{2} \right) - \Delta_z m_\tau \left(\tau + \frac{1}{2}, i, z + \frac{1}{2} \right) \right] \right), \quad (3.8)$$

¹⁰This is because $W_z(a)^{r_a} = \prod_{i,z} U_z(i, z + \frac{1}{2})^{r_a(Q^{-1})_{ai}} = \prod_{i,j,z} U_z(i, z + \frac{1}{2})^{P_{aj}L_{ji}} = \prod_j i \prod_z U(j, z + 1)U(j, z)^{\dagger P_{aj}} = 1$, where we used the facts that $RQ^{-1} = PL$, and $F = 1$ on the ground states.

¹¹The power of 2 in (3.7) is related to the fact that the anisotropic \mathbb{Z}_N Laplacian model is the anisotropic extension of the \mathbb{Z}_N Laplacian model of Appendix D, whose GSD is $|\text{Jac}(\Gamma, N)|$ (D14). This is similar to the relation $\text{GSD}_{3+1\text{D}} \text{anisotropic } \mathbb{Z}_N \text{ lineon model} = (\text{GSD}_{2+1\text{D}} \mathbb{Z}_N \text{ Ising plaquette model})^2$.

where the integer fields (m_τ, m, m_z) have a gauge symmetry

$$m_\tau \sim m_\tau + \Delta_\tau k + Nq_\tau,$$

$$m \sim m + \Delta_L k + Nq,$$

$$m_z \sim m_z + \Delta_z k + Nq_z, \quad (3.9)$$

where k and (q_τ, q, q_z) are integers, and similarly for $(\tilde{m}_\tau, \tilde{m}, \tilde{m}_z)$. [Note that, when working modulo N , the second line of (3.9) is exactly the equivalence relation discussed in (2.3).]

The theory is self-dual under the map $(m_\tau, m, m_z) \rightarrow (\tilde{m}_\tau, \tilde{m}, \tilde{m}_z)$ and $(\tilde{m}_\tau, \tilde{m}, \tilde{m}_z) \rightarrow -(m_\tau, m, m_z)$.

The integer BF action (3.8) describes the ground states of a stabilizer code given by the Hamiltonian (3.1). Here we will not elaborate on the relation between the Euclidean and Hamiltonian presentations. We refer the readers to Appendix C.2 of [27] for an analogous discussion of the relation between the 2+1D \mathbb{Z}_N toric code and the 2+1D \mathbb{Z}_N gauge theory in the integer BF presentation.

1. Ground-state degeneracy

We can count the number of ground states by counting the number of solutions to the “equations of motion” of $(\tilde{m}_\tau, \tilde{m}, \tilde{m}_z)$ modulo gauge transformations:

$$\Delta_z m - \Delta_L m_z = 0 \pmod{N},$$

$$\Delta_\tau m - \Delta_L m_\tau = 0 \pmod{N}, \quad (3.10)$$

$$\Delta_\tau m_z - \Delta_z m_\tau = 0 \pmod{N}.$$

A gauge field (m_τ, m, m_z) that satisfies (3.10) is a flat \mathbb{Z}_N gauge field. We can use the gauge freedom in k to set $m_\tau(\tau + \frac{1}{2}, i, z)|_{\tau \neq 0} = 0 \pmod{N}$, and $m_z(\tau, i, z + \frac{1}{2})|_{z \neq 0} = 0 \pmod{N}$. In this gauge choice, the last line of (3.10) implies that

$$\Delta_\tau m_z(\tau + \frac{1}{2}, i, z + \frac{1}{2})|_{z=0} = 0 \pmod{N},$$

$$\Delta_z m_\tau(\tau + \frac{1}{2}, i, z + \frac{1}{2})|_{\tau=0} = 0 \pmod{N}. \quad (3.11)$$

The first two lines of (3.10) then imply that

$$\Delta_\tau m(\tau + \frac{1}{2}, i, z) = 0 \pmod{N},$$

$$\Delta_z m(\tau, i, z + \frac{1}{2}) = 0 \pmod{N}, \quad (3.12)$$

which in turn imply that

$$\Delta_L m_z(i, z + \frac{1}{2})|_{z=0} = 0 \pmod{N},$$

$$\Delta_L m_\tau(\tau + \frac{1}{2}, i)|_{\tau=0} = 0 \pmod{N}. \quad (3.13)$$

The remaining τ and z -independent gauge freedom, $m(i) \sim m(i) + \Delta_L k(i)$, is exactly the equivalence relation in (2.3). So, after gauge fixing, we can set $m(i)$ to be of the form (2.6), i.e., there are $|\text{Jac}(\Gamma, N)|$ independent holonomies in $m(i)$. Since $m_z(i, z + \frac{1}{2})|_{z=0}$ and $m_\tau(\tau + \frac{1}{2}, i)|_{\tau=0}$ satisfy (3.13), which is exactly the discrete Laplace equation (2.2), they are of the form (2.5). So there are $|\text{Jac}(\Gamma, N)|$ independent holonomies in both of them. Finally, the set of gauge transformations $k(\tau, i, z)$ that do not act on (m_τ, m, m_z) satisfy $\Delta_\tau k = \Delta_z k = \Delta_L k = 0 \pmod{N}$. In other words, $k(\tau, i, z) = k(i)$ is independent of τ, z , and $k(i)$ satisfies the discrete Laplace equation (2.2). So such gauge transformations are of

the form (2.5), and there are $|\text{Jac}(\Gamma, N)|$ of them. Therefore, the ground-state degeneracy is

$$\text{GSD} = \frac{|\text{Jac}(\Gamma, N)|^3}{|\text{Jac}(\Gamma, N)|} = |\text{Jac}(\Gamma, N)|^2 = \prod_{a=1}^N \text{gcd}(N, r_a)^2. \quad (3.14)$$

2. Global symmetry

There is an electric symmetry associated with the shift of (m_τ, m, m_z) by a flat \mathbb{Z}_N gauge field. By the analysis following (3.10), up to gauge transformations, the electric (spacelike) symmetry acts as¹²

$$\begin{aligned} m(\tau, i, z) &\rightarrow m(\tau, i, z) + \lambda(i), \\ \lambda(i) &= \sum_{a=1}^N p_a (Q^{-1})_{ai}, \\ m_z\left(\tau, i, z + \frac{1}{2}\right) &\rightarrow m_z\left(\tau, i, z + \frac{1}{2}\right) + \delta_{z,0} \lambda_z(i), \\ \lambda_z(i) &= \sum_{a=1}^N \frac{N Q_{ia} p_{z,a}}{\text{gcd}(N, r_a)}, \end{aligned} \quad (3.15)$$

where p_a and $p_{z,a}$ are both integers modulo $\text{gcd}(N, r_a)$ for $a = 1, \dots, N$. There is also a magnetic (spacelike) symmetry which acts on \tilde{m} and \tilde{m}_z in a similar way.

The electric (spacelike) symmetry is generated by the Wilson operators of $(\tilde{m}_\tau, \tilde{m}, \tilde{m}_z)$:

$$\begin{aligned} \tilde{W}_z(a) &= \exp \left[\frac{2\pi i}{N} \sum_{i,z} (Q^{-1})_{ai} \tilde{m}_z\left(\tau + \frac{1}{2}, i, z\right) \right], \\ \tilde{W}\left(a; z + \frac{1}{2}\right) &= \exp \left[\frac{2\pi i}{\text{gcd}(N, r_a)} \right. \\ &\quad \left. \times \sum_i \tilde{m}\left(\tau + \frac{1}{2}, i, z + \frac{1}{2}\right) Q_{ia} \right], \end{aligned} \quad (3.16)$$

for $a = 1, \dots, N$ and $z = 0, \dots, L_z - 1$. The electrically charged operators are the Wilson operators of (m_τ, m, m_z) , i.e., $W(a; z)$ and $W_z(a)$. Similarly, the magnetic (spacelike) symmetry is generated by $W(a; z)$ and $W_z(a)$, while the magnetically charged operators are $\tilde{W}_z(a)$ and $\tilde{W}(a; z + \frac{1}{2})$. These are the operators in (3.4) and (3.5) in the low-energy limit. The commutation relation (3.6) can now be understood as a mixed 't Hooft anomaly between electric and magnetic spacelike symmetries.

3. Timelike symmetry and lineons

The integer BF action has defects, which extend in the time direction, such as

$$W_\tau(i, z) = \exp \left[\frac{2\pi i}{N} \sum_\tau m_\tau\left(\tau + \frac{1}{2}, i, z\right) \right]. \quad (3.17)$$

¹²These are symmetries of the action (3.8) because $\Delta_\tau m$, $\Delta_z m$, and $\Delta_\tau m_z$ are clearly unaffected by the shifts, whereas $\Delta_L m_z$ is shifted by $\delta_{z,0} \Delta_L \lambda_z(i) = 0 \pmod N$ because $\lambda_z(i)$ is a \mathbb{Z}_N -valued discrete harmonic function (2.5).

This describes the world-line of an infinitely heavy particle of unit charge at position (i, z) . It also represents the low-energy limit of an electric excitation at position (i, z) in the stabilizer code (3.1). We can deform the defect to

$$\begin{aligned} &\exp \left[\frac{2\pi i}{N} \sum_{\tau < 0} m_\tau\left(\tau + \frac{1}{2}, i, z\right) \right] \\ &\times \exp \left[\frac{2\pi i}{N} \sum_{z \leq z'' < z'} m_z\left(0, i, z'' + \frac{1}{2}\right) \right] \\ &\times \exp \left[\frac{2\pi i}{N} \sum_{\tau \geq 0} m_\tau\left(\tau + \frac{1}{2}, i, z'\right) \right]. \end{aligned} \quad (3.18)$$

This configuration describes a particle moving along the z direction.

Next, we discuss the mobility of the particle along the graph Γ . Such a motion is constrained by the timelike global symmetry, which acts on extended defects rather than the operators or states of the Hilbert space (see [35] for more discussions on timelike global symmetries). Up to gauge transformation, the electric timelike symmetry acts as¹³

$$\begin{aligned} m_\tau\left(\tau + \frac{1}{2}, i, z\right) &\rightarrow m_\tau\left(\tau + \frac{1}{2}, i, z\right) + \delta_{\tau,0} \lambda_\tau(i), \\ \lambda_\tau(i) &= \sum_{a=1}^N \frac{N Q_{ia} p_{\tau,a}}{\text{gcd}(N, r_a)}, \end{aligned} \quad (3.19)$$

where $p_{\tau,a} = 0, \dots, \text{gcd}(N, r_a) - 1$. Hence, the group of electric timelike symmetry is $\text{Jac}(\Gamma, N)$.

Two defects at sites (i, z) and (i', z') carry the same timelike charges, or equivalently, a particle can hop from (i, z) to (i', z') , if and only if¹⁴

$$Q_{ia} = Q_{i'a} \pmod{\text{gcd}(N, r_a)}, \quad \forall a = 1, \dots, N. \quad (3.21)$$

In other words, the timelike charges Q_{ia} encode the superselection sector of a defect.

¹³This is a symmetry of the action (3.8) because $\Delta_z m_\tau$ is clearly unaffected by the shift, and $\Delta_L m_\tau$ is shifted by $\delta_{\tau,0} \Delta_L \lambda_\tau(i) = 0 \pmod N$ because $\lambda_\tau(i)$ is a \mathbb{Z}_N -valued discrete harmonic function (2.5).

¹⁴Indeed, when this condition holds, the defect that ‘‘moves’’ a particle from (i, z) to (i', z') at time $\tau = 0$ is given by

$$\begin{aligned} &\exp \left[\frac{2\pi i}{N} \sum_{\tau < 0} m_\tau\left(\tau + \frac{1}{2}, i, z\right) \right] \\ &\times \exp \left[-\frac{2\pi i}{N} \sum_{a,j} \left(\frac{Q_{ia} - Q_{i'a}}{\text{gcd}(N, r_a)} \right) \tilde{r}_a P_{aj} m(0, j, z) \right] \\ &\times \exp \left[\frac{2\pi i}{N} \sum_{z \leq z'' < z'} m_z\left(0, i', z'' + \frac{1}{2}\right) \right] \\ &\times \exp \left[\frac{2\pi i}{N} \sum_{\tau \geq 0} m_\tau\left(\tau + \frac{1}{2}, i', z'\right) \right], \end{aligned} \quad (3.20)$$

where for each a , \tilde{r}_a is the integer solution of the equation $\tilde{r}_a r_a = \text{gcd}(N, r_a) \pmod N$.

Similarly, there are defects of \tilde{m}_τ which represent the low-energy limit of magnetic excitations of the stabilizer code (3.1). By the self-duality, similar mobility restrictions apply to the defects of \tilde{m}_τ due to a $\text{Jac}(\Gamma, N)$ dual magnetic timelike symmetry.

While this selection rule (3.21) is not very intuitive, we will give strong mobility constraints in the special case where the spatial lattice is a cubic lattice (i.e., Γ is a 2D torus graph $C_{L_x} \times C_{L_y}$) in Sec. IV. In particular, under some mild conditions, the particles can move only along the z direction, i.e., they are lineons.

4. Robustness

Let us examine the robustness of the low-energy theory. Typically, in order to address this question we should map local operators in the UV theory to local operators in the IR theory. However, if Γ is a general graph, it has no notion of locality and we cannot discuss local operators. Therefore, the usual discussion of robustness does not apply. Instead, we will restrict Γ to be a regular lattice (such as square lattice, honeycomb lattice, cubic lattice, etc.), where there is an unambiguous notion of locality and we can consider localized operators. (One might be able to extend the discussion to the case of an infinite graph Γ with some restrictions on its connectivity. We will not attempt to do it here.)

The only operators that act nontrivially on the ground states are $\tilde{W}(a; z)$ and $\tilde{W}_z(a)$ of (3.16), and similarly, $W(a; z)$ and $W_z(a)$. It is clear that $W_z(a)$ and $\tilde{W}_z(a)$ are supported over L_z sites in the z direction, so they are not finitely supported in the infinite volume limit. Now, we show that $W(a; z)$ is not finitely supported when Γ is a regular lattice. Assume to the contrary that it is finitely supported. It generates a $\text{Jac}(\Gamma, N)$ magnetic symmetry that shifts the gauge field $\tilde{m}(\tau + \frac{1}{2}, i, z')$ by $\delta_{z', z} f(i)$, where $f(i)$ is a \mathbb{Z}_N -valued discrete harmonic function. The support of $f(i)$ is precisely the support of the operator $W(a; z)$, so $f(i)$ is also finitely supported. However, on a regular lattice, there is no nontrivial finitely supported discrete harmonic function.¹⁵ Therefore, $W(a; z)$ cannot be finitely supported. Similarly, $\tilde{W}(a; z + \frac{1}{2})$ is also not finitely supported.

Since there are no finitely supported operators that act nontrivially in the space of ground states, the anisotropic \mathbb{Z}_N Laplacian model is robust. We can deform the microscopic model with finitely supported operators. As long as their coefficients are small enough, they map to localized deformations of the low-energy theory. However, since there are no local pointlike operators acting in the low-energy theory, it cannot change.

¹⁵For example, on a square lattice with coordinates (x, y) , let $f(x, y)$ be a finitely supported discrete harmonic function. Consider a large rectangular region R that contains the support of $f(x, y)$, i.e., $f(x, y) = 0$ for (x, y) outside R . Using the discrete Laplace equation $\Delta_L f(x, y) = 0$ at the points immediately outside R , one can show that $f(x, y) = 0$ at the points immediately inside R . By induction, $f(x, y) = 0$ everywhere inside R . This argument extends to any regular lattice. It also extends to a more general class of graphs, but we do not discuss this here.

IV. 3+1D ANISOTROPIC \mathbb{Z}_N LAPLACIAN MODEL ON A TORUS

In this section, we analyze the GSD and restricted mobility of the anisotropic \mathbb{Z}_N Laplacian model on an $L_x \times L_y \times L_z$ cubic lattice with periodic boundary condition, i.e., Γ is a 2D torus graph $C_{L_x} \times C_{L_y}$. On $\Gamma = C_{L_x} \times C_{L_y}$, we have the following identification between the lattice points:

$$(x, y) \sim (x + L_x, y) \sim (x, y + L_y). \quad (4.1)$$

Throughout this section, we use (x, y) to denote a vertex of the 2D torus graph, and reserve i to denote a vertex of a general graph Γ . Then the discrete Laplacian operator Δ_L takes the more familiar form $\Delta_x^2 + \Delta_y^2$ in the xy plane.

A. Upper bound on \log_N GSD

It is clear from (3.7) that the GSD depends only on properties of Γ and is independent of L_z . Here we give an upper bound on how fast \log_N GSD can grow with L_x, L_y .

Recall that the GSD of the anisotropic \mathbb{Z}_N Laplacian model is $|\text{Jac}(\Gamma, N)|^2$ (3.7). As we showed in Sec. II A, $|\text{Jac}(\Gamma, N)|$ is also the number of equivalence classes under the equivalence relation “ \sim ” in (2.3). Combining these facts, we have

$$\text{GSD} = \left| \frac{\{g(i)\}}{g(i) \sim g(i) + \Delta_L f(i)} \right|^2, \quad (4.2)$$

where $g(i)$ and $f(i)$ are \mathbb{Z}_N -valued functions on the graph Γ .

When Γ is the 2D torus graph $C_{L_x} \times C_{L_y}$, interpreting the equivalence relation as a gauge symmetry, we can gauge fix $g(x, y) = 0 \pmod N$ everywhere except at $x = 0, 1$, or at $y = 0, 1$. In other words, the number of sites where $g(x, y) \neq 0$ after gauge fixing is at most $2 \min(L_x, L_y)$. Since $g(x, y)$ is \mathbb{Z}_N -valued, it follows that the number of nontrivial configurations of $g(x, y)$ is at most $N^{2 \min(L_x, L_y)}$. Therefore,

$$\log_N \text{GSD} \leq 4 \min(L_x, L_y). \quad (4.3)$$

B. $N = 2$

When $N = 2$, the stabilizer terms (3.2) simplify to

$$\begin{aligned} G(x, y, z) &= V_z \left(x, y, z + \frac{1}{2} \right) V_z \left(x, y, z - \frac{1}{2} \right) \\ &\quad \times \prod_{\epsilon_x, \epsilon_y = \pm 1} V(x + \epsilon_x, y + \epsilon_y, z), \\ F \left(x, y, z + \frac{1}{2} \right) &= U(x, y, z + 1) U(x, y, z) \\ &\quad \times \prod_{\epsilon_x, \epsilon_y = \pm 1} U_z \left(x + \epsilon_x, y + \epsilon_y, z + \frac{1}{2} \right). \end{aligned} \quad (4.4)$$

Here the product of the four V 's in G and the product of the four U_z 's in F both involve only the four sites around the “ $\frac{\pi}{4}$ -tilted plaquette” centered at (x, y) . This is illustrated in Fig. 4. Therefore, each stabilizer term of the 3+1D anisotropic \mathbb{Z}_2 Laplacian model is equivalent to a stabilizer term of the 3+1D anisotropic \mathbb{Z}_2 lineon model of [13, 26] on a “ $\frac{\pi}{4}$ -tilted” lattice.




FIG. 4. The stabilizer terms (4.4) of the 3+1D anisotropic \mathbb{Z}_2 Laplacian model are equivalent to those of the 3+1D anisotropic \mathbb{Z}_2 lineon model of [13,26] on a “ $\frac{\pi}{4}$ -tilted” lattice (the z direction is suppressed in this figure). In particular, the product of V ’s in G and the product of U_z ’s in F involve only the four sites around the “ $\frac{\pi}{4}$ -tilted plaquette” centered at (x, y) . Such a plaquette consists of only red sites or only blue sites but not both. One such red tilted plaquette is shown. The red and blue sublattices might or might not give independent copies of the 3+1D anisotropic \mathbb{Z}_2 lineon model depending on the parities of L_x and L_y in the identifications (4.1). We use the coordinates (x', y') for the tilted lattice, which are related to the original coordinates (x, y) as $x' = \frac{x+y}{2}$ and $y' = \frac{x-y}{2}$. They are integers on the blue sublattice, and half-integers on the red sublattice.

Let us define the coordinates $(x', y') = (\frac{x+y}{2}, \frac{y-x}{2})$ for the tilted lattice. The tilted lattice decomposes into two sublattices: those with integral (x', y') and those with half-integral (x', y') . These are shown in blue and red in Fig. 4. Observe that any tilted plaquette consists of sites from only one of the sublattices. Therefore, locally, there are two copies of the 3+1D anisotropic \mathbb{Z}_2 lineon model, one on each sublattice. On an infinite lattice, these two copies are independent. However, the identifications (4.1) can couple them: when L_x and L_y are both even, the two sublattices are decoupled and there are two copies of the 3+1D anisotropic \mathbb{Z}_2 lineon model, whereas when L_x or L_y is odd, the two sublattices are identified, so there is only one copy of the 3+1D anisotropic \mathbb{Z}_2 lineon model. In all these cases, the identifications on the tilted lattice for the 3+1D anisotropic \mathbb{Z}_2 lineon model are given in (B6), (B9), and (B13).

We present the GSD and mobility restrictions in this model for $L_x = L_y = L$ and refer the readers to Appendix B on results for arbitrary L_x and L_y . The ground-state degeneracy is given by

$$\text{GSD} = \begin{cases} 2^{4L}, & L \text{ even,} \\ 2^{4L-2}, & L \text{ odd.} \end{cases} \quad (4.5)$$

This is in agreement with the plot for $N = 2$ in Fig. 2, and it saturates the bound in (4.3) when L is even. Furthermore, a z -lineon cannot hop between different sites in the xy plane. In contrast, a dipole of z -lineons separated in the $(1, \pm 1)$ direction can move in the $(1, \mp 1)$ direction in the xy plane. These mobility restrictions follow from the relation between the 3+1D anisotropic \mathbb{Z}_2 Laplacian model and the 3+1D anisotropic \mathbb{Z}_2 lineon model on the tilted lattice.

To conclude, the $N = 2$ anisotropic Laplacian model is made out of the known anisotropic lineon model of [13,26], with a relatively simple GSD (4.5). The next subsection discusses the anisotropic \mathbb{Z}_p Laplacian model with p an odd prime, which is a genuinely new model and has a much more intricate GSD.

C. $N = p$ prime larger than 2

When $N = p$ is a prime larger than 2 we can follow [47] and use techniques from commutative algebra to compute the ground-state degeneracy. We show that the GSD is given by (1.2). See Appendix C 1 for the meaning and derivation of this expression.

The expression in (1.2) can be simplified in some special cases. Let $q \neq p$ be another odd prime such that p is a primitive root modulo q^m , where $m \geq 1$, i.e., p is the generator of the multiplicative group of integers modulo q^m , denoted as $\mathbb{Z}_{q^m}^\times$.¹⁶ Then for $L_x = p^{k_x} q^m$ and $L_y = p^{k_y} q^m$, where $k_x, k_y, m \geq 0$, we show that

$$\log_p \text{GSD} = 2[2p^{\min(k_x, k_y)} - \delta_{k_x, k_y}]. \quad (4.6)$$

We see that the bound (4.3) is saturated whenever $k_x \neq k_y$ and $m = 0$, i.e., there are infinitely many L_x, L_y for which $\log_p \text{GSD}$ scales as $O(L_x, L_y)$. On the other hand, when $k_x = k_y = 0$, we have $\log_p \text{GSD} = 2$ for any m , i.e., there are also infinitely many L_x, L_y for which $\log_p \text{GSD}$ remains finite.

The last statement relies on the existence of an odd prime q such that p is a primitive root modulo q^m for all $m \geq 1$. A sufficient condition for this is that p is a primitive root modulo q^2 [[48], Sec. 2.8]. For example, 3 is a primitive root modulo 5^2 , so for $p = 3$, we can choose $q = 5$. Similarly, for $p = 5, 7$, we can choose $q = 7, 11$ respectively. In fact, one can verify numerically that for all $p \lesssim 10^9$, there is such a q . However, there is no proof of existence of such q for arbitrary p .

Interestingly, Artin’s conjecture on primitive roots [49] states that there are infinitely many prime q such that p is a primitive root modulo q .¹⁷ Whenever $L_x = L_y = q$ for any such q , we find that $\log_p \text{GSD} = 2$. This gives another infinite family of L_x, L_y for which $\log_p \text{GSD}$ remains finite. However,

¹⁶For any positive integer n , the set of all integers a such that $1 \leq a < n$ and $\text{gcd}(a, n) = 1$ form a group under multiplication, known as the multiplicative group of integers modulo n , and denoted as \mathbb{Z}_n^\times . It is cyclic exactly when $n = 1, 2, 4, q^m$, or $2q^m$, where q is an odd prime and $m \geq 1$ [[48], Sec. 2.8]. Whenever \mathbb{Z}_n^\times is cyclic, it has a single generator, and the notion of “primitive root modulo n ” is well defined.

¹⁷Note that p being a primitive root modulo q does not imply that p is a primitive root modulo q^2 .

Artin's conjecture is still unproven, except under the assumption of the *generalized Riemann hypothesis* [50], which is also unproven.

We can apply similar techniques to determine the mobility of z -lineons in the xy plane as well. (See Appendix C 2 for more details.) There exist certain special values of L_x, L_y (e.g., $L_x = L_y = q^m$, where q is an odd prime such that p is a primitive root modulo q^m) for which the z -lineons are completely mobile in the xy plane. However, on an infinite square lattice, any finite set of z -lineons is completely immobile (unless they can be annihilated), assuming that their charges and the separations between them are fixed during the motion, i.e., they cannot move “rigidly.”

It is surprising that the set of L_x, L_y for which $\log_p \text{GSD}$ remains finite and the z -lineons are completely mobile is intimately related to well-known open problems in number theory.

Note added. Recently [51] appeared on arXiv, which studies the same anisotropic \mathbb{Z}_N Laplacian model using its stabilizer code.

ACKNOWLEDGMENTS

We are grateful to H. Ebisu, E. Studnia, and S. Velusamy for helpful discussions. P.G. was supported by the Physics Department of Princeton University. H.T.L. was supported in part by a Croucher fellowship from the Croucher Foundation, the Packard Foundation, and the Center for Theoretical Physics at MIT. The work of N.S. was supported in part by DOE Grant No. DE-SC0009988 and by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651440). The work of S.H.S. was supported in part by NSF Grant No. PHY-2210182. The authors of this paper are ordered alphabetically.

APPENDIX A: \mathbb{Z}_N -VALUED FUNCTIONS ON A GRAPH

In this Appendix, we analyze the space of \mathbb{Z}_N -valued harmonic functions and the equivalence classes of \mathbb{Z}_N -valued functions on a general graph Γ . We use the Smith decomposition (2.4) of the Laplacian matrix L to give complete answers of these two questions mentioned in Sec. II A.

Recall that the Smith normal form of L is given by $R = PLQ$, where $P, Q \in GL_N(\mathbb{Z})$, and $R = \text{diag}(r_1, \dots, r_N)$. Here r_a 's are nonnegative integers such that r_a divides r_{a+1} for $a = 1, \dots, N-1$. While R is uniquely determined by L , the matrices P and Q are not.

In the index notation of (2.4), we have $R_{ab} = r_a \delta_{ab} = \sum_{i,j} P_{ai} L_{ij} Q_{jb}$. While all the indices here run from 1 to N , only i, j have a natural interpretation as vertices of the graph Γ .

We can now answer the first question raised in Sec. II A: find all the \mathbb{Z}_N -valued discrete harmonic functions. We first transform the \mathbb{Z}_N -valued function $h(i)$ to a new basis:

$$h'_a = \sum_i (Q^{-1})_{ai} h(i) \pmod{N}, \quad (\text{A1})$$

In this basis, the discrete Laplace equation (2.2) is “diagonal”:

$$r_a h'_a = 0 \pmod{N}, \quad a = 1, \dots, N. \quad (\text{A2})$$

We can solve this equation independently for each a . The most general solution is

$$h'_a = \frac{N p_a}{\text{gcd}(N, r_a)} \pmod{N}, \quad a = 1, \dots, N, \quad (\text{A3})$$

where $p_a = 0, \dots, \text{gcd}(N, r_a) - 1$. Transforming back to the original basis, the most general \mathbb{Z}_N -valued discrete harmonic function is

$$h(i) = \sum_{a=1}^N \frac{N Q_{ia} p_a}{\text{gcd}(N, r_a)} \pmod{N}. \quad (\text{A4})$$

Let us now address the second question raised in Sec. II A: find all the equivalence classes under the equivalence relation “ \sim .” Since the Laplacian matrix L is symmetric, taking the transpose of $R = PLQ$ gives another Smith decomposition $R = Q^T L P^T$. Using this, we transform the \mathbb{Z}_N -valued function $g(i)$ to a (different) new basis

$$g''_a = \sum_i g(i) Q_{ia} \pmod{N}. \quad (\text{A5})$$

We define \tilde{g}''_a similarly for another function $\tilde{g}(i)$ in the same equivalence class. In this basis, the equivalence relation (2.3) is “diagonal”:

$$\tilde{g}''_a - g''_a = r_a \hat{f}_a \pmod{N}, \quad (\text{A6})$$

where $\hat{f}_a = \sum_i f(i) P_{ia}^{-1} \pmod{N}$. Therefore, the equivalence class of $g(i)$ is completely determined by N congruence classes:

$$g''_a \pmod{\text{gcd}(N, r_a)}, \quad a = 1, \dots, N. \quad (\text{A7})$$

Going back to the original basis, a representative of the equivalence class “ $p_a \pmod{\text{gcd}(N, r_a)}$ ” is

$$g(i) = \sum_{a=1}^N p_a (Q^{-1})_{ai} \pmod{N}, \quad (\text{A8})$$

where $p_a = 0, \dots, \text{gcd}(N, r_a) - 1$ for $a = 1, \dots, N$.

APPENDIX B: 3+1D ANISOTROPIC \mathbb{Z}_2 LAPLACIAN MODEL

In this Appendix, we use the relation between the 3+1D anisotropic \mathbb{Z}_2 Laplacian model of Sec. IV B and the 3+1D anisotropic \mathbb{Z}_2 lineon model to compute the GSD and restricted mobility of the former.

1. Ground-state degeneracy

Recall that the stabilizer terms of the 3+1D anisotropic \mathbb{Z}_2 Laplacian model are given by (4.4), which are equivalent to those of the 3+1D anisotropic \mathbb{Z}_2 lineon model of [13,26] on a tilted lattice. Moreover, the latter is an anisotropic extension of the 2+1D \mathbb{Z}_2 Ising plaquette model [25] on the tilted lattice.

Now, the identifications on the original lattice are (4.1)

$$(x, y) \sim (x + L_x, y) \sim (x, y + L_y). \quad (\text{B1})$$

In the new coordinates $(x', y') = (\frac{x+y}{2}, \frac{y-x}{2})$, the identifications on the tilted lattice take the schematic form

$$(x', y') \sim (x' + L_x^u, y' + L_y^u) \sim (x' + L_x^v, y' + L_y^v). \quad (\text{B2})$$




FIG. 5. The minimal identifications on the blue sublattice of Fig. 4 for different parities of L_x and L_y . The black circles represent the identifications (4.1), and the shaded regions represent the fundamental domains of the blue sublattice under these identifications.

The authors of [32] analyzed the 2+1D \mathbb{Z}_2 Ising plaquette model on a 2D spatial torus with such identifications. Their strategy was to reduce the identifications to the form

$$(x', y') \sim (x' + ML_{x'}^{\text{eff}}, y') \sim (x' + KL_{x'}^{\text{eff}}, y' + L_{y'}^{\text{eff}}), \quad (\text{B3})$$

where $\text{gcd}(M, K) = 1$. Then they showed that

$$\text{GSD}_{2+1\text{D } \mathbb{Z}_2 \text{ Ising plaq}} = \text{gcd}(2, M)2^{L_{x'}^{\text{eff}} + L_{y'}^{\text{eff}} - 1}. \quad (\text{B4})$$

It follows that (see footnote 11)

$$\text{GSD}_{3+1\text{D aniso } \mathbb{Z}_2 \text{ lineon}} = [\text{gcd}(2, M)2^{L_{x'}^{\text{eff}} + L_{y'}^{\text{eff}} - 1}]^2. \quad (\text{B5})$$

Let us use these results to compute the GSD of the 3+1D anisotropic \mathbb{Z}_2 Laplacian model:

(1) L_x, L_y even: In this case, the identifications in (4.1) do not couple the two sublattices. So there are two independent copies of the 3+1D anisotropic \mathbb{Z}_2 lineon model, and we can work with one copy at a time, say, the blue sublattice in Fig. 4. In the new coordinates (x', y') , the minimal identifications on the blue sublattice are [see Fig. 5(a)]

$$(x', y') \sim \left(x' + \frac{L_x}{2}, y' - \frac{L_x}{2}\right) \sim \left(x' + \frac{L_y}{2}, y' + \frac{L_y}{2}\right). \quad (\text{B6})$$

Let $(\tilde{L}_x, \tilde{L}_y)$ be the integer solution of the equation $\tilde{L}_x L_x + \tilde{L}_y L_y = \text{gcd}(L_x, L_y)$. Then, in the notation of (B3), we have

$$\begin{aligned} ML_{x'}^{\text{eff}} &= \text{lcm}(L_x, L_y), \\ KL_{x'}^{\text{eff}} &= \frac{1}{2}(\tilde{L}_x L_x - \tilde{L}_y L_y), \\ L_{y'}^{\text{eff}} &= \frac{1}{2} \text{gcd}(L_x, L_y), \end{aligned} \quad (\text{B7})$$

and hence,

$$\text{GSD} = [\text{gcd}(2, M)2^{L_{x'}^{\text{eff}} + L_{y'}^{\text{eff}} - 1}]^4. \quad (\text{B8})$$

Here the power is 4 rather than 2 because there are two copies of the 3+1D anisotropic \mathbb{Z}_2 lineon model, one on each sublattice.

(2) L_x odd, L_y even: In this case, the identifications in (4.1) couple the two sublattices so that effectively there is only one sublattice, say the blue sublattice in Figure 4. Hence, there is only one copy of the 3+1D anisotropic \mathbb{Z}_2 lineon model. The minimal identifications on the blue sublattice are [see Fig. 5(b)]

$$(x, y) \sim (x + 2L_x, y) \sim (x, y + L_y), \quad (\text{B9})$$

which can be written as

$$(x', y') \sim (x' + L_x, y' - L_x) \sim \left(x' + \frac{L_x}{2}, y' + \frac{L_y}{2}\right), \quad (\text{B10})$$

in the new coordinates (x', y') . Let $(\tilde{L}_x, \tilde{L}_y)$ be the integer solution of the equation $\tilde{L}_x L_x + \tilde{L}_y (\frac{L_y}{2}) = \text{gcd}(L_x, \frac{L_y}{2})$. Then, in the notation of (B3), we have

$$\begin{aligned} ML_{x'}^{\text{eff}} &= 2 \text{lcm}\left(L_x, \frac{L_y}{2}\right), \\ KL_{x'}^{\text{eff}} &= \tilde{L}_x L_x - \frac{1}{2} \tilde{L}_y L_y, \\ L_{y'}^{\text{eff}} &= \text{gcd}\left(L_x, \frac{L_y}{2}\right), \end{aligned} \quad (\text{B11})$$

and hence,

$$\text{GSD} = [\text{gcd}(2, M)2^{L_{x'}^{\text{eff}} + L_{y'}^{\text{eff}} - 1}]^2. \quad (\text{B12})$$

(3) L_x, L_y odd: In this case, once again, the identifications in (4.1) couple the two sublattices so that effectively there is only one sublattice, say the blue sublattice in Fig. 4. Hence, there is only one copy of the 3+1D anisotropic \mathbb{Z}_2 lineon model. The minimal identifications on the blue sublattice are [see Fig. 5(c)]

$$(x, y) \sim (x + 2L_x, y) \sim (x + L_x, y + L_y), \quad (\text{B13})$$

which can be written as

$$(x', y') \sim (x' + L_x, y' - L_x) \sim \left(x' + \frac{L_x + L_y}{2}, y' - \frac{L_x - L_y}{2}\right), \quad (\text{B14})$$

in the new coordinates (x', y') . Let $(\tilde{L}_x, \tilde{L}_y)$ be the integer solution of the equation $\tilde{L}_x L_x + \tilde{L}_y (\frac{L_x - L_y}{2}) = \text{gcd}(L_x, \frac{L_x - L_y}{2})$. Then, in the notation of (B3), we have

$$\begin{aligned} ML_{x'}^{\text{eff}} &= \frac{L_x L_y}{\text{gcd}\left(L_x, \frac{L_x - L_y}{2}\right)}, \\ KL_{x'}^{\text{eff}} &= \tilde{L}_x L_x + \tilde{L}_y \left(\frac{L_x + L_y}{2}\right), \\ L_{y'}^{\text{eff}} &= \text{gcd}\left(L_x, \frac{L_x - L_y}{2}\right), \end{aligned} \quad (\text{B15})$$

and hence,

$$\text{GSD} = [\text{gcd}(2, M)2^{L_{x'}^{\text{eff}}+L_{y'}^{\text{eff}}-1}]^2. \quad (\text{B16})$$

When $L_x = L_y = L$, we have

$$L_{x'}^{\text{eff}} = L_{y'}^{\text{eff}} = \frac{L}{M}, \quad K = 1, \quad M = \begin{cases} 2, & L \text{ even,} \\ 1, & L \text{ odd.} \end{cases} \quad (\text{B17})$$

Then the above expressions for the ground-state degeneracy simplify to

$$\text{GSD} = \begin{cases} 2^{4L}, & L \text{ even,} \\ 2^{4L-2}, & L \text{ odd.} \end{cases} \quad (\text{B18})$$

2. Mobility restrictions

Let us now discuss the mobility of the z -lineons in the xy plane in the 3+1D anisotropic \mathbb{Z}_2 Laplacian model. First, note that in the 2+1D \mathbb{Z}_2 Ising plaquette model on a 2D spatial torus given by the identifications (B3), the defect describing a single particle of unit charge can “hop” from (x', y') to $(x' + \text{gcd}(2, M)L_{x'}^{\text{eff}}, y')$ [35]. This motion is nontrivial if and only if $\text{gcd}(2, M) = 1$ and $M > 1$. It follows that the z -lineons of the 3+1D anisotropic \mathbb{Z}_2 lineon model, in addition to moving along the z direction, can “hop” in the xy plane in the same way.

In particular, consider $L_x = L_y = L$. It follows from (B17) that the two conditions $\text{gcd}(2, M) = 1$ and $M > 1$ cannot be simultaneously satisfied for any L . Therefore, the z -lineons cannot hop in the xy plane, but can only move in the z direction (and hence the name lineon).

A dipole of z -lineons at (x, y) and $(x + s, y \pm s)$, where $s \in \mathbb{Z}$, can move in the $(1, \mp 1)$ direction. This follows from the motion of a dipole of fractons in the 2+1D \mathbb{Z}_2 Ising plaquette model in the direction orthogonal to their separation.

APPENDIX C: 3+1D ANISOTROPIC \mathbb{Z}_p LAPLACIAN MODEL

In this Appendix, we use techniques from commutative algebra to compute the ground-state degeneracy and analyze the restricted mobility of the 3+1D anisotropic \mathbb{Z}_p Laplacian model of Sec. IV C when $p > 2$ is prime. Such techniques were used to analyze translationally invariant Pauli stabilizer codes [47]. All the mathematical facts used here can be found in standard textbooks on the subject, such as [52,53].

1. Ground-state degeneracy

Recall the relation (4.2) between the GSD of the anisotropic \mathbb{Z}_N Laplacian model and the number of equivalence classes of \mathbb{Z}_N -valued functions under the equivalence relation “ \sim ” of (2.3). Here we set $N = p > 2$, where p is prime,¹⁸ and $\Gamma = C_{L_x} \times C_{L_y}$, the 2D torus graph. We first find an exact expression for the \log_p GSD in terms of commutative-algebraic quantities using the relation (4.2), then explain how to compute this expression in general using a *Gröbner basis*, and finally compute it explicitly for some

special values of L_x, L_y . In particular, we show that there are infinitely many L_x, L_y for which \log_p GSD is $O(L_x, L_y)$, and also infinitely many L_x, L_y for which \log_p GSD is finite.

a. Exact expression for \log_p GSD

Since there are $L_x L_y$ points in the $\Gamma = C_{L_x} \times C_{L_y}$ torus and since we are interested in \mathbb{Z}_p -valued functions on that space, it is clear that there are $p^{L_x L_y}$ such functions. As in (2.3), they fall into equivalence classes $g(x, y) \sim \tilde{g}(x, y)$ when $g(x, y) - \tilde{g}(x, y) = \Delta_L f(x, y) \pmod p$. We would like to find the number of such equivalence classes.

As a first step, we give a more abstract description of these $p^{L_x L_y}$ functions. Let $\mathcal{R} = \mathbb{Z}_p[X, Y]$ be the ring of polynomials with coefficients in \mathbb{Z}_p , and $\mathfrak{j} = (\mathbf{Q}_x, \mathbf{Q}_y)$ be the *ideal* of \mathcal{R} generated by the polynomials $\mathbf{Q}_x(X, Y) = X^{L_x} - 1$ and $\mathbf{Q}_y(X, Y) = Y^{L_y} - 1$. Given two polynomials $F, G \in \mathcal{R}$, we write¹⁹

$$F(X, Y) = G(X, Y) \pmod{\mathfrak{j}}, \quad (\text{C1})$$

if and only if $F(X, Y) - G(X, Y)$ is a polynomial in \mathfrak{j} . The set of equivalence classes modulo \mathfrak{j} is the quotient ring \mathcal{R}/\mathfrak{j} .

Any equivalence class of \mathcal{R}/\mathfrak{j} is represented by a unique polynomial that is a \mathbb{Z}_p -linear combination of the monomials $X^a Y^b$ with $0 \leq a < L_x$ and $0 \leq b < L_y$. (Here we used the equivalence relations to remove higher powers of X or Y . This is a special case of a more general procedure, called complete reduction, which we will describe below.) Therefore, the number of equivalence classes is $p^{L_x L_y}$. In fact, since \mathbb{Z}_p is a field, \mathcal{R}/\mathfrak{j} can be thought of as a vector space over \mathbb{Z}_p . The above monomials form a basis of this vector space, so

$$\dim_{\mathbb{Z}_p} \mathcal{R}/\mathfrak{j} = L_x L_y. \quad (\text{C2})$$

Here “ $\dim_{\mathbb{Z}_p}$ ” denotes the dimension of a vector space over \mathbb{Z}_p .

It is convenient to represent a \mathbb{Z}_p -valued function $f(x, y)$ on the 2D torus graph $\Gamma = C_{L_x} \times C_{L_y}$ as a polynomial representing an equivalence class of the quotient ring \mathcal{R}/\mathfrak{j} as follows:

$$\hat{f}(X, Y) = \sum_{x=0}^{L_x-1} \sum_{y=0}^{L_y-1} f(x, y) X^{L_x-x-1} Y^{L_y-y-1} \pmod{\mathfrak{j}}. \quad (\text{C3})$$

$\hat{f}(X, Y)$ can be thought of as a lattice Fourier transform of $f(x, y)$ with $X = e^{ik_x}$ and $Y = e^{ik_y}$, which depend on the momenta k_x and k_y .

Observe that, for any integer $0 \leq k < L_x$, we have

$$\begin{aligned} X^k \hat{f}(X, Y) &= \left[\sum_{x=0}^{L_x-k-1} \sum_{y=0}^{L_y-1} f(x+k, y) X^{L_x-x-1} Y^{L_y-y-1} \right. \\ &\quad \left. + \sum_{x=L_x-k}^{L_x-1} \sum_{y=0}^{L_y-1} f(x-L_x+k, y) X^{L_x-x-1} Y^{L_y-y-1} \right] \pmod{\mathfrak{j}}, \end{aligned} \quad (\text{C4})$$

¹⁸Actually, all of the following discussion up to (C36) works even for $p = 2$. The discussion after that does not work for $p = 2$ for reasons we will explain later.

¹⁹We do not write the “mod p ” explicitly because we are working in $\mathbb{Z}_p[X, Y]$.

so X can be interpreted as the generator of translations in the x direction.²⁰ The fact that translating in x by L_x takes the graph C_{L_x} back to itself is related to the trivial equation

$$X^{L_x} \hat{f}(X, Y) = \hat{f}(X, Y) \pmod{j}. \quad (\text{C5})$$

Also, the difference operator Δ_x is associated with the polynomial $X - 1$. For convenience, we define the *displaced* discrete Laplacian operator, denoted by $\tilde{\Delta}_L$, as

$$\tilde{\Delta}_L f(x, y) = (\Delta_x^2 + \Delta_y^2) f(x + 1, y + 1). \quad (\text{C6})$$

[Here we extended $f(x, y)$ to a periodic function on \mathbb{Z}^2 .] It is associated with the polynomial

$$\tilde{P}(X, Y) = Y(X - 1)^2 + X(Y - 1)^2. \quad (\text{C7})$$

In general, any *local* difference operator in the xy plane, after an appropriate displacement, is associated with a polynomial $S \in \mathcal{R}$ satisfying $S(1, 1) = 0$.

Let $\mathfrak{i} = (\tilde{P})$ be the ideal of \mathcal{R} generated by the polynomial $\tilde{P}(X, Y)$. Then $\mathfrak{i}/(\mathfrak{i} \cap \mathfrak{j}) \cong (\mathfrak{i} + \mathfrak{j})/\mathfrak{j}$ is an ideal of the quotient ring \mathcal{R}/\mathfrak{j} .²¹ In fact, it is the subspace of the vector space \mathcal{R}/\mathfrak{j} that corresponds to the \mathbb{Z}_p -valued functions of the form $\tilde{\Delta}_L f(x, y)$.

It is then clear that the quotient

$$\frac{\mathcal{R}/\mathfrak{j}}{(\mathfrak{i} + \mathfrak{j})/\mathfrak{j}} \cong \mathcal{R}/(\mathfrak{i} + \mathfrak{j}) \quad (\text{C8})$$

corresponds to the set of equivalence classes of \mathbb{Z}_p -valued functions under the equivalence relation “ \sim ” of (2.3) on the 2D torus graph $C_{L_x} \times C_{L_y}$. It follows that

$$\log_p \text{GSD} = 2 \dim_{\mathbb{Z}_p} \mathcal{R}/(\mathfrak{i} + \mathfrak{j}), \quad (\text{C9})$$

or more explicitly,

$$\log_p \text{GSD} = 2 \dim_{\mathbb{Z}_p} \frac{\mathbb{Z}_p[X, Y]}{(Y(X - 1)^2 + X(Y - 1)^2, X^{L_x} - 1, Y^{L_y} - 1)}. \quad (\text{C10})$$

Below we give a general procedure to compute this quantity.

b. Computing \log_p GSD using Gröbner basis

One way to compute the (vector space) dimension of $\mathcal{R}/(\mathfrak{i} + \mathfrak{j})$ is by first computing the *Gröbner basis* of the ideal $\mathfrak{i} + \mathfrak{j}$. Before defining a Gröbner basis, we need an ordering on all the monomials. For our purposes, it is sufficient to define a *lexicographic monomial ordering*: $X^m Y^n > X^k Y^l$ if and only if $m > k$, or $m = k$ and $n > l$. Then, for any polynomial

$F(X, Y)$, we define its *leading term* as the term that has the largest monomial among all the terms.

Now, we say a polynomial $F(X, Y)$ is *reducible* with respect to a set of polynomials $\mathcal{G} = \{G_1, \dots, G_n\}$ if some term of $F(X, Y)$ is a multiple of the leading term of one of the $G_i(X, Y)$'s. We say it is *irreducible* otherwise.

Given an ideal \mathfrak{J} of \mathcal{R} and a polynomial $F \in \mathcal{R}$, one can ask what the equivalence class of $F(X, Y)$ in \mathcal{R}/\mathfrak{J} is. One way to answer this question is to *reduce* $F(X, Y)$ with respect to a generating set \mathcal{B} of \mathfrak{J} repeatedly until we are left with a polynomial $H(X, Y)$ that is irreducible with respect to \mathcal{B} . [Such a procedure is known as a *complete reduction* of $F(X, Y)$ with respect to \mathcal{B} .] One might hope that $H(X, Y)$ uniquely specifies the equivalence class of $F(X, Y)$. However, for a generic \mathcal{B} , the $H(X, Y)$ so obtained depends on the choices made in the repeated reductions, and hence, may not be unique.

A Gröbner basis $\mathcal{G} = \{G_1, \dots, G_n\}$ is a special generating set of \mathfrak{J} such that $F(X, Y)$ can be written as

$$F(X, Y) = H(X, Y) + \sum_{i=1}^n H_i(X, Y) G_i(X, Y), \quad (\text{C11})$$

where $H(X, Y)$ is *uniquely* determined by the requirement that it is irreducible with respect to \mathcal{G} .²² In this case, we write

$$F(X, Y) = H(X, Y) \pmod{\mathcal{G}}. \quad (\text{C12})$$

It follows that there is a one-one correspondence between \mathcal{R}/\mathfrak{J} and the set of all polynomials that are irreducible with respect to \mathcal{G} . Indeed, the set of all monomials that are irreducible with respect to \mathcal{G} forms a basis of the vector space \mathcal{R}/\mathfrak{J} . From this we conclude that $\dim_{\mathbb{Z}_p} \mathcal{R}/\mathfrak{J}$ equals the number of monomials that are irreducible with respect to \mathcal{G} .

While there is an algorithm, known as *Buchberger's algorithm*, to compute a Gröbner basis of an ideal given its generators, it is not always easy to compute it analytically.²³ Nonetheless, for fixed values of p, L_x, L_y , the Gröbner basis, and therefore the GSD, can be readily computed with the help of computer programs. Let us do an explicit calculation for the GSD when $p = 3$ and $L_x = L_y = 4$ as an example. Using the `GroebnerBasis` command with `Modulus` $\rightarrow 3$ in *Mathematica*, we can compute the Gröbner basis for the ideal $(\tilde{P}, Q_x, Q_y) = (Y(X - 1)^2 + X(Y - 1)^2, X^4 - 1, Y^4 - 1)$ in this case. We find $\mathcal{G} = \{Y^4 - 1, XY + X - Y^3 - Y^2, X^2 + Y^3 - Y^2 + Y + 1\}$ under the lexicographic ordering $X > Y$. The leading terms of \mathcal{G} are $\{Y^4, XY, X^2\}$, and the five irreducible monomials with respect to \mathcal{G} are $1, Y, Y^2, Y^3, X$. We conclude that $\dim_{\mathbb{Z}_3} \mathcal{R}/(\mathfrak{i} + \mathfrak{j}) = 5$, and hence $\log_3 \text{GSD} = 10$. In fact, the plot of $\log_N \text{GSD}$ as a function of $L_x = L_y = L$ in Fig. 2 was obtained exactly in this way.

²⁰Our functions $f(x, y)$ are defined on $L_x L_y$ points. They can be extended to periodic functions on \mathbb{Z}^2 . Then it is straightforward to apply k translations. The expression (C4) corresponds to applying such k translation and then expressing the result in terms of the function $f(x, y)$ in the fundamental domain $0 \leq x \leq L_x - 1, 0 \leq y \leq L_y - 1$. Equivalently, we can use the periodicity of f to write (C4) as $X^k \hat{f}(X, Y) = \sum_{x=0}^{L_x-1} \sum_{y=0}^{L_y-1} f(x+k, y) X^{L_x-x-1} Y^{L_y-y-1} \pmod{j}$.

²¹Here $\mathfrak{i} + \mathfrak{j} = (\tilde{P}, Q_x, Q_y)$ is the ideal of \mathcal{R} generated by the three polynomials $\tilde{P}(X, Y)$, $Q_x(X, Y)$, and $Q_y(X, Y)$. It is known as the *sum of ideals* \mathfrak{i} and \mathfrak{j} .

²²Note that $H_i(X, Y)$'s are not uniquely determined by this procedure. Indeed, shifting $H_1(X, Y)$ by $G_2(X, Y)$ and $H_2(X, Y)$ by $-G_1(X, Y)$ gives another complete reduction of $F(X, Y)$ with respect to \mathcal{G} .

²³For the ideal $\mathfrak{j} = (Q_x, Q_y)$, the generating set $\{Q_x, Q_y\}$ is already a Gröbner basis, and moreover, the irreducible monomials are $X^x Y^y$ for $0 \leq x < L_x$ and $0 \leq y < L_y$, which form a basis of \mathcal{R}/\mathfrak{j} . [This fact was used in the analysis leading to (C2).] However, for the ideal $\mathfrak{i} + \mathfrak{j} = (\tilde{P}, Q_x, Q_y)$, the generating set $\{\tilde{P}, Q_x, Q_y\}$ is not always a Gröbner basis.

Fortunately, for the problem at hand, it is possible to simplify the expression (C9) for the GSD further analytically. To prepare for the simplified expression, we first define a few things:

A field F is *algebraically closed* if any polynomial in $F[X]$ can be factorized completely into linear factors in $F[X]$. For example, \mathbb{R} is not algebraically closed because the polynomial $X^2 + 1$, which is in $\mathbb{R}[X]$, cannot be factorized into linear factors in $\mathbb{R}[X]$. In contrast, it is well known that \mathbb{C} is algebraically closed. Every field, by definition, contains the multiplicative identity 1.

The *characteristic* of a field is the smallest positive integer n such that $1 + \dots + 1$ (n times) = 0. It is clear that \mathbb{Z}_p is a field of characteristic p . By convention, the characteristics of \mathbb{Q} , \mathbb{R} , and \mathbb{C} are all defined to be 0. It is known that the characteristic of any field is either 0 or a prime number. As we see, there are several fields with the same characteristic p .

For *finite fields*, the field is uniquely specified by the number of its elements and the characteristic. In particular, for any integer $k \geq 1$ and prime p , \mathbb{F}_{p^k} denotes the unique (up to field isomorphisms) finite field of order p^k and characteristic p . Furthermore, \mathbb{F}_{p^∞} denotes the unique algebraically closed field of characteristic p .²⁴

Let $\mathcal{S} = \mathbb{F}_{p^\infty}[X, Y]$ be the ring of polynomials in X, Y with coefficients in \mathbb{F}_{p^∞} . We use the same symbols \mathfrak{i} and \mathfrak{j} for the ideals of \mathcal{S} generated by the polynomials $\tilde{P}(X, Y)$ and $\tilde{Q}_x(X, Y)$, $\tilde{Q}_y(X, Y)$ respectively. The *algebraic set* of $\mathfrak{i} + \mathfrak{j}$, denoted as $V(\mathfrak{i} + \mathfrak{j})$, is the set of distinct solutions $(X_0, Y_0) \in \mathbb{F}_{p^\infty}^2$ of the system of polynomial equations

$$X^{L_x} - 1 = Y^{L_y} - 1 = \tilde{P}(X, Y) = 0. \tag{C13}$$

We will find it convenient to parametrize L_x and L_y in terms of the order of our group \mathbb{Z}_p as

$$L_x = p^{k_x} L'_x, \quad L_y = p^{k_y} L'_y, \quad \gcd(p, L'_x) = \gcd(p, L'_y) = 1. \tag{C14}$$

Now, for each solution $(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})$, we define the polynomials

$$\tilde{Q}_{x, X_0}(X, Y) = (X - X_0)^{p^{k_x}}, \quad \tilde{Q}_{y, Y_0}(X, Y) = (Y - Y_0)^{p^{k_y}}, \tag{C15}$$

and the ideal $\mathfrak{i}'_{X_0, Y_0} = (\tilde{P}, \tilde{Q}_{x, X_0}, \tilde{Q}_{y, Y_0})$ of \mathcal{S} .

With these preparations, the simplified expression for the GSD (C9) is

$$\log_p \text{GSD} = 2 \sum_{(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})} \dim_{\mathbb{F}_{p^\infty}} \mathcal{S}/\mathfrak{i}'_{X_0, Y_0}. \tag{C16}$$

It is obtained using techniques from commutative algebra (see e.g., [52,53]), that were used in [47]. For readers who are familiar with such techniques, a derivation of (C16) is given below. Others, who are willing to accept it, can skip directly to Appendix C 1 d, where we compute the GSD for some special values of L_x, L_y explicitly.

²⁴More explicitly, $\mathbb{F}_{p^\infty} = \bigcup_{n=1}^\infty \mathbb{F}_{p^n}$. Here we used the fact that \mathbb{F}_{p^k} is a subfield of \mathbb{F}_{p^m} if and only if k divides m .

c. Derivation of (C16)

Since $\mathcal{R}/(\mathfrak{i} + \mathfrak{j})$ is a finite-dimensional vector space over \mathbb{Z}_p , it is an *Artinian ring*,²⁵ and hence, it has finitely many *maximal ideals*.²⁶ Moreover, for an Artinian ring, it is known that

$$\mathcal{R}/(\mathfrak{i} + \mathfrak{j}) \cong \bigoplus_{\mathfrak{m}} [\mathcal{R}/(\mathfrak{i} + \mathfrak{j})]_{\mathfrak{m}}, \tag{C17}$$

where the sum is over all maximal ideals of $\mathcal{R}/(\mathfrak{i} + \mathfrak{j})$, and $[\mathcal{R}/(\mathfrak{i} + \mathfrak{j})]_{\mathfrak{m}}$ denotes the *localization* of $\mathcal{R}/(\mathfrak{i} + \mathfrak{j})$ at \mathfrak{m} .²⁷ It follows that

$$\log_p \text{GSD} = 2 \sum_{\mathfrak{m}} \dim_{\mathbb{Z}_p} [\mathcal{R}/(\mathfrak{i} + \mathfrak{j})]_{\mathfrak{m}}. \tag{C18}$$

So we can compute dimension for each term in the sum and then add them up. However, the maximal ideals of $\mathcal{R}/(\mathfrak{i} + \mathfrak{j})$ are a bit complicated to work with. Instead, we proceed as follows.

We can replace \mathbb{Z}_p by \mathbb{F}_{p^∞} in (C9) and get the same answer for $\log_p \text{GSD}$. More concretely, we have²⁸

$$\dim_{\mathbb{Z}_p} \mathcal{R}/(\mathfrak{i} + \mathfrak{j}) = \dim_{\mathbb{F}_{p^\infty}} \mathcal{S}/(\mathfrak{i} + \mathfrak{j}), \tag{C19}$$

where, on the right-hand side, \mathfrak{i} and \mathfrak{j} are ideals of \mathcal{S} generated by the same polynomials as before. Since $\mathcal{S}/(\mathfrak{i} + \mathfrak{j})$ is also Artinian, we have

$$\log_p \text{GSD} = 2 \sum_{\mathfrak{m}} \dim_{\mathbb{F}_{p^\infty}} [\mathcal{S}/(\mathfrak{i} + \mathfrak{j})]_{\mathfrak{m}}, \tag{C20}$$

where the sum is over all maximal ideals of $\mathcal{S}/(\mathfrak{i} + \mathfrak{j})$. We now characterize the maximal ideals of $\mathcal{S}/(\mathfrak{i} + \mathfrak{j})$.

²⁵A ring \mathcal{R} is Artinian if it satisfies the *descending chain condition*, i.e., if $\mathfrak{I}_1 \supseteq \mathfrak{I}_2 \supseteq \mathfrak{I}_3 \supseteq \dots$ is a descending chain of ideals, then there is a $k \geq 1$ such that $\mathfrak{I}_k = \mathfrak{I}_{k+1} = \mathfrak{I}_{k+2} = \dots$. For example, the ring of integers \mathbb{Z} is not Artinian because $(2) \supset (4) \supset (8) \supset \dots$, where (n) denotes the ideal generated by the integer n . On the other hand, for any integer n , the ring of integers modulo n , $\mathbb{Z}/(n) = \mathbb{Z}_n$, is Artinian. Moreover, any ring that is also a finite dimensional vector space over a field is always Artinian, which is exactly the case here.

²⁶A proper ideal is an ideal that is not the ring itself. For example, in \mathbb{Z} , the ideal (4) is proper because it does not contain 1. (The only ideal containing 1 is the entire ring itself.) A maximal ideal is a proper ideal that is not contained in any other proper ideal except itself. For example, in \mathbb{Z} , (4) is not a maximal ideal because it is contained in the proper ideal (2). The latter is maximal; in fact, the ideal (n) is maximal if and only if n is prime.

²⁷Intuitively, given a multiplicatively closed subset S of a ring \mathcal{R} , the localization of \mathcal{R} with respect to S , denoted as $S^{-1}\mathcal{R}$, means “formally adding multiplicative inverses” for all the elements of S . For example, in \mathbb{Z} , the subset of nonzero integers is multiplicatively closed, and localizing with respect to this set gives the rationals \mathbb{Q} . In any ring \mathcal{R} , given a maximal ideal \mathfrak{m} , the set $\mathcal{R} \setminus \mathfrak{m}$ is always multiplicatively closed. So we define the localization of \mathcal{R} at \mathfrak{m} , denoted by $\mathcal{R}_{\mathfrak{m}}$, as the localization of \mathcal{R} with respect to $\mathcal{R} \setminus \mathfrak{m}$.

²⁸Given a field F , one can talk about extending it to a larger field F' such that F is a subfield of F' . For example, \mathbb{R} is a subfield of \mathbb{C} , or equivalently, \mathbb{C} is an extension of \mathbb{R} . Now, let V be a vector space over F . One can “extend the base field” from F to F' by tensoring V with F' , denoted as $V \otimes_F F'$. Then $\dim_{F'} V = \dim_F(V \otimes_F F')$. Since \mathbb{F}_{p^∞} is an extension of \mathbb{Z}_p , the result in (C19) follows.

By the *correspondence theorem for quotient rings*, the maximal ideals of $\mathcal{S}/(\mathfrak{i} + \mathfrak{j})$ are in one-one correspondence with the maximal ideals of \mathcal{S} that contain the ideal $\mathfrak{i} + \mathfrak{j}$. Moreover, since \mathbb{F}_{p^∞} is algebraically closed, by *Hilbert's Nullstellensatz*, the maximal ideals of \mathcal{S} are in one-one correspondence with ideals of the form $(X - X_0, Y - Y_0)$, where $(X_0, Y_0) \in \mathbb{F}_{p^\infty}^2$. Now, the ideal $(X - X_0, Y - Y_0)$ contains $\mathfrak{i} + \mathfrak{j}$ if and only if (X_0, Y_0) is a root of all the polynomials in $\mathfrak{i} + \mathfrak{j}$. It follows that the maximal ideals of $\mathcal{S}/(\mathfrak{i} + \mathfrak{j})$ are in one-one correspondence with ideals of \mathcal{S} of the form $(X - X_0, Y - Y_0)$, where (X_0, Y_0) is a solution of the system of polynomial equations

$$X^{L_x} - 1 = Y^{L_y} - 1 = \tilde{P}(X, Y) = 0, \quad (\text{C21})$$

i.e., $(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})$, the algebraic set of $\mathfrak{i} + \mathfrak{j}$. Then

$$\log_p \text{GSD} = 2 \sum_{(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})} \dim_{\mathbb{F}_{p^\infty}} [\mathcal{S}/(\mathfrak{i} + \mathfrak{j})]_{(X - X_0, Y - Y_0)}. \quad (\text{C22})$$

We can simplify it further. Let $L_i = p^{k_i} L'_i$ with $\text{gcd}(p, L'_i) = 1$. Recall that since \mathbb{F}_{p^∞} is algebraically, any polynomial in $\mathbb{F}_{p^\infty}[X]$ can be factorized completely into linear factors in $\mathbb{F}_{p^\infty}[X]$. Using this fact, we have

$$X^{L_x} - 1 = (X^{L'_x} - 1)^{p^{k_x}} = \prod_{\xi \in \mathbb{F}_{p^\infty} : \xi^{L'_x} = 1} (X - \xi)^{p^{k_x}}; \quad (\text{C23})$$

that is, there are L'_x distinct ξ 's in \mathbb{F}_{p^∞} that satisfy $\xi^{L_x} = 1$, each with multiplicity p^{k_x} .

Let $(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})$. It is clear that X_0 is one of the ξ 's in (C23). Consider the localization $\mathcal{S}_{(X - X_0, Y - Y_0)}$. Recall that every element outside the ideal $(X - X_0, Y - Y_0)$ becomes a *unit* in the localization $\mathcal{S}_{(X - X_0, Y - Y_0)}$.²⁹ In particular, the polynomial $\mathbf{Q}_x(X, Y) = X^{L_x} - 1$ generates the same ideal in $\mathcal{S}_{(X - X_0, Y - Y_0)}$ as the polynomial $\tilde{\mathbf{Q}}_{x, X_0}(X, Y) = (X - X_0)^{p^{k_x}}$ because the other linear factors in (C23) associated with $\xi \neq X_0$ all have inverses.³⁰ Similarly, $\mathbf{Q}_y(X, Y) = Y^{L_y} - 1$ generates the same ideal as $\tilde{\mathbf{Q}}_{y, Y_0}(X, Y) = (Y - Y_0)^{p^{k_y}}$. It follows that

$$(\mathfrak{i} + \mathfrak{j})_{(X - X_0, Y - Y_0)} = (\tilde{P}, \tilde{\mathbf{Q}}_{x, X_0}, \tilde{\mathbf{Q}}_{y, Y_0})_{(X - X_0, Y - Y_0)}, \quad (\text{C24})$$

in the localization $\mathcal{S}_{(X - X_0, Y - Y_0)}$. Here we use the notation \mathfrak{J}_m to denote the ideal in \mathcal{S}_m generated by the image of the ideal $\mathfrak{J} \subseteq \mathcal{S}$ under the *localization map* $\mathcal{S} \rightarrow \mathcal{S}_m$, where m is a maximal ideal of \mathcal{S} .

Defining $i'_{X_0, Y_0} = (\tilde{P}, \tilde{\mathbf{Q}}_{x, X_0}, \tilde{\mathbf{Q}}_{y, Y_0})$, we then have

$$\begin{aligned} [\mathcal{S}/(\mathfrak{i} + \mathfrak{j})]_{(X - X_0, Y - Y_0)} &\cong \mathcal{S}_{(X - X_0, Y - Y_0)}/(\mathfrak{i} + \mathfrak{j})_{(X - X_0, Y - Y_0)} \\ &\cong \mathcal{S}_{(X - X_0, Y - Y_0)}/(i'_{X_0, Y_0})_{(X - X_0, Y - Y_0)} \\ &\cong (\mathcal{S}/i'_{X_0, Y_0})_{(X - X_0, Y - Y_0)}, \end{aligned} \quad (\text{C25})$$

where in the first and third lines, we used the slogan ‘‘localization commutes with quotienting,’’ and the second line

²⁹A unit is an element of the ring that has a multiplicative inverse. In any ring, the multiplicative identity 1 is its own inverse, so it is always a unit. In particular, in \mathbb{Z} , ± 1 are the only units, whereas in \mathbb{Q} , any nonzero rational is a unit.

³⁰More abstractly, if r is one of the generators of an ideal \mathfrak{J} of \mathcal{R} , then we can replace it by ur , where u is a unit.

follows from (C24). Now, the quotient $\mathcal{S}/i'_{X_0, Y_0}$ is also Artinian, and by Hilbert's Nullstellensatz, its only maximal ideal is $(X - X_0, Y - Y_0)$. Hence, by a result similar to the one in (C17), we have

$$(\mathcal{S}/i'_{X_0, Y_0})_{(X - X_0, Y - Y_0)} \cong \mathcal{S}/i'_{X_0, Y_0}, \quad (\text{C26})$$

and therefore,

$$\log_p \text{GSD} = 2 \sum_{(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})} \dim_{\mathbb{F}_{p^\infty}} \mathcal{S}/i'_{X_0, Y_0}. \quad (\text{C27})$$

d. Computation of log_p GSD for special values of L_x, L_y

Given the expression (C16) for the GSD, all we need to do now is to compute a Gröbner basis of i'_{X_0, Y_0} in \mathcal{S} for each $(X_0, Y_0) \in V(\mathfrak{i} + \mathfrak{j})$. First, we note certain ‘‘symmetries’’ in the set $V(\mathfrak{i} + \mathfrak{j})$. Recall that $V(\mathfrak{i} + \mathfrak{j})$ is the set of $(X_0, Y_0) \in \mathbb{F}_{p^\infty}^2$ that solve the system of polynomial equations

$$X^{L'_x} - 1 = Y^{L'_y} - 1 = \tilde{P}(X, Y) = 0, \quad (\text{C28})$$

where we used the facts that $X^{L_x} - 1 = (X^{L'_x} - 1)^{p^{k_x}}$ and $Y^{L_y} - 1 = (Y^{L'_y} - 1)^{p^{k_y}}$ in $\mathbb{F}_{p^\infty}[X, Y]$. [Here we used the parametrization (C14).] Given a solution (X_0, Y_0) of (C28), we can generate three more solutions using the transformations

$$X_0 \rightarrow X_0^{-1}, \quad Y_0 \rightarrow Y_0^{-1}, \quad (\text{C29})$$

because the equations (C28) are invariant under these transformations. (These transformations are well defined because $X_0 \neq 0$ and $Y_0 \neq 0$.) Furthermore, if $L'_x = L'_y$, we can generate four more solutions using the exchange

$$X_0 \leftrightarrow Y_0. \quad (\text{C30})$$

It is clear that (1,1) is the only solution that is invariant under all these transformations. We will exploit these transformations in our analysis below.

In general, it is hard to compute a Gröbner basis of i'_{X_0, Y_0} for arbitrary $(X_0, Y_0) \in \mathbb{F}_{p^\infty}^2$ except when $(X_0, Y_0) = (1, 1)$. So below we specialize to those values of L_x, L_y for which $(X_0, Y_0) = (1, 1)$ is the only solution of (C28). These special values of L_x, L_y contain infinite families of L_x, L_y with interesting behaviors of GSD.

(1) Consider the special case $L'_x = L'_y = 1$, where (C28) becomes

$$X - 1 = Y - 1 = \tilde{P}(X, Y) = 0. \quad (\text{C31})$$

Clearly, $(X_0, Y_0) = (1, 1)$ is the only solution of (C31). In this case, changing the variables X and Y to $\tilde{X} = X - 1$ and $\tilde{Y} = Y - 1$, we have $i'_{1,1} = (\tilde{X}^2 \tilde{Y} + \tilde{X} \tilde{Y}^2 + \tilde{X}^2 + \tilde{Y}^2, \tilde{X}^{p^{k_x}}, \tilde{Y}^{p^{k_y}})$. We can assume that $k_x \geq k_y$ without loss of generality. Then, with a lexicographic monomial order on \tilde{X}, \tilde{Y} with $\tilde{X} \succ \tilde{Y}$, a Gröbner basis of $i'_{1,1}$ is given by the following:

(1) When $k_y > 0$,

$$\begin{aligned} \mathbf{G}_1(\tilde{X}, \tilde{Y}) &= \tilde{Y}^{p^{k_y}}, \\ \mathbf{G}_2(\tilde{X}, \tilde{Y}) &= \tilde{X} \tilde{Y}^{p^{k_y} - 1} \delta_{k_x, k_y}, \end{aligned} \quad (\text{C32})$$

$$\mathbf{G}_3(\tilde{X}, \tilde{Y}) = \tilde{X}^2 + (\tilde{X} + 1) \tilde{Y}^2 \sum_{i=0}^{p^{k_y} - 3} (-\tilde{Y})^i - \tilde{X} \tilde{Y}^{p^{k_y} - 1} \delta_{k_x, k_y}.$$

(2) When $k_y = 0$,

$$\begin{aligned} \mathbf{G}_1(\tilde{X}, \tilde{Y}) &= \tilde{Y}, \\ \mathbf{G}_2(\tilde{X}, \tilde{Y}) &= \tilde{X} \delta_{k_x, 0} + \tilde{X}^2(1 - \delta_{k_x, 0}). \end{aligned} \quad (\text{C33})$$

The set of monomials that are irreducible with respect to this Gröbner basis are

$$\{\tilde{Y}^i : 0 \leq i < p^{k_y}\} \cup \{\tilde{X}\tilde{Y}^j : 0 \leq j < p^{k_y} - \delta_{k_x, k_y}\}. \quad (\text{C34})$$

They form a basis of $\mathcal{S}/i'_{1,1}$, so we have

$$\dim_{\mathbb{F}_{p^\infty}} \mathcal{S}/i'_{1,1} = 2p^{k_y} - \delta_{k_x, k_y}. \quad (\text{C35})$$

By (C16), we conclude that

$$\log_p \text{GSD} = 2(2p^{k_y} - \delta_{k_x, k_y}). \quad (\text{C36})$$

(2) Next, we generalize the previous special case to $L'_x = L'_y = q$ with $q > 2$ a prime such that p is a *primitive root* modulo q .³¹ Then (C28) becomes

$$X^q - 1 = Y^q - 1 = \tilde{\mathbf{P}}(X, Y) = 0. \quad (\text{C37})$$

We will show that $X_0 = Y_0 = 1$ is the only solution of (C37). We argue by contradiction. We assume that there is a solution with $X_0 \neq 1$. Then any other solution (X_0, Y_0) of (C37) is obtained from a solution of the form (X_0, X_0^s) for some $1 \leq s \leq (q-1)/2$ using the transformations (C29) and (C30).³² Since $X_0^q = 1$ and $X_0 \neq 1$, X_0 is a root of the *cyclotomic polynomial* $\Phi_q(X) = \sum_{j=0}^{q-1} X^j$. Moreover, since (X_0, X_0^s) satisfies $\tilde{\mathbf{P}}(X, Y) = 0$, X_0 is also a root of the polynomial $\tilde{\mathbf{P}}(X, X^s)$. We can write

$$\tilde{\mathbf{P}}(X, X^s) = X(X^s - 1)^2 + X^s(X - 1)^2 = X(X - 1)^2 \hat{\mathbf{P}}_s(X), \quad (\text{C38})$$

where

$$\hat{\mathbf{P}}_s(X) = \left(\sum_{i=0}^{s-1} X^i \right)^2 + X^{s-1}. \quad (\text{C39})$$

$\hat{\mathbf{P}}_s(X)$ is a nonzero polynomial in $\mathbb{Z}_p[X]$ because $\hat{\mathbf{P}}_s(0) = 1$ mod p for $s > 1$, and $2 \pmod p$ for $s = 1$.³³ Since X_0 is a root of $\tilde{\mathbf{P}}(X, X^s)$ and $X_0 \neq 0, 1$, it must be a root of $\hat{\mathbf{P}}_s(X)$. We now use the fact that $\Phi_q(X)$ is the *minimal polynomial* of X_0 in $\mathbb{Z}_p[X]$ because p is a primitive root modulo q [[54],

³¹In other words, p is a generator of the multiplicative group of integers modulo q , denoted as \mathbb{Z}_q^\times . For any positive integer n , the order of the group \mathbb{Z}_n^\times is known as the *Euler totient function* of n , denoted as $\varphi(n)$. It is easy to see that $\varphi(q^m) = q^m - q^{m-1}$ for any odd prime q and any $m \geq 1$.

³²Since $X_0 \neq 1$, $X_0^q = 1$, and q is prime, X_0 is a primitive q th root of unity, i.e., powers of X_0 generates all the q th roots of unity. Since $Y_0^q = 1$, it is a q th root of unity, and hence, $Y_0 = X_0^s$ for some $0 \leq s < q$. But $s \neq 0$ because there is no solution of (C37) of the form $(X_0, 1)$ for $X_0 \neq 1$. Moreover, using the transformation (C29), (X_0, X_0^{q-s}) is also a solution, so we can restrict s to the range $1 \leq s \leq (q-1)/2$.

³³This is not true for $p = 2$ because $\hat{\mathbf{P}}_s(X) = 0$ identically for $s = 1$, so there is always a solution of (C28) of the form (X_0, X_0) even for $X_0 \neq 1$.

Sec. 11.2.B]. This means $\Phi_q(X)$ divides $\hat{\mathbf{P}}_s(X)$. But this is impossible because

$$\deg_X \hat{\mathbf{P}}_s(X) = 2s - 2 \leq q - 3 < q - 1 = \deg_X \Phi_q(X), \quad (\text{C40})$$

so there is no such X_0 . In other words, when p is a primitive root modulo q , the only solution of (C37) is $(1, 1)$. Then, by the analysis in point 1 above, $\log_p \text{GSD}$ is again given by (C36).

(3) We now generalize as follows. Let $q > 2$ be a prime such that p is a primitive root modulo q^m for some $m \geq 2$. Set $L'_x = L'_y = q^m$, so that (C28) becomes

$$X^{q^m} - 1 = Y^{q^m} - 1 = \tilde{\mathbf{P}}(X, Y) = 0. \quad (\text{C41})$$

Again, we argue by contradiction that the only solution of (C41) $(X_0, Y_0) \in \mathbb{F}_{p^\infty}^2$ is $(1, 1)$. We assume that $X_0 \neq 1$. Then any other solution (X_0, Y_0) of (C41) is obtained from a solution of the form (X_0, X_0^s) for some $1 \leq s \leq (q^m - 1)/2$ using the transformations (C29) and (C30). Actually, the range of s can be smaller than this. Let q^r be the order of X_0 for some $0 \leq r \leq m$, i.e., $X_0^{q^r} = 1$, but $X_0^{q^{r'}} \neq 1$ for any $r' < r$. Since $r = 0$ corresponds to the trivial solution $(1, 1)$, we have $r > 0$. Then $1 \leq s \leq (q^r - 1)/2$. We consider two cases:

(3) $s \leq \varphi(q^r)/2$: [See footnote 31 for the definition of $\varphi(q^r)$.] Since the order of X_0 is q^r , it is a root of the cyclotomic polynomial $\Phi_{q^r}(X) = \sum_{j=0}^{q^r-1} X^{jq^{r-1}}$. Since $X_0 \neq 0, 1$, it is also a root of the polynomial $\hat{\mathbf{P}}_s(X)$ in (C39). We now use the fact that $\Phi_{q^r}(X)$ is the *minimal polynomial* of X_0 in $\mathbb{Z}_p[X]$ because p is a primitive root modulo q^r [[54], Sec. 11.2.B].³⁴ This means $\Phi_{q^r}(X)$ must divide $\hat{\mathbf{P}}_s(X)$. But this is impossible because

$$\deg_X \hat{\mathbf{P}}_s(X) = 2s - 2 \leq \varphi(q^r) - 2 < \varphi(q^r) = \deg_X \Phi_{q^r}(X), \quad (\text{C42})$$

so there is no such X_0 .

(4) $\varphi(q^r)/2 < s \leq (q^r - 1)/2$: Let $X_0 = Z_0^2$, where Z_0 also has order q^r . [Such a Z_0 exists because $\gcd(2, q) = 1$ for $q > 2$] Then the solution (X_0, Y_0) is of the form (Z_0^2, Z_0^{2s}) . By the transformation (C29), (Z_0^2, Z_0^t) is also a solution, where $t = q^r - 2s$. Then

$$\frac{\varphi(q^r)}{2} < s \leq \frac{q^r - 1}{2} \Rightarrow 1 \leq t < q^{r-1} \leq \frac{q^r - q^{r-1}}{2} = \frac{\varphi(q^r)}{2}, \quad (\text{C43})$$

where the rightmost inequality holds for $q > 2$. Clearly, Z_0 is a root of $\Phi_{q^r}(X)$. Since (Z_0^2, Z_0^t) satisfies $\tilde{\mathbf{P}}(X, Y) = 0$, Z_0 is a root of $\tilde{\mathbf{P}}(X^2, X^t)$. We can write

$$\begin{aligned} \tilde{\mathbf{P}}(X^2, X^t) &= X^2(X^t - 1)^2 + X^t(X^2 - 1)^2 \\ &= \begin{cases} X(X - 1)^2 \check{\mathbf{P}}_1(X), & t = 1, \\ X^2(X - 1)^2 \check{\mathbf{P}}_t(X), & t > 1, \end{cases} \end{aligned} \quad (\text{C44})$$

³⁴Here we used the fact that p is a primitive root modulo $q^m \Rightarrow p$ is a primitive root modulo q^r for all $r \leq m$. In fact, for $m \geq 2$, p is a primitive root modulo $q^m \Rightarrow p$ is a primitive root modulo q^{m+1} . Combining these facts, p is a primitive root modulo $q^2 \Rightarrow p$ is a primitive root modulo q^m for all $m \geq 1$ [[48], Sec. 2.8].

where

$$\check{P}_t(X) = \begin{cases} X + (X + 1)^2, & t = 1, \\ (\sum_{i=0}^{t-1} X^i)^2 + X^{t-2}(X + 1)^2, & t > 1. \end{cases} \quad (\text{C45})$$

$\check{P}_t(X)$ is a nonzero polynomial because $\check{P}_t(0) = 1 \pmod p$ for $t \neq 2$, and $2 \pmod p$ for $t = 2$.³⁵ Since Z_0 is a root of $\check{P}(X^2, X^t)$ and $Z_0 \neq 0, 1$, it is a root of $\check{P}_t(X)$ as well. We now use the fact that $\Phi_{q^r}(X)$ is the minimal polynomial of Z_0 in $\mathbb{Z}_p[X]$ because p is a primitive root modulo q^r [[54], Sec. 11.2.B]. This means $\Phi_{q^r}(X)$ must divide $\check{P}_t(X)$. But this is impossible because

$$\begin{aligned} \deg_X \check{P}_t(X) &= t + \max(t, 2) - 2 + \delta_{t,1} \leq 2t < \varphi(q^r) \\ &= \deg_X \Phi_{q^r}(X), \end{aligned} \quad (\text{C46})$$

where in the third line, we used (C43). So there is no such Z_0 .

Therefore, when p is a primitive root modulo q^m , then (1,1) is the only solution of (C41), and hence, \log_p GSD is still given by (C36).

To conclude, when $L_x = p^{k_x} q^m$ and $L_y = p^{k_y} q^m$, where q is an odd prime such that p is a primitive root modulo q^m , and $k_x, k_y, m \geq 0$, the ground-state degeneracy of the 3+1D anisotropic \mathbb{Z}_p Laplacian model is given by

$$\log_p \text{GSD} = 2[2p^{\min(k_x, k_y)} - \delta_{k_x, k_y}]. \quad (\text{C47})$$

When $m = 0$, we see that \log_p GSD scales as $4 \min(L_x, L_y)$. This gives an infinite family of L_x, L_y for which \log_p GSD is $O(L_x, L_y)$.

Say q is such that p is a primitive root modulo q^2 . Then p is a primitive root modulo q^m for all $m \geq 1$ (see footnote 34). Then, for $k_x = k_y = 0$ and any $m \geq 0$, we see that \log_p GSD = 2, a finite number. This gives an infinite family of L_x, L_y for which \log_p GSD remains finite.³⁶

Note that the last conclusion relies on the existence of an odd prime q such that p is a primitive root modulo q^2 . However, we do not know of a proof for general p . Another interesting possibility is the following. By *Artin's conjecture on primitive roots* [49],³⁷ there are infinitely many prime q such that p is a primitive root modulo q . (Recall from footnote 34 that this does not imply that p is a primitive root modulo q^2 .) Then choosing $L_x = L_y = q$ for all such q gives another infinite family of L_x, L_y for which \log_p GSD = 2. However, Artin's conjecture is still unproven, except under the assumption of the *generalized Riemann hypothesis* [50], which is also unproven.

2. Mobility restrictions

We now discuss the mobility of z -lineons in the xy plane in the 3+1D anisotropic \mathbb{Z}_p Laplacian model. The lineons

³⁵Once again, this is not true for $p = 2$ because $\check{P}_t(X) = 0$ identically when $t = 2$, so there is always a solution of (C28) of the form (Z_0^2, Z_0^2) even for $Z_0 \neq 1$.

³⁶In contrast, when $N = 2$, \log_2 GSD in (B18) always scales as $4L$ for any L . This is because, when $p = 2$, the above arguments do not go through, as explained in footnotes 33 and 35.

³⁷The conjecture is actually stronger: the set of such q has positive asymptotic density inside the set of all primes.

are represented as defects in the low-energy theory and their motion is implemented by operators acting at fixed time.

These operators fall into two kinds. First, there are operators supported in a small region, e.g., the line joining the two points. Second, there are also situations where the operator spans over $O(L_x, L_y)$ sites. Operators of the second kind exist only for certain special values of L_x, L_y depending on some number-theoretic properties of L_x, L_y , whereas the first kind exist for all L_x, L_y . In particular, only the first kind exist on an infinite square lattice. (See the discussion in [23,35].)

As an example of the second kind of operator, consider $L_x = L_y = q^m$, where $q > 2$ is a prime such that p is a primitive root modulo q^m . In Appendix C 1, we showed that for $L_x = L_y = q^m$, where $q > 2$ is a prime such that p is a primitive root modulo q^m , the ground-state degeneracy is given by $\log_p \text{GSD} = 2$. It follows that $|\text{Jac}(C_{q^m} \times C_{q^m}, p)| = p$, or equivalently, $\text{Jac}(C_{q^m} \times C_{q^m}, p) = \mathbb{Z}_p$. Therefore, in this case, the only selection imposed by the $\text{Jac}(C_{q^m} \times C_{q^m}, p)$ timelike symmetry is that the total charge of the defects is conserved modulo p . In particular, a z -lineon can move anywhere within the xy plane when $L_x = L_y = q^m$. However, we show below that a z -lineon is completely immobile when Γ is an infinite square lattice. This means, the operator that moves a z -lineon on the 2D torus graph $C_{q^m} \times C_{q^m}$ must be of the second kind.

We now show that a z -lineon is completely immobile on an infinite square lattice. In fact, we show that any finite configuration of z -lineons is completely immobile (except in some trivial cases) as long as their charges and the separations between them are fixed during the motion, i.e., we allow only “rigid” motion. Without this restriction, the groups of lineons can move. We will not discuss this motion.

Our analysis will be similar to the analogous discussion in [23]. The main difference between them is that here our variables are in \mathbb{Z}_p and therefore various properties of the polynomials will depend on p .

Consider n z -lineons, with charges q_i and positions (x_i, y_i) for $i = 1, \dots, n$, described by the defect

$$\exp \left[\frac{2\pi i}{p} \sum_{\tau} \sum_{i=1}^n q_i m_{\tau} \left(\tau + \frac{1}{2}, x_i, y_i \right) \right]. \quad (\text{C48})$$

(Since they are z -lineons, we can assume without loss of generality that they all have the same z coordinate, and omit writing it.) They can move “rigidly” by $(x_0, y_0) \neq (0, 0)$ if there is a defect of the form

$$\begin{aligned} & \exp \left[\frac{2\pi i}{p} \sum_{\tau < 0} \sum_{i=1}^n q_i m_{\tau} \left(\tau + \frac{1}{2}, x_i, y_i \right) \right] \\ & \times \exp \left[\frac{2\pi i}{p} \sum_{j=1}^l s_j m(0, x_j, y_j) \right] \\ & \times \exp \left[\frac{2\pi i}{p} \sum_{\tau \geq 0} \sum_{i=1}^n q_i m_{\tau} \left(\tau + \frac{1}{2}, x_i + x_0, y_i + y_0 \right) \right]. \end{aligned} \quad (\text{C49})$$

This defect is gauge invariant if and only if

$$\begin{aligned} & \sum_{i=1}^n q_i [k(0, x_i + x_0, y_i + y_0) - k(0, x_i, y_i)] \\ &= \sum_{j=1}^l s_j (\Delta_x^2 + \Delta_y^2) k(0, x_j, y_j) \pmod{p}, \end{aligned} \quad (\text{C50})$$

for any integer gauge parameter k in (3.9).

Using a formal Laurent power series

$$\hat{k}(X, Y) = \sum_{(x,y) \in \mathbb{Z}^2} k(0, x, y) X^{-x} Y^{-y}, \quad (\text{C51})$$

associated with the gauge parameter $k(0, x, y)$, the condition (C50) can be written as

$$(X^{x_0} Y^{y_0} - 1) \mathbf{Q}(X, Y) = \mathbf{S}(X, Y) \mathbf{P}(X, Y) \pmod{p}, \quad (\text{C52})$$

where

$$\mathbf{P}(X, Y) = (X - 2 + X^{-1}) + (Y - 2 + Y^{-1}) \quad (\text{C53})$$

is the *Laurent polynomial* (i.e., an element of $\mathbb{Z}_p[X, X^{-1}, Y, Y^{-1}]$) associated with the discrete Laplacian operator $\Delta_x^2 + \Delta_y^2$, and

$$\mathbf{Q}(X, Y) = \sum_{i=1}^n q_i X^{x_i} Y^{y_i}, \quad \mathbf{S}(X, Y) = \sum_{j=1}^l s_j X^{x_j} Y^{y_j} \quad (\text{C54})$$

are also Laurent polynomials. The coefficients and monomials in $\mathbf{Q}(X, Y)$ and $\mathbf{S}(X, Y)$ are obtained from the defect (C49).

If there is a Laurent polynomial $\mathbf{R}(X, Y)$ such that $\mathbf{Q}(X, Y) = \mathbf{R}(X, Y) \mathbf{P}(X, Y)$, then (C52) can be trivially satisfied by choosing $\mathbf{S}(X, Y) = (X^{x_0} Y^{y_0} - 1) \mathbf{R}(X, Y)$. However, in this case, the defect (C48) can end at $\tau = 0$ as follows:

$$\begin{aligned} & \exp \left[\frac{2\pi i}{p} \sum_{\tau < 0} \sum_{i=1}^n q_i m_{\tau} \left(\tau + \frac{1}{2}, x_i, y_i \right) \right] \\ & \times \exp \left[-\frac{2\pi i}{p} \sum_{j'=1}^{l'} r_{j'} m(0, x_{j'}, y_{j'}) \right], \end{aligned} \quad (\text{C55})$$

where $r_{j'}$'s and $(x_{j'}, y_{j'})$'s are obtained from $\mathbf{R}(X, Y) = \sum_{j'=1}^{l'} r_{j'} X^{x_{j'}} Y^{y_{j'}}$. Therefore, in this case, the defect (C49) describes the annihilation of the n z -lineons at their original positions and their creation at positions displaced by (x_0, y_0) at time $\tau = 0$.

A more interesting situation occurs for a defect like (C49) when $\mathbf{Q}(X, Y)$ cannot be written as $\mathbf{R}(X, Y) \mathbf{P}(X, Y)$ for any Laurent polynomial $\mathbf{R}(X, Y)$. Then, to satisfy (C52), $\mathbf{P}(X, Y)$ and $X^{x_0} Y^{y_0} - 1$ must share a nontrivial factor.³⁸ Let us show that this cannot happen. In the following, it is crucial that $(x_0, y_0) \neq (0, 0)$.

³⁸By *nontrivial factor* we mean a nonconstant Laurent factor that is not a Laurent monomial.

First, note that $\mathbf{P}(X, Y)$ is nonconstant and irreducible up to a monomial in $\mathbb{Z}_p[X, X^{-1}, Y, Y^{-1}]$ for any odd prime p .³⁹ So all we need to show is that $X^{x_0} Y^{y_0} - 1$ is not a multiple of $\mathbf{P}(X, Y)$ in $\mathbb{Z}_p[X, X^{-1}, Y, Y^{-1}]$.

Let p^k be the largest power of p that divides both x_0 and y_0 , i.e., $x'_0 = x_0/p^k$ and $y'_0 = y_0/p^k$ are integers and $d = \gcd(x'_0, y'_0)$ is not divisible by p . Then, in $\mathbb{Z}_p[X, X^{-1}, Y, Y^{-1}]$, we have

$$X^{x_0} Y^{y_0} - 1 = (X^{x'_0} Y^{y'_0} - 1)^{p^k} = [(X^{x''_0} Y^{y''_0} - 1) \mathbf{T}(X, Y)]^{p^k}, \quad (\text{C56})$$

where $x''_0 = x'_0/d$, $y''_0 = y'_0/d$, and $\mathbf{T}(X, Y) = \sum_{c=0}^{d-1} (X^{x''_0} Y^{y''_0})^c$. Now, $\mathbf{T}(1, 1) = d \neq 0 \pmod{p}$, whereas $\mathbf{P}(1, 1) = 0$, so $\mathbf{T}(X, Y)^{p^k}$ is not a multiple of $\mathbf{P}(X, Y)$. Since $\gcd(x''_0, y''_0) = 1$, the factor $X^{x''_0} Y^{y''_0} - 1$ is nonconstant and irreducible up to a monomial for any p [56]. So $(X^{x''_0} Y^{y''_0} - 1)^{p^k}$ is also not a multiple of $\mathbf{P}(X, Y)$. Therefore, $X^{x_0} Y^{y_0} - 1$ is not a multiple of $\mathbf{P}(X, Y)$.

To conclude, a finite set of z -lineons cannot move “rigidly” in the xy plane in the 3+1D anisotropic \mathbb{Z}_p Laplacian model, unless they can be annihilated.

APPENDIX D: \mathbb{Z}_N LAPLACIAN MODEL ON A GRAPH

In this Appendix, we analyze a gapped fracton model on a simple, connected, undirected spatial graph Γ . We refer to it as the \mathbb{Z}_N Laplacian model because the theory is defined using the discrete Laplacian operator Δ_L on the graph Γ . The anisotropic \mathbb{Z}_N Laplacian model in Sec. III is an anisotropic extension of this \mathbb{Z}_N Laplacian model by adding another direction.

1. Hamiltonian

In the Hamiltonian formulation of the \mathbb{Z}_N Laplacian model, there are a \mathbb{Z}_N variable U_i and its conjugate variable V_i , i.e., $U_i V_i = e^{2\pi i/N} V_i U_i$, on every site of the graph Γ where i labels the sites. The Hamiltonian is

$$H = -\gamma_1 \sum_i G_i + \text{H.c.}, \quad (\text{D1})$$

where

$$G_i = \prod_{j:(i,j) \in \Gamma} V_i V_j^\dagger. \quad (\text{D2})$$

³⁹A polynomial in $\mathbb{Z}_p[X, Y]$ is said to be *irreducible* if it cannot be written as a product of two nonconstant polynomials. A Laurent polynomial $\mathbf{F}(X, Y)$ in $\mathbb{Z}_p[X, X^{-1}, Y, Y^{-1}]$ is said to be *irreducible up to a monomial* if $X^a Y^b \mathbf{F}(X, Y)$ is an irreducible polynomial for some $a, b \in \mathbb{Z}$. For example, $\mathbf{P}(X, Y)$ is irreducible up to a monomial because $\tilde{\mathbf{P}}(X, Y) = XY \mathbf{P}(X, Y)$, given by (C7), is an irreducible polynomial. The irreducibility of $\tilde{\mathbf{P}}(X, Y)$ for any prime $p > 6$ follows from [[55], Corollary 3]. It is easy to verify by hand, or in *Mathematica*, that it is irreducible even for $p = 3, 5$. It is, however, not irreducible for $p = 2$ because $\tilde{\mathbf{P}}(X, Y) = (XY + 1)(X + Y) \pmod{2}$. This is one way of seeing why a dipole of z -lineons separated in $(1, \pm 1)$ direction can move in $(1, \mp 1)$ direction in the 3+1D anisotropic \mathbb{Z}_2 Laplacian model.

Here $\langle i, j \rangle$ means i and j are connected by an edge in the graph Γ .

Since all the G_i s commute, the ground states satisfy $G_i = 1$ for all i and the excitations are violations of $G_i = 1$. We could take the limit $\gamma_1 \rightarrow \infty$, in which case, the Hilbert space consists of only the ground states and the Hamiltonian is trivial. The Euclidean presentation of this model in this limit will be discussed later in Appendix D 2.

We are particularly interested in those operators that commute with the Hamiltonian (D1) and act nontrivially on its ground states. They are the global symmetry operators of the model in the low-energy limit.

The electric symmetry operators are V_i , which trivially commute with the Hamiltonian. Since the ground states satisfy $G_i = 1$, some of these operators are equivalent when acting on the ground states. The independent symmetry operators are

$$\tilde{W}_\lambda = \prod_i V_i^{\lambda(i)}, \quad (\text{D3})$$

where $\lambda(i)$ takes the form (2.6)

$$\lambda(i) = \sum_{a=1}^N p_a (Q^{-1})_{ai}, \quad p_a \sim p_a + \text{gcd}(N, r_a). \quad (\text{D4})$$

Let us explain the identification on p_a . We have $p_a \sim p_a + N$ because $V_i^N = 1$. We also have $p_a \sim p_a + r_a$ because $r_a (Q^{-1})_{ai} = \sum_j L_{ij} P_{aj}$ and $\prod_i V_i^{L_{ij}} = 1$ when acting on the ground states. Combining the two identifications, we get $p_a \sim p_a + \text{gcd}(N, r_a)$. The symmetry operators generate a $\text{Jac}(\Gamma, N)$ electric symmetry.

The magnetic symmetry operators are

$$W_{\tilde{\lambda}} = \prod_i U_i^{\tilde{\lambda}(i)}, \quad (\text{D5})$$

where $\tilde{\lambda}(i)$ obeys $\Delta_L \tilde{\lambda}(i) = 0 \pmod{N}$, and the most general solution takes the form (2.5)

$$\tilde{\lambda}(i) = \sum_{a=1}^N \frac{N Q_{ia} \tilde{p}_a}{\text{gcd}(N, r_a)}, \quad \tilde{p}_a \sim \tilde{p}_a + \text{gcd}(N, r_a). \quad (\text{D6})$$

The symmetry operators generate a $\text{Jac}(\Gamma, N)$ magnetic symmetry.

A convenient basis of electric and magnetic spacelike symmetry operators is given by

$$\begin{aligned} \tilde{W}(a) &= \prod_i V_i^{(Q^{-1})_{ai}}, \\ W(a) &= \prod_i U_i^{\frac{N}{\text{gcd}(N, r_a)} Q_{ia}} \end{aligned} \quad (\text{D7})$$

for $a = 1, \dots, N$. Both $W(a)$ and $\tilde{W}(a)$ are $\mathbb{Z}_{\text{gcd}(N, r_a)}$ operators. They satisfy the commutation relations

$$\begin{aligned} W(a) \tilde{W}(b) &= \exp \left[\frac{2\pi i \delta_{ab}}{\text{gcd}(N, r_a)} \right] \tilde{W}(b) W(a), \\ a, b &= 1, \dots, N. \end{aligned} \quad (\text{D8})$$

For each a , there is only one b that has nontrivial commutation relation. So for each a , there is an independent $\mathbb{Z}_{\text{gcd}(N, r_a)}$ Heisenberg algebra generated by $W(a)$ and $\tilde{W}(a)$, leading to

a ground-state degeneracy

$$\text{GSD} = \prod_{a=1}^N \text{gcd}(N, r_a) = |\text{Jac}(\Gamma, N)|. \quad (\text{D9})$$

2. Euclidean presentation

We now discuss the Euclidean presentation of the \mathbb{Z}_N Laplacian model. We place the theory on a Euclidean spacetime lattice $C_{L_\tau} \times \Gamma$, where Γ is the spatial slice. We use (τ, i) to label a site in the spacetime lattice. The integer BF action of the \mathbb{Z}_N Laplacian theory is

$$S = \frac{2\pi i}{N} \sum_{\tau, i} \tilde{m} \left(\tau + \frac{1}{2}, i \right) \left[\Delta_\tau m(\tau, i) - \Delta_L m_\tau \left(\tau + \frac{1}{2}, i \right) \right], \quad (\text{D10})$$

where the integer fields \tilde{m} and (m_τ, m) have an integer gauge symmetry

$$\begin{aligned} \tilde{m} &\sim \tilde{m} + N\tilde{k}, \\ m_\tau &\sim m_\tau + \Delta_\tau k + Nq_\tau, \\ m &\sim m + \Delta_L k + Nq, \end{aligned} \quad (\text{D11})$$

where k, \tilde{k} , and (q_τ, q) are integers. [Note that, when working modulo N , the last line of (D11) is precisely the equivalence relation discussed in (2.3).] The integer BF action (D10) describes the ground states of the Hamiltonian (D1).

a. Ground-state degeneracy

We can count the number of ground states by counting the number of solutions to the ‘‘equations of motion’’ of (m_τ, m) :

$$\Delta_\tau \tilde{m} = 0 \pmod{N}, \quad \Delta_L \tilde{m} = 0 \pmod{N}. \quad (\text{D12})$$

The first equation implies that $\tilde{m}(\tau, i)$ is independent of τ . Then, as discussed in Sec. II A, the general solution is

$$\tilde{m}(i) = \sum_{a=1}^N \frac{N Q_{ia} p_a}{\text{gcd}(N, r_a)}, \quad p_a \sim p_a + \text{gcd}(N, r_a). \quad (\text{D13})$$

Therefore, the ground-state degeneracy is

$$\text{GSD} = \prod_{a=1}^N \text{gcd}(N, r_a) = |\text{Jac}(\Gamma, N)|. \quad (\text{D14})$$

b. Global symmetry

The above ground-state degeneracy can also be obtained from the (spacelike) global symmetry. There are electric (spacelike and timelike) and magnetic (spacelike) global symmetries, whose groups are both

$$\text{Jac}(\Gamma, N) = \prod_{a=1}^N \mathbb{Z}_{\text{gcd}(N, r_a)}. \quad (\text{D15})$$

The electric global symmetry acts as

$$(m_\tau, m) \rightarrow (m_\tau, m) + (\lambda_\tau, \lambda), \quad (\text{D16})$$

where (λ_τ, λ) is a flat \mathbb{Z}_N gauge field, i.e., $\Delta_\tau \lambda - \Delta_L \lambda_\tau = 0 \pmod{N}$. Using k , we can set $\lambda_\tau(\tau + \frac{1}{2}, i)|_{\tau \neq 0} = 0 \pmod{N}$. Then, by flatness, we have $\Delta_\tau \lambda = 0 \pmod{N}$. This in turn

implies that $\Delta_L \lambda_\tau(\tau + \frac{1}{2}, i)|_{\tau=0} = 0 \pmod{N}$, which is the discrete Laplace equation (2.2).

The remaining time-independent gauge freedom, $\lambda(i) \sim \lambda(i) + \Delta_L k(i)$, is precisely the equivalence relation in (2.3). Therefore, we can gauge fix $\lambda(i)$ to

$$\lambda(i) = \sum_{a=1}^N p_a (Q^{-1})_{ai}, \quad (\text{D17})$$

where $p_a = 0, \dots, \text{gcd}(N, r_a) - 1$. Since $\lambda_\tau(\tau + \frac{1}{2}, i)|_{\tau=0}$ satisfies the discrete Laplace equation, the most general solution is (2.5)

$$\lambda_\tau\left(\tau + \frac{1}{2}, i\right)\Big|_{\tau=0} = \sum_{a=1}^N \frac{NQ_{ia} p_{\tau,a}}{\text{gcd}(N, r_a)}, \quad (\text{D18})$$

where $p_{\tau,a} = 0, \dots, \text{gcd}(N, r_a) - 1$. The parameters $p_{\tau,a}$ and p_a generate the electric timelike and spacelike global symmetries respectively.

The magnetic spacelike global symmetry acts as

$$\begin{aligned} \tilde{m}\left(\tau + \frac{1}{2}, i\right) &\rightarrow \tilde{m}\left(\tau + \frac{1}{2}, i\right) + \tilde{\lambda}(i), \quad \tilde{\lambda}(i) \\ &= \sum_{a=1}^N \frac{NQ_{ia} \tilde{p}_a}{\text{gcd}(N, r_a)}, \end{aligned} \quad (\text{D19})$$

and $\tilde{p}_a = 0, \dots, \text{gcd}(N, r_a) - 1$.

A convenient basis of electric and magnetic spacelike symmetry operators is given by

$$\begin{aligned} \tilde{W}(a) &= \exp\left[\frac{2\pi i}{N} \sum_i (Q^{-1})_{ai} \tilde{m}\left(\tau + \frac{1}{2}, i\right)\right], \\ W(a) &= \exp\left[\frac{2\pi i}{\text{gcd}(N, r_a)} \sum_i m(\tau, i) Q_{ia}\right] \end{aligned} \quad (\text{D20})$$

for $a = 1, \dots, N$. These operators are the low-energy counterpart of the operators in (D7). The commutation relation (D8) can now be understood as a mixed 't Hooft anomaly between the electric and magnetic spacelike symmetries.

c. Timelike symmetry and fractons

The \mathbb{Z}_N Laplacian model has defects that extend in the time direction, such as

$$W_\tau(i) = \exp\left[\frac{2\pi i}{N} \sum_\tau m_\tau\left(\tau + \frac{1}{2}, i\right)\right]. \quad (\text{D21})$$

This describes the world-line of an infinitely heavy particle of unit charge at position $i \in \Gamma$.

Below we discuss the timelike global symmetry that acts on these defects. The electric timelike symmetry acts as

$$m_\tau\left(\tau + \frac{1}{2}, i\right) \rightarrow m_\tau\left(\tau + \frac{1}{2}, i\right) + \delta_{\tau,0} \sum_{a=1}^N \frac{NQ_{ia} p_{\tau,a}}{\text{gcd}(N, r_a)}. \quad (\text{D22})$$

Therefore, two defects at sites i and i' carry the same timelike charges, or equivalently, a particle can hop from i to i' , if and only if

$$Q_{ia} = Q_{i'a} \pmod{\text{gcd}(N, r_a)}, \quad a = 1, \dots, N. \quad (\text{D23})$$

Indeed, when this condition holds, the defect that ‘‘moves’’ a particle from i to i' at time $\tau = 0$ is given by

$$\begin{aligned} &\exp\left[\frac{2\pi i}{N} \sum_{\tau < 0} m_\tau\left(\tau + \frac{1}{2}, i\right)\right] \\ &\times \exp\left[-\frac{2\pi i}{N} \sum_{a,j} \left(\frac{Q_{ia} - Q_{i'a}}{\text{gcd}(N, r_a)}\right) \tilde{r}_a P_{aj} m(0, j)\right] \\ &\times \exp\left[\frac{2\pi i}{N} \sum_{\tau \geq 0} m_\tau\left(\tau + \frac{1}{2}, i'\right)\right], \end{aligned} \quad (\text{D24})$$

where \tilde{r}_a is an integer such that $\tilde{r}_a r_a = \text{gcd}(N, r_a) \pmod{N}$.

While the selection rule (D23) is not very intuitive, it leads to strong mobility constraints in the special case where the spatial lattice is a square lattice (i.e., Γ is a 2D torus graph $C_{L_x} \times C_{L_y}$). This will be shown in Appendix D 3 b. In particular, under some mild conditions, the particles are completely immobile, i.e., they are fractons.

d. Robustness

We now discuss the robustness of the low-energy limit of the \mathbb{Z}_N Laplacian model. The only operators that act nontrivially on the ground states are $W(a)$ and $\tilde{W}(a)$ of (D20). $W(a)$ is an extended operator with support spanning over the entire graph. In contrast, $\tilde{W}(a)$ can be written as a product of local operators of the form $e^{\frac{2\pi i}{N} \tilde{m}(\tau + \frac{1}{2}, i)}$. [In (D20) we defined $\tilde{W}(a)$ in such a way that its commutation relation (D8) with the extended operator $W(a)$ is simple.] Since these local operators act nontrivially in the space of ground states, the low-energy limit of the model is not robust.

3. Examples

a. $\Gamma = C_{L_x}$

Let Γ be a cycle graph, i.e., $\Gamma = C_{L_x}$, where L_x is the number of sites in the cycle. The operator Δ_L associated with the Laplacian matrix of Γ is the same as the standard Laplacian operator Δ_x^2 in the x direction.

In this case, the \mathbb{Z}_N Laplacian model simplifies to the 1+1D rank-2 \mathbb{Z}_N tensor gauge theory of [35]. Indeed, the diagonal entries in the Smith normal form of L are

$$r_a = \begin{cases} 1, & 1 \leq a < L_x - 1, \\ L_x, & a = L_x - 1, \\ 0, & a = L_x. \end{cases} \quad (\text{D25})$$

To be more concrete, we can write $R = PLQ$, where

$$\begin{aligned} R &= \begin{pmatrix} I_{L_x-2} & \mathbf{0} & \mathbf{0} \\ \mathbf{0}^T & L_x & 0 \\ \mathbf{0}^T & 0 & 0 \end{pmatrix}, \quad P = \begin{pmatrix} \tilde{P} & \mathbf{0} \\ \mathbf{1}^T & 1 \end{pmatrix}, \\ Q &= \begin{pmatrix} \tilde{Q} & \mathbf{1} \\ \mathbf{0}^T & 1 \end{pmatrix}, \end{aligned} \quad (\text{D26})$$

where \tilde{P} and \tilde{Q} are $(L_x - 1) \times (L_x - 1)$ integer matrices given by

$$\begin{aligned}\tilde{P}_{a,x+1} &= \min\{a, x + 1\}, \\ \tilde{Q}_{x+1,a} &= \delta_{x+1,a} - (x + 1)(1 - \delta_{x,L_x-2})\delta_{a,L_x-1},\end{aligned}\quad (\text{D27})$$

where $a = 1, \dots, L_x$, and $x = 0, \dots, L_x - 1$. For example, for $L_x = 5$, the 4×4 matrices \tilde{P} and \tilde{Q} are

$$\tilde{P} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{pmatrix}, \quad \tilde{Q} = \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & -3 \\ 0 & 0 & 0 & 1 \end{pmatrix}.\quad (\text{D28})$$

The electric (timelike and spacelike) and magnetic (space-like) symmetries of the \mathbb{Z}_N Laplacian gauge theory are $\mathbb{Z}_N \times \mathbb{Z}_{\text{gcd}(N, L_x)}$. These are in agreement with the 1+1D rank-2 \mathbb{Z}_N tensor gauge theory of [35].

b. $\Gamma = C_{L_x} \times C_{L_y}$

Let Γ be a torus graph, i.e., $\Gamma = C_{L_x} \times C_{L_y}$, where L_x and L_y are the number of sites in the x -cycle and y -cycle. The operator Δ_L associated with the Laplacian matrix of Γ is the same as the standard Laplacian operator $\Delta_x^2 + \Delta_y^2$ on a square lattice.

In this case, the \mathbb{Z}_N Laplacian model can be viewed as the \mathbb{Z}_N version of the Laplacian ϕ -theory or the \mathbb{Z}_N version of the $U(1)$ Laplacian gauge theory discussed in [22,23]. Its ground-state degeneracy is the square root of that of the anisotropic \mathbb{Z}_N Laplacian model, which is computed in Appendix C 1. When N is a prime, the GSD of the \mathbb{Z}_N Laplacian model is thus

$$\begin{aligned}\log_N \text{GSD} &= \dim_{\mathbb{Z}_N} \frac{\mathbb{Z}_N[X, Y]}{(Y(X-1)^2 + X(Y-1)^2, X^{L_x} - 1, Y^{L_y} - 1)}.\end{aligned}\quad (\text{D29})$$

The GSD depends on L_x, L_y in an erratic way. There exists a sequence of L_x, L_y where the $\log_N \text{GSD} \sim O(L_x, L_y)$, but

there also exists a sequence where the GSD stays at order 1 if $N > 2$.

Like the GSD, the mobility of a particles depends on number-theoretic properties of L_x, L_y . Since the mobility of these particles is the same as the mobility of the z -lineons of the 3+1D anisotropic \mathbb{Z}_N Laplacian model in the xy plane, the analysis of Appendix C 2 applies here. In particular, when N is an odd prime, there are infinitely many values of L_x, L_y for which a single particle is completely mobile. In contrast, on an infinite square lattice, any finite set of particles is completely immobile (unless they can be annihilated), assuming they move ‘‘rigidly.’’

When $N = 2$, the \mathbb{Z}_N Laplacian model is equivalent to two copies of a known model, the \mathbb{Z}_2 Ising plaquette model [25], when both L_x and L_y are even, and only one copy when L_x or L_y is odd. Therefore, the GSD and mobility restrictions of the \mathbb{Z}_2 Laplacian model are relatively simple in this case, and follow from the analysis in Appendix B.

Let us contrast the \mathbb{Z}_N Laplacian model with the 2+1D rank-2 \mathbb{Z}_N tensor gauge theory discussed in [57–61], which is another 2+1D generalization of the 1+1D rank-2 \mathbb{Z}_N tensor gauge theory. These two models differ in several aspects:

(1) The \mathbb{Z}_N tensor gauge theory has a ground-state degeneracy of

$$N^3 \text{gcd}(N, L_x) \text{gcd}(N, L_y) \text{gcd}(N, L_x, L_y).\quad (\text{D30})$$

In particular, the GSD of \mathbb{Z}_N tensor gauge theory is always bounded by N^6 , whereas there are infinitely many L_x, L_y for which $\log_N \text{GSD}$ of the \mathbb{Z}_N Laplacian model scales as $O(L_x, L_y)$, at least when N is prime.

(2) Relatedly, the low-energy limit of the \mathbb{Z}_N tensor gauge theory is robust, whereas the low-energy limit of the \mathbb{Z}_N Laplacian model is not.

(3) A particle in the \mathbb{Z}_N tensor gauge theory can always hop by N sites on an infinite square lattice,⁴⁰ whereas a particle in the \mathbb{Z}_N Laplacian model is completely immobile on an infinite square lattice, at least when N is prime.

⁴⁰More precisely, this is true for an electrically charged particle. Magnetically charged particles come in two flavors: x -lineons, which can move anywhere in the x direction but can only hop by N sites in the y direction, and similarly y -lineons.

[1] C. Chamon, Quantum Glassiness in Strongly Correlated Clean Systems: An Example of Topological Overprotection, *Phys. Rev. Lett.* **94**, 040402 (2005).
[2] J. Haah, Local stabilizer codes in three dimensions without string logical operators, *Phys. Rev. A* **83**, 042330 (2011).
[3] S. Vijay, J. Haah, and L. Fu, Fracton topological order, generalized lattice gauge theory and duality, *Phys. Rev. B* **94**, 235157 (2016).
[4] J. Haah, A degeneracy bound for homogeneous topological order, *SciPost Phys.* **10**, 011 (2021).
[5] R. M. Nandkishore and M. Hermele, Fractons, *Annu. Rev. Condens. Matter Phys.* **10**, 295 (2019).

[6] M. Pretko, X. Chen, and Y. You, Fracton phases of matter, *Int. J. Mod. Phys. A* **35**, 2030003 (2020).
[7] K. T. Grosvenor, C. Hoyos, F. Peña Benitez, and P. Surówka, Space-dependent symmetries and fractons, *Front. Phys.* **9**, 792621 (2022).
[8] T. Brauner, S. A. Hartnoll, P. Kovtun, H. Liu, M. Mezei, A. Nicolis, R. Penco, S.-H. Shao, and D. T. Son, Snowmass white paper: Effective field theories for condensed matter systems, *arXiv:2203.10110*.
[9] J. McGreevy, Generalized symmetries in condensed matter, *arXiv:2204.03045*.

- [10] C. Cordova, T. T. Dumitrescu, K. Intriligator, and S.-H. Shao, Snowmass white paper: Generalized symmetries in quantum field theory and beyond, [arXiv:2205.09545](https://arxiv.org/abs/2205.09545).
- [11] K. Slagle and Y. B. Kim, X-cube model on generic lattices: Fracton phases and geometric order, *Phys. Rev. B* **97**, 165106 (2018).
- [12] W. Shirley, K. Slagle, Z. Wang, and X. Chen, Fracton Models on General Three-Dimensional Manifolds, *Phys. Rev. X* **8**, 031051 (2018).
- [13] W. Shirley, K. Slagle, and X. Chen, Fractional excitations in foliated fracton phases, *Ann. Phys.* **410**, 167922 (2019).
- [14] W. Shirley, K. Slagle, and X. Chen, Foliated fracton order in the checkerboard model, *Phys. Rev. B* **99**, 115123 (2019).
- [15] W. Shirley, K. Slagle, and X. Chen, Foliated fracton order from gauging subsystem symmetries, *SciPost Phys.* **6**, 041 (2019).
- [16] K. Slagle, A. Prem, and M. Pretko, Symmetric tensor gauge theories on curved spaces, *Ann. Phys.* **410**, 167910 (2019).
- [17] K. Slagle, D. Aasen, and D. Williamson, Foliated field theory and string-membrane-net condensation picture of fracton order, *SciPost Phys.* **6**, 043 (2019).
- [18] W. Shirley, K. Slagle, and X. Chen, Twisted foliated fracton phases, *Phys. Rev. B* **102**, 115103 (2020).
- [19] K. Slagle, Foliated Quantum Field Theory of Fracton Order, *Phys. Rev. Lett.* **126**, 101603 (2021).
- [20] P.-S. Hsin and K. Slagle, Comments on foliated gauge theories and dualities in 3+1d, *SciPost Phys.* **11**, 032 (2021).
- [21] H. Geng, S. Kachru, A. Karch, R. Nally, and B. C. Rayhaun, Fractons and exotic symmetries from branes, *Fortschr. Phys.* **69**, 2100133 (2021).
- [22] P. Gorantla, H. T. Lam, and S.-H. Shao, Fractons on graphs and complexity, *Phys. Rev. B* **106**, 195139 (2022).
- [23] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, 2+1d compact Lifshitz theory, tensor gauge theory, and fractons, [arXiv:2209.10030](https://arxiv.org/abs/2209.10030).
- [24] N. Manoj and V. B. Shenoy, Arboreal topological and fracton phases, [arXiv:2109.04259](https://arxiv.org/abs/2109.04259).
- [25] N. Seiberg and S.-H. Shao, Exotic symmetries, duality, and fractons in 2+1-dimensional quantum field theory, *SciPost Phys.* **10**, 027 (2021).
- [26] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, fcc lattice, checkerboards, fractons, and quantum field theory, *Phys. Rev. B* **103**, 205116 (2021).
- [27] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, A modified Villain formulation of fractons and other exotic theories, *J. Math. Phys.* **62**, 102301 (2021).
- [28] N. Seiberg, Field theories with a vector global symmetry, *SciPost Phys.* **8**, 050 (2020).
- [29] N. Seiberg and S.-H. Shao, Exotic $U(1)$ symmetries, duality, and fractons in 3+1-dimensional quantum field theory, *SciPost Phys.* **9**, 046 (2020).
- [30] N. Seiberg and S.-H. Shao, Exotic \mathbb{Z}_N symmetries, duality, and fractons in 3+1-dimensional quantum field theory, *SciPost Phys.* **10**, 003 (2021).
- [31] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, More exotic field theories in 3+1 dimensions, *SciPost Phys.* **9**, 073 (2020).
- [32] T. Rudelius, N. Seiberg, and S.-H. Shao, Fractons with twisted boundary conditions and their symmetries, *Phys. Rev. B* **103**, 195113 (2021).
- [33] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, Low-energy limit of some exotic lattice theories and UV/IR mixing, *Phys. Rev. B* **104**, 235116 (2021).
- [34] F. J. Burnell, T. Devakul, P. Gorantla, H. T. Lam, and S.-H. Shao, Anomaly inflow for subsystem symmetries, *Phys. Rev. B* **106**, 085113 (2022).
- [35] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, Global dipole symmetry, compact Lifshitz theory, tensor gauge theory, and fractons, *Phys. Rev. B* **106**, 045112 (2022).
- [36] M. Pretko, The fracton gauge principle, *Phys. Rev. B* **98**, 115134 (2018).
- [37] D. J. Williamson, Z. Bi, and M. Cheng, Fractonic matter in symmetry-enriched $U(1)$ gauge theory, *Phys. Rev. B* **100**, 125150 (2019).
- [38] C. Castelnovo and C. Chamon, Topological quantum glassiness, *Philos. Mag.* **92**, 304 (2012).
- [39] B. Yoshida, Exotic topological order in fractal spin liquids, *Phys. Rev. B* **88**, 125122 (2013).
- [40] F. Chung, *Spectral Graph Theory*, CBMS Regional Conference Series in Mathematics Vol. 92 (American Mathematical Society, Providence, RI, 1997).
- [41] K. A. Berman, Bicycles and spanning trees, *SIAM J. Algebraic Discrete Methods* **7**, 1 (1986).
- [42] H. J. S. Smith, XV. On systems of linear indeterminate equations and congruences, *Philos. Trans. R. Soc. London* **151**, 293 (1861).
- [43] D. Lorenzini, Smith normal form and Laplacians, *J. Comb. Theory, Ser. B* **98**, 1271 (2008).
- [44] D. Dhar, Self-Organized Critical State of Sandpile Automaton Models, *Phys. Rev. Lett.* **64**, 1613 (1990).
- [45] D. J. Lorenzini, A finite group attached to the Laplacian of a graph, *Discrete Math.* **91**, 277 (1991).
- [46] N. L. Biggs, Chip-firing and the critical group of a graph, *J. Algebraic Combinatorics* **9**, 25 (1999).
- [47] J. Haah, Commuting Pauli Hamiltonians as maps between free modules, *Commun. Math. Phys.* **324**, 351 (2013).
- [48] I. Niven, H. Zuckerman, and H. Montgomery, *An Introduction to the Theory of Numbers*, 5th ed. (Wiley, New York, 1991).
- [49] P. Moree, Artin's primitive root conjecture—A survey, *Integers* **12**, 1305 (2012).
- [50] C. Hooley, On Artin's conjecture, *J. Reine Angew. Math.* **225**, 209 (1967).
- [51] H. Ebisu and B. Han, Anisotropic higher rank \mathbb{Z}_N topological phases on graphs, [arXiv:2209.07987](https://arxiv.org/abs/2209.07987).
- [52] D. S. Dummit and R. M. Foote, *Abstract Algebra*, 3rd ed. (Wiley, New York, 2004).
- [53] D. Eisenbud, *Commutative Algebra: With a View toward Algebraic Geometry*, Graduate Texts in Mathematics Vol. 150 (Springer, New York, 2013).
- [54] D. A. Cox, *Galois Theory*, 2nd ed. (John Wiley & Sons, New York, 2012).
- [55] S. Gao and V. M. Rodrigues, Irreducibility of polynomials modulo p via Newton polytopes, *J. Number Theory* **101**, 32 (2003).
- [56] S. Gao, Absolute irreducibility of polynomials via Newton polytopes, *J. Algebra* **237**, 501 (2001).
- [57] D. Bulmash and M. Barkeshli, The Higgs mechanism in higher-rank symmetric $U(1)$ gauge theories, *Phys. Rev. B* **97**, 235112 (2018).

- [58] H. Ma, M. Hermele, and X. Chen, Fracton topological order from the Higgs and partial-confinement mechanisms of rank-two gauge theory, *Phys. Rev. B* **98**, 035111 (2018).
- [59] Y.-T. Oh, J. Kim, E.-G. Moon, and J. H. Han, Rank-2 toric code in two dimensions, *Phys. Rev. B* **105**, 045128 (2022).
- [60] Y.-T. Oh, J. Kim, and J. H. Han, Effective field theory of dipolar braiding statistics in two dimensions, *Phys. Rev. B* **106**, 155150 (2022).
- [61] S. D. Pace and X.-G. Wen, Position-dependent excitations and UV/IR mixing in the \mathbb{Z}_N rank-2 toric code and its low-energy effective field theory, *Phys. Rev. B* **106**, 045145 (2022).