Erratum: Impact of dark excitons on Förster-type resonant energy transfer between dye molecules and atomically thin semiconductors [Phys. Rev. B 107, 035304 (2023)]

Manuel Katzer D, Sviatoslav Kovalchuk, Kyrylo Greben, Kirill I. Bolotin, Malte Selig, and Andreas Knorr

(Received 6 November 2023; published 17 November 2023)

DOI: 10.1103/PhysRevB.108.199903

In our paper, we gave an incorrect definition for the acronym DCM, which should refer instead to the laser dye molecule 4dicyanomethylene-2-methyl-6-(p(dimethylamino)styryl)-4H-pyran [1] (empirical formula $C_{19}H_{17}N_3O$ [2]). The sketch in Fig. 1 of the paper depicts a simplification of this molecule and is not an accurate display of its structure. The used optical matrix element of the dye molecule in our calculation corresponds to a lifetime of 26 ns [3]. All results and conclusions of our work remain valid.

We are grateful to Professor Hasenstab-Riedel and Professor Haag from FU Berlin for pointing out this error, and to L. Greten and J. Grumm from TU Berlin for the careful reading and assistance in this correction process.

- [2] P. Hammond, Laser dye DCM, its spectral properties, synthesis and comparison with other dyes in the red, Opt. Commun. 29, 331 (1979).
- [3] M. Meyer, J. Mialocq, and M. Rougée, Fluorescence lifetime measurements of the two isomers of the laser dye DCM, Chem. Phys. Lett. 150, 484 (1988).

^[1] S. Bondarev, V. Knyukshto, V. Stepuro, A. Stupak, and A. Turban, Fluorescence and electronic structure of the laser dye dcm in solutions and in polymethylmethacrylate, J. Appl. Spectrosc. **71**, 194 (2004).