

Model of a non-Fermi liquid with power-law resistivity: Strange metal with a not-so-strange originPatrick A. Lee *Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

(Received 25 March 2024; revised 26 August 2024; accepted 27 August 2024; published 13 September 2024)

We construct a model that exhibits resistivity going as a power law in temperature T , as T^α down to the lowest temperature. There is no residual resistivity because we assume the absence of disorder and momentum relaxation is due to umklapp scattering. Our model consists of a quantum spin liquid state with a spinon Fermi surface and a hole Fermi surface made out of doped holes. The key ingredient is a set of singular $2k_F$ modes living on a ring in momentum space. Depending on parameters, α may be unity (strange metal) or even smaller. The model may be applicable to a doped organic compound, which has been found to exhibit linear T resistivity. We conclude that it is possible to obtain strange metal behavior starting with a model that is not so strange.

DOI: [10.1103/PhysRevB.110.L121110](https://doi.org/10.1103/PhysRevB.110.L121110)

Introduction. The notion of strange metals emerged out of transport measurements from the early days of cuprate superconductivity, and it has been applied broadly over the years to instances where linear in temperature resistivity is observed, whether it is low temperature or high temperature, and with or without a finite residual resistivity due to disorder. In this broad sense, strange metal behavior has been seen in many materials and has attracted a great deal of attention from the community, as summarized nicely in recent reviews [1,2]. The strange metal is usually associated with the violation of Landau's theory of Fermi liquid, and it is believed to be driven by strong correlation physics. In this paper, we will restrict the use of the term "strange metal" to low temperatures and where disorder is either absent or does not play a dominant role. The high-temperature linear resistivity anomaly is associated with violations of the Mott-Ioffe-Regel limit and clearly requires a separate set of physical input [2]. On the opposite end, while there are examples where linear T resistivity survives to low temperatures, in many cases the linear regime does not extend above the residual resistivity beyond a value comparable to the residual resistivity itself. Examples of this behavior include the overdoped [3] and electron doped cuprates [4]. While this phenomenon is not understood and is of great interest, the effect of disorder is likely to be strongly relevant, and a different set of explanations may be required. Furthermore, as a matter of principle, Landau's Fermi liquid theory refers to the clean case. In this paper, we will not discuss models where disorder plays a key role [5–8]. We will focus on the situation in which a power-law resistivity extrapolates to a small residual value, so that the power extends over a range much larger than the residual resistivity, and disorder may be considered unimportant. Our goal is to produce a model that produces a power-law resistivity T^α with $\alpha < 2$ which is valid down to zero temperature and which include the linear resistivity as a special case $\alpha = 1$. As explained below (see also [6]), there are barriers towards accomplishing this goal, which may explain the *paucity* of such models. While our model is unlikely to be applicable to cuprate

superconductors, it may find application in a doped organic compound [9].

Why is it difficult to construct such a model? In the absence of disorder, the total momentum is conserved under scattering unless umklapp scattering is allowed. Therefore, in the absence of both disorder and umklapp scattering, the conductivity $\sigma(\omega)$ is proportional to $\delta(\omega)$, which is incompatible with any power-law function $\omega^{-\alpha}$. Hence the coefficient of $\delta(\omega)$ must be zero. This case was considered by Else and Senthil [10], who show that a power-law resistivity requires the divergence of a certain kind of fluctuations in order to kill the δ function. We shall not pursue this route here and instead consider models where umklapp scattering is allowed down to $T = 0$. Such models immediately rule out scattering from critical modes at $q = 0$ such as those from ferromagnetic or nematic order or from emergent gauge fields [6,11]. This leaves soft modes that are associated with a critical point involving ordering at a finite momentum such as antiferromagnets or charge density waves. Such scatterings are limited to "hot spots" on the Fermi surface. Without scatterings which rapidly equilibrate the other momentum states on the Fermi surface, the resistivity is dominated by scattering rates away from the hot spots [5,12]. This problem can be brought under control with the introduction of disorder scattering [5,6,11]. This indeed gives rise to a linear T regime with a coefficient that is independent of the disorder. However, the price one pays is that this regime is limited to an increase of resistivity which is equal to or less than the residual resistivity. Hence the zero disorder limit cannot be taken without losing the linear T regime [5,11].

It turns out that these difficulties can be circumvented in a model of doping holes into a quantum spin liquid with a spinon Fermi surface. The spinon Fermi surface has a singular self-energy and violates the Landau criterion for quasiparticles. On a triangular lattice umklapp scattering is possible if the density of doped holes is large enough. We shall show that in this case the hot spot becomes a hot region on the Fermi surface, which may even cover the entire Fermi surface. So

the bottleneck problem mentioned earlier does not arise, and the power-law resistivity survives to low temperature. We also emphasize that if the power law α is equal to or less than unity, the Landau criterion is violated and quasiparticles as defined by Landau do not exist. Nevertheless, it was shown long ago by Prange and Kadanoff [13] that even without Landau's quasiparticles, a Boltzmann equation that describes transport properties can be derived. Taking advantage of their insight, the resistivity can be calculated in a simple way even in the "non-Fermi-liquid" case. While a quantum spin liquid with a spinon Fermi surface is an exotic state of matter, it is no longer considered very strange. In fact, these state may be realized in certain organic compounds [14] and in monolayer $1T$ -TaSe₂ [15]. In any case, as a matter of principle, it is useful to have an example that can exhibit strange metal behavior based on a model that is not so strange.

The model. We consider a Hubbard model on the triangular lattice with nearest-neighbor hopping t and on-site U . We assume U/t is such that we are on the insulator side of the Mott transition, but not deep inside, so the charge gap is finite but relatively small. $S = 1/2$ local moments are formed on the sites, and we assume that they do not order but form a spin liquid state. We further assume that the electrons have fractionalized into fermionic spinons carrying $S = 1/2$ but no charge, and relativistic bosonic chargons. Both are coupled to emergent U(1) gauge fields [16]. This state was proposed to characterize the organic spin liquids compounds, the ET, and dmit salts [14]. Recently, the ET salt was found to have a spin gap below a phase transition at 6 K, so if there is a Fermi surface, at least part of it is gapped out at low temperatures [17]. We note that with doping the 6 K transition is absent and it is reasonable to assume that the spinon Fermi surface is intact [9]. More recently, the spinon Fermi surface state was also proposed to be realized in monolayer $1T$ -TaSe₂ and $1T$ -TaS₂, and there is evidence of such a state from the appearance of incommensurate modulations at wave vectors given by $2k_F$ expected in a Fermi liquid [15]. Further evidence comes from Kondo screening of adsorbed magnetic impurities [18]. We note that recent density matrix renormalization group (DMRG) calculations found a chiral spin liquid in the vicinity of the Mott transition and not a spinon Fermi surface [19,20]. On the other hand, there is no experimental evidence for time reversal breaking in the two systems mentioned above. So we continue to assume that the Fermi surface state is either realized with additional coupling or is a competing state that may arise with carrier doping.

Now we consider doping this spin liquid with carriers that can be electrons or holes. For concreteness we shall use the hole notation. So far the $1T$ -TaSe₂ and $1T$ -TaS₂ has not been systematically doped, even though significant charge transfer is known to occur from the $1T$ to the $1H$ layer in $4Hb$ -TaS₂ [21]. On the other hand, in the case of the ET salt, it has been possible to introduce Hg chains between the layers, resulting in a doped hole concentration p of 11% [9]. At the mean-field level, the doped hole will occupy the gapped holon band. On the triangular lattice, the band minimum may occur at the Γ point or the K points depending on the sign of t . Here we make a further assumption. We consider a strong attraction between the holon and the spinon so they recombine to form physical holes that carry both charge and spin. These holes




FIG. 1. (a) The dashed circle shows the spinon Fermi surface in the hexagonal Brillouin zone for a triangular lattice. The solid blue circle shows the hole Fermi surface holding p doped holes. The two Fermi surfaces can exchange momentum \mathbf{P} via an umklapp process where \mathbf{P} is the sum of a $2k_F$ vector (dashed orange line) and a reciprocal-lattice vector \mathbf{G} (green line). (b) An expanded view of the region between the two spinon Fermi surfaces shown in (a). \mathbf{P}_0 is the shortest umklapp vector connecting the two spinon Fermi surfaces. The hole Fermi surface is positioned so that its Fermi surface is spanned by \mathbf{P}_0 . An initial hole state (green dot) at angle $-\phi_0$ is scattered to a final hole state (orange dot) at angle $\phi_0 + \phi$ in an umklapp process. The short red arrow denotes the vector $\Delta\mathbf{q}$ [defined above Eq. (2)] which measures the deviation of the collective mode momentum \mathbf{Q} from $2k_F$.

form a Fermi surface containing p holes if the band bottom is located at Γ , or they form two Fermi surfaces containing $p/2$ holes each if the band bottoms are at the zone corner K points. We shall refer to such bands as hole bands. Alternatively, there may be an additional band that happens to be located below the charge gap, and the doped holes enter that band to form conventional hole pockets. There is in fact evidence for such a band in $1T$ -TaSe₂ [22]. The situation is illustrated in Fig. 1. The spinon band is half-filled and is almost circular. For concreteness, the hole band is assumed to be at the Γ point and has an area corresponding to p spinful holes.

Note that the hole band containing p holes violates the Luttinger theorem, which states that the Fermi surfaces should enclose a total of $1 - p$ fermions. This situation has been dubbed FL* by Senthil, Sachdev, and Vojta [23]. They explained that one can get around the nonperturbative proof of Luttinger theorem [24] if the spinon sector is in a gapped topological state with degenerate ground states upon flux insertion in a torus geometry. In our case, the spinon sector is gapless and has massive degeneracy. As a result, the non-perturbative proof that relies on returning to the same ground state after a flux insertion in a torus also fails [25]. While topology does not play a role, we follow [23] and denote this as an FL* state. The properties of the hole pocket can be quite conventional, apart from its small Fermi surface area. In contrast, the spinon Fermi surface is strongly coupled to a U(1) gauge field which the holes do not see. This problem has been well studied and one key result is that the self-energy goes as $\omega^{2/3}$ [27]. The spinon decay rate exceeds ω , which violates Landau's criterion for the existence of quasiparticles. The origin of the strong decay rate is that the gauge field is strongly overdamped, with a propagator that goes as $(|\omega|/q + q^2)^{-1}$. The excitation ω scales as q^3




FIG. 2. A plot of the exponent α which characterizes the resistivity going as T^α vs the exponent σ . The dashed line indicates the breakdown of the linear relation given by Eq. (5) for $\sigma > 5/6$. The Fermi liquid behavior $\alpha = 2$ is recovered for $\sigma < 1/6$, and linear T resistivity is obtained for $\sigma = 2/3$, as indicated by the short-dashed line. σ is defined as the divergent exponent of the vertex function $\gamma(\omega) \propto \omega^{-\sigma}$. As shown in the inset, this divergence is due to repeated exchange of the gauge field (dotted line). Also shown is the diagram for the spinon polarization function Π_{2k_F} . The intermediate spinon lines denote excitations of spinon particle-hole pairs, which are restricted to low energy, hence two factors of the vertex functions appear and contribute the factor $\tilde{\omega}^{-2\sigma}$ in Eq. (1). For details, see Ref. [26].

resulting in a copious amount of low-energy excitations. For a number of years it was believed that this strong coupled problem can be controlled with a $1/N$ expansion, where N is the number of fermion species. The low-energy physics is found to be described by a nontrivial fixed point, which is exactly marginal and described by scaling functions [26,28]. Of particular interest to us is that the polarizability function for the system comprising spinons coupled to gauge fields, $\Pi(\omega, \mathbf{Q})$, is singular for $|\mathbf{Q}|$ near $2k_F$ [26]. Importantly, note that this results in a ring of low energy excitations in momentum space with radius $2k_F$. Unfortunately, upon further scrutiny it was shown by Sung-Sik Lee that the $1/N$ is not controlled even in the large N limit [29]. So the nature of the low energy physics is not known even for large N , let alone the physical case of $N = 2$. Nevertheless, it is still-possible that the low-energy physics remains to be described by a nontrivial and marginal fixed point. There is support for this possibility by studies involving more expansion parameters [30]. We will adopt this point of view and assume a scaling form for $\Pi(\omega, \mathbf{Q})$ which is the same for the large- N expansion as given by Altshuler, Ioffe, and Millis [26], except that we will treat the scaling exponent σ as an unknown parameter. [σ is the exponent that characterizes the divergence of the $2k_F$ vertex function γ due to coupling to the gauge fluctuations as shown in Fig. 2, namely, $\gamma(\omega) \propto |\omega|^{-\sigma}$.] Since $\Pi(\omega, \mathbf{Q})$ is independent of the direction of \mathbf{Q} , we introduce the variable $\Delta\mathbf{q} = \mathbf{Q} - 2k_F\hat{\mathbf{Q}}$ to denote the distance of \mathbf{Q} from the Fermi surface. We write $\Pi(\omega, \mathbf{Q})$ as $\Pi_{2k_F}(\omega, \Delta\mathbf{q})$ and denote the imaginary part of the retarded function after analytic continuation to real frequency Ω as Π'' . We define the scaled variable \tilde{q} as $\tilde{q}\hat{\mathbf{Q}} = \Delta\mathbf{q}/k_F$, together with $\tilde{\omega} = \omega/E_F$. We have (for

details, see Appendix A)

$$\Pi''_{2k_F}(\omega, \Delta\mathbf{q}) \propto \frac{1}{E_F} \tilde{\omega}^{2/3-2\sigma}, \quad \tilde{\omega}^{2/3} > |\tilde{q}|, \quad (1)$$

$$\Pi''_{2k_F}(\omega, \Delta\mathbf{q}) \propto \begin{cases} \frac{1}{E_F} \tilde{\omega} |\tilde{q}|^{-1/2-3\sigma}, & \tilde{\omega}^{2/3} < |\tilde{q}|, \tilde{q} < 0, \\ \frac{1}{E_F} \tilde{\omega}^{5/3} |\tilde{q}|^{-3/2-3\sigma}, & \tilde{\omega}^{2/3} < |\tilde{q}|, \tilde{q} > 0. \end{cases} \quad (2)$$

In these equations, the σ -dependent part comes from the vertex function γ and the rest comes from particle-hole excitations including the self-energy correction. Equation (1) gives the limit $\tilde{\omega}^{2/3} > |\tilde{q}|$ and is given in Ref. [26]. The limit $\tilde{\omega}^{2/3} < |\tilde{q}|$ is given in Eq. (2). The first line is applicable for $\tilde{q} < 0$ or $|\mathbf{Q}| < 2k_F$ and apart from the σ -dependent factor, it is the same as the familiar form for free fermions [31]. The only difference is that the condition of validity is changed from $\tilde{\omega} < |\tilde{q}|$ to $\tilde{\omega}^{2/3} < |\tilde{q}|$. The second line in Eq. (2) gives the case $|\mathbf{Q}| > 2k_F$ which is zero for free fermions because we are outside of the particle-hole continuum. In our case, there is a finite contribution due to a self-energy that goes as $\omega^{2/3}$. Note that Eqs. (1) and (2) satisfy the scaling form $\Pi''_{2k_F} \propto \tilde{\omega}^{2/3-2\sigma} F(\tilde{\omega}/\tilde{q}^{3/2})$ so that $\tilde{\omega}$ scales as $\tilde{q}^{3/2}$. [$F(x)$ goes to 1 for x small.]

We note that while Eqs. (1) and (2) are often derived assuming a circular Fermi surface, it is generally applicable to any Fermi surface shape as long as opposite k points on the Fermi surface have parallel tangents. In this case, k_F is a function of angle θ , and Eqs. (1) and (2) remain valid.

Resistivity. Now we are ready to compute the resistivity of the hole band due to scattering by the soft $2k_F$ mode of the spinon Fermi surface. We note that after shifting by a reciprocal-lattice vector \mathbf{G} , the $2k_F$ vectors are equivalent to a set of vectors \mathbf{P} centered at the M point that connect the Fermi surfaces on neighboring extended Brillouin zones, as shown in Fig. 1(a). If the hole Fermi surface is large enough, these vectors can connect points on the hole Fermi surface and give rise to umklapp scattering, which relaxes the momentum and current. The condition on the size of the hole Fermi surface is the following. With nearest-neighbor hopping the spinon Fermi surface is nearly a circle with radius $k_F \approx 0.375|\mathbf{b}_1|$, where \mathbf{b}_1 is the reciprocal-lattice vector along \hat{x} [15]. The length of the shortest vector that connects the two Fermi surfaces is $0.25|\mathbf{b}_1|$. Let us denote the Fermi momentum of the hole pocket by p_F . The condition for umklapp is that $p_F > 0.125|\mathbf{b}_1|$, which is $1/3$ of k_F for one spinon per unit cell. Hence we conclude that umklapp scattering begins for $p > 1/9$ in the case of a single circular hole pocket centered at Γ . This condition can be relaxed if the Fermi surface deviates from a circle, which is likely the case if the doped holes occupy a separate trivial band. Below we will assume that this condition is satisfied, and we will calculate the hole lifetime due to umklapp scattering. Since the momentum transfer is large and umklapp, the same lifetime will enter the resistivity and all transport phenomena.

Unlike the spinon Fermi surface, the hole Fermi surface is not coupled to the gauge field, and it does not have the anomalous large self-energy and damping rate. The major source of damping at low temperatures comes from the umklapp scattering channel under consideration. If this rate is smaller than linear in T or its frequency Ω , the quasiparticle is well

defined and a Fermi liquid description is valid in the Landau sense, even if the Luttinger theorem is not obeyed. Here I remark that even if the decay rate ends up with a power law smaller than unity and Landau quasiparticles do not exist, it is still possible to treat the transport problem using the Boltzmann equation, as long as the self-energy has only a frequency dependence and no singular momentum dependence, which is the case here. This was shown by Prange and Kadanoff [13], and their idea has been applied to the fermion coupled to the gauge field problem [11,32]. The idea is that at low frequency, the electron spectral function is a sharp peak in momentum space, and a Fermi surface can be defined by the crossing of this sharp peak across p_F . This is familiar in the angle-resolved photoemission spectroscopy (ARPES) literature in that the momentum distribution curve (MDC) can be sharp while the energy distribution curve (EDC) is broad. This allows us to use the Boltzmann equation approach to calculate the resistivity, and the result will remain valid even if the exponent α ends up being less than unity, which will happen in a certain parameter range.

In its simplest form, the solution of the Boltzmann equation is just the computation of the scattering rate $1/\tau$ using Fermi's Golden Rule. As shown in Fig. 1(b), we label the state on the hole Fermi surface by an angle ϕ . For simplicity of exposition, we consider the scattering from an initial state at $-\phi_0$ to a final state at the Fermi surface near ϕ_0 such that they are connected by the shortest umklapp vector, which lies along \hat{x} . Different ϕ_0 's will have longer spanning umklapp wave vectors, but they will have a similar scattering rate with the same power law. The important point is to note that in general a finite region of ϕ_0 values will satisfy this umklapp condition and will have similar umklapp scattering rates. Furthermore, there are six minimal spanning vectors in total which replicate the one shown in Fig. 1(b). Therefore, it is quite likely that these regions of initial states will cover the entire Fermi surface. This situation is totally different from the scattering from a critical mode at a finite momentum, such as that due to antiferromagnetic instability. This results in the so called "hot spots," and the problem is that the rapid relaxation is limited to these hot spots and there is a bottleneck to relax momenta from the rest of the Fermi surface [5,12]. To overcome the bottleneck constraint, one needs to introduce disorder scattering, which gives a finite resistivity at zero temperature, leading to the difficulties described earlier. An important feature of the current model is that the entire Fermi surface or large fragments of it is hot, which allows us to reach the clean limit. The umklapp scattering rate is given by [11]

$$1/\tau = V_0^2 \int_0^\Omega d\omega \int d\phi \Pi_{2k_F}''(\omega, \Delta\mathbf{q}(\phi)), \quad (3)$$

where V_0 is a short-range interaction constant between the holes and the spinons. Since the imaginary part of Π_{2k_F} represents the excitation of a particle-hole pair of spinons, this equation captures the scattering of holes by spinons described in Fig. 1. The integral is over a final state located at $\phi_0 + \phi$ on the Fermi surface, and $\mathbf{Q}(\phi)$ denotes a vector in the direction connecting this point and the center of the spinon Fermi surface circle, which is close to \hat{x} for small ϕ . We are interested in $\mathbf{Q}(\phi)$ near $2k_F$, so the integration over ϕ can be converted to a one-dimensional integral over the length of $\Delta\mathbf{q}$, where $\Delta\mathbf{q}$ is

the deviation of \mathbf{Q} from the $2k_F$ vector as defined earlier. It is easy to see that $|\Delta\mathbf{q}| \approx k_F \phi \cos(\phi_0)$. Hence the integral over ϕ is converted to an integral over $\tilde{q}/\cos(\phi_0)$. We have

$$1/\tau = \frac{V_0^2}{\cos(\phi_0)} \int_0^\Omega d\omega \int_{-\Lambda}^\Lambda d\tilde{q} \Pi_{2k_F}''(\omega, \Delta\mathbf{q}(\phi)), \quad (4)$$

where $\Lambda \approx 1$ is an ultraviolet (UV) cutoff in the \tilde{q} integration. We can divide the \tilde{q} integral into two regions. Region (1) is for $\tilde{\omega}^{2/3} > |\tilde{q}|$ as given by Eq. (1). The \tilde{q} integral gives a factor $\tilde{\omega}^{2/3}$ and we find $1/\tau \propto \Omega^{7/3-2\sigma}$. Region (2) is for $\tilde{\omega}^{2/3} < |\tilde{q}|$ as given by Eq. (2). The contribution from this region is the same as that from region (1), as long as $\sigma > 1/6$. In this case, all the integrals are ultraviolet-convergent and the integral can be obtained by power counting using the scaling property described after Eq. (2). (For details, see Appendix B.) Hence we find the exponent for the scattering rate $1/\tau \propto \Omega^\alpha$ or T^α , where

$$\alpha = 7/3 - 2\sigma, \quad \sigma > 1/6. \quad (5)$$

For $\sigma < 1/6$ the region $q < 0$ (corresponding to the region inside the particle-hole continuum) dominates because the q integral is ultraviolet-divergent and scaling fails. As a result, we find that $\alpha = 2$ and we recover the standard Landau Fermi liquid result. We note that the spinon self-energy is given by ω^α due to scattering by the holes. When $\alpha < 2/3$, the exponent will need to be determined self-consistently. Hence the validity of Eq. (5) is limited to the range $2/3 < \alpha < 7/3$. This final result is summarized in Fig. 2.

We do not know what the value of σ is, but as a matter of principle, this model gives a power-law behavior of the resistivity down to the lowest temperature, which can include the linear T resistivity. In fact, the linear T case $\alpha = 1$ does not play any special role, and there is no obvious restriction that α cannot be smaller than unity.

The effect of disorder can easily be included in the Boltzmann approach by adding a scattering rate $1/\tau_0$, which simply adds a residual resistivity. Since the umklapp scattering already relaxes momentum, disorder does not modify the umklapp rate given by Eq. (3) in an essential way, in contrast with the hot spot models [5,6].

Discussion. We have constructed a model based on doping of a spin liquid with a spinon Fermi surface, forming a hole Fermi surface with an area that corresponds to the doped hole concentration p . As such it violates Luttinger theorem and belongs to the classification FL* [23]. We show that for p large enough, umklapp scattering between the hole Fermi surface and the spinon Fermi surface becomes possible, resulting in a resistivity that goes as T^α , where α is given by Eq. (5) and can vary over a wide range, depending on the strength of the $2k_F$ singularity of the spinon. In principle, α equal to unity or even smaller is possible. In these cases, the Landau criterion for the existence of his quasiparticle is violated, and this state can be called a non-Fermi liquid. Our calculation relies on the formulation of Prange and Kadanoff [13], who showed that quasiparticles in the Landau sense are not necessary to derive a Boltzmann equation to describe transport. Therefore, our result should remain valid for α equal to or less than unity. On the other hand, since the transport properties are based on the Boltzmann equation, other properties such as the




FIG. 3. Diagram for $\Pi_{2k_F}(\omega, \mathbf{Q})$. Solid triangles represent the vertex function $\gamma(\omega, \Delta \mathbf{q})$ shown in the inset of Fig. 2.

Hall constant and magnetoresistance should be conventional. In particular, the magnetoresistance should obey Koehler's rule, which states that the correction to resistivity goes as $(B\tau)^2$. In cuprates, a linear in B magnetoresistance is often associated with a linear in T resistivity. It is unlikely that our model or something similar is relevant to the cuprates. The most promising material candidate is the doped organic system where 11% doping has been achieved, and linear or close to linear in T resistivity, which extends over several times the residual resistivity, has been observed over some range of pressure [9]. The Hall constant is given by the hole doping in this pressure range, consistent with small Fermi pockets of total area p . Interestingly, 11% is on the border of applicability of our model, which requires $p > 1/9$.

We note that the key ingredient of our model is a critical mode that is soft along a line in momentum space. This gives rise to low-energy scattering in finite regions on the Fermi surface, which allows us to circumvent the hot spot problem associated with scattering by a mode that is critical at one momentum. We employ a model with a spinon Fermi surface because that represents a critical state that exists over a range in parameter space. Therefore, the power-law resistivity exists over a range of parameters such as doping or pressure. Furthermore, a circular spinon or hole Fermi surface is not needed, because the singularity of Π_{2k_F} is a consequence of a two-patch model, which relies on the fact that opposite k points in the spinon Fermi surface have parallel tangents.

There have been a lot of discussions concerning the so-called Planckian bound, $\hbar/\tau < k_B T$ [33]. Attempts to support this conjecture with more rigorous derivations have centered on the ability of the system to reach local equilibrium within some relaxation time τ_{eq} . There is no clear connection of τ_{eq} to the transport time [2] so that at present there is no clear connection between transport lifetime and the Planckian bound [2]. Our model is relatively simple and can serve as a counterexample to the Planckian bound. The fact that phonon scattering at high temperature is given by $2\pi\lambda kT$ and can exceed kT has been used to argue against the Planckian bound [4]. Our model has the advantage that it is valid down to zero temperature.

Acknowledgments. I thank Andrey Chubukov and T. Senthil for enlightening discussions, and I acknowledge support by the U.S. Department of Energy, Office of Basic Sciences Grant No. DE-FG02-03ER46076.

Appendix A: The imaginary part of the polarization bubble $\Pi'_{2k_F}(\omega, \Delta \mathbf{q})$. In this Appendix, we derive the expression for the imaginary part of the polarization bubble $\Pi'_{2k_F}(\omega, \Delta \mathbf{q})$ given by Eqs. (1) and (2). The diagram is shown in Fig. 3.

The vertex function is given by Altshuler *et al.* [26] as

$$\gamma(\omega, \Delta \mathbf{q}) \propto \begin{cases} \tilde{\omega}^{-\sigma}, & \tilde{\omega}^{2/3} > |\tilde{q}|, \\ |\tilde{q}|^{-3\sigma/2}, & \tilde{\omega}^{2/3} < |\tilde{q}|, \end{cases} \quad (\text{A1})$$

where a tilde denotes the scaled variables defined above Eq. (1) in the main text. From Fig. 3 we see that $\Pi'_{2k_F}(\omega, \Delta \mathbf{q})$ is given by the product of two factors of the vertex function γ and the bubble Π_0 given the product of two fermion Green functions G ,

$$\Pi_0(\omega, \mathbf{Q}) \propto \int d\nu \int d\mathbf{p} G(\mathbf{p}, \nu) G(\mathbf{p} + \mathbf{Q}, \nu + \omega). \quad (\text{A2})$$

We work at $T = 0$, and ω and ν are Matsubara frequencies. The Green function is given by $G(\mathbf{p}, \nu) = 1/[i\Sigma(\nu) - \xi_p]$, where $\xi_p = p^2/2m - k_F^2/2m$, and the self-energy $\Sigma(\nu) \propto \text{sgn}(\nu)|\nu|^{2/3}$ due to gauge fluctuations and dominates over the usual linear ν term, which has been dropped. We write $\mathbf{Q} = 2\mathbf{k}_F + \mathbf{q}$ and pick the directions of $\mathbf{Q}, \mathbf{k}_F, \mathbf{q}$ to be along the x axis. For small q the important \mathbf{p} is near $-\mathbf{k}_F$ and it is natural to define $\mathbf{p} = -k_F \hat{x} + \mathbf{k}$. It follows that $\xi_p = -v_F k_x + k_y^2/2m$ and $\xi_{p+Q} = v_F(q + k_x) + k_y^2/2m$. Equation (A2) becomes

$$\Pi_0(\omega, \mathbf{Q}) \propto \int d\nu \int dk_x dk_y \frac{1}{i\Sigma(\nu) - k_y^2/2m + v_F k_x} \times \frac{1}{i\Sigma(\nu + \omega) - v_F q - k_y^2/2m - v_F k_x}. \quad (\text{A3})$$

The k_x integral can be done. Note that the poles cannot be both in the upper or lower half-plane. This results in a factor with the θ function $\theta(\text{sgn}(\nu)\text{sgn}(\nu + \omega))$ and

$$\Pi_0(\omega, \mathbf{Q}) \propto \int d\nu \int dk_y \frac{2\pi i \theta(\text{sgn}(\nu)\text{sgn}(\nu + \omega))}{i\Sigma(\nu) + i\Sigma(\nu + \omega) - v_F q - k_y^2/2m}. \quad (\text{A4})$$

The k_y integral can now be done, resulting in

$$\Pi_0(\omega, \mathbf{Q}) \propto \int d\nu \frac{(2\pi i)^2 \theta(\text{sgn}(\nu)) \theta(\text{sgn}(\nu)\text{sgn}(\nu + \omega))}{\sqrt{-v_F q + i\Sigma(\nu) + i\Sigma(\nu + \omega)}}. \quad (\text{A5})$$

Keeping the region in the ν integral that is not canceled by the θ functions, we have for positive ω

$$\Pi_0(\omega, \mathbf{Q}) \propto \int_0^\omega d\nu \frac{1}{\sqrt{-v_F q + i|\nu|^{2/3} + i|\nu + \omega|^{2/3}}}. \quad (\text{A6})$$

Now we can consider various limits. We note that in order to get the imaginary part of Π_0 , we need to analytically continue the Matsubara ω in Eq. (A6) to the real axis. In the equations below, ω is on the real axis. First for $\tilde{\omega}^{2/3} > |\tilde{q}|$, we can drop q in the denominator and obtain, in agreement with [26],

$$\Pi_0(\omega, \mathbf{Q}) \propto e^{-i\pi/4} |\omega|^{2/3}, \quad \tilde{\omega}^{2/3} > |\tilde{q}|. \quad (\text{A7})$$

For $\tilde{\omega}^{2/3} < |\tilde{q}|$ and $q < 0$, i.e., $Q < 2k_F$, we are inside the particle-hole continuum. For the leading contribution, it suffices to keep the $\sqrt{-q}$ term in the denominator, resulting in

$$\text{Im } \Pi_0(\omega, \mathbf{Q}) \propto \omega / \sqrt{|q|}, \quad \tilde{\omega}^{2/3} < |\tilde{q}| \text{ and } q < 0. \quad (\text{A8})$$

This is a well-known result for free fermions. On the other hand, for $q > 0$ we are outside the particle-hole continuum, and the free-fermion polarization bubble has no imaginary part. The presence of the self-energy Σ gives a finite contribution in our case, and we need to keep the next order by expanding the ω -dependent term in the denominator, resulting in

$$\text{Im } \Pi_0(\omega, \mathbf{Q}) \propto \frac{\omega^{5/3}}{|q|^{3/2}}, \quad \tilde{\omega}^{2/3} < |\tilde{q}| \text{ and } q > 0. \quad (\text{A9})$$

Restoring the scaling factors and multiplying by two factors of the vertex function γ given in Eq. (A1), we obtain Eqs. (1) and (2) for $\Pi''_{2k_F}(\omega, \Delta\mathbf{q})$ in the text.

Appendix B: The scattering rate $1/\tau$. We calculate the scattering rate $1/\tau$ using the Fermi Golden Rule, which is appropriate for a solution of the Boltzmann equation. Since momentum scattering is large, there is no difference between the scattering rate and the transport scattering rate. As explained in the text, the final state is parametrized by the angle ϕ shown in Fig. 1. Importantly, for a given ϕ and for a final state on the spinon Fermi surface, the momentum transfer is given by the deviation of \mathbf{Q} from $2\mathbf{k}_F$ where these two vectors are parallel. Therefore, the integral over ϕ can be converted to a *one-dimensional* integral over \tilde{q} as shown in Eq. (4), and we can use Eqs. (1) and (2) for $\Pi''_{2k_F}(\omega, \Delta\mathbf{q})$. We now break up the integral into three regions, corresponding to the different limits given in Eqs. (1) and (2). The first region corresponds to $\tilde{\omega}^{2/3} > |\tilde{q}|$ and the integral takes the form

$$\int_0^\Omega d\omega \int_0^{\omega^{2/3}} dq \omega^{2/3-2\sigma} \propto \Omega^{7/3-2\sigma}. \quad (\text{B1})$$

The second region corresponds to $\tilde{\omega}^{2/3} < |\tilde{q}|$ and $q > 0$, i.e., the region outside the particle-hole continuum in the free-

fermion case. The integral takes the form

$$\int_0^\Omega d\omega \int_{\omega^{2/3}}^\Lambda dq \frac{\omega^{5/3}}{q^{3/2+3\sigma}}. \quad (\text{B2})$$

Note that the q integral is ultraviolet convergent as long as $\sigma > -1/2$, which is the case for a divergent vertex function. The upper cutoff can be extended to infinity, and this integral gives the same result as in Eq. (B1). This is a consequence of the scaling between $\tilde{\omega}^{2/3}$ and $|\tilde{q}|$, and the result can be obtained by power counting. The final region corresponds to $\tilde{\omega}^{2/3} < |\tilde{q}|$ and $q < 0$, i.e., the region inside the particle-hole continuum in the free-fermion case. The integral takes the form

$$\int_0^\Omega d\omega \int_{-\Lambda}^{-\omega^{2/3}} dq \frac{\omega|q|^{-3\sigma}}{\sqrt{|q|}}. \quad (\text{B3})$$

The important difference is that the q integral is ultraviolet-convergent only for $\sigma > 1/6$, in which case the contribution is the same as Eq. (B1) again. For $\sigma < 1/6$ the q integral gives a constant proportional to $\omega\Lambda$, and we recover the standard Fermi liquid result $1/\tau \propto \Omega^2$ and the power counting argument fails. This becomes the dominant contribution, and we conclude that Fermi liquid is restored for $\sigma < 1/6$, as shown in Fig. 2.

Finally, these results can easily be extended for finite temperature by noting that since everything is a power law, T simply scales as Ω if we include the usual Fermi factors, which depend on ω/T . Therefore, we simply replace Ω by T in the finite result for the scattering rate.

-
- [1] P. W. Phillips, N. E. Hussey, and P. Abbamonte, Stranger than metals, *Science* **377**, eabh4273 (2022).
- [2] S. A. Hartnoll and A. P. Mackenzie, Colloquium: Planckian dissipation in metals, *Rev. Mod. Phys.* **94**, 041002 (2022).
- [3] C. Putzke, S. Benhabib, W. Tabis, J. Ayres, Z. Wang, L. Malone, S. Licciardello, J. Lu, T. Kondo, T. Takeuchi *et al.*, Reduced hall carrier density in the overdoped strange metal regime of cuprate superconductors, *Nat. Phys.* **17**, 826 (2021).
- [4] R. L. Greene, P. R. Mandal, N. R. Poniatowski, and T. Sarkar, The strange metal state of the electron-doped cuprates, *Annu. Rev. Condens. Matter Phys.* **11**, 213 (2020).
- [5] A. Rosch, Magnetotransport in nearly antiferromagnetic metals, *Phys. Rev. B* **62**, 4945 (2000).
- [6] D. L. Maslov, V. I. Yudson, and A. V. Chubukov, Resistivity of a non-Galilean-Invariant Fermi liquid near Pomeranchuk quantum criticality, *Phys. Rev. Lett.* **106**, 106403 (2011).
- [7] P. Cha, N. Wentzell, O. Parcollet, A. Georges, and E.-A. Kim, Linear resistivity and Sachdev-Ye-Kitaev (SYK) spin liquid behavior in a quantum critical metal with spin-1/2 fermions, *Proc. Natl. Acad. Sci. USA* **117**, 18341 (2020).
- [8] A. A. Patel, H. Guo, I. Esterlis, and S. Sachdev, Universal theory of strange metals from spatially random interactions, *Science* **381**, 790 (2023).
- [9] Y. Suzuki, K. Wakamatsu, J. Ibuka, H. Oike, T. Fujii, K. Miyagawa, H. Taniguchi, and K. Kanoda, Mott-driven bec-bcs crossover in a doped spin liquid candidate κ -(BEDT-TTF)₄Hg_{2.89}Br₈, *Phys. Rev. X* **12**, 011016 (2022).
- [10] D. V. Else and T. Senthil, Strange metals as Ersatz Fermi liquids, *Phys. Rev. Lett.* **127**, 086601 (2021).
- [11] P. A. Lee, Low-temperature T -linear resistivity due to umklapp scattering from a critical mode, *Phys. Rev. B* **104**, 035140 (2021).
- [12] R. Hlubina and T. M. Rice, Resistivity as a function of temperature for models with hot spots on the Fermi surface, *Phys. Rev. B* **51**, 9253 (1995).
- [13] R. E. Prange and L. P. Kadanoff, Transport theory for electron-phonon interactions in metals, *Phys. Rev.* **134**, A566 (1964).
- [14] Y. Zhou, K. Kanoda, and T.-K. Ng, Quantum spin liquid states, *Rev. Mod. Phys.* **89**, 025003 (2017).
- [15] W. Ruan, Y. Chen, S. Tang, J. Hwang, H.-Z. Tsai, R. L. Lee, M. Wu, H. Ryu, S. Kahn, F. Liou *et al.*, Evidence for quantum spin liquid behaviour in single-layer 1T-TaSe₂ from scanning tunnelling microscopy, *Nat. Phys.* **17**, 1154 (2021).
- [16] S.-S. Lee and P. A. Lee, U (1) gauge theory of the Hubbard model: Spin liquid states and possible application to κ -(BEDT-TTF)₂Cu₂(CN)₃, *Phys. Rev. Lett.* **95**, 036403 (2005).

- [17] B. Miksch, A. Pustogow, M. J. Rahim, A. A. Bardin, K. Kanoda, J. A. Schlueter, R. Hübner, M. Scheffler, and M. Dressel, Gapped magnetic ground state in quantum spin liquid candidate κ -(BEDT-TTF)₂Cu₂(CN)₃, *Science* **372**, 276 (2021).
- [18] Y. Chen, W.-Y. He, W. Ruan, J. Hwang, S. Tang, R. L. Lee, M. Wu, T. Zhu, C. Zhang, H. Ryu *et al.*, Evidence for a spinon Kondo effect in cobalt atoms on single-layer 1T-TaSe₂, *Nat. Phys.* **18**, 1335 (2022).
- [19] A. Szasz, J. Motruk, M. P. Zaletel, and J. E. Moore, Chiral spin liquid phase of the triangular lattice Hubbard model: A density matrix renormalization group study, *Phys. Rev. X* **10**, 021042 (2020).
- [20] B.-B. Chen, Z. Chen, S.-S. Gong, D. N. Sheng, W. Li, and A. Weichselbaum, Quantum spin liquid with emergent chiral order in the triangular-lattice Hubbard model, *Phys. Rev. B* **106**, 094420 (2022).
- [21] A. Kumar Nayak, A. Steinbok, Y. Roet, J. Koo, I. Feldman, A. Almoalem, A. Kanigel, B. Yan, A. Rosch, N. Avraham *et al.*, First-order quantum phase transition in the hybrid metal–Mott insulator transition metal dichalcogenide 4Hb-TaS₂, *Proc. Natl. Acad. Sci. USA* **120**, e2304274120 (2023).
- [22] Y. Chen, W. Ruan, M. Wu, S. Tang, H. Ryu, H.-Z. Tsai, R. L. Lee, S. Kahn, F. Liou, C. Jia *et al.*, Strong correlations and orbital texture in single-layer 1T-TaSe₂, *Nat. Phys.* **16**, 218 (2020).
- [23] T. Senthil, S. Sachdev, and M. Vojta, Fractionalized Fermi liquids, *Phys. Rev. Lett.* **90**, 216403 (2003).
- [24] M. Oshikawa, Topological approach to Luttinger’s theorem and the Fermi surface of a Kondo lattice, *Phys. Rev. Lett.* **84**, 3370 (2000).
- [25] T. Senthil, M. Vojta, and S. Sachdev, Weak magnetism and non-Fermi liquids near heavy-fermion critical points, *Phys. Rev. B* **69**, 035111 (2004).
- [26] B. L. Altshuler, L. B. Ioffe, and A. J. Millis, Low-energy properties of fermions with singular interactions, *Phys. Rev. B* **50**, 14048 (1994).
- [27] P. A. Lee and N. Nagaosa, Gauge theory of the normal state of high- T_c superconductors, *Phys. Rev. B* **46**, 5621 (1992).
- [28] J. Polchinski, Low-energy dynamics of the spinon-gauge system, *Nucl. Phys. B* **422**, 617 (1994).
- [29] S.-S. Lee, Low-energy effective theory of Fermi surface coupled with U(1) gauge field in 2 + 1 dimensions, *Phys. Rev. B* **80**, 165102 (2009).
- [30] D. F. Mross, J. McGreevy, H. Liu, and T. Senthil, Controlled expansion for certain non-Fermi-liquid metals, *Phys. Rev. B* **82**, 045121 (2010).
- [31] S.-S. Zhang, E. Berg, and A. V. Chubukov, Free energy and specific heat near a quantum critical point of a metal, *Phys. Rev. B* **107**, 144507 (2023).
- [32] Y. B. Kim, P. A. Lee, X.-G. Wen, and P. C. E. Stamp, Influence of gauge-field fluctuations on composite fermions near the half-filled state, *Phys. Rev. B* **51**, 10779 (1995).
- [33] J. Zaanen, Why the temperature is high, *Nature (London)* **430**, 512 (2004).