

## ERRATUM

Nuclear Spin-Lattice Relaxation in Pure and Impure Indium. I. Normal State, D. E. MacLaughlin, J. D. Williamson, and J. Butterworth [Phys. Rev. B **4**, 60 (1971)].

- (i) The lowest number on the ordinate of Fig. 1 should be 0.001 instead of 0.00.  
 (ii) Equation (34) should read

$$a_i^{(1)}(t) = -A_{ii}^{(Q)} a_i^{(0)} t - \sum_{j \neq i} A_{ij}^{(Q)} a_j^{(0)} \frac{(e^{(\lambda_i^M - \lambda_j^M)t} - 1)}{(\lambda_i^M - \lambda_j^M)}.$$

- (iii) In Eq. (36), for  $(\lambda_i^M - \lambda_i^M)$  read  $(\lambda_i^M - \lambda_j^M)$ .

(iv) Page 64, column 2, line 9: for "...adequately spanned by the Fourier transition..." read "...adequately spanned by the Fourier transform..."

Throughout the article, solute concentrations given in at. % are erroneous, and should be taken to be in wt. %. The conversion factors for the dilute alloys studied are approximately  $c(\text{at. \%})/c(\text{wt. \%}) = 1.022, 0.562, 0.967,$  and  $0.554$  for Cd, Tl, Sn, and Pb solutes, respectively. No major conclusions are affected by this error, although details of the discussion in Sec. IV are inaccurate.