

# PHYSICAL REVIEW B

## CONDENSED MATTER

THIRD SERIES, VOLUME 52, NUMBER 7

15 AUGUST 1995-I

### BRIEF REPORTS

*Brief Reports are accounts of completed research which, while meeting the usual Physical Review B standards of scientific quality, do not warrant regular articles. A Brief Report may be no longer than four printed pages and must be accompanied by an abstract. The same publication schedule as for regular articles is followed, and page proofs are sent to authors.*

#### Conductivity of a metal with inverse-power-law correlated impurities

M. C. Varriale

*Instituto de Matemática, Universidade Federal do Rio Grande do Sul, Caixa Postal 15040, 91501-970 Porto Alegre, Rio Grande do Sul, Brazil*

Alba Theumann

*Instituto de Física, Universidade Federal do Rio Grande do Sul, Caixa Postal 15051, 91501-970 Porto Alegre, Rio Grande do Sul, Brazil*

(Received 28 March 1995)

We calculate the cooperon for a system of noninteracting electrons in the presence of random potentials with correlations  $W(\mathbf{r}) = W_0\delta(\mathbf{r}) + W_1 r^{-(d+\sigma)}$  in  $d$  dimensions, for arbitrary values  $\sigma$  and  $d$ , to first order in  $W_1$ . Our detailed results confirm the exactitude of the universal form for the cooperon proposed earlier by other authors based on general principles. We find that the system will have short-range (long-range) behavior according to  $\sigma$  being positive (negative).

#### I. INTRODUCTION

The critical behavior of interacting systems depends on general features of the interactions that may be classified in universality classes.<sup>1</sup> One important class is that of short-range (SR) interactions with the Fourier transform expressible as a series in  $q^2$ , while a different class constitutes the long-range (LR) inverse power-law interactions decaying like  $r^{-(d+\sigma)}$ , where  $d$  is the space dimensionality and  $\sigma$  the range parameter, with a Fourier transform in  $q^\sigma$ . Crossover from LR to SR critical behavior occurs when  $\sigma > \sigma_c$ , where the exact value of  $\sigma_c$ , either 2 or  $2 - \eta_{SR}$ , depends on the renormalization procedure.<sup>1,2</sup>

Interesting phenomena occur in systems with random interactions  $V(\mathbf{r})$ . Within a renormalization-group calculation, the only relevant cumulants<sup>3</sup> are the mean and variance

$$M(\mathbf{r}) = \langle V(\mathbf{r}) \rangle, \quad (1)$$
$$W(|\mathbf{r}-\mathbf{r}'|) = \langle V(\mathbf{r})V(\mathbf{r}') \rangle - \langle V(\mathbf{r}) \rangle \langle V(\mathbf{r}') \rangle,$$

which can be of either the SR or the LR type. The case of LR correlations in random spin systems,  $W(r) = r^{-(d+\sigma)}$  in Eq. (1), was studied in Ref. 4. The critical value of  $\sigma$  that separates SR and LR behavior is in this case  $\sigma_c = 0$ .

In a previous work<sup>5</sup> we investigated the Anderson lo-

calization transition in the presence of random potentials with inverse power-law correlations. We followed Ref. 4 to perform a renormalization-group calculation in a double expansion in  $\epsilon = 4 - d$  and  $\sigma$ . We found a fixed point for  $\sigma < 0$  and  $2\sigma < \epsilon < \sigma$  that we interpreted to indicate a LR-induced localization transition at  $d > 4$ , with the corresponding scaling law for the conductivity.

A microscopic calculation of the conductivity was performed in Ref. 6 for the case of scattering by Yukawa-like potentials in two dimensions. The main result in this work was that, in spite of the mathematical intricacies of the problem, the cooperon in Fig. 1 could be cast into a geometric series with the familiar result for SR correlations

$$\Gamma_c(\mathbf{Q}; \omega) = \frac{1}{2\pi\rho_F\tau^2} \frac{1}{-i\omega + Q^2 D_{tr}} \quad (2)$$

for small values of the cooperon momentum  $\mathbf{Q}$ , where  $D_{tr} = k_F^2 \tau_{tr}/d$  for  $d = 2$  and  $\rho_F$  is the density of states at the Fermi energy. All the details of the interaction were absorbed into the elastic and transport lifetimes  $\tau$  and  $\tau_{tr}$ , respectively. In a subsequent publication<sup>7</sup> it was shown that, provided the cooperon exhibits a diffusion pole as in Eq. (2), the conductivity takes the universal scaling form of SR interactions for arbitrary dimensionality  $d$ .

The question still remains, however, whether the valid-




FIG. 1. (a) Series of maximally crossed diagrams for the cooperon  $\Gamma_c[\mathbf{k}, \mathbf{k}', \mathbf{Q}; \omega]$ . A full line of momentum  $\mathbf{k}$  stands for the Green function  $G(\mathbf{k})$  while a broken line stands for the correlation  $W(\mathbf{k})$ . (b) Green function and self-energy.

ity of Eq. (2) is not restricted to potentials belonging to the SR class,<sup>1</sup> such as the Yukawa potential. Following the formal theory in Ref. 8, it can be established that the relation between the retarded  $K^R(\mathbf{k}; \omega + i\delta)$  and causal  $K^C(\mathbf{k}; \epsilon + \omega + i\delta; \epsilon - i\delta)$  density-density correlation functions is given by

$$K^R(\mathbf{k}; \omega + i\delta) \approx \frac{1}{2\pi i} \int_{-\infty}^{+\infty} d\epsilon \frac{\partial n_F(\epsilon)}{\partial \epsilon} \{ \omega K^C(\mathbf{k}; \epsilon + \omega + i\delta; \epsilon - i\delta) - 2\pi i \rho(\epsilon) \}, \quad (3)$$

where  $n_F(\epsilon)$  is the Fermi function and  $\rho(\epsilon)$  the density of states. In the limit  $\mathbf{k} = 0$  we obtain

$$K^R(\mathbf{k} = 0; t_1 - t_2) = \theta(t_1 - t_2) \langle [\rho_{\mathbf{k}=0}(t_1), \rho_{\mathbf{k}=0}(t_2)] \rangle = 0 \quad (4)$$

because the density operator is given by

$$\rho_{\mathbf{k}=0} = \int d\mathbf{p} n(\mathbf{p}) = N, \quad (5)$$

where  $n(\mathbf{p})$  is the particle number operator for momentum  $\mathbf{p}$  and  $N$  is the conserved total number of particles. From Eqs. (4) and Eq. (5), the exact relation<sup>8</sup> follows:

$$K^C(\mathbf{k} = 0; \epsilon + \omega + i\delta; \epsilon - i\delta) = \frac{2\pi i \rho(\epsilon)}{\omega}. \quad (6)$$

The causal density-density correlation function corresponds to the vertex part shown in Fig. 1. Then Eq. (6) demonstrates that  $\Gamma(\mathbf{Q} = 0; \omega)$  in Eq. (2) is exact but the

theory does not provide rigorously for the first correction in  $Q^2$  in the case of an arbitrary potential. The derivation of Eq. (2) is straightforward for a contact<sup>8</sup> potential, but in the Yukawa case<sup>6</sup> the calculations were far from trivial. Hence we consider interesting the calculation of the cooperon in Fig. 1 for random impurities when the correlations  $W(\mathbf{r})$  in Eq. (1) are a combination of a contact potential plus a LR part, with the Fourier transform

$$W(\mathbf{q}) = W_0 + W_1 q^\sigma. \quad (7)$$

This particular form is dictated by a renormalization-group procedure<sup>5</sup> that automatically generates  $W_0$  starting from  $W_1$ .

In Sec. II we present results that are obtained to first-order perturbation theory in  $W_1$  by following the method of Ref. 9 and that confirm the exactitude of Eq. (2) also for the LR correlations in Eq. (7). We discuss the results for the localization part of the conductivity<sup>7</sup> and we obtain, according to the range parameter  $\sigma$  being positive or negative, that the system will exhibit SR or LR behavior, as predicted in Ref. 5.

## II. CALCULATION OF THE COOPERON AND CONCLUSIONS

We consider a gas of noninteracting electrons in  $d$  dimensions, in the presence of random potentials  $V(\mathbf{r})$  with mean and variance defined in Eq. (1). Here we consider  $M = 0$  and  $W(\mathbf{r})$  is defined by its Fourier transform in Eq. (7). We work in units  $\hbar = m = c = 1$ .

The sum of maximally crossed diagrams for the particle-hole vertex part of cooperon shown in Fig. 1 is obtained by solving the Bethe-Salpeter equation<sup>9</sup>

$$\Gamma_c(\mathbf{k}, \mathbf{k}', \mathbf{Q}; \omega) = W(\mathbf{k} - \mathbf{k}') + \int \frac{d\mathbf{p}}{(2\pi)^d} W(\mathbf{k} - \mathbf{p}) G_+(\mathbf{p}) G_-(\mathbf{Q} - \mathbf{p}) \times \Gamma_c(\mathbf{p}, \mathbf{k}', \mathbf{Q}; \omega), \quad (8)$$

where the electron Green function is given by

$$G_\pm(\mathbf{p}) = [\frac{1}{2}(p^2 - k_F^2) - \epsilon_\pm]^{-1} \quad (9)$$

and  $\epsilon_0$  and  $\epsilon_+$  stand for  $-i/2\tau$  and  $\omega + i/2\tau$ , respectively, while we indicate by  $k_F$  the Fermi momentum. The pole in the Green function ensures that all relevant momenta will be at the Fermi surface. Then, in Eq. (8) we have, from Eq. (7),

$$W(\mathbf{k} - \mathbf{k}') = W_0 + W_1 [2k_F \sin(\frac{1}{2}\theta_{kk'})]^\sigma, \quad (10)$$

where  $\theta_{kk'}$  is the angle between the vectors  $\mathbf{k}$  and  $\mathbf{k}'$ . Also we perform the integrals within the standard approximation

$$\int \frac{d\mathbf{p}}{(2\pi)^d} = k_F^{d-2} \int_{-\infty}^{+\infty} d\xi d\Omega_p, \quad (11)$$

where  $\xi = (p^2 - k_F^2)/2$ ,  $d\Omega_p$  is the angular differential on the unit sphere divided by  $(2\pi)^d$ , and  $\int d\Omega_p = \Omega_d = 2^{1-d}/\pi^{d/2} \Gamma(d/2)$ . The inverse lifetime in Eq. (9) is calculated in the Born approximation from the self-

energy in Fig. 1, with the result

$$\frac{1}{\tau} = \frac{1}{\tau_0} \left[ 1 + \frac{W_1}{W_0} f_\tau \right], \quad (12)$$

where

$$\frac{1}{\tau_0} = 2\pi\rho_F W_0, \quad (13)$$

$$f_\tau = 2^{\sigma+d-2} \frac{\Gamma^{(d/2)} \Gamma \left[ \frac{d-1+\sigma}{2} \right]}{\sqrt{\pi} \Gamma \left[ d-1 + \frac{\sigma}{2} \right]} k_F^\sigma, \quad (14)$$

and  $\rho_F = k_F^{d-2} \Omega_d$  is the density of states at the Fermi energy.

The solution of Eq. (8) to first order in  $W_1$  is obtained by following the method of Béal-Monod and Forgacs.<sup>9</sup> We obtain

$$\Gamma_c(\mathbf{k}, \mathbf{k}', \mathbf{Q}; \omega) = \frac{W_0}{1-W_0B} + W_1 \left[ |\mathbf{k}-\mathbf{k}'|^\sigma + 2 \frac{W_0 C}{1-W_0B} + \frac{W_0^2 E}{(1-W_0B)^2} \right], \quad (15)$$

where

$$\begin{aligned} B &= \int \frac{d\mathbf{p}}{(2\pi)^d} G_+(\mathbf{p}) G_-(\mathbf{Q}-\mathbf{p}), \\ C &= \int \frac{d\mathbf{p}}{(2\pi)^d} G_+(\mathbf{p}) G_-(\mathbf{p}) [2k_F \sin(\frac{1}{2}\theta_{pk})]^\sigma, \\ E &= \int \frac{d\mathbf{p}}{(2\pi)^d} \frac{d\mathbf{q}}{(2\pi)^d} G_+(\mathbf{p}) G_-(\mathbf{Q}-\mathbf{p}) \\ &\quad \times [2k_F \sin(\frac{1}{2}\theta_{pq})]^\sigma G_+(\mathbf{q}) G_-(\mathbf{Q}-\mathbf{q}). \end{aligned} \quad (16)$$

As we are looking for the singular contributions to  $\Gamma_c$  when  $Q \rightarrow 0$ , we must expand  $B$  and  $E$  in Eq. (16) to  $O(Q^2)$ . Also the expressions for  $C$  and  $E$  should be evaluated for  $W_1=0$ . The results for  $B$  and  $C$  are straightforward<sup>10</sup> and read

$$B = 2\pi\rho_F \tau \left[ 1 + i\omega\tau - Q^2 \frac{k_F^2 \tau^2}{d} \right], \quad (17)$$

$$C = 2\pi\rho_F \tau \omega f_\tau. \quad (18)$$

The expression for  $E$  is more complicated

$$E = (2\pi\rho_F \tau_0)^2 \{ f_\tau [1 - 2Q^2 \tau_0 D_0] - Q^2 k_F^2 \tau_0^2 2^\sigma k_F^\sigma J \}, \quad (19)$$

where  $D_0 = k_F^2 \tau_0 / d$  is the diffusion coefficient and

$$J = \frac{1}{(\Omega_d)^2} \int d\Omega_p \int d\Omega_q \cos\theta_{pQ} \cos\theta_{qQ} [\sin(\frac{1}{2}\theta_{pq})]^\sigma. \quad (20)$$

We introduce  $d$ -dimensional polar coordinates in momentum space

$$\begin{aligned} q_i &= q \cos\theta_{i-1} \prod_{j=i}^{d-1} \sin\theta_j, \quad i=1, \dots, d-1 \\ q_d &= q \cos\theta_{d-1}, \quad \theta_0=0, \end{aligned} \quad (21)$$

with Jacobian  $q^{d-1} \prod_{j=1}^{d-1} (\sin\theta_j)^{j-1}$ . Then the angle  $\theta_{qQ}$  between two vectors  $\mathbf{q}(\theta_1, \dots, \theta_{d-1})$  and  $\mathbf{Q}(\varphi_1, \dots, \varphi_{d-1})$  is given by

$$\cos\theta_{qQ} = \frac{\mathbf{q} \cdot \mathbf{Q}}{qQ} = \sum_{i=1}^{d-1} \cos\theta_i \cos\varphi_i \prod_{j=i}^{d-1} \sin\theta_j \sin\varphi_j. \quad (22)$$

To perform the integral over  $d\Omega_q$  in Eq. (19), we set the  $d$  axis along  $\mathbf{p}$ ; then  $\theta_{pq} = \theta_{d-1}$  and  $\theta_{pQ} = \varphi_{d-1}$  and we obtain<sup>10</sup>

$$J = \frac{1}{(\Omega_d)^2} \int d\Omega_p \cos\theta_{pQ} K(\theta_{pQ}), \quad (23)$$

where, from Eq. (22),

$$\begin{aligned} K(\theta_{pQ}) &= \frac{1}{(2\pi)^d} \int \prod_{j=1}^{d-1} [d\theta_j (\sin\theta_j)^{j-1}] \\ &\quad \times \cos\theta_{qQ} \left[ \sin \left[ \frac{\theta_{d-1}}{2} \right] \right]^\sigma \\ &= -\frac{\Omega_{d-1}}{2\pi} 2^{d-3} \sigma \frac{\Gamma \left[ \frac{d+\sigma-1}{2} \right] \Gamma \left[ \frac{d-1}{2} \right]}{\Gamma \left( d + \frac{\sigma}{2} \right)} \\ &\quad \times \cos\varphi_{d-1} \end{aligned} \quad (24)$$

because only the term with  $i=d-1$  in Eq. (22) gives a nonzero contribution to the integral in Eq. (24). It follows by introducing Eq. (24) into Eq. (23), where now  $d\Omega_p = 1/(2\pi)^d \prod_{j=1}^{d-1} (\sin\varphi_j)^{j-1} d\varphi_j$ , that

$$J = -\frac{2^{d-3}}{d} \sigma \frac{\Gamma \left[ \frac{d+\sigma-1}{2} \right] \Gamma \left[ \frac{d}{2} \right]}{\sqrt{\pi} \Gamma \left[ d + \frac{\sigma}{2} \right]} \quad (25)$$

and, from Eqs. (25) and (19),

$$E = (2\pi\rho_F \tau_0)^2 f_\tau \left[ 1 - Q^2 \tau_0 D_0 \frac{4(d-1)+\sigma}{2(d-1)+\sigma} \right]. \quad (26)$$

Introducing Eqs. (17), (18), and (26) into Eq. (15) and expanding the terms originating in  $(1-W_0B)^{-1}$  to first order in  $W_1$ , we obtain the singular part

$$\begin{aligned} [\Gamma_c(\mathbf{Q})]_{\omega=0} &= \frac{1}{2\pi\rho_F} \frac{1}{\tau_0^2 D_0 Q^2} \\ &\quad \times \left[ 1 + \frac{W_1}{W_0} f_\tau 2 \frac{3(d-1)+2\sigma}{2(d-1)+\sigma} \right], \\ &\quad Q \rightarrow 0. \end{aligned} \quad (27)$$

In Ref. 7, it was proposed, from first-principles arguments based on the Ward identity, that the universal form for the cooperon should be

$$\Gamma_c(\mathbf{Q}) = \frac{1}{2\pi\rho_F \tau^2} \frac{1}{DQ^2}, \quad Q \rightarrow 0, \quad (28)$$

where

$$D = D_0 \frac{\tau_{tr}}{\tau_0} \quad (29)$$

and  $\tau_{tr}$  is the transport<sup>7</sup> lifetime, which in our case gives

$$\frac{\tau_0}{\tau_{tr}} = \frac{\tau_0}{\tau} + \frac{W_1}{W_0} f_\tau \frac{\sigma}{2(d-1)+\sigma}, \quad (30)$$

with  $\tau$  and  $f_\tau$  defined in Eqs. (12) and (14). The expansion of Eq. (28) to first order in  $W_1$  yields exactly the result obtained in Eq. (27); hence we conclude that our calculation provides a rigorous proof of the validity of Eq. (28) also for inverse power-law correlations.

The calculation of the conductivity now proceeds as indicated in Ref. 7, where it was shown rigorously that *all* the relevant contributions combine into the following expression, which we quote for completeness:

$$\sigma_{dc} = \sigma_0 \frac{\tau_r}{\tau_0} - \frac{\Omega_d}{2\pi} \int_{L^{-1}}^{Q_{max}} Q^{d-3} dQ = \sigma_{Drude} + \sigma_{Loc}, \quad (31)$$

where  $L^{-1}=0$  for  $d > 2$ ,  $\sigma_0 = e^2 \rho_F D_0$ , and  $\sigma_{Drude}$  ( $\sigma_{Loc}$ ) indicates the first (second) term in Eq. (31), while

$$Q_{max} = \begin{cases} \frac{1}{k_F \tau} & \text{if } \frac{1}{\tau} < \frac{1}{\tau_{tr}} \\ \frac{1}{k_F \sqrt{\tau \tau_{tr}}} & \text{if } \frac{1}{\tau} > \frac{1}{\tau_{tr}} \end{cases}. \quad (32a) \quad (32b)$$

We can see, from the last term in Eq. (31), that the localization contribution to the conductivity depends on the potential *only* through  $Q_{max}$ : if  $\sigma > 0$  then we are in the case (32a) and  $\sigma_{Loc}$  depends solely on the elastic lifetime  $\tau$ , *just as for a contact potential*, while for  $\sigma < 0$  we are in the case (32b) and  $\sigma_{Loc}$  exhibits a long-range behavior through  $\tau_{tr}$ . These results confirm the renormalization-group prediction in Ref. 5 that the localization transition will have a LR or SR character according to the range parameter  $\sigma$  being negative (LR) or positive (SR). We consider these results to be valid for all values of  $d$  and  $\sigma$  that ensure well defined  $\Gamma$  functions by analytic continuation.

#### ACKNOWLEDGMENTS

This work was partially supported by Financiadora de Estudos e Projetos and Conselho Nacional de Desenvolvimento Científico e Tecnológico.

<sup>1</sup>M. E. Fisher, Rev. Mod. Phys. **46**, 597 (1974); M. E. Fisher, S. K. Ma, and B. G. Nickel, Phys. Rev. Lett. **29**, 917 (1972); M. A. Gusmão and W. K. Theumann, Phys. Rev. B **28**, 6545 (1983).

<sup>2</sup>J. Sak, Phys. Rev. B **8**, 281 (1973).

<sup>3</sup>T. C. Lubensky, Phys. Rev. B **11**, 3573 (1975).

<sup>4</sup>A. Weinrib and B. I. Halperin, Phys. Rev. B **27**, 413 (1983).

<sup>5</sup>M. C. Varriale and A. Theumann, J. Phys. A **23**, L719 (1990).

<sup>6</sup>M. T. Béal-Monod, A. Theumann, and G. Forgacs, Phys. Rev.

B **46**, 15 726 (1992).

<sup>7</sup>M. T. Béal-Monod, Phys. Rev. B **47**, 3495 (1993).

<sup>8</sup>S. V. Maleev and B. T. Toperverg, Zh. Eksp. Teor. Fiz. **69**, 1440 (1975) [Sov. Phys. JETP **42**, 734 (1976)].

<sup>9</sup>M. T. Béal-Monod and G. Forgacs, Phys. Rev. B **37**, 6646 (1988).

<sup>10</sup>I. S. Gradshteyn and I. M. Ryzhik, in *Table of Integrals, Series and Products*, edited by A. Jeffrey (Academic, New York, 1980).