X-ray magnetic circular dichroism at the iron *K* edge in rare-earth-transition-metal intermetallics: Experimental probe of the rare-earth magnetic moment

J. Chaboy

Instituto de Ciencia de Materiales de Aragón, 50009 Zaragoza, Spain

H. Maruyama

Department of Physics, Faculty of Science, Okayama University, Okayama, Japan

L. M. García and J. Bartolomé Instituto de Ciencia de Materiales de Aragón, 50009 Zaragoza, Spain

K. Kobayashi and N. Kawamura Department of Physics, Faculty of Science, Okayama University, Okayama, Japan

A. Marcelli

INFN, Laboratori Nazionali di Frascati, C.P. 13, 00044 Frascati, Italy

L. Bozukov

Department Solid State Physics, Faculty of Physics, Sofia University, 1126 Sofia, Bulgaria (Received 16 September 1996)

We present a systematic x-ray magnetic-circular-dichroism (XMCD) study performed at the Fe *K* edge on the R_2 Fe₁₄B series (*R*=rare earth and Y). The magnetic XMCD signal has been identified as due to the addition of two components associated with the magnetic contributions from the iron and the rare-earth sublattices. The contribution of the rare-earth sublattice has been extracted from the total XMCD signal. The integrated XMCD signal of the *R* component is proven to be directly correlated with the *R* magnetic moment. [S0163-1829(97)50202-5]

Since the discovery of Nd₂Fe₁₄B,¹ worldwide efforts have been conducted to find novel materials among the ironrich ternary compounds based on rare earths (R) showing high hard-magnet performance. Moreover, the interstitial doping with light elements such as hydrogen or nitrogen has been profusely developed as to become a new route to improve the permanent-magnet technological performance of previously well-known *R*-Fe compounds.

Previous x-ray absorption spectroscopy investigations have addressed the existence of strong electronic effects induced upon light atoms charging.² They conclude that the interplay between electronic charge transfer to the conduction bands and the change of the R(5d)-Fe(3d) hybridization play a major role in governing the modification of the electronic and magnetic properties of these systems. According to Campbell's model, the R-Fe interaction is determined by the 5d-3d hybridization through intra-atomic 4f-5d exchange interactions and direct interatomic 3d-5d spin-spin interactions, so that the R and Fe magnetic moments are ferromagnetically (antiferromagnetically) coupled for light (heavy) rare earths.³ Therefore, it is of paramount interest to characterize magnetically the R(5d) states, which tune the magnetic properties of the *R*-Fe intermetallic compounds, to obtain a further verification of these results.

This information can be gained by means of the x-ray magnetic-circular-dichroism (XMCD) technique. Indeed, one of the most striking capabilities of XMCD applied to R-Fe intermetallics lies in the possibility of directly probing

the spin polarization of the 5d empty states of the rare-earth $L_{2,3}$ edges. Moreover, it offers, in principle, the capability to separate the spin and orbital contributions to the magnetic moments by use of theoretical sum rules derived within an atomic framework.⁴ This model is able to describe also in a rather satisfactory way the XMCD signal coming from localized electronic states [Fe(3d) $L_{2,3}$ edges, $R(4f) M_{4,5}$ edges]. However, recent works have shown the limitations of these sum rules,⁵ so that their validity when applied to $L_{2,3}$ -edge spectra of rare-earth metals is strongly questioned. In addition, Wang *et al.* have shown that the conventional two-step description of the XMCD process fails to account for the R $L_{2,3}$ -edge spectra because it neglects the spin dependence of the transition matrix elements induced by the 5d-4fexchange.⁶ Indeed, systematic studies of the spin-dependent L_{23} absorption in *R*-Fe intermetallics indicate that the sign of the L_{2.3} XMCD signals cannot be directly correlated to the mutual orientation of the rare-earth and iron magnetic moments.^{7,8} In the case of the iron K edge, where the 4pempty states of Fe are probed, the interpretation of the XMCD spectra is still far from being solved. The K-Fe XMCD profile has been described by taking into account the spin splitting of the final *p*-projected states at the Fe site and the hybridization of the 4p and 3d states⁹ so that in some cases the behavior of the XMCD amplitude has been shown to be correlated to variations in the local magnetization.^{9,10} However, to date the relationship between the Fe K-edge XMCD signals and the local magnetic moments is not welldefined.

© 1996 The American Physical Society

In this paper we report a systematic Fe *K*-edge XMCD study at the *K* edge on R_2 Fe₁₄B systems. Our results show that the study of the Fe *K*-edge XMCD signal yields an experimental way to overcome the limitations of the $L_{2,3}$ -edge analysis to characterize the *R* 5*d*-states. Our study shows how the Fe *K*-edge XMCD signal is due to the addition of magnetic contributions from both the iron and the rare-earth sublattices, so that they can be isolated and correlated to the iron and rare-earth magnetic moments.

XMCD experiments have been performed at the iron K edge in R_2 Fe₁₄B (R = Y, La, Ce, Pr, Nd, Sm, Gd, Tb, Dy, Er, Ho, Tm, Yb, and Lu). Several polycrystalline samples were measured in different experimental runs at the Photon Factory (PF) synchrotron radiation facility of the National Laboratory for High Energy Physics (KEK) in Tsukuba. The magnetic XMCD experiments were performed at room temperature in the transmission mode using left circularly polarized x rays on beamline 28B at PF-KEK. The PF storage ring was operated with a positron beam energy of 2.5 GeV and a maximum stored current of 360 mA. The x-ray radiation was monochromatized using a Si(220) doublecrystal monochromator, being the degree of circular polarization after monochromatization $P_c \sim 0.35$.¹¹ The XMCD spectra were recorded by reversing the sample magnetization for a fixed polarization of the incoming radiation. In our experimental setup a magnetic field of 0.6 T was applied parallel to the plane of the sample at 45° to the incident beam and reversed twice for each energy value.

The spin-dependent absorption coefficient has been obtained as the difference of the absorption coefficient $\mu_c = (\mu^- - \mu^+)$ for antiparallel, μ^- , and parallel, μ^+ , orientation of the photon helicity and the magnetic field applied to the sample. The spectra were normalized to the averaged absorption coefficient at high energy, μ_0 , in order to eliminate the dependence of the absorption on the sample thickness, so that $\mu_c(E)/\mu_0 = [\mu^-(E) - \mu^+(E)]/\mu_0$ corresponds to the dimensionless spin-dependent absorption coefficient. The origin of the energy scale was chosen at the inflection point of the absorption edge.

In Fig. 1 we show the normalized Fe K-edge XMCD signals for the measured R_2 Fe₁₄B compounds. In the case of the nonmagnetic rare earths (La, Ce) and Y, the XMCD spectra exhibit a characteristic two-peak structure at the absorption edge, similar to that of pure iron. The positive peak is very narrow, of about 4 eV, whereas the negative dip at higher energy covers an energy width of about 12 eV. In the case of the light rare-earth compounds (R = Pr, Nd, and Sm), the XMCD signals exhibit a strong modification with respect to the nonmagnetic R based compounds: (i) the negative dip gets narrower (~8.5 eV) and its intensity becomes ~2.5 times higher; (ii) an additional positive ($\sim 8 \text{ eV}$ wide) contribution arises at the high-energy side of the negative dip. This behavior is not exhibited by the heavy rare-earth compounds: the XMCD signal of Gd₂Fe₁₄B resembles that of Fe metal or R_2 Fe₁₄B compounds in which R is a nonmagnetic rare earth, see Fig. 1, except for a small positive contribution arising just in the center of the negative dip. The intensity of this feature is found to increase across the series from Gd to Ho, for which the maximum intensity is reached, followed by a progressive decrease till near disappearance for Yb. For $Lu_2Fe_{14}B$ the normal behavior for nonmagnetic R is re-




FIG. 1. Normalized x-ray magnetic circular dichroism spectra at the Fe K-edge for the R_2 Fe₁₄B compounds (R=rare earth and Y). The scale (×10³) is referred to the R=Y (left panel) and R=Gd (right panel) case. For sake of clarity, the normalized Fe K-edge XAS spectrum for Y₂Fe₁₄B is also shown.

gained. This result is an indication of the influence of the particular rare-earth magnetic atom on the *K*-Fe XMCD profile of each compound. The observed experimental behavior across the series indicates that this influence strongly depends on the magnetic state of the rare earth, i.e., nonmagnetic *R*, light *R* ($\mathbf{J}=\mathbf{L}-\mathbf{S}$ coupling), and heavy *R* ($\mathbf{J}=\mathbf{L}+\mathbf{S}$ coupling).

In order to obtain a deeper insight on the origin of such behavior we have performed a further analysis in the framework of a two-magnetic sublattice model. In this model, it is assumed that: (i) the total magnetization of the R_2 Fe₁₄B compounds is the addition of the magnetization of the iron and the rare-earth sublattices, i.e., $M_T = M_R + M_{Fe}$; and (ii) the contribution of the Fe sublattice to the magnetization, $M_{\rm Fe}$, is identical to that of Y₂Fe₁₄B. For each compound in the lanthanide series, the subtraction of this contribution to the magnetization allows one to sort out the magnetic contribution coming from the rare-earth sublattice. In identical manner, in our model we assume the hypothesis that the total Fe K-edge XMCD signal results from the addition of two components, associated with the magnetic contribution from the Fe and R sublattices, respectively. To analyze our data under this model, we have subtracted first the Fe K-edge XMCD signal of Y_2 Fe₁₄B from the Fe K-edge XMCD spectra of the R₂Fe₁₄B series as shown in Fig. 2, then, the remaining signal in the XMCD spectra is assigned to the rareearth sublattice magnetic contribution. The validity of such an assumption is supported by the fact that in the case of the




FIG. 2. Normalized x-ray magnetic circular dichroism spectra at the Fe *K*-edge for the R_2 Fe₁₄B compounds (R=rare earth and Y), after the subtraction of the Y₂Fe₁₄B XMCD signal. The scale (×10³) is referred to the R=Y (left panel) and R=Gd (right panel) case. For sake of clarity, the normalized Fe *K*-edge XAS spectrum for Y₂Fe₁₄B is also shown.

nonmagnetic *R*-based compounds, (La, Ce, and Lu), the signals obtained by substraction of the Y₂Fe₁₄B signal are zero within the signal-to-noise ratio, i.e., the Fe 4*p* moment is nearly constant in the series. Indeed, despite that under substitution of yttrium by a magnetic lanthanide the mean magnetic contribution of iron to the magnetization is increased up to 8%, Mössbauer effect measurements on ⁵⁷Fe in R_2 Fe₁₄B compounds suggest that the electronic configuration of iron atoms is nearly the same for all these compounds, and that the iron 3*d* shell dominates the contribution due to the conduction electron polarization to the Fe hyperfine fields.¹²

After applying the subtraction procedure the remaining signal for all the investigated light-*R* compounds (R=Pr, Nd, and Sm) exhibits a negative peak centered at ($\sim 7 \text{ eV}$) above E_0 with a width that ranges between 6.5 and 9 eV, and a positive peak at ~ 13 eV above the edge, its width ranging between 7.4 and 8.2 eV. In the case of the heavy rare-earths compounds the negative peak is strongly depressed, the main feature on the signal being a positive peak located at about 7 eV above the edge, with a width that varies from 5.7 to 7 eV across the series.

Within a one-electron framework and by introducing the difference of the density of empty states, $\Delta(E)$, the XMCD at the Fe *K* edge is proportional to $\Delta\rho(E) = \rho^{\downarrow}(E) - \rho^{\uparrow}(E)$, where \downarrow (\uparrow) holds for the majority (minority) spin density.^{13,14} In the case of iron metal, the XMCD profile has

been accounted for by considering the spin splitting of the final *p*-projected states at the Fe site and the hybridization of the 4*p* and 3*d* states, which modulates the weight of the spin-dependent absorption cross-section for transitions towards majority (minority) 4*p* states.¹⁴ In the R_2 Fe₁₄B compounds, the Fe sublattice (3*d* \downarrow , majority) determines the sign of the magnetization. In the *R*-Fe intermetallic compounds a strong hybridization between the rare earth 5*d* and Fe-3*d* orbitals is present, so one can expect an additional influence of the *R* magnetic state on the Fe *K*-edge XMCD signal. The subtraction of the Y₂Fe₁₄B Fe *K*-edge XMCD signal, (Fig. 2), would lead to isolation of the *R* contribution to the Fe *K*-edge XMCD.

In this way, the two-peak profile of the R contribution found in the case of the light-R compounds (see Fig. 2) can be associated with the exchange splitting of the 5d bands which occurs when there is a nonzero 4f moment at the R site, because of the intra-atomic 4f-5d overlap. This raises the energy of the *R*-minority 5d band of the *R* sublattice with respect to that of its majority band. Hence, the negative sign of the XMCD signal indicates that the spin of the 5d band just above the Fermi level is antiparallel $(5d^{\uparrow})$ to that of the Fe atoms $(3d^{\downarrow})$. So, the experimental XMCD signal identifies the dominant 5d levels close to the Fermi level as those of the minority-spin band, while the change of sign in the Rcontribution to the XMCD signal for increasing energy implies that above a certain energy the density of 5d-majority band becomes dominant. This result is in agreement with previous electronic structure calculations showing that the highly localized 4f peak is found to be located near the Fermi level, and that the center of the $5d^{\downarrow}$ ($5d^{\uparrow}$) band is located at about 4.7 (6.7) eV above E_F .¹⁵ On the other hand, when a heavy-R (Gd) is involved, the occupied 4f level lies below the Fermi level and the empty 4f orbitals lies at about 3.1 eV above the Fermi level.¹⁶ The XMCD data, showing a prominent positive peak only indicating that the dominant 5dempty states correspond to the majority 5d band, are in agreement to these electronic calculations.

The above results indicate unambiguously that the extracted contribution to the Fe K-edge XMCD signal shown in Fig. 2 reflects the magnetic state of the R atoms in these systems. In a simple model we propose that the Fe K-edge XMCD signals are directly related to the Fe(4p,3d)-R(5d)hybridized band, and that the extracted signals resemble the magnetic state of the rare earth through the splitting of the 5dcomponent due to the exchange interaction with the 4f magnetic moments. If so, the extracted signals should be related to the magnitude of the R magnetic moments. To verify this point we present in Fig. 3 the comparison between the rareearth 4f magnetic moments according to Russel-Saunders coupling and the absolute value of Fe K-edge integrated XMCD signal for the R_2 Fe₁₄B compounds after subtracting the Y₂Fe₁₄B spectrum. The maximum integrated XMCD signal, corresponding to the Dy-based compound, has been scaled to the Dy magnetic moment, and the proportionality factor found was then applied to the whole XMCD signals. The remarkable proportionality between the integrated *R*-component signal and $g_J J$ supports the above hypothesis and thus the validity of our model of additive contributions to the XMCD. Moreover, the *R*-component integrated signal changes sign from the light-R compounds (negative) to the heavy-R compounds (positive). This result agrees with the




FIG. 3. Comparison between the rare-earth 4f magnetic moments, gJ, according to Russel-Saunders coupling (\Box) and the Fe *K*-edge integrated XMCD signal for the R_2 Fe₁₄B compounds, (\diamond), after subtracting the Y₂Fe₁₄B spectrum (see text for details). The values of **S** (solid line) and **L** (dashed line) for the ground state of rare-earth ions along the lanthanide series are also shown, where the sign refers to the spin of the Fe atoms, according to Campbell's model (Ref. 3).

known fact that the R and Fe magnetic moments are parallel(antiparallel) in the light-R (heavy-R) R-Fe intermetallic compounds. We conclude that the extracted signal reflects both the direction and magnitude of the rare-earth magnetic moment. This result is rather surprising in view of the expected dependence of the *R*-*M* interaction on the 5*d* spin value, according to the current models for R(4f-5d) spin-spin coupling.¹⁷ Our results point towards the existence of non-negligible orbital contribution of the 5*d* magnetic moment, arising from the overlap of the 5*d* and 4*f* shells, that is in agreement with the model proposed by Jo and Imada to account for the electron distribution in the 5*d* state due to the 4*f*-5*d* exchange interaction.¹⁸ A similar conclusion has been reached from the correlation found between the branching ratio of the XMCD effect at $L_{2,3}$ edges and the orbital angular momentum of the rare earth (L_z^{4f}) in *R*Fe₂ Laves phases.¹⁹

Our study identifies the influence of the rare-earth magnetic state into the Fe *K*-edge XMCD signals in R_2 Fe₁₄B intermetallic compounds. This signal results from the addition of two components, each one being due to the magnetic contribution of both the iron and the rare-earth sublattices. The contribution of the *R* sublattice to the XMCD signal has been extracted yielding a direct correlation to the *R* magnetic moment. Therefore we provide an approach to characterize magnetically the R(5d) states and thus, the *R*-Fe interaction, in *R*-Fe intermetallics materials.

We are indebted to T. Iwazumi for the experimental support at KEK. This work was partially supported by INFN-CICYT agreement and Spanish DGICYT MAT93-0240C04 and MAT96-0448 grants. This work has been performed with the approval of the Photon Factory Program Advisory Committee (Proposal No. 94-G161).

- ¹J.J. Croat, J.F. Herbst, R.W. Lee, and F.E. Pinkerton, Appl. Phys. Lett. **44**, 148 (1984); M. Sagawa, S. Fujimura, N. Togawa, H. Yamamoto, and Y. Matsuura, J. Appl. Phys. **55**, 2083 (1984).
- ²J. Chaboy, A. Marcelli, and L. Bozukov, J. Phys., Condens. Matter**7**, 8197 (1995); J. Chaboy, A. Marcelli, L. Bozukov, F. Baudelet, E. Dartyge, A. Fontaine, and S. Pizzini, Phys. Rev. B **51**, 9005 (1995).
- ³I.A. Campbell, J. Phys. F 2, L47 (1972).
- ⁴B.T. Thole, P. Carra, F. Sette, and G. van der Laan, Phys. Rev. Lett. **68**, 1943 (1992); P. Carra, B.T. Thole, M. Altarelli, and X. Wang, *ibid.* **70**, 694 (1993).
- ⁵R. Wu, D.S. Wang, and A.J. Freeman, Phys. Rev. Lett. **71**, 3581 (1993); R. Wu and A.J. Freeman, *ibid.* **73**, 1994 (1994); J. Stöhr and H. König, *ibid.* **75**, 3748 (1995); W.L. O'Brien and B.P. Tonner, Phys. Rev. B **50**, 12 672 (1994).
- ⁶X. Wang, T.C. Lueng, B.N. Harmon, and P. Carra, Phys. Rev. B 47, 9087 (1993).
- ⁷P. Fischer, G. Schütz, and G. Wiesinger, Solid State Commun. **76**, 777 (1990).
- ⁸F. Baudelet, C. Brouder, E. Dartyge, A. Fontaine, J.P. Kappler, and G. Krill, Europhys. Lett. **13**, 751 (1990).
- ⁹S. Stähler, G. Schütz, and H. Ebert, Phys. Rev. B 47, 818 (1993).

- ¹⁰S. Pizzini, A. Fontaine, E. Dartyge, C. Giorgetti, F. Baudelet, J.P. Kappler, P. Boher, and F. Giron, Phys. Rev. B **50**, 3779 (1994).
- ¹¹T. Iwazumi, A. Koyama, and Y. Sakurai, Rev. Sci. Instrum. 66, 1691 (1995); K. Hirano, T. Ishikawa, A. Koyama, and T. Iwazumi, Photon Factory Activity Report 11, 412 (1993).
- ¹²E. Burzo and H.R. Kirchmayr, in *Handbook on the Physics and Chemistry of Rare Earths*, edited by K.A. Gschneidner, Jr. and L. Eyring (Elsevier, Amsterdam, 1989), Vol. 12, and references therein.
- ¹³G. Schütz, W. Wagner, W. Wilhelm, P. Kienle, R. Zeller, R. Frahm, and G. Materlik, Phys. Rev. Lett. 58, 737 (1987).
- ¹⁴C. Brouder and M. Hikam, Phys. Rev. B 43, 3809 (1991).
- ¹⁵Z.Q. Gu and W.Y. Ching, Phys. Rev. B **36**, 8546 (1987).
- ¹⁶W.Y. Ching and Z.Q. Gu, J. Appl. Phys. **63**, 3716 (1988).
- ¹⁷M.S.S. Brooks and B. Johansson, *Handbook of Magnetic Materials*, edited by K.H.J. Buschow (Elsevier, Amsterdam, 1993), Vol. 7.
- ¹⁸T. Jo and S. Imada, J. Phys. Soc. Jpn. 62, 3271 (1993).
- ¹⁹ F. Baudelet, Ch. Giorgetti, S. Pizzini, Ch. Brouder, E. Dartyge, A. Fontaine, J.P. Kappler, and G. Krill, J. Electron Spectrosc. Relat. Phenom. **62**, 153 (1993).