

Ordered phase in the two-dimensional randomly coupled ferromagnet

A. K. Hartmann

Institut für Theoretische Physik, Bunsenstrasse 9, 37073 Göttingen, Germany

I. A. Campbell

*Laboratoire des Verres, Université Montpellier II, Place Eugène Bataillon, 34095 Montpellier Cedex 05, France**and Physique des Solides, Université Paris Sud, 91405 Orsay, France*

(Received 25 May 2000; published 12 February 2001)

True ground states are evaluated for a $2d$ Ising model with random near-neighbor interactions and ferromagnetic second-neighbor interactions (the randomly coupled ferromagnet). The spin-glass stiffness exponent is positive when the absolute value of the random interaction is weaker than the ferromagnetic interaction. This result demonstrates that in this parameter domain the spin-glasslike ordering temperature is nonzero for these systems, in strong contrast to the $2d$ Edwards-Anderson spin glass.

DOI: 10.1103/PhysRevB.63.094423

PACS number(s): 75.10.Nr, 02.10.De, 75.40.Mg

I. INTRODUCTION

For more than two decades the intensive numerical work on the spin-glass (SG) problem has been concentrated almost exclusively on the Edwards-Anderson Ising spin glass: Ising spins on a regular (hyper)cubic lattice with random near-neighbor Gaussian or binomial interaction distributions.¹ The many possible alternative Ising systems with a randomness ingredient have hardly been touched on and such results as exist have been largely ignored.

One such family of alternative systems was proposed by Lemke and Campbell.² It consists of a $2d$ square lattice of $L \times L$ Ising spins $\sigma_i = \pm 1$ with uniform ferromagnetic second near-neighbor interactions of strength J , plus random near neighbor interactions $J_{ij} = \pm \lambda J$; we will refer to it as the RCF (randomly coupled ferromagnet) model.³ It is described by the following Hamiltonian:

$$H = - \sum_{\langle i,j \rangle} J_{ij} \sigma_i \sigma_j - \sum_{[i,j]} J \sigma_i \sigma_j. \quad (1)$$

For each realization of the randomness, the J_{ij} are drawn with the constraint $\sum_{\langle i,j \rangle} J_{ij} = 0$ to reduce fluctuations.

As the spins are coupled through the ferromagnetic second-near-neighbor interactions, the system can be partitioned into two interpenetrating sublattices in checkerboard-like fashion. In the limit $\lambda = 0$ the two sublattices order ferromagnetically and independently below the Onsager temperature $T = 2.27J$. Because each sublattice can order up or down, there are four degenerate ground states. As was pointed out in Ref. 2, for nonzero λ the near-neighbor interactions can be considered in terms of effective random fields exerted by each sublattice on the other, so that for finite λ and large enough L the ferromagnetic sublattice ordering is expected to be broken up, as in the $2d$ random-field Ising (RFI) model.⁴ The ground state will consist of coexisting domains of each of the four types: up/up, up/down, down/up, and down/down. The question is: is the break-up accompanied by paramagnetic order down to $T = 0$?

A number of Monte Carlo simulations were performed, and it was concluded on the basis of standard numerical

criteria^{2,3} that when the ratio λ is less than about 1, the RCF systems show spin-glasslike ordering at a *finite* temperature, whereas $2d$ Edwards Anderson ISG's are paramagnetic down to $T = 0$.⁵ The finite-ordering temperature interpretation was strongly questioned by Parisi *et al.*⁶ who criticized the initial work² on the grounds that the results were restricted to relatively small sample sizes L . On the basis of Monte Carlo data obtained on rather larger samples, Parisi *et al.* suggested that the RCF systems are always paramagnetic down to $T = 0$, like the Edwards Anderson (EA) ISG's. Further large sample Monte Carlo results,³ however, indicated finite-temperature ordering.

Here we present data from ground-state configuration evaluations which gives evidence that RCF systems indeed exhibit finite temperature SG-like ordering for λ less than about 1. This opens up intriguing possibilities for the testing of fundamental properties of complex ordered systems at finite temperatures in a $2d$ context.

II. ALGORITHM

In the present work, ground-state configurations have been found for periodic boundary conditions, and for the case where in one direction the boundary conditions are switched to antiperiodic. By comparing the ground-state energies of the different boundary conditions for each realization conclusions on the ordering behavior can be obtained. Similar studies were performed for simple d -dimensional EA spin glasses in two,⁵ three,⁷ and four⁸ dimensions.

For readers not familiar with the calculation of spin-glass ground states now a short introduction to the subject and a description of the algorithm used here are given. A detailed overview can be found in Ref. 9.

The concept of *frustration*¹⁰ is important for understanding the behavior of $\pm J$ Ising spin glasses. The simplest example of a frustrated system is a triplet of spins where all pairs are connected by antiferromagnetic bonds, see Fig. 1. A bond is called *satisfied* if it contributes with a negative value to the total energy by choosing the values of its adjacent spins properly. For the triangle, it is not possible to find a spin-configuration where all bonds are satisfied. In general, a




FIG. 1. The simplest frustrated system: a triple of spins, each pair of spins connected by antiferromagnetic bonds (dashed lines). It is not possible to satisfy all bonds.

system is frustrated if closed loops of bonds exists, where the product of these bond values is negative. For square and cubic systems, the smallest closed loops consist of four bonds. They are called (elementary) *plaquettes*.

As we will see later, the presence of frustration makes the calculation of exact ground states of such systems computationally hard. Only for the special case of the two-dimensional system with periodic boundary conditions in no more than one direction and without external field a polynomial-time algorithm is known.¹¹ In general, only methods with exponential running times are known, one says the problem is *NP hard*.¹² Now for the general case three basic methods are briefly reviewed and the largest system sizes which can be treated are given for three-dimensional systems, the standard spin-glass model, where data are available for comparison.

The simplest method works by enumerating all 2^N possible states and has obviously an exponential running time. Even a system size of 4^3 is too large. The basic idea of the so-called *Branch-and-Bound* algorithm¹³ is to exclude the parts of the state space, where no low-lying states can be found, so that the complete low-energy landscape of systems of size 4^3 can be calculated.¹⁴

A more sophisticated method called *Branch-and-Cut*^{15,16} works by rewriting the quadratic energy function as a linear function with an additional set of inequalities which must hold for the feasible solutions. Since not all inequalities are known *a priori*, the method iteratively solves the linear problem, looks for inequalities which are violated, and adds them to the set until the solution is found. Since the number of inequalities grows exponentially with the system size, the same holds for the computation time of the algorithm. With Branch-and-Cut anyway, small systems up to 8^3 are feasible.

The method used here is able to calculate true ground states⁷ up to size 14^3 . For two-dimensional systems, as considered in this paper, sizes up to 50^2 can be treated. The method is based on a special genetic algorithm^{17,18} and on *cluster-exact approximation*.¹⁹ CEA is an optimization method designed specially for spin glasses. Its basic idea is to transform the spin glass in a way that graph-theoretical methods can be applied, which work only for systems exhibiting no frustrations. Next a description of the genetic CEA is given.

Genetic algorithms are inspired by the evolution of populations in biology. An optimal solution is found by treating many instances of the problem in parallel, keeping only better instances and replacing bad ones by new ones (survival of




FIG. 2. Example of the cluster-exact approximation method. A part of a spin glass is shown. The circles represent lattice sites/spins. Straight lines represent ferromagnetic bonds the jagged lines antiferromagnetic interactions. The left half shows the initial situation. The construction starts with the spin at the center. The right half displays the final stage. The spins which belong to the cluster carry a plus or minus sign which indicates how each spin is transformed, so that only ferromagnetic interactions remain inside the cluster. All other spins cannot be added to the cluster because it is not possible to multiply them by ± 1 to make all adjacent bonds positive. Please note that many other combinations of spins can be used to build a cluster without frustration.

the fittest). The genetic algorithm starts with an initial population of M_i randomly initialized spin configurations (= *individuals*), which are linearly arranged using an array. The last one is also a neighbor of the first one. Then $n_o \times M_i$ times two neighbors from the population are taken (called *parents*) and two configurations called *offspring* are created. For that purpose, the *triadic crossover* is used which turned out to be very efficient for spin glasses: a mask is used which is a third randomly chosen (usually distant) member of the population with a fraction of 0.1 of its spins reversed. In a first step, the offspring are created as copies of the parents. Then those spins are selected, where the orientations of the first parent and the mask agree.²⁰ The values of these spins are swapped between the two offspring. Then a *mutation* with a rate of p_m is applied to each offspring, i.e., a fraction p_m of the spins is reversed.

Next for both offspring the energy is reduced by applying CEA: The method constructs iteratively and randomly a nonfrustrated cluster of spins. Spins adjacent to many unsatisfied bonds are more likely to be added to the cluster. During the construction of the cluster, a local gauge-transformation of the spin variables is applied so that all interactions between cluster spins become ferromagnetic.

Figure 2 shows an example of how the construction of the cluster works using a small spin-glass system. For $2d \pm J$ spin glasses each cluster contains typically 70% of all spins. The noncluster spins act like local magnetic fields on the cluster spins, so the ground state of the cluster is not trivial. Since the cluster has only ferromagnetic interactions, an energetic minimum state for its spins can be calculated in polynomial time by using graph theoretical methods:^{21,22} an equivalent network is constructed,²³ the maximum flow^{24,25} is calculated and the spins of the cluster are set to their orientations leading to a minimum in energy. This minimization step is performed n_{\min} times for each offspring.

Afterwards each offspring is compared with one of its parents. The pairs are chosen in the way that the sum of the

TABLE I. Simulation parameters: L =system size, M_i =initial size of population, n_o =average number of offsprings per configuration, n_{\min} =number of CEA minimization steps per offspring, p_m =mutation rate, and n_R =number of independent runs per realization.

L	M_i	n_o	m_{\min}	p_m	n_R
5	8	1	1	0.05	5
10	16	1	2	0.05	5
14	16	2	2	0.05	5
20	32	8	2	0.05	5
28	128	16	2	0.05	5
40	512	16	2	0.05	5
56	1024	16	2	0.05	5

phenotypic differences between them is minimal. The phenotypic difference is defined here as the number of spins where the two configurations differ. Each parent is replaced if its energy is not lower (i.e., not better) than the corresponding offspring. After this whole step is done $n_o \times M_i$ times, the population is halved: from each pair of neighbors, the configuration which has the higher energy is eliminated. If more than four individuals remain the process is continued; otherwise it is stopped and the best individual is taken as a result of the calculation.

The representation in the appendix summarizes the algorithm.

The whole algorithm is performed n_R times and all configurations which exhibit the lowest energy are stored, resulting in n_g statistical independent ground state configurations. The running time of the algorithm with suitable parameters chosen (see Table I) grows exponentially with the system size. On an 80 Mhz PowerPC processor, a typical $L=40$ instance takes 3 h (15 h for $L=56$).

III. RESULTS

In this work, ground states of the RCF are studied for values of $\lambda=0.5, 0.7, 0.9$, and 1.1 and system sizes up to $L=56$ ($\lambda=0.7$, and some realizations for 0.9), respectively $L=40$ ($\lambda=0.5, 1.1$). Usually 1000 different realizations were treated, each submitted to periodic (pbc) and antiperiodic (apbc) boundary conditions in one direction and always pbc in the other direction. The apbc are realized by inverting one line of bonds in the system with pbc. Because of the enormous computational effort, for the largest system sizes only realizations with $\lambda=0.7$ were considered with large statistics (and about 100 realizations with $L=56, \lambda=0.9$).

The periodic ground states give a direct measurement of the $T=0$ break up length L_b at each value of λ , which is defined as follows: for small enough L and finite value of λ , the ground states will always be such that there is a full ferromagnetic ordering within each sublattice. With increasing L , more and more samples will be found with ground states having at least one of the sublattices incompletely ferromagnetic. The break up length L_b is defined²⁶ as the value of L above which more than half the samples do not have full ferromagnetic order in each sublattice. The breakup lengths

TABLE II. Breakup length L_b for different values of the interaction strength λ .

λ	L_b
0.5	45
0.7	10
0.9	8
1.1	5

for the values of lambda considered here are presented in Table II. With the wisdom of hindsight, it can be seen that the measurements done in^{2,3,6} for $\lambda=0.5$ were mainly in the regime $L < L_b$ while for $\lambda=0.7$, the larger samples were well in the regime $L > L_b$.

A ‘‘typical’’ ground state for $\lambda=0.7$ and $L=56$ is shown in Fig. 3. All four possible types of domains occur. Because of the discrete structure of the interaction, usually the ground state is degenerate. But in contrast to the EA spin glasses with only $\pm J$ near-neighbor interactions, where a complex ground-state landscape exists, the structure of the degeneracy is trivial for $\lambda \leq 1$: the whole system may be flipped, sometimes it is possible to flip both sublattices independently, and usually some small clusters occur which can take two orientations.

But for studying whether the model exhibits long range order or not, it is sufficient to concentrate on the ground-state energies E_P, E_{AP} for periodic and antiperiodic boundary conditions. The energy differences $\Delta = E_P - E_{AP}$ give information about whether a system exhibits some kind of stiffness against perturbations of the boundary, i.e., about the presence of order. Δ is called the *stiffness energy*. For samples with the same set of interactions, the stiffness can be analyzed in terms of the size dependence of the average $\langle \Delta \rangle$ and of the width $W \equiv \sqrt{\sigma^2(\Delta)}$ of the distribution $P(\Delta)$. For $\lambda=0.7$, the distribution is presented in Fig. 4. In Fig. 5, the




FIG. 3. Typical ground state of one RCF realization ($L=56$) for $\lambda=0.70$ with periodic boundary conditions in all directions. Two different symbols (white/black square), (unfilled/filled diamond) are used to represent the orientations on the different sublattices.




FIG. 4. Distribution of stiffness energies Δ at $\lambda = 0.7$ for different system sizes $L = 14, 28,$ and 56 .

behavior of the average stiffness energy as a function of L is shown for all four values of λ . For system sizes larger than the breakup length of $\lambda \geq 0.7$, the stiffness energy decreases, indicating that no ferromagnetic long-range order is present in the system. For $\lambda = 0.5$, the breakup length is very large, so the asymptotic behavior is not visible, but the results are included for completeness. A better result for small values of λ can be obtained from direct evaluation of the magnetization, see Fig. 6 and Fig. 7. We determine a threshold value λ_{TH} for each system size from $m(\lambda_{TH}) = 0.9$. The value of 0.9 is somewhat arbitrary, so the results give just a rough impression of what happens. It is not possible to use much smaller thresholds, because the magnetization remains large for small system sizes even for strong spin-glass interaction. Via a fit $\lambda_{TH}(L) = \lambda_{\infty} + eL^{-f}$ we obtain $\lambda_{\infty} = 0.27(8)$. Therefore, there is strong evidence that also for $\lambda = 0.5$, no ferromagnetic order is present. For much smaller values of λ , nothing can be concluded from our data. Furthermore, for smaller values of λ it remains possible that the ground states of the RCF model do not exhibit ferromagnetic ordering for




FIG. 5. The mean stiffness energy $\langle \Delta \rangle$ as a function of L for $\lambda = 0.5, 0.7, 0.9,$ and 1.1 .




FIG. 6. Average magnetization m as a function of the strength λ of the spin-glass couplings for different system sizes $5 \leq L \leq 56$.

any finite value of the relative coupling constant λ .

Now we turn to the question whether some kind of spin-glass order is present in the system. This can be investigated by analyzing the dependence of the variance $\sigma^2(\Delta)$ of the stiffness-energy distributions on the system size, the result is shown in Fig. 8. For small system sizes, the variance grows for all values of the coupling constant λ . In order to exclude finite-size effects, only systems larger than the breakup length $L_b(\lambda)$ should be taken into account. Please note that for $L = 56, \lambda = 0.9$, only results for few realizations of the disorder were available due to the huge computational effort. Above L_b , there is a good linear size dependence of $\log W(L)$ against $\log L$, with $\theta_{SG} = 0.59(8), 0.29(1), 0.09(5),$ and $-0.16(2)$, respectively, for $\lambda = 0.5, 0.7, 0.9,$ and 1.1 . The values were obtained for $\lambda = 0.7, 0.9,$ and 1.1 from fits using only system sizes $L > L_b$ and are represented by straight lines in the figure. For $\lambda = 0.5$, also smaller system sizes were included, so the results for θ_{SG} is only included for completeness.




FIG. 7. Threshold value λ_{TH} as a function of the system size L . The threshold value is determined from $m(\lambda_{TH}) = 0.9$. The solid line shows a fit to a function $\lambda_{TH}(L) = \lambda_{\infty} + eL^{-f}$, resulting in $\lambda_{\infty} = 0.27(8), f = 0.53(10)$.




FIG. 8. Width $\sigma^2(\Delta)$ of the distribution of stiffness energies as a function of the system size $L \geq 10$ for $\lambda = 0.5, 0.7, 0.9$, and 1.1 . The solid lines are fits to algebraic functions of the form $\sigma^2(L) = gL^{\theta_{SG}}$. The inset shows the values of the exponent for different values of λ . The value for $\lambda = 0.5$ is included for completeness, since system sizes smaller than the breakup length have been included into the estimate. For the same reason, the corresponding fit is not shown.

The values of θ_{SG} against λ are shown in the inset of Fig. 8. The result for $\lambda = 0.5$ is not very reliable, because the largest system size is of the order of the breakup length. In the log-log plot, the datapoints for $\lambda = 0.5$ exhibit a negative curvature, thus the asymptotic value of θ_{SG} may be smaller than 0.59 . For the other systems, the breakup length is quite small, so the results give good evidence for spin-glasslike ordering in the large size limit, with a nonzero ordering temperature. Please note that for $\lambda = 1.1$, a change in the behavior is visible around $L = 14$ which is three times L_b . That means especially for $\lambda = 0.7$, where $L_b \approx 10$, the result $\theta_{SG} > 0$ is very reliable. This is the main result of this work, while the specific values obtained for θ_{SG} may be only rough guesses.

Thus, it seems indeed not necessary to carry out further calculations with larger systems to prove the fact, that there are values of the coupling constant giving rise to an ordered spin-glass phase in the RCF. The limiting value λ_c above which θ_{SG} is negative is very close to 1.0 ; λ_c would correspond to the highest value at which the ordering temperature is nonzero, in good agreement with the initial estimate from the Monte Carlo work.²

IV. CONCLUSION

We have calculated ground states of the randomly coupled ferromagnet for different values of the spin-glass coupling constant λ and with periodic as well as antiperiodic boundary conditions. By using the genetic cluster-exact approximation algorithm, we were able to treat system sizes up to $N = 56 \times 56$. The breakup length was calculated for each value of λ . From the calculation of the $T = 0$ stiffness energy, it could be concluded that below $\lambda_c \approx 1$, the RCF exhibits an ordered spin-glasslike phase at finite temperature. It should

be stressed again that for $\lambda = 0.7, 0.9$, and 1.1 , the largest system sizes are well beyond the breakup length, so no changes are to be expected when treating larger system sizes. This holds even for $\lambda = 0.7$, where the maximum system size is five times larger than the breakup length, which should be sufficient. For $\lambda \leq 0.5$, especially if one likes to test whether the model exhibits ferromagnetic ordering, ground-states calculations of larger systems are needed to study the behavior in more detail. Unfortunately, these studies are beyond the power of current computers and algorithms.

Although the zero-temperature stiffness exponent values give no direct information on the ordering temperatures, the present results are consistent with the conclusions drawn in Refs. 2 and 3, where Monte Carlo estimates of the critical temperatures were made using the finite-size scaling of the spin glass susceptibility and the form of the time dependence of the autocorrelation function relaxation. Ordering temperatures were estimated to be close to 2.0 for $\lambda = 0.5$ and 0.7 , dropping to zero near $\lambda = 1$. Rather remarkably, the $T = 0$ crossover as a function of L at L_b appears to have little effect on the behavior of the SG susceptibility as a function of size in the temperature region close to T_g^3 . However, for $\lambda = 0.5$, Parisi *et al.*⁶ observed weakly nonmonotonic behavior of the Binder parameter with L for sizes that we now know to be in the region of the crossover.

Since the existence of a spin-glasslike phase seems to be likely, at least for intermediate values of λ , it would be instructive to carry out further careful Monte Carlo measurements for sample sizes well in the regime $L > L_b$ and over a range of λ values. Is the physics of the $2d$ RCF above, at and below the ordering temperature strictly analogous to that of the standard Edwards Anderson ISG at dimensions where there is finite-temperature ordering? To what extent could the RCF enlighten us concerning problems which in the Edwards Anderson ISG context have remained conflictual for more than twenty years? The fact that the RCF lives on a $2d$ lattice rather than in a higher dimension should facilitate understanding of the fundamental physics of ordering in complex systems.

Finally, there may even be possible experimental realizations of systems where quasi-two-dimensional magnets form short-range clusters with local ferromagnetic or antiferromagnetic order, with random frustrated interactions linking these clusters together. Examples of promising behavior of this sort are Fe compounds with halogens,²⁷ where it might be interesting to look at the data again in view of the present results.

ACKNOWLEDGMENTS

A.K.H. was supported by the Graduiertenkolleg ‘‘Modellierung und Wissenschaftliches Rechnen in Mathematik und Naturwissenschaften’’ at the Interdisziplinäres Zentrum für Wissenschaftliches Rechnen in Heidelberg and the Paderborn Center for Parallel Computing by the allocation of computer time. A.K.H. obtained financial support from the DFG (Deutsche Forschungs Gemeinschaft) under Grant No. Zi209/6-1. I.A.C. gratefully acknowledges very helpful discussions with Dr. Lemke, and thanks Professor T. Shirakura

for having shown him very interesting unpublished data. Meetings organized by the Monbusho collaboration “Statistical physics of fluctuations in glassy systems” and by the ESF network “Sphinx” played an essential role for the present work.

APPENDIX: GENETIC CLUSTER-EXACT APPROXIMATION

algorithm genetic CEA($\{J_{ij}\}, M_i, n_o, p_m, n_{\min}$)

begin

create M_i configurations randomly

while ($M_i > 4$) *do*

begin

for $i = 1$ *to* $n_o \times M_i$ *do*

begin

select two neighbors

create two offspring using triadic crossover

do mutations with rate p_m

for both offspring *do*

begin

for $j = 1$ *to* n_{\min} *do*

begin

construct unfrustrated cluster of spins

construct equivalent network

calculate maximum flow

construct minimum cut

set new orientations of cluster spins

end

if offspring is not worse than related parent

then

replace parent with offspring

end

end

half population; $M_i = M_i/2$

end

return one configuration with lowest energy

end

¹For reviews on spin glasses see K. Binder and A. P. Young, *Rev. Mod. Phys.* **58**, 801 (1986); M. Mezard, G. Parisi, M. A. Virasoro, *Spin Glass Theory and Beyond* (World Scientific, Singapore 1987); K. H. Fisher and J. A. Hertz, *Spin Glasses* (Cambridge University Press, 1991).

²N. Lemke and I. A. Campbell, *Phys. Rev. Lett.* **76**, 4616 (1996).

³N. Lemke and I. A. Campbell, *J. Phys. A* **32**, 7851 (1999).

⁴Y. Imry and S.-K. Ma, *Phys. Rev. Lett.* **35**, 1399 (1975), M. Aizenman and J. Wehr, *ibid.* **62**, 2503 (1989).

⁵N. Kawashima and H. Rieger, *Europhys. Lett.* **39**, 85 (1997).

⁶G. Parisi, J. J. Ruiz-Lorenzo, and D. A. Stariolo, *J. Phys. A* **31**, 4657 (1998).

⁷A. K. Hartmann, *Phys. Rev. E* **59**, 84 (1999).

⁸A. K. Hartmann, *Phys. Rev. E* **60**, 5135 (1999).

⁹H. Rieger, in *Advances in Computer Simulation*, Vol. 501 of *Lecture Notes in Physics*, edited by J. Kertesz and I. Kondor (Springer-Verlag, Heidelberg, 1998).

¹⁰G. Toulouse, *Commun. Phys.* **2**, 115 (1977).

¹¹F. Barahona, R. Maynard, R. Rammal, and J. P. Uhry, *J. Phys. A* **15**, 673 (1982).

¹²F. Barahona, *J. Phys. A* **15**, 3241 (1982).

¹³A. Hartwig, F. Daske, and S. Kobe, *Comput. Phys. Commun.* **32**, 133 (1984).

¹⁴T. Klotz and S. Kobe, *J. Phys. A* **27**, L95 (1994).

¹⁵C. De Simone, M. Diehl, M. Jünger, P. Mutzel, G. Reinelt, and G. Rinaldi, *J. Stat. Phys.* **80**, 487 (1995).

¹⁶C. De Simone, M. Diehl, M. Jünger, P. Mutzel, G. Reinelt, and G. Rinaldi, *J. Stat. Phys.* **84**, 1363 (1996).

¹⁷K. F. Pál, *Physica A* **223**, 283 (1996).

¹⁸Z. Michalewicz, *Genetic Algorithms + Data Structures = Evolution Programs* (Springer, Berlin, 1992).

¹⁹A. K. Hartmann, *Physica A* **224**, 480 (1996).

²⁰K. F. Pál, *Biol. Cybern.* **73**, 335 (1995).

²¹J. D. Claiborne, *Mathematical Preliminaries for Computer Networking* (Wiley, New York, 1990).

²²M. N. S. Swamy and K. Thulasiraman, *Graphs, Networks and Algorithms* (Wiley, New York, 1991).

²³J.-C. Picard and H. D. Ratliff, *Network* **5**, 357 (1975).

²⁴J. L. Träff, *Eur. J. Oper. Res.* **89**, 564 (1996).

²⁵R. E. Tarjan, *Data Structures and Network Algorithms* (Society for Industrial and Applied Mathematics, Philadelphia, 1983).

²⁶E. T. Seppälä, V. Petäjä, and M. J. Alava, *Phys. Rev. E* **58**, 5217 (1998).

²⁷J. Vetel, M. Yahiaoui, D. Bertrand, A. R. Fert, J. P. Redoules, and J. Ferre, *J. Phys. Colloq.* **8**, **49**, 1067 (1988), D. Bertrand, F. Bensamka, A. R. Fert, J. Gelard, J. P. Redoules, and S. Legrand, *J. Phys. C* **17**, 1725 (1984).