

Thermal transport in the Falicov-Kimball model

J. K. Freericks¹ and V. Zlatić²

¹*Department of Physics, Georgetown University, Washington, DC 20057-0995*

²*Institute of Physics, Zagreb, Croatia*

(Received 29 August 2001; published 10 December 2001)

We prove the Jonson-Mahan theorem for the thermopower of the Falicov-Kimball model by solving explicitly for correlation functions in the large dimensional limit. We prove a similar result for the thermal conductivity. We separate the results for thermal transport into the pieces of the heat current that arise from the kinetic energy and those that arise from the potential energy. Our method of proof is specific to the Falicov-Kimball model, but illustrates the near cancellations between the kinetic- and potential-energy pieces of the heat current implied by the Jonson-Mahan theorem.

DOI: 10.1103/PhysRevB.64.245118

PACS number(s): 72.15.Jf, 72.20.Pa, 71.30.+h, 71.10.Fd

I. INTRODUCTION

The Jonson-Mahan^{1,2} theorem shows that there is a simple relation between the transport coefficient for the electrical conductivity and that needed for the thermopower. The relation is that the integral for the L_{12} coefficient has one more power of frequency in the integrand than the L_{11} coefficient. This result has been known for many years³ for a noninteracting system—the Jonson-Mahan theorem generalizes this result for a wide class of many-body systems (including the Falicov-Kimball model, the Holstein model, the periodic Anderson model with on-site hybridization, and the Hubbard model).

We use the exact solution of the Falicov-Kimball model in the large-dimensional limit to provide an alternate derivation of the Jonson-Mahan theorem by explicitly evaluating all relevant correlation functions needed for the thermal transport. Our exact analysis also allows us to separate the contributions to thermal transport that arise from the kinetic- and potential-energy pieces of the heat current. These results provide an interesting interpretation of thermal transport in correlated systems.

In Sec. II we develop a formalism for deriving the dc conductivity, the thermopower, and the thermal conductivity. We derive exact results for the relevant correlation functions, and use them to prove the Jonson-Mahan theorem and its generalization to the thermal conductivity. In Sec. III we provide numerical results for the thermal transport illustrating the different contributions to the thermal coefficients for a number of illustrative cases. Conclusions are presented in Sec. IV.

II. FORMALISM FOR THE THERMAL TRANSPORT

The Hamiltonian for our system is the spin- $\frac{1}{2}$ Falicov-Kimball model⁴

$$H = -\frac{t^*}{2\sqrt{d}} \sum_{\langle i,j \rangle \sigma} c_{i\sigma}^\dagger c_{j\sigma} + E_f \sum_i w_i + U \sum_{i\sigma} w_i c_{i\sigma}^\dagger c_{i\sigma}, \quad (1)$$

where $c_{i\sigma}^\dagger$ ($c_{i\sigma}$) is the electron creation (annihilation) operator for an electron at site i with spin σ , E_f is the energy level

of the localized electrons, w_i is a variable that is equal to zero or one and corresponds to the localized electron number, and U is the interaction strength. The hopping integral is scaled with the spatial dimension d so as to have a finite result in the limit⁵ $d \rightarrow \infty$; we measure all energies in units of $t^* = 1$. We work on a hypercubic lattice where the noninteracting density of states is a Gaussian $\rho(\epsilon) = \exp(-\epsilon^2)/\sqrt{\pi}$.

The Falicov-Kimball model can be solved exactly by employing dynamical mean-field theory^{6,7}. Because the self-energy $\Sigma(z)$ is local, the local Green's function satisfies

$$G(z) = \int d\epsilon \rho(\epsilon) \frac{1}{z + \mu - \Sigma(z) - \epsilon}, \quad (2)$$

with z anywhere in the complex plane (we suppress the spin index here). The self-energy, the local Green's function, and the effective medium G_0 are related by

$$G_0^{-1}(z) - G^{-1}(z) = \Sigma(z), \quad (3)$$

and the Green's function also satisfies

$$G(z) = (1 - w_1)G_0(z) + w_1 \frac{1}{G_0^{-1}(z) - U}. \quad (4)$$

Here w_1 is the average concentration of localized electrons,

$$w_1 = 2 \exp[-\beta(E_f - \mu)] Z_\uparrow(\mu - U) Z_\downarrow(\mu - U) / Z, \quad (5)$$

with $Z = Z_\uparrow(\mu) Z_\downarrow(\mu) + 2 \exp[-\beta(E_f - \mu)] Z_\uparrow(\mu - U) Z_\downarrow(\mu - U)$ and

$$Z_\sigma(\mu) = 2 e^{\beta\mu/2} \prod_n \frac{i\omega_n + \mu - \lambda_\sigma(i\omega_n)}{i\omega_n}. \quad (6)$$

The factor of 2 in Eq. (5) arises from the spin degeneracy of the f electrons, and the constraint that no more than one f electron is allowed on any site. The symbol $\lambda_\sigma(z)$ is defined from the effective medium via $\lambda_\sigma(i\omega_n) = i\omega_n + \mu - G_{0\sigma}^{-1}(i\omega_n)$. $\omega_n = \pi T(2n + 1)$ is the fermionic Matsubara frequency, and $\beta = 1/T$. The algorithm for determining the Green's function begins with the self energy set equal to zero. Then Eq. (2) is used to find the local Green's function. The effective medium is found from Eq. (3), and the localized electron filling from Eq. (5). The new local Green's

function is then found from Eq. (4), and the new self-energy from Eq. (3). This algorithm is repeated until it converges.

Transport properties are calculated within a Kubo-Greenwood formalism.⁸ This relates the transport coefficients to correlation functions of the corresponding transport current operators. We will deal with two current operators here—the particle current⁹

$$\mathbf{j} = \sum_{q\sigma} \mathbf{v}_q c_{q\sigma}^\dagger c_{q\sigma} \quad (7)$$

(where the velocity operator is $\mathbf{v}_q = \nabla_q \epsilon(q)$ and the Fourier transform of the creation operator is $c_q^\dagger = \sum_j \exp[iq \cdot \mathbf{R}_j] c_j^\dagger / N$) and the heat current^{1,9}

$$\begin{aligned} \mathbf{j}_Q = & \sum_{q\sigma} (\epsilon_q - \mu) \mathbf{v}_q c_{q\sigma}^\dagger c_{q\sigma} + \frac{U}{2} \\ & \times \sum_{qq'\sigma} W(q - q') [\mathbf{v}_q + \mathbf{v}_{q'}] c_{q\sigma}^\dagger c_{q'\sigma}, \end{aligned} \quad (8)$$

[where $W(q) = \sum_j \exp(-iq \cdot \mathbf{R}_j) w_j / N$]. The heat current can be broken into two pieces: (i) a kinetic-energy piece \mathbf{j}_Q^K , which is the first term in Eq. (8); and (ii) a potential-energy piece \mathbf{j}_Q^P , which is the second term in Eq. (8).

The particle current is defined by the commutator of the Hamiltonian with the polarization operator.⁹ When evaluated on a lattice with nearest-neighbor hopping, one finds factors that involve the weighted summation of the nearest-neighbor translation vectors δ weighted by phase factors $\exp(iq\delta)$, which yield the velocity operator terms above. The definition of the heat-current operator is more involved, and requires the Hamiltonian to be separated into operators h_i that involve the site i (in decomposing the kinetic-energy operator into “localized” pieces, one symmetrically assigns half of the $c_{i\sigma}^\dagger c_{j\sigma} + c_{j\sigma}^\dagger c_{i\sigma}$ term to site i and half to site j). These operators can be combined with the position operator to construct an “energy” polarization operator $\sum_i \mathbf{R}_i h_i$, which is commuted with the Hamiltonian to determine the energy current operator; the heat current operator is just this energy current operator shifted by the chemical potential multiplied by the number current operator. Important operator relations between the heat-current operator and the particle-current operator are described fully below [see the discussion around Eq. (60)].

The dc conductivity σ , thermopower S , and electronic thermal conductivity κ can all be determined from relevant correlation functions of the current operators. We define three transport coefficients L_{11} , $L_{12} = L_{21}$, and L_{22} . Then

$$\sigma = \frac{e^2}{T} L_{11}, \quad (9)$$

$$S = - \frac{k_B}{e|T|} \frac{L_{12}}{L_{11}}, \quad (10)$$

and

$$\kappa = k_B^2 \left[L_{22} - \frac{L_{12} L_{21}}{L_{11}} \right]. \quad (11)$$

The transport coefficients are found from the analytical continuation of the relevant “polarization operators” at zero frequency,

$$\begin{aligned} L_{11} &= \lim_{\nu \rightarrow 0} \text{Re} \frac{i}{\nu} \bar{L}_{11}(\nu), \\ \bar{L}_{11}(i\nu_l) &= \pi T \int_0^\beta d\tau e^{i\nu_l \tau} \langle T_\tau j_\alpha^\dagger(\tau) j_\beta(0) \rangle, \end{aligned} \quad (12)$$

where $\nu_l = 2\pi T l$ is the bosonic Matsubara frequency, the τ dependence of the operator is with respect to the full Hamiltonian in Eq. (1), and we must analytically continue $\bar{L}_{11}(i\nu_l)$ to the real axis $\bar{L}_{11}(\nu)$ before taking the limit $\nu \rightarrow 0$. Similar definitions hold for the other transport coefficients:

$$\begin{aligned} L_{12} &= \lim_{\nu \rightarrow 0} \text{Re} \frac{i}{\nu} \bar{L}_{12}(\nu), \\ \bar{L}_{12}(i\nu_l) &= \pi T \int_0^\beta d\tau e^{i\nu_l \tau} \langle T_\tau j_\alpha^\dagger(\tau) j_{Q\beta}(0) \rangle, \end{aligned} \quad (13)$$

and

$$\begin{aligned} L_{22} &= \lim_{\nu \rightarrow 0} \text{Re} \frac{i}{\nu} \bar{L}_{22}(\nu), \\ \bar{L}_{22}(i\nu_l) &= \pi T \int_0^\beta d\tau e^{i\nu_l \tau} \langle T_\tau j_{Q\alpha}^\dagger(\tau) j_{Q\beta}(0) \rangle. \end{aligned} \quad (14)$$

In all of these equations, the subscripts α and β denote the respective spatial indexes of the current vectors.

We begin with a derivation that shows the analytic continuation for the conductivity. Substituting the definition of the particle current operator of Eq. (7) into Eq. (12) for \bar{L}_{11} yields

$$\begin{aligned} \bar{L}_{11}(i\nu_l) &= \pi T \int_0^\beta d\tau e^{i\nu_l \tau} \sum_{qq'\sigma\sigma'} \mathbf{v}_{q\alpha} \mathbf{v}_{q'\beta} \\ & \times \langle T_\tau c_{q\sigma}^\dagger(\tau) c_{q\sigma}(\tau) c_{q'\sigma'}^\dagger(0) c_{q'\sigma'}(0) \rangle. \end{aligned} \quad (15)$$

The correlation function can be determined from Dyson’s equation, which relates the dressed correlation function to the bare correlation function via the irreducible charge vertex. Since the charge vertex is local in the infinite-dimensional limit, it is an even function of momentum, and any sum over momentum that is weighted by just one factor of \mathbf{v}_q will vanish. Hence the dressed correlation function is equal to just the bare correlation function¹⁰ (note that the contractions of the operators at equal times also vanish when summed over momentum). This produces




FIG. 1. Contours used in the analytical continuation: (a) contour needed for the Matsubara frequency summation and (b) deformed contour to lines parallel to the real axis.

$$\bar{L}_{11}(i\nu_l) = -\pi T \int_0^\beta d\tau e^{i\nu_l\tau} \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} G_{q\sigma}(\tau) G_{q\sigma}(-\tau). \quad (16)$$

Now we introduce the Fourier transform of the Green's function $G(\tau) = T \sum_n \exp(-i\omega_n\tau) G_n$ with $G_n = \int_0^\beta d\tau \exp(i\omega_n\tau) G(\tau)$. Substituting into Eq. (16) allows us to perform the integral over τ . This finally produces

$$\bar{L}_{11}(i\nu_l) = -\pi T^2 \sum_n \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} G_{q\sigma}(i\omega_n) G_{q\sigma}(i\omega_{n+l}). \quad (17)$$

The next step is to perform an analytical continuation from the imaginary axis to the real axis. The procedure is standard.⁹ We first write the summation over Matsubara frequencies as an integral over the contour C shown in Fig. 1(a), which has contributions at the poles of the Fermi function $f(\omega) = 1/[1 + \exp(\beta\omega)]$ which lie at the fermionic Matsubara frequencies. The contour is then deformed to lines

parallel to the real axis, with the Green's functions evaluated with either retarded (R) or advanced (A) functions. The result is

$$\begin{aligned} \bar{L}_{11}(i\nu_l) &= -\frac{T}{2i} \int_C d\omega f(\omega) \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} G_{q\sigma}(\omega) G_{q\sigma}(\omega + i\nu_l) \\ &= -\frac{T}{2i} \int_{-\infty}^{\infty} d\omega f(\omega) \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} \\ &\quad \times [G_{q\sigma}^R(\omega) - G_{q\sigma}^A(\omega)] G_{q\sigma}^R(\omega + i\nu_l) - \frac{T}{2i} \\ &\quad \times \int_{-\infty}^{\infty} d\omega f(\omega - i\nu_l) \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} G_{q\sigma}^A(\omega - i\nu_l) \\ &\quad \times [G_{q\sigma}^R(\omega) - G_{q\sigma}^A(\omega)]. \end{aligned} \quad (18)$$

The analytical continuation is performed by first rewriting $f(\omega - i\nu_l) = f(\omega)$, then taking $i\nu_l \rightarrow \nu + i\delta$ and shifting the integration variable $\omega \rightarrow \omega + \nu$ in the second integral. Then, using the definition for L_{11} , we finally arrive at

$$\begin{aligned} L_{11} &= \lim_{\nu \rightarrow 0} -\frac{T}{2\nu} \int_{-\infty}^{\infty} d\omega \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} \text{Re}\{f(\omega) G_{q\sigma}(\omega) \\ &\quad \times G_{q\sigma}(\omega + \nu) - f(\omega + \nu) G_{q\sigma}^*(\omega) G_{q\sigma}^*(\omega + \nu) \\ &\quad - [f(\omega) - f(\omega + \nu)] G_{q\sigma}^*(\omega) G_{q\sigma}(\omega + \nu)\}. \end{aligned} \quad (19)$$

Since $G_{q\sigma}(\omega) = 1/[\omega + \mu - \Sigma(\omega) - \epsilon(q)]$, we can perform a summation over q directly. Because $\epsilon(q)$ is an even function of q and \mathbf{v}_q is odd, we must have $\alpha = \beta$. Converting the fraction into the integral of an exponential then allows the summation over q to be performed directly.¹¹ The summation over q can be written as an integral over energy with a weighting factor of $\rho(\epsilon) t^{*2}/d$. This yields

$$\begin{aligned} L_{11} &= \lim_{\nu \rightarrow 0} -\frac{T t^{*2}}{2\nu d} \delta_{\alpha\beta} \int_{-\infty}^{\infty} d\omega \sum_{\sigma} \int_{-\infty}^{\infty} d\epsilon \rho(\epsilon) \\ &\quad \times \text{Re}\{f(\omega) G_{q\sigma}(\omega) G_{q\sigma}(\omega + \nu) \\ &\quad - f(\omega + \nu) G_{q\sigma}^*(\omega) G_{q\sigma}^*(\omega + \nu) \\ &\quad - [f(\omega) - f(\omega + \nu)] G_{q\sigma}^*(\omega) G_{q\sigma}(\omega + \nu)\}. \end{aligned} \quad (20)$$

If we define $\sigma_0 = e^2 t^{*2}/(2d)$, and we perform the integral over ϵ , we arrive at

$$\begin{aligned} L_{11} &= \lim_{\nu \rightarrow 0} \frac{T\sigma_0}{e^2} \delta_{\alpha\beta} \int_{-\infty}^{\infty} d\omega \frac{f(\omega) - f(\omega + \nu)}{\nu} \\ &\quad \times \text{Re} \left[-\frac{G(\omega) - G(\omega + \nu)}{\nu + \Sigma(\omega) - \Sigma(\omega + \nu)} \right. \\ &\quad \left. + \frac{G^*(\omega) - G(\omega + \nu)}{\nu + \Sigma^*(\omega) - \Sigma(\omega + \nu)} \right]. \end{aligned} \quad (21)$$

The final step is to take the limit of $\nu \rightarrow 0$. Using the facts that

$$\lim_{\nu \rightarrow 0} \frac{f(\omega) - f(\omega + \nu)}{\nu} = -\frac{df(\omega)}{d\omega} \quad (22)$$

and

$$\lim_{\nu \rightarrow 0} \frac{G(\omega) - G(\omega + \nu)}{\nu + \Sigma(\omega) - \Sigma(\omega + \nu)} = -2 + 2[\omega + \mu - \Sigma(\omega)]G(\omega) \quad (23)$$

produces our final result for L_{11} ,

$$L_{11} = \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) \tau(\omega), \quad (24)$$

with the relaxation time $\tau(\omega)$ defined by

$$\tau(\omega) = \frac{\text{Im} G(\omega)}{\text{Im} \Sigma(\omega)} + 2 - 2 \text{Re} \{ [\omega + \mu - \Sigma(\omega)] G(\omega) \}. \quad (25)$$

This result appears different from that originally derived for the hypercubic lattice,¹² but Eqs. (24) and (25) do yield the same result; the form presented here has the integral over the noninteracting density of states performed exactly.

The derivation for the transport coefficient L_{12} , needed for the thermopower, proceeds in a similar fashion. We can divide the heat current into two pieces; one corresponding to the kinetic energy, and one corresponding to the potential energy. This allows us to write $L_{12} = L_{12}^K + L_{12}^P$. The derivation for the kinetic-energy piece follows exactly like the derivation for L_{11} except there is an extra factor of $\epsilon - \mu$ that appears in Eq. (20). The integral over ϵ can then be performed straightforwardly, producing

$$L_{12}^K = \lim_{\nu \rightarrow 0} \frac{T\sigma_0}{e^2} \delta_{\alpha\beta} \int_{-\infty}^{\infty} d\omega \frac{f(\omega) - f(\omega + \nu)}{\nu} \text{Re} \left\{ -\frac{[\omega - \Sigma(\omega)]G(\omega) - [\omega + \nu - \Sigma(\omega + \nu)]G(\omega + \nu)}{\nu + \Sigma(\omega) - \Sigma(\omega + \nu)} + \frac{[\omega - \Sigma^*(\omega)]G^*(\omega) - [\omega + \nu - \Sigma(\omega + \nu)]G(\omega + \nu)}{\nu + \Sigma^*(\omega) - \Sigma(\omega + \nu)} \right\}. \quad (26)$$

Evaluating the limit $\nu \rightarrow 0$ is simple. The final result is

$$L_{12}^K = \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) \{ [\omega - \text{Re} \Sigma(\omega)] \tau(\omega) - 2 \text{Im} \Sigma(\omega) \text{Im} [(\omega + \mu - \Sigma(\omega))G(\omega)] \}, \quad (27)$$

with $\tau(\omega)$ defined in Eq. (25).

The derivation of the potential-energy piece is much more involved. The first step is to replace the momentum-dependent operator $W(q - q')$ by its Fourier transform. Simplifying the expression for \bar{L}_{12}^P yields

$$\begin{aligned} \bar{L}_{12}^P(i\nu_l) &= \frac{\pi T U}{2} \int_0^\beta d\tau e^{i\nu_l \tau} \sum_{qq'\sigma\sigma'} \frac{1}{N} \sum_j \mathbf{v}_{q\alpha} \mathbf{v}_{q'\beta} \\ &\times [e^{-iq' \cdot \mathbf{R}_j} \langle T_\tau w_j c_{q\sigma}^\dagger(\tau) c_{q\sigma}(\tau) c_{q'\sigma'}^\dagger(0) c_{j\sigma'}(0) \rangle \\ &+ e^{iq' \cdot \mathbf{R}_j} \langle T_\tau w_j c_{q\sigma}^\dagger(\tau) c_{q\sigma}(\tau) c_{j\sigma'}^\dagger(0) c_{q'\sigma'}(0) \rangle]. \end{aligned} \quad (28)$$

Noting that the w_j operator commutes with the fermionic operators allows us to use Wick's theorem to rewrite the terms in the square bracket as

$$\begin{aligned} \frac{1}{N} \sum_j [-e^{-iq' \cdot \mathbf{R}_j} \langle T_\tau w_j c_{q\sigma}^\dagger(\tau) c_{j\sigma}(0) \rangle G_{q\sigma}(\tau) \\ + e^{iq' \cdot \mathbf{R}_j} \langle T_\tau w_j c_{q\sigma}(\tau) c_{j\sigma}^\dagger(0) \rangle G_{q\sigma}(-\tau)] \delta_{qq'} \delta_{\sigma\sigma'}, \end{aligned} \quad (29)$$

where we have taken the appropriate contractions (note the velocity operators guarantee that we need not worry about any vertex corrections). The correlation functions in Eq. (29) can be evaluated by taking the derivative with respect to the components of an infinitesimal field $-\sum_j h_j w_j$. These correlation functions have a factor of $\exp[-\beta H]$ in the numerator and a factor of Z in the denominator. In addition, the τ dependence of the operators arises from factors of $\exp[\pm \tau H]$. Since the operator w_j commutes with all fermionic operators, it is easy to verify that the expression in Eq. (29) becomes

$$\begin{aligned} -\frac{1}{N} \sum_j \left[G_{q\sigma}(\tau) \left(T \frac{\partial}{\partial h_j} + \langle w_j \rangle \right) G_{q\sigma}(-\tau) + G_{q\sigma}(-\tau) \right. \\ \left. \times \left(T \frac{\partial}{\partial h_j} + \langle w_j \rangle \right) G_{q\sigma}(\tau) \right] \delta_{qq'} \delta_{\sigma\sigma'}, \end{aligned} \quad (30)$$

which follows by first removing the w_j operator through the derivative, then expressing the fermionic operator at site j through a Fourier transform, and finally evaluating the fermionic averages. Substituting this result into Eq. (28) then yields

$$\begin{aligned} \bar{L}_{12}^P(i\nu_l) &= -\frac{\pi T^2 U}{2N} \sum_n \sum_{q\sigma} \sum_j \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} \\ &\times \left(\left[T \frac{\partial}{\partial h_j} + \langle w_j \rangle \right] G_{q\sigma}(i\omega_n) \right) G_{q\sigma}(i\omega_{n+l}) \\ &+ G_{q\sigma}(i\omega_n) \left[T \frac{\partial}{\partial h_j} + \langle w_j \rangle \right] G_{q\sigma}(i\omega_{n+l}). \end{aligned} \quad (31)$$

The derivatives need to be computed. Writing the momentum-dependent Green's function as a Fourier transform,

$$G_{q\sigma}(i\omega_n) = \frac{1}{N} \sum_{i-j} e^{iq \cdot (\mathbf{R}_i - \mathbf{R}_j)} G_{ij\sigma}(i\omega_n), \quad (32)$$

and using the identity

$$G_{ij\sigma}(i\omega_n) = \sum_{kl} G_{ik\sigma}(i\omega_n) G_{kl\sigma}^{-1}(i\omega_n) G_{lj\sigma}(i\omega_n) \quad (33)$$

(with G^{-1} the matrix inverse of G), allows us to compute the derivative as

$$\begin{aligned} \frac{\partial}{\partial h_j} G_{q\sigma}(i\omega_n) &= \frac{1}{N} \sum_{i-j} e^{iq \cdot (\mathbf{R}_i - \mathbf{R}_j)} \\ &\times G_{ij\sigma}(i\omega_n) \frac{\partial \Sigma_{jj\sigma}(i\omega_n)}{\partial h_j} G_{jj\sigma}(i\omega_n). \end{aligned} \quad (34)$$

But in a homogeneous phase, the derivative of the local self-energy with respect to the local field and the local Green's function are both independent of the site j , so we finally arrive at

$$\frac{\partial G_{q\sigma}(i\omega_n)}{\partial h} = G_{n\sigma} \frac{\partial \Sigma_{n\sigma}}{\partial h} G_{q\sigma}(i\omega_n). \quad (35)$$

Since the self-energy depends only on G_n and w_1 , the derivative can be computed by taking partial derivatives and using the chain rule

$$\frac{\partial \Sigma_{n\sigma}}{\partial h} = \frac{\frac{\partial \Sigma_{n\sigma}}{\partial w_1} \frac{\partial w_1}{\partial h}}{1 - G_{n\sigma}^2 \frac{\partial \Sigma_{n\sigma}}{\partial G_{n\sigma}}}. \quad (36)$$

Each of the derivatives in Eq. (36) can be found directly:^{6,7}

$$\frac{\partial \Sigma_{n\sigma}}{\partial w_1} = \frac{U}{1 + G_{n\sigma}(2\Sigma_{n\sigma} - U)}, \quad (37)$$

$$\frac{\partial w_1}{\partial h} = \frac{w_1(1 - w_1)}{T}, \quad (38)$$

and

$$1 - G_{n\sigma}^2 \frac{\partial \Sigma_{n\sigma}}{\partial G_{n\sigma}} = \frac{(1 + G_{n\sigma} \Sigma_{n\sigma})(1 + G_{n\sigma}[\Sigma_{n\sigma} - U])}{1 + G_{n\sigma}(2\Sigma_{n\sigma} - U)}. \quad (39)$$

Substituting these derivatives into Eq. (36), and performing some straightforward simplifications that involve the quadratic equation that the self-energy satisfies,⁶ finally yields

$$\left[T \frac{\partial}{\partial h} + \langle w \rangle \right] G_{q\sigma}(i\omega_n) = \frac{\Sigma_{n\sigma}}{U} G_{q\sigma}(i\omega_n). \quad (40)$$

Now we are ready to perform the analytical continuation. First we substitute Eq. (40), into Eq. (31) and we note that the sum over j cancels the factor of $1/N$:

$$\begin{aligned} \bar{L}_{12}^P(i\nu_l) &= -\frac{\pi T^2}{2} \sum_n \sum_{q\sigma} \mathbf{v}_{q\alpha} \mathbf{v}_{q\beta} [\Sigma_{\sigma}(i\omega_n) \\ &+ \Sigma_{\sigma}(i\omega_{n+l})] G_{q\sigma}(i\omega_n) G_{q\sigma}(i\omega_{n+l}). \end{aligned} \quad (41)$$

Next, we rewrite the sum over Matsubara frequencies as a contour integral and perform the analytical continuation in the exact same way as before. If we then evaluate L_{12}^P , we find

$$\begin{aligned} L_{12}^P &= \lim_{\nu \rightarrow 0} -\frac{T\sigma_0}{2e^2\nu} \int_{-\infty}^{\infty} d\omega \int_{-\infty}^{\infty} d\epsilon \rho(\epsilon) \text{Re}(f(\omega)) \\ &\times \{ [\Sigma(\omega) + \Sigma(\omega + \nu)] G_q(\omega) G_q(\omega + \nu) \\ &- [\Sigma^*(\omega) + \Sigma(\omega + \nu)] G_q^*(\omega) G_q(\omega + \nu) \\ &+ f(\omega + \nu) \{ [\Sigma^*(\omega) + \Sigma(\omega + \nu)] G_q^*(\omega) G_q(\omega + \nu) \\ &- [\Sigma^*(\omega) + \Sigma^*(\omega + \nu)] G_q^*(\omega) G_q^*(\omega + \nu) \}. \end{aligned} \quad (42)$$

Now the integral over ϵ can be performed, and the limit $\nu \rightarrow 0$ can be taken. It becomes

$$\begin{aligned} L_{12}^P &= \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) (\text{Re} \Sigma(\omega) \tau(\omega) \\ &+ 2 \text{Im} \Sigma(\omega) \text{Im}[\{\omega + \mu - \Sigma(\omega)\} G(\omega)]). \end{aligned} \quad (43)$$

Adding together Eqs. (27) and (43) yields the Jonson-Mahan result of

$$L_{12} = \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) \tau(\omega) \omega. \quad (44)$$

Our final derivation is for the thermal conductivity coefficient L_{22} . Like before, we separate this into pieces corresponding to the kinetic energy and the potential energy: $L_{22} = L_{22}^{KK} + L_{22}^{KP} + L_{22}^{PK} + L_{22}^{PP}$. Due to the symmetry of the terms, we have $L_{22}^{KP} = L_{22}^{PK}$. The kinetic-energy piece is simple to calculate. Like in our derivation for L_{12}^K , the steps are identical to the derivation for L_{11} except we have an extra factor of $(\epsilon - \mu)^2$ in Eq. (20). Performing the integration over ϵ and collecting terms finally yields

$$\begin{aligned} L_{22}^{KK} &= \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) \{ [\omega - \text{Re} \Sigma(\omega)]^2 \tau(\omega) \\ &+ \text{Im} G(\omega) \text{Im} \Sigma(\omega) - 2[\text{Im} \Sigma(\omega)]^2 + 2[\text{Im} \Sigma(\omega)]^2 \\ &\times \text{Re}\{[\omega + \mu - \Sigma(\omega)] G(\omega)\} - 4[\omega - \text{Re} \Sigma(\omega)] \\ &\times \text{Im} \Sigma(\omega) \text{Im}\{[\omega + \mu - \Sigma(\omega)] G(\omega)\} \}. \end{aligned} \quad (45)$$

The derivation for $L_{22}^{KP} = L_{22}^{PK}$ is identical to that of L_{12}^P except we have an extra factor of $(\epsilon - \mu)$ in Eq. (42). Performing the integration over ϵ then produces

$$\begin{aligned}
L_{22}^{KP} = & \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) ([\omega - \text{Re} \Sigma(\omega)] \text{Re} \Sigma(\omega) \tau(\omega) \\
& - \text{Im} G(\omega) \text{Im} \Sigma(\omega) + 2[\text{Im} \Sigma(\omega)]^2 - 2[\text{Im} \Sigma(\omega)]^2 \\
& \times \text{Re}\{[\omega + \mu - \Sigma(\omega)] G(\omega)\} + 2[\omega - 2\text{Re} \Sigma(\omega)] \\
& \times \text{Im} \Sigma(\omega) \text{Im}\{[\omega + \mu - \Sigma(\omega)] G(\omega)\}). \quad (46)
\end{aligned}$$

The final term we must evaluate is L_{22}^{PP} . This is the most complicated term to evaluate, and we are unable to do so following the same strategy as employed in the L_{12}^P derivation. Instead, we proceed by an alternate method based on the equation of motion (EOM) technique. The EOM's for the fermionic creation and annihilation operators (in the momentum basis) are

$$\frac{\partial}{\partial \tau} c_{q\sigma}^\dagger(\tau) = [\epsilon(q) - \mu] c_{q\sigma}^\dagger(\tau) + U \sum_k W(k) c_{q+k\sigma}^\dagger(\tau) \quad (47)$$

and

$$\frac{\partial}{\partial \tau} c_{q\sigma}(\tau) = -[\epsilon(q) - \mu] c_{q\sigma}(\tau) - U \sum_k W(k) c_{q-k\sigma}(\tau). \quad (48)$$

These EOM's can be employed to express the correlation function of the heat-current operators in terms of derivatives with respect to imaginary time as shown below:

$$\begin{aligned}
\bar{L}_{22}^{PP}(i\nu_l) = & \frac{\pi T U^2}{4} \int_0^\beta d\tau e^{i\nu_l \tau} \sum_{qq'q''\sigma\sigma'} (\mathbf{v}_{q\alpha} + \mathbf{v}_{q''\alpha}) \\
& \times (\mathbf{v}_{q'\beta} + \mathbf{v}_{q''\beta}) \langle T_\tau W(q - q'') W(q' - q'') \\
& \times c_{q\sigma}^\dagger(\tau) c_{q''\sigma}(\tau) c_{q'\sigma'}^\dagger(0) c_{q''\sigma'}(0) \rangle \\
= & \pi T \int_0^\beta d\tau e^{i\nu_l \tau} \sum_{qq'\sigma\sigma'} \mathbf{v}_{q\alpha} \mathbf{v}_{q'\beta} \lim_{\tau' \rightarrow \tau^-} \lim_{\tau'' \rightarrow \tau'' - \rightarrow 0^+} \\
& \times \left\langle T_\tau \left[\frac{1}{2} (\partial_\tau - \partial_{\tau'}) - (\epsilon_{q'} - \mu) \right] \right. \\
& \times c_{q'\sigma}^\dagger(\tau) c_{q'\sigma}(\tau') \left. \left[\frac{1}{2} (\partial_{\tau''} - \partial_{\tau'''}) - (\epsilon_q - \mu) \right] \right. \\
& \times c_{q\sigma}^\dagger(\tau'') c_{q\sigma}(\tau''') \left. \right\rangle. \quad (49)
\end{aligned}$$

Now each of the operator averages can be expressed in terms of Green's functions, since the velocity factors guarantee there will be no vertex corrections. Noting further that the integrals will only contribute if $\alpha = \beta$ finally yields

$$\begin{aligned}
\bar{L}_{22}^{PP}(i\nu_l) = & \pi T \int_0^\beta d\tau e^{i\nu_l \tau} \sum_{q\sigma} v_{q\alpha}^2 \left[\frac{1}{2} \partial_\tau G_{q\sigma}(\tau) \partial_\tau G_{q\sigma}(-\tau) \right. \\
& - \frac{1}{4} \partial_\tau^2 G_{q\sigma}(\tau) G_{q\sigma}(-\tau) - \frac{1}{4} G_{q\sigma}(\tau) \partial_\tau^2 G_{q\sigma}(-\tau) \\
& + (\epsilon_q - \mu) \{ G_{q\sigma}(\tau) \partial_\tau G_{q\sigma}(-\tau) - \partial_\tau G_{q\sigma}(\tau) G_{q\sigma} \\
& \times (-\tau) \} - (\epsilon_q - \mu)^2 G_{q\sigma}(\tau) G_{q\sigma}(-\tau) \left. \right]. \quad (50)
\end{aligned}$$

We need to be able to produce expressions for the derivatives of the Green's functions. We do so by first writing the Green's function as a Fourier series over the Matsubara frequencies, and then taking the derivative into the Matsubara summation. This may appear to be mathematically unsound, but we do so by adding and subtracting the known $\delta(\tau)$ behavior of the derivative, to regularize the summation. Since we are interested only in $0 < \tau < \beta$, this procedure has no convergence issues. Likewise, one is also able to take the second derivative in this fashion. For $0 < \tau < \beta$ we find

$$\begin{aligned}
\partial_\tau G_{q\sigma}(\tau) = & -(\epsilon_q - \mu) G_{q\sigma}(\tau) \\
& - T \sum_m e^{-i\omega_m \tau} \frac{\sum_{m\sigma} \omega_m}{i\omega_m + \mu - \sum_{m\sigma} \omega_m - \epsilon_q}, \quad (51)
\end{aligned}$$

$$\begin{aligned}
\partial_\tau^2 G_{q\sigma}(\tau) = & +(\epsilon_q - \mu)^2 G_{q\sigma}(\tau) + (\epsilon_q - \mu) T \\
& \times \sum_m e^{-i\omega_m \tau} \frac{\sum_{m\sigma} \omega_m^2}{i\omega_m + \mu - \sum_{m\sigma} \omega_m - \epsilon_q} \\
& + T \sum_m e^{-i\omega_m \tau} \frac{i\omega_m \sum_{m\sigma} \omega_m}{i\omega_m + \mu - \sum_{m\sigma} \omega_m - \epsilon_q},
\end{aligned}$$

with similar formulas for $G_{q\sigma}(-\tau)$. Substituting the derivatives from Eq. (51) into Eq. (50), and then simplifying the result, finally produces

$$\begin{aligned}
\bar{L}_{22}^{PP}(i\nu_l) = & -\frac{\pi T^2}{4} \sum_n \sum_{q\sigma} v_{q\alpha}^2 (\{ \sum_\sigma(i\omega_n) \\
& + \sum_\sigma(i\omega_{n+1}) \}^2 G_{q\sigma}(i\omega_n) G_{q\sigma}(i\omega_{n+1}) \\
& + \sum_\sigma(i\omega_{n+1}) G_{q\sigma}(i\omega_n) + \sum_\sigma(i\omega_n) G_{q\sigma}(i\omega_{n+1})). \quad (52)
\end{aligned}$$

It is easy to understand the first terms in this expression, as they are what one would naively recover when following the same Wick analysis that was done previously for L_{12}^P . We have not been able to discover a direct operator-based derivation of the second terms, but they are critical for providing the right answer for L_{22}^{PP} . Performing an analytical continuation and simplifying yields our final result

$$\begin{aligned}
L_{22}^{PP} &= \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) ([\text{Re}\Sigma(\omega)]^2 \tau(\omega) \\
&\quad - 2[\text{Im}\Sigma(\omega)]^2 + \text{Im}\Sigma(\omega)\text{Im}G(\omega) \\
&\quad + 2[\text{Im}\Sigma(\omega)]^2 \text{Re}[\omega + \mu - \Sigma(\omega)]G(\omega) \\
&\quad + 4\text{Re}\Sigma(\omega)\text{Im}\Sigma(\omega)\text{Im}[\omega + \mu - \Sigma(\omega)]G(\omega)]. \quad (53)
\end{aligned}$$

Summing together Eq. (45), twice Eq. (46), and Eq. (53) gives the Mott form

$$L_{22} = \frac{T\sigma_0}{e^2} \int_{-\infty}^{\infty} d\omega \left(-\frac{df(\omega)}{d\omega} \right) \tau(\omega) \omega^2. \quad (54)$$

We can also generalize the original Jonson-Mahan argument to prove relations between L_{21} and L_{11} and between L_{22} and L_{12} . Our method is different from their proof, and relies on the infinite-dimensional limit, but one could proceed in their fashion if desired. We begin with the generalized two-particle correlation function

$$\begin{aligned}
F_{\alpha\beta}(\tau, \tau', \tau'', \tau''') &= \sum_{qq'\sigma\sigma'} \mathbf{v}_{q\alpha} \mathbf{v}_{q'\beta} \langle T_{\tau} c_{q\sigma}^{\dagger}(\tau) \\
&\quad \times c_{q\sigma}(\tau') c_{q'\sigma'}^{\dagger}(\tau'') c_{q'\sigma'}(\tau''') \rangle. \quad (55)
\end{aligned}$$

In the infinite-dimensional limit, the two-particle correlation function is expressed by just its bare bubble because the irreducible charge vertex has a different symmetry than \mathbf{v}_q . Hence we immediately learn that

$$F_{\alpha\beta}(\tau, \tau', \tau'', \tau''') = - \sum_{q\sigma} \mathbf{v}_{q\alpha}^2 \delta_{\alpha\beta} G_{q\sigma}(\tau''' - \tau) G_{q\sigma}(\tau' - \tau''). \quad (56)$$

But

$$G_{q\sigma}(\tau) = \int d\omega A(q, \omega) e^{-\omega\tau} [1 - f(\omega)] \quad (57)$$

for $\tau > 0$, and

$$G_{q\sigma}(\tau) = \int d\omega A(q, \omega) e^{-\omega\tau} [-f(\omega)] \quad (58)$$

for $\tau < 0$. Substituting into Eq. (56) then yields

$$\begin{aligned}
F_{\alpha\beta}(\tau, \tau', \tau'', \tau''') &= \frac{\delta_{\alpha\beta}}{2d} \int d\epsilon \rho(\epsilon) \int d\omega \int d\omega' A(\epsilon, \omega) A(\epsilon, \omega') \\
&\quad \times e^{\omega(\tau - \tau''') - \omega'(\tau' - \tau'')} f(\omega) [1 - f(\omega')]. \quad (59)
\end{aligned}$$

Using this function we can construct the relevant ‘‘polarization operators.’’ Recalling the EOM in Eqs. (47) and (48) shows that

$$\lim_{\tau' \rightarrow \tau^-} \frac{1}{2} \left(\frac{\partial}{\partial \tau} - \frac{\partial}{\partial \tau'} \right) \sum_{q\sigma} \mathbf{v}_q c_{q\sigma}^{\dagger}(\tau) c_{q\sigma}(\tau') = \mathbf{j}_Q(\tau). \quad (60)$$

The Jonson-Mahan theorem will hold for any Hamiltonian that satisfies Eq. (60) (which follows from the relevant commutators and equations of motion). This identity holds true for the Falicov-Kimball model, the Holstein model, the periodic Anderson model with on-site hybridization, and the Hubbard model. The ‘‘polarization operators’’ then become

$$\bar{L}_{11} = \pi T \int_0^{\beta} e^{i\nu\tau} F(\tau, \tau^-, 0, 0) \quad (61)$$

for the conductivity,

$$\bar{L}_{12} = \pi T \int_0^{\beta} e^{i\nu\tau} \frac{1}{2} \left(\frac{\partial}{\partial \tau} - \frac{\partial}{\partial \tau'} \right) F(\tau, \tau', 0, 0) \quad (62)$$

(in the limit where $\tau' \rightarrow \tau^-$) for the thermopower, and

$$\bar{L}_{22} = \pi T \int_0^{\beta} e^{i\nu\tau} \frac{1}{4} \left(\frac{\partial}{\partial \tau} - \frac{\partial}{\partial \tau'} \right) \left(\frac{\partial}{\partial \tau''} - \frac{\partial}{\partial \tau'''} \right) F(\tau, \tau', \tau'', \tau''') \quad (63)$$

(in the limit where $\tau' \rightarrow \tau^-$, $\tau'' \rightarrow \tau''^-$, and $\tau''' \rightarrow 0^+$) for the thermal conductivity. Because of Eq. (59), the analytical continuation is trivial (one first converts from imaginary time to Matsubara frequencies and then performs the Wick rotation to the real frequency axis), and if we note the identity

$$f(\omega) - f(\omega + \nu) = -f(\omega) [1 - f(\omega + \nu)] [e^{-\beta\nu} - 1], \quad (64)$$

then we can easily compute that

$$L_{11} = \frac{T\sigma_0}{e^2} \int d\epsilon \rho(\epsilon) \int d\omega \left(-\frac{df(\omega)}{d\omega} \right) A^2(\epsilon, \omega) \quad (65)$$

for the conductivity,

$$L_{12} = \frac{T\sigma_0}{e^2} \int d\epsilon \rho(\epsilon) \int d\omega \left(-\frac{df(\omega)}{d\omega} \right) A^2(\epsilon, \omega) \omega \quad (66)$$

for the thermopower, and

$$L_{22} = \frac{T\sigma_0}{e^2} \int d\epsilon \rho(\epsilon) \int d\omega \left(-\frac{df(\omega)}{d\omega} \right) A^2(\epsilon, \omega) \omega^2 \quad (67)$$

for the thermal conductivity. This proves Mott’s form for the thermal transport [since $\int d\epsilon \rho(\epsilon) A^2(\epsilon, \omega) = \tau(\omega)$].

III. NUMERICAL RESULTS

We first present results for a case where the filling of the localized electrons is a constant. We choose the symmetric case of $\langle w \rangle = 1/2$ and half-filling $\rho_e = 1$ for the electrons. We perform calculations for two cases: (i) $U = 1$ which is a moderately correlated metal (see Figs. 2 and 3), and (ii) $U = 2$ which is a strongly correlated insulator (see Figs. 4 and 5).

The thermopower in Fig. 2 behaves as expected—it van-




FIG. 2. Thermopower for the case $U=1$, $w_1=0.5$, and $\rho_d=1$. The main panel shows the two different contributions from the kinetic- and potential-energy pieces of the heat current, and the inset shows the net thermopower. Note how the two pieces are large and nearly cancel to produce S , and how there is a sign change near $T \approx 1.4$.

ishes for large and small temperature, it has an electronlike peak at lower temperatures, and it has a sign change at a temperature on the order of half the bandwidth. What is surprising is that there is such a large compensation between the kinetic- and potential-energy pieces of the thermopower to produce the net thermopower (note the three order of magnitude difference in the scales for the main figure and the inset). The thermal conductivity also appears as expected. We can see that while the contributions from the potential energy are critical in determining the right thermopower, they have a relatively mild effect in the thermal conductivity




FIG. 3. Thermal conductivity for the case $U=1$, $w_1=0.5$, and $\rho_d=1$. The plot shows the different contributions from the kinetic- and potential-energy pieces of the heat current. Note how the thermal conductivity is essentially described by the kinetic-energy-only piece for moderate correlation strength.




FIG. 4. Thermopower for the case $U=2$, $w_1=0.5$, and $\rho_d=1$. The main panel shows the two different contributions from the kinetic- and potential-energy pieces of the heat current, and the inset shows the net thermopower. Note how the two pieces are large and nearly cancel to produce S , and how the thermopower appears to diverge as T becomes small (calculations run into numerical problems as $T \rightarrow 0$).

for a moderately correlated metal (note how close the total thermal conductivity is to the kinetic-energy-only contribution) (see Fig. 3).

As we increase the correlation strength, so that the interacting density of states has a gap and the system is a correlated insulator, the behavior of the thermal transport changes. The thermopower in Fig. 4 has the characteristic insulating behavior here, with what appears to be a divergence as $T \rightarrow 0$. The divergence arises from the presence of a gap in the single-particle spectrum—both L_{11} and L_{12} approach zero exponentially in T (with the same exponent), but the ratio is




FIG. 5. Thermal conductivity for the case $U=2$, $w_1=0.5$, and $\rho_d=1$. The plot shows the different contributions from the kinetic- and potential-energy pieces of the heat current. Note how the potential-energy terms become increasingly important.

proportional to the size of the gap, so $S \rightarrow C/T + B$ as $T \rightarrow 0$ (similar results have been found when applying scaling theory to the Anderson transition¹³). Here this is “allowed” thermodynamically, because the “ground state” of the insulating phase has a nonzero entropy, since we forced the system into the paramagnetic insulating phase. In a real system, however, there must be a transition to a ground state where the entropy is quenched. In such a case, one would expect to see a large peak in the thermopower at low energies, with the thermopower ultimately going to zero as $T \rightarrow 0$. We know of no real correlated insulator that has a diverging thermopower as $T \rightarrow 0$.

The computation at very small values of T is difficult because the electrical conductivity (or equivalently L_{11}) approaches zero, and there are numerical difficulties associated with properly calculating the conductivity in this regime due to the cancellation of two large and nearly equal numbers. Since the thermopower requires the electrical conductivity, if that cannot be calculated accurately then spurious behavior will be seen in the thermopower. The thermal conductivity in Fig. 5 looks similar to the weaker correlated case, except that it appears to go to zero at a nonzero temperature which is the expected behavior for a correlated insulator with a gap and the potential-energy pieces become increasingly more important (particularly at low temperature).

In both of these half-filled cases, that of a correlated metal and a correlated insulator, the low-temperature thermopower is determined by a slightly larger contribution from the kinetic-energy piece of the heat current than from the potential-energy piece of the heat current. The thermal conductivity, on the other hand, has an evolution of going from a result nearly completely determined by the kinetic-energy-only piece of the heat current correlation functions to one where the potential-energy pieces of the heat current contribute progressively more and more to the total thermal conductivity.

Next we present results for the case where the total filling $\rho_e + \langle w \rangle = 1.5$ is a constant but the electrons can change from localized to itinerant (i.e., we fix the total electron concentration not the individual electron concentrations). We choose values of the parameters¹⁴ where the system has a sharp transition from a state at high temperature that has large f occupancy ($\langle w \rangle \approx 0.36$ for $0.2 < T < 0.8$), to a state at low temperature with no f electrons (the crossover occurs near $T = 0.04$). We find that the results do not depend too strongly on the parameters in this regime, and choose $E_f = -0.7$ and $U = 4$ as a canonical system that is similar to YbInCu_4 . The main difference from the symmetric case studied above is that the localized electron filling goes to zero as $T \rightarrow 0$. Hence both the kinetic and potential contributions to the thermopower become small in this limit, and there is no large cancellation between two nearly equal numbers to determine the thermopower (see Fig. 6). We see that in the thermal conductivity in Fig. 7, the contributions from the mixed kinetic- and potential-energy pieces are the most important, which is an indication of the strengthening of the correlations in the system. This feature is hard to see from the shape of the thermal conductivity itself.

The thermopower has an interesting exchange of impor-




FIG. 6. Thermopower for the case $U=4$, $w_1 + \rho_d = 1.5$, and $E_f = -0.7$. The main panel shows the two different contributions from the kinetic- and potential-energy pieces of the heat current, and the inset shows the net thermopower. Note the absolute scale for the thermopower is much larger here.

tance of the different pieces of the heat current as temperature is varied. For high temperatures, the thermopower is determined by both parts of the heat current and a compensation effect is important. In the moderate-temperature regime, the thermopower is dominated by kinetic-energy pieces, which then give way to the potential-energy domination at low temperature, that eventually shrinks as $T \rightarrow 0$ and the thermopower vanishes.

Note that one would need to reverse the sign of the thermopower to describe YbInCu_4 , since its charge carriers are holes rather than electrons. We should also remark that the thermopower of the Falicov-Kimball model does not have a low-energy peak associated with the “valence-change” transition—such a sharp peak occurs in YbInCu_4 -like systems because of hybridization effects not included in this




FIG. 7. Thermal conductivity for the case $U=4$, $w_1 + \rho_d = 1.5$, and $E_f = -0.7$. The plot shows the different contributions from the kinetic- and potential-energy pieces of the heat current.

model. Consequently, the thermopower also does not have a low-temperature sign changes as seen in the experimental data¹⁵ (a sign change occurs at $T \approx 2$), but we see that if one could reduce the potential-energy piece of the thermal current then the kinetic-energy contributions to the thermopower could cause a sign change to occur. This cannot happen in a pure Falicov-Kimball model, however, because of the Jonson-Mahan theorem. The sign change for the pure Falicov-Kimball model generically occurs at much larger values of temperature, on the order of the bandwidth. As regards the thermal conductivity, we find κ is dominated by the potential-energy piece as $T \rightarrow 0$ and the mixed piece yields a negative contribution over a wide temperature range.

IV. CONCLUSIONS

We have examined thermal transport in the spin- $\frac{1}{2}$ Falicov-Kimball model. We chose this model because the transport properties can be solved exactly, and they provide an alternate proof of the Jonson-Mahan theorem for the thermopower. We provide the proof in two different ways. The first is a brute-force application of the dynamical mean field theory to calculate all relevant correlation functions, and combine all terms to yield final expressions for the thermal transport coefficients. The second is based largely on the techniques of Jonson and Mahan, but one can determine the important “generalized polarization” functions exactly in the large dimensional limit. Here we extend the Jonson-Mahan

arguments to show analogous results hold for the thermal conductivity as well.

Our formulation also allows us to decompose the contributions to the thermopower and the thermal conductivity into the respective contributions from the kinetic-energy piece and the potential energy piece of the thermal current. We find that, generically, these pieces are large and opposite in sign for the thermopower, so that thermal transport carried by the kinetic heat current is almost completely compensated for by the potential heat current, producing a small net thermopower. For the thermal conductivity, we see an evolution of the transport being dominated first by kinetic-energy terms and then potential-energy terms as the strength of the correlations increase. We note that, because the kinetic-energy contribution to the thermopower can be straightforwardly determined for a number of models, any Hamiltonian that satisfies the Jonson-Mahan theorem can be separated into its kinetic and potential pieces for the thermopower by simply subtracting the kinetic-energy piece from the Jonson-Mahan result.

ACKNOWLEDGMENTS

We would like to acknowledge stimulating discussions with G. Czycholl, B. Letfulov, G. Mahan, R. Römer, G. Kotliar, and C. Villagonzalo. This work was supported by the National Science Foundation under Grant No. DMR-9973225.

-
- ¹M. Jonson and G. D. Mahan, Phys. Rev. B **21**, 4223 (1980); **42**, 9350 (1990).
- ²G. D. Mahan, Phys. Rev. B **56**, 11 833 (1997); Solid State Phys. **51**, 81 (1998).
- ³G. V. Chester and A. Thellung, Proc. Phys. Soc. London **77**, 1005 (1961); N. F. Mott and E. A. Davis, *Electronic Processes in Non-Crystalline Materials* (Clarendon, Oxford, 1971), p. 47.
- ⁴L. M. Falicov and J. C. Kimball, Phys. Rev. Lett. **22**, 997 (1969); R. Ramirez, L. M. Falicov, and J. C. Kimball, Phys. Rev. B **2**, 3383 (1970).
- ⁵W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989).
- ⁶U. Brandt and C. Mielsch, Z. Phys. B: Condens. Matter **75**, 365 (1989); **75**, 295 (1990); U. Brandt, A. Fledderjohann, and G. Hülsenbeck, *ibid.* **81**, 409 (1990); U. Brandt and C. Mielsch, *ibid.* **82**, 37 (1991).
- ⁷J. K. Freericks and V. Zlatić, Phys. Rev. B **58**, 322 (1998); V. Zlatić, J. K. Freericks, R. Lemański, and G. Czycholl, Philos. Mag. B **81**, 1443 (2001).
- ⁸R. Kubo, J. Phys. Soc. Jpn. **12**, 570 (1957); D. A. Greenwood, Proc. Phys. Soc. London **71**, 585 (1958).
- ⁹G. D. Mahan, *Many-Particle Physics* (Plenum, New York, 1990).
- ¹⁰A. Khurana, Phys. Rev. Lett. **64**, 1990 (1990).
- ¹¹E. Müller-Hartmann, Z. Phys. B: Condens. Matter **74**, 507 (1989); **76**, 211 (1989).
- ¹²Th. Pruschke, D.L. Cox, and M. Jarrell, Phys. Rev. B **47**, 3553 (1993); Th. Pruschke, D. L. Cox, and M. Jarrell, Europhys. Lett. **21**, 593 (1993).
- ¹³C. Villagonzalo and R. Römer, Ann. Phys. (Leipzig) **7**, 394 (1998); C. Villagonzalo, R. Römer, and M. Schreiber, Eur. Phys. J. B **12**, 179 (1999); special issue of Ann. Phys. (Leipzig) **8**, (1999), edited by M. Schreiber, p. S1-269; C. Villagonzalo, R. Römer, M. Schreiber, and A. MacKinnon, Phys. Rev. B **62**, 16 446 (2000).
- ¹⁴V. Zlatić and J. K. Freericks, in *Open Problems in Strongly Correlated Electron Systems*, edited by J. Bonča, P. Prelovšek, A. Ramšak, and S. Sarkar (Kluwer, Dordrecht, 2001) p. 371; V. Zlatić and J. K. Freericks, Acta Phys. Pol. B **32**, 3253 (2001).
- ¹⁵M. Očko, D. Drobac, J. L. Sarrao, and Z. Fisk, Phys. Rev. B **64**, 085103 (2001); M. Očko and J. L. Sarrao, Physica B (to be published).