

## Self-trapping of polarons in the Rashba-Pekar model

A. S. Mishchenko,<sup>1,2</sup> N. Nagaosa,<sup>1,3</sup> N. V. Prokof'ev,<sup>4</sup> A. Sakamoto,<sup>3</sup> and B. V. Svistunov<sup>2</sup>

<sup>1</sup>*Correlated Electron Research Center, AIST, Tsukuba Central 4, Tsukuba 305-8562, Japan*

<sup>2</sup>*Russian Research Center "Kurchatov Institute," 123182, Moscow, Russia*

<sup>3</sup>*Department of Applied Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113, Japan*

<sup>4</sup>*Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003*

(Received 7 May 2002; published 24 June 2002)

We performed a quantum Monte Carlo study of the exciton-polaron model which features the self-trapping phenomenon when the coupling strength and/or particle momentum is varied. Accurate data for energy, effective mass, the structure of the polaronic cloud, dispersion law, and spectral function became available throughout the crossover region. We observed that self-trapping cannot be reduced to hybridization of two states with different lattice deformation, and that at least three states are involved in the crossover from light- to heavy-mass regimes.

DOI: 10.1103/PhysRevB.66.020301

PACS number(s): 71.38.Ht, 02.70.Ss, 71.38.Fp

Properties of particles strongly coupled to their environment are of importance in many fields of physics and are attracting constant attention given extreme diversity of what may be called a "particle" and "environment," and how they interact with each other. In most general terms, self-trapping (ST) means a dramatic transformation of particle properties when system parameters are slightly changed. Landau<sup>1</sup> showed that the "trapped" (T) particle state with strong lattice deformation around it and the weakly perturbed "free-" (F) particle state may have the same energy at some critical value of the coupling strength,  $\alpha_c$ .

Of course, the resonance between F and T states is not infinitely sharp since the matrix element hybridizing them is nonzero, i.e., the ST phenomenon is a crossover, rather than a transition, and all polaron properties are analytic in  $\alpha$ —see Ref. 2 for explicit proof. This theorem makes the notion of ST rather vague since there is always some admixture of one state in another. Moreover, it challenges the adopted opinion that only two states, namely, F and T, are in competition. If there are more than two states within the energy scale of the hybridization matrix element then all of them are mixed and the F-T classification fails. In fact, the two-states assumption on which the current theory is based is not supported by experiments and rather complex spectra are usually observed instead.<sup>3,4</sup>

According to the standard criterion,<sup>5,6</sup> ST takes place if there is a barrier  $U_B$  in the adiabatic potential between the bare-particle and polaron states. It occurs, almost by definition, in the intermediate coupling regime where perturbation theory is not applicable. Hence, the existence of a barrier—if the very notion of the adiabatic potential is not ill defined—and the ST phenomenon can be addressed only by an exact method, because in the intermediate coupling regime an analytic solution is hardly available, and even sophisticated variational treatments often give misleading results.<sup>7</sup>

In this paper, we consider a typical model in which a particle couples to the environment of gapped dispersionless optical phonons. For this model it is possible to define ST in a mathematically rigorous way and proceed with its quantitative study. We show how various particle properties (energy, effective mass, dispersion law, and the structure of the

polaronic cloud) change between weak- and strong-coupling limits, and provide detailed information about the ST of polarons, which is not based on any approximations. In addition, we show that there are at least three states involved in mixing in the critical region and, thus the commonly accepted concept of only F and T states mixing at  $\alpha_c$  appears to be oversimplified. In fact, we are not aware of any other numerical study testing how accurate are existing treatments of the ST problem.

The Hamiltonian of the system consists of the free-particle term [we consider a continuum three-dimensional case with the dispersion relation  $\varepsilon(\mathbf{k}) = k^2/2m$ ]

$$H_e = \sum_{\mathbf{k}} \varepsilon(\mathbf{k}) a_{\mathbf{k}}^\dagger a_{\mathbf{k}}, \quad (1)$$

the Hamiltonian of the phonon bath

$$H_{\text{ph}} = \sum_{\mathbf{q}} \omega_{\mathbf{q}} b_{\mathbf{q}}^\dagger b_{\mathbf{q}} = \omega_0 \sum_{\mathbf{q}} b_{\mathbf{q}}^\dagger b_{\mathbf{q}}, \quad (2)$$

and the standard density-displacement interaction<sup>8</sup>

$$H_{e-\text{ph}} = \sum_{\mathbf{k}, \mathbf{q}} V(\mathbf{q}) (b_{\mathbf{q}}^\dagger - b_{-\mathbf{q}}) a_{\mathbf{k}-\mathbf{q}}^\dagger a_{\mathbf{k}}. \quad (3)$$

In Eqs. (1)–(3),  $a_{\mathbf{k}}$  and  $b_{\mathbf{q}}$  are the particle and phonon annihilation operators in momentum space, correspondingly.

Our study is based on the quantum Monte Carlo simulation of the polaron Green function in imaginary time at  $T = 0$  and subsequent analytic continuation to the real frequencies.<sup>9–11</sup> The method developed in Refs. 9–11 is free from approximations and systematic errors. It is particularly suited for the study of the ST problem where several  $\delta$  functional peaks are expected below the spectral continuum in the Lehman expansion

$$S^{(\mathbf{k})}(\omega) = \sum_{\nu} \delta[\omega - E_{\nu}(\mathbf{k})] |\langle \nu | a_{\mathbf{k}}^\dagger | \text{vac} \rangle|^2. \quad (4)$$

Here  $\{|\nu\rangle\}$  is a complete set of eigenstates of  $H$  in the momentum sector  $\mathbf{k}$ , i.e.,  $H|\nu(\mathbf{k})\rangle = E_{\nu}(\mathbf{k})|\nu(\mathbf{k})\rangle$ .




FIG. 1. The ground-state energy, average number of phonons, and effective mass as functions of  $\alpha$  (points connected by solid lines). Relative statistic errors are less than  $10^{-3}$  and  $10^{-2}$  for the energy and  $\langle N \rangle$ , respectively. The relative statistic errors for the mass are of order  $10^{-2}$  for  $\alpha < 18.5$  and around  $5 \times 10^{-2}$  for larger coupling constants. Dashed lines show results of the perturbation theory while the dotted line corresponds to the strong-coupling limit.

Separating stable quasiparticle states (labeled by the index  $i$ ) from the continuum, we rewrite Eq. (4) as

$$S^{(\mathbf{k})}(\omega) = \sum_i Z_i^{(\mathbf{k})}(0) \delta(\omega - E_i^{(\mathbf{k})}) + \int_{\omega_c} d\omega s^{(\mathbf{k})}(\omega), \quad (5)$$

where  $Z_i^{(\mathbf{k})}(0)$  and  $E_i^{(\mathbf{k})}$  are  $Z$  factors and energies of stable states, and the continuum threshold is given by  $\omega_c = E_0^{(\mathbf{k}=0)} + \omega_0$ . Any state with  $E > \omega_c$  is unstable against the single- ( $n=1$ ) or multiphonon ( $n>1$ ) emission process  $E \rightarrow E_i^{(\mathbf{p})} + n\omega_0$ , where momentum  $\mathbf{p}$  is selected only by the energy-conservation law since phonons are dispersionless.

Speaking rigorously, by self-trapping one understands the existence of such a region in the parameter space of  $H$  where more than one stable polaron state, differing by the degree of polarization of the lattice, coexist. This definition implies three critical points in the coupling constant (keeping other parameters fixed for simplicity),  $\alpha_{c1}(\mathbf{k}) < \alpha_c(\mathbf{k}) < \alpha_{c2}(\mathbf{k})$ . The ST “transition point”  $\alpha_c$  is understood as the point of avoided crossing between the two lowest polaron states. At this point the ground state of the polaron is a hybrid of states with substantially different degrees of lattice polarization.




FIG. 2. Partial weights of  $n$ -phonon states in the polaron ground state ( $\mathbf{k}=0$ ) at  $\alpha=18$  (circles),  $\alpha=18.35$  (squares), and  $\alpha=19$  (diamonds). Statistical error bars of order  $3 \times 10^{-3}$  are less than the symbol size.

The critical points  $\alpha_{c1}$  and  $\alpha_{c2}$  correspond to the appearance and disappearance of the extra stable state(s), respectively. [By definition, the energy difference  $\Delta E^{(\mathbf{k})}(\alpha)$  between the ground and first stable excited state, which has its minimum at  $\alpha_c(\mathbf{k})$ , ought to be less than  $\omega_c - E_0^{(\mathbf{k})}$ .] Critical couplings introduced above are consistent with previous considerations<sup>1,5,6</sup> and have an advantage of being unambiguous even when the minimal gap  $\Delta E^{(\mathbf{k})}[\alpha_c(\mathbf{k})]$  is not small.<sup>12</sup>

A typical system that is believed to feature ST is the so-called Rashba-Pekar model,<sup>13,14</sup> which describes Wannier exciton in the  $1s$  state interacting with optical vibrations via electrostatic potential,<sup>6</sup>

$$V(\mathbf{q}) = \gamma(\mathbf{q}) \left\{ \frac{1}{[1 + (\xi_e a_B q)^2]^2} - \frac{1}{[1 + (\xi_h a_B q)^2]^2} \right\}, \quad (6)$$

$$\gamma(\mathbf{q}) = i(2\sqrt{2}\alpha\pi)^{1/2} q^{-1}. \quad (7)$$

Here  $\alpha$  is the standard dimensionless coupling constant,  $a_B$  is the Bohr radius, and  $\xi_{e,h} = m_{e,h}/[2(m_e + m_h)]$  is given in terms of electron ( $m_e$ ) and hole ( $m_h$ ) masses, respectively.

In this paper, we focus on the parameters corresponding to the curve 2 of Ref. 13, which is believed to describe ST in the strong-coupling regime. Setting the total bare mass of the exciton  $m = m_e + m_h$ , phonon frequency  $\omega_0$ , electric charge, and Planck constant to unity, one finds that  $m_e = 0.065$ . The Bohr radius can be used to change the degree of adiabaticity in the model. Below we shall thoroughly consider an “almost adiabatic” case (i) with  $U_B/\omega_0 = 2$  (which is realized for  $a_B = 0.467$ ) and outline some peculiar features of the “nonadiabatic” situation (ii) with  $U_B/\omega_0 = 0.5$  ( $a_B = 0.934$ ). The critical coupling constants, determined within the approach of Ref. 13, are then  $\alpha_c^{\text{ad}} \approx 14.3$  and  $\alpha_c^{\text{ad}} \approx 7.2$ , respectively.

In Fig. 1 we show how the ground-state properties ( $\mathbf{k} = 0$ ) depend on the coupling strength. The ground-state en-




FIG. 3. The Lehman spectral function  $S^{(\mathbf{k}=0)}$  at coupling constants  $\alpha = 18.35$  (upper panel) and  $\alpha = 18.75$  (lower panel).

ergy, the effective mass  $m^*$ , and the average number of phonons in the polaronic cloud

$$\langle N \rangle = \left\langle \mathbf{k}=0 \left| \sum_{\mathbf{q}} b_{\mathbf{q}}^{\dagger} b_{\mathbf{q}} \right| \mathbf{k}=0 \right\rangle, \quad (8)$$

clearly indicate drastic changes around  $\alpha_c \approx 18.35$ . At this point the energy derivative changes very fast, and both  $\langle N \rangle$




FIG. 4. Energies of the ground (circles) and excited stable (squares, diamonds, and triangles) states vs interaction constant. The dashed line is the threshold of incoherent continuum. Typical error bars for the first, second, and third excited states are  $10^{-2}$ ,  $3 \times 10^{-2}$ , and  $4 \times 10^{-2}$ , respectively.




FIG. 5. The wave-vector dependence of energy and average number of phonons for  $\alpha = 17.75$  (circles connected by solid lines). The error bars are  $3 \times 10^{-3}$  and  $10^{-2}$  for energy and  $\langle N \rangle$ , respectively. The dashed curve is the effective-mass approximation  $E^{(\mathbf{k})} = E_0 + \mathbf{k}^2/2m^*$  with  $E_0(\alpha = 17.75) = -3.7946$  and  $m^*(\alpha = 17.75) = 2.258$  obtained from direct Monte Carlo estimators. The dotted curve is the parabolic dispersion law fitted to the last four points in the energy plot with parameters  $E_1(\alpha = 17.75) = -3.5273$  and  $m_1^* = 195$ . The open rectangle is the energy obtained from spectral analysis for the first excited state.

and  $m^*$  undergo a stepwise increase visible even on the logarithmic plot for  $m^*$ . In a narrow region between  $\alpha = 17.5$  and  $\alpha = 19$  the effective mass increases by two orders of magnitude. The remarkable fact is that at  $\alpha_c$  the strong-coupling approach is still far from being accurate (dotted line<sup>14</sup> in the upper panel of Fig. 1). In addition, the adiabatic critical constant  $\alpha_c^{\text{ad}} \approx 21$  (Ref. 14) differs significantly from our value  $\alpha_c \approx 18.35$ . For the “nonadiabatic” case (ii) the behavior of the ground state properties is qualitatively the same, but quantitative deviations from the strong-coupling limit are larger.

Next, we study how the phonon cloud evolves throughout the ST critical region. Partial  $n$ -phonon contributions to the polaron ground state  $Z_0^{\mathbf{k}}(n)$  are the probabilities of finding exactly  $n$  phonons in the cloud, and the average number of phonons introduced earlier is just  $\langle N \rangle = \sum_n n Z(n)$ . Figure 2 shows  $Z_0^{\mathbf{k}=0}(n)$  distributions at  $\alpha = 18$  (below the crossover region),  $\alpha = \alpha_c = 18.35$ , and  $\alpha = 19$  (trapped state). We see that the distribution at  $\alpha_c$  has two peaks and is half way between the two limiting cases. However, in the “nonadiabatic” case (ii) the structure with two maxima in  $Z(n)$  is missing. Therefore, the peculiar behavior presented in Fig. 1

is a general feature of the ST phenomenon whereas the two-peak structure of the phonon distribution is specific for the adiabatic limit.

The spectral function around the critical point<sup>15</sup> reveals up to three stable excited states below the continuum threshold (see examples of the Lehman function in Fig. 3). We observe in Fig. 4 that three polaronic states (in the energy range comparable with the hybridization strength) participate in the ST crossover. We emphasize that all three states have large  $Z$  factors ( $>0.1$ ). Therefore, for the given set of parameters more than two states are mixed at the crossover point and the standard picture of F-T hybridization at the tip of the ST crossover fails. One can speculate that extra stable states in the gap, which standard theory puts into the spectral continuum, are due to excited levels of highly nonlinear ST potential in the resonating region. However, this interpretation is essentially qualitative since the concept of adiabatic potential breaks down in the crossover region.

So far we have considered ST crossover at zero momentum, i.e., for the ground state. However, same considerations apply to finite momentum states, as long as  $k < k_c$  where  $k_c$  is defined as the point where  $E_0^{(k)} = \omega_c$  and the polaron spectrum has an end point.<sup>9,16</sup> For  $\alpha < \alpha_c$  the ground state is characterized by small lattice distortion and light effective mass and thus the dispersion law associated with this state rises steeply with  $\mathbf{k}$ . On another hand, the dispersion curve of the excited heavy-mass state is nearly flat which means that light and heavy branches have to intersect at some wave vector. In Fig. 5 we plot the ground-state dispersion law and

the average number of phonons in the ground state for  $\alpha = 17.75$  which we interpret as the level crossing picture (see the figure caption). Notice the agreement between the spectral analysis at  $\mathbf{k}=0$  and the parabolic fit of the heavy-mass branch.

By all accounts, the existence of more than one stable polaron state is a highly nontrivial qualitative property of the model whether the hybridization gap between these states is small, or not. Moreover, the ST crossover is not necessarily limited to hybridization of only two states. Recent studies of the Holstein polaron in one dimension strongly support the universality of the latter statement since more than two stable states were found there as well.<sup>17</sup>

We found that the dependence of  $E_0$  and  $\langle N \rangle$  (but not the structure of the cloud) on coupling can be used as an indirect indication of the self-trapping phenomenon with a well-defined “transition point” (the point of minimal gap between the ground and first stable excited state). It was considered previously as a theorem, that is, ST may occur only in dimensions  $d > 2$ ;<sup>5,6</sup> however, the definition of what has to be counted as a ST transition was not given in quantitative terms. We believe that it is more appropriate to use a less “radical,” but unambiguous, definition adopted in our paper. Recently, the second stable polaron state was found to exist for the Holstein polaron in a one-dimensional lattice<sup>18</sup> and in an infinite-dimension approximation.<sup>19</sup>

This work was supported by the National Science Foundation under Grant No. DMR-0071767 and RFBR 01-02-16508.

<sup>1</sup>L.D. Landau, *Phys. Z. Sowjetunion* **3**, 664 (1933).

<sup>2</sup>B. Gerlach and H. Löwen, *Rev. Mod. Phys.* **63**, 63 (1991).

<sup>3</sup>A. Matsui, K. Mizuno, and M. Ikemura, *J. Phys. Soc. Jpn.* **51**, 1871 (1982).

<sup>4</sup>T. Koda and Y. Kaneko, in *Luminescence: Phenomena, Materials and Devices*, edited by R.P. Rao (Nova Science Publishers, New York, 1992), p. 175.

<sup>5</sup>E.I. Rashba, *Self-Trapping of Excitons*, in *Modern Problems in Condensed Matter Sciences*, Edited by V.M. Agranovich and A.A. Maradudin (North-Holland, Amsterdam, 1982), Vol. 2, p. 543; A.S. Ioselevich and E.I. Rashba, *Theory of Nonradiative Trapping in Crystals* (North-Holland, Amsterdam, 1992), Vol. 34, p. 347.

<sup>6</sup>M. Ueta, H. Kanzaki, K. Kobayashi, Y. Toyozawa, and E. Hanamura, *Exciton Processes in Solids* (Springer-Verlag, Berlin, 1986).

<sup>7</sup>Self-trapping was predicted to occur in the Fröhlich polaron model in various papers, but recently it has been proven to be an artifact of the variational scheme (Ref. 10).

<sup>8</sup>J. Appel, in *Solid State Physics*, edited by H. Ehrenreich, F. Seitz, and D. Turnbull (Academic, New York, 1968), Vol. 21.

<sup>9</sup>N.V. Prokof'ev and B.V. Svistunov, *Phys. Rev. Lett.* **81**, 2514 (1998).

<sup>10</sup>A.S. Mishchenko, N.V. Prokof'ev, A. Sakamoto, and B.V. Svistunov, *Phys. Rev. B* **62**, 6317 (2000).

<sup>11</sup>A.S. Mishchenko and N. Nagaosa, *Phys. Rev. Lett.* **86**, 4624 (2001).

<sup>12</sup>With nonzero coupling to acoustic modes the excited polaron states have finite lifetimes. However, when the energy gap between the ground and excited states gets small, the lifetime diverges due to vanishing probability of emitting a long-wavelength phonon.

<sup>13</sup>S.I. Pekar, E.I. Rashba, and V.I. Sheka, *Zh. Éksp. Teor. Fiz.* **76** 251 (1979) [*Sov. Phys. JETP* **49**, 129 (1979)].

<sup>14</sup>I.M. Dykman and S.I. Pekar, *Dokl. Akad. Nauk SSSR* **83**, 825 (1952).

<sup>15</sup>The specific features of the spectral analysis in the present case will be presented in a subsequent paper.

<sup>16</sup>D.M. Larsen, *Phys. Rev.* **144**, 697 (1966).

<sup>17</sup>A. Macridin, A.S. Mishchenko, N.V. Prokof'ev, G.A. Sawatzky, and B.V. Svistunov (unpublished).

<sup>18</sup>J. Bonča, S.A. Trugman, and I. Batistić, *Phys. Rev. B* **60**, 1633 (1999).

<sup>19</sup>S. Ciuchi, F. de Pasquale, and D. Feinberg, *Europhys. Lett.* **30**, 151 (1995).