

Domain wall renormalization group study of the XY model with quenched random phase shifts

N. Akino¹ and J. M. Kosterlitz²¹*Institut für Physik, WA 331, Johannes Gutenberg-Universität, D-55099 Mainz, Germany*²*Department of Physics, Brown University, Box 1843, Providence, Rhode Island 02912*

(Received 15 March 2002; published 29 August 2002)

The XY model with quenched random disorder is studied by a zero-temperature domain wall renormalization group method in two dimensions (2D) and three dimensions (3D). Instead of the usual phase representation we use the charge (vortex) representation to compute the domain wall, or defect, energy. For the gauge glass corresponding to the maximum disorder we reconfirm earlier predictions that there is no ordered phase in 2D but an ordered phase can exist in 3D at low temperature. However, our simulations yield spin stiffness exponents $\theta_s \approx -0.36$ in 2D and $\theta_s \approx +0.31$ in 3D, which are considerably larger than previous estimates and strongly suggest that the lower critical dimension is less than three. For the $\pm J$ XY spin glass in 3D, we obtain a spin stiffness exponent $\theta_s \approx +0.10$ which supports the existence of spin glass order at finite temperature in contrast with previous estimates which obtain $\theta_s < 0$. Our method also allows us to study renormalization group flows of both the coupling constant and the disorder strength with a length scale L . Our results are consistent with recent analytic and numerical studies suggesting the absence of a reentrant transition in 2D at low temperature. Some possible consequences and connections with real vortex systems are discussed.

DOI: 10.1103/PhysRevB.66.054536

PACS number(s): 05.40.-a, 74.60.Ge, 75.50.Lk

I. INTRODUCTION

The XY model with quenched random phase shifts as a model for a superconducting glass has been intensively investigated over the last decade, focusing on the so-called gauge glass model which corresponds to the case with maximal disorder. Since a transport current exerts a force on a flux lattice, it tends to move in response which causes dissipation of the current. The existence of disorder, which destroys the flux lattice structure, is essential to pin the vortices in order for a superconducting phase to exist in a high- T_c superconductor.¹⁻³ Although there exists no proof whether or not the gauge glass and vortex glass are in the same universality class, it is of interest as the simplest model of a disordered superconductor and is still not understood despite all the effort expended on it.

From numerical⁴⁻⁹ and experimental¹⁰ studies, it is believed that the gauge glass has no ordered phase at any finite temperature in two dimensions (2D). In three dimensions (3D), numerical domain wall renormalization group (DWRG) studies^{11,12} indicate that the lower critical dimension seems to be close to three. However the situation is less conclusive, since the simulations are limited to small system sizes. Finite temperature Monte Carlo studies yield a transition temperature $T_c/J \sim \mathcal{O}(1)$,^{4,5,13-15} which is difficult to reconcile with DWRG studies^{5,6,8,9} as these studies imply that the lower critical dimension for superconducting glass order is close to three. Experimentally there is also some evidence for a finite temperature phase transition to a superconducting glass phase.^{16,17}

The Hamiltonian of the XY model with random quenched disorder can be written as

$$H = \sum_{\langle ij \rangle} V(\theta_i - \theta_j - A_{ij}), \quad (1)$$

where $V(\phi)$ is an even, 2π periodic function of ϕ with a maximum at $\phi = \pi$ and minimum at $\phi = 0$, usually taken as

$V(\phi_{ij}) = -J_{ij} \cos(\phi_{ij})$. The sum is over all nearest-neighbor pairs of sites and the coupling constants, J_{ij} , are assumed uniform, $J_{ij} = J > 0$. The random bond variables A_{ij} , which are responsible for the randomness and frustration, are taken to be independent and uniformly distributed in $(-\alpha\pi, \alpha\pi]$ with $0 \leq \alpha \leq 1$. For a gauge glass, θ_i is the phase of the superconducting order parameter at site i of a square lattice in 2D and a simple cubic lattice in 3D. The random bond variables A_{ij} are taken to correspond to maximal disorder with $\alpha = 1$. An external field applied to an extreme type-II superconductor induces a uniform component $A_{ij}^0 = (2\pi/\Phi_0) \int_i^j \mathbf{A} \cdot d\mathbf{l}$ where \mathbf{A} is the vector potential of the applied field and $\Phi_0 = hc/2e$ is the quantum of flux. In this work, we take $A_{ij}^0 = 0$, corresponding to zero applied field. Unless explicitly stated, we consider an unscreened system with $\alpha = 1$ corresponding to maximal disorder. The Hamiltonian of Eq. (1) also describes the XY magnet with random Dzyaloshinski-Moriya interactions¹⁸ and also a Josephson-junction array with positional disorder.^{19,20} These studies¹⁸⁻²⁰ showed that the existence of weak disorder ($\alpha \ll 1$) does not destroy an ordered phase at intermediate temperature but predicts a reentrant transition to a disordered phase at low temperature in two dimensions. However, recent analytic²¹⁻²⁵ and numerical⁸ studies suggest the absence of a reentrant transition and that there exists an ordered phase for $T < T_c(\alpha)$ when $\alpha < \alpha_c$.

When the random bond variables A_{ij} are restricted to 0 or π with equal probabilities, this model reduces to the $\pm J$ XY spin glass, which is believed to be in a different universality class due to the additional reflection symmetry²⁶ which is absent in the case of uniformly or Gaussian distributed A_{ij} . An XY spin glass may have both spin- and chiral glass order associated with rotational and reflection symmetries, respectively. It has been suggested that, in 2D and 3D, spin and chiral variables decouple at long distances and order independently,²⁷⁻³⁰ and the lower critical dimensions are d_l

≥ 4 for spin-glass order and $d_l < 3$ for chiral glass order. However, the decoupling scenario contradicts the analytic studies on a ladder lattice,³¹ on a tube lattice,³² and on a 2D lattice with a special choice of disorder.³³ Recent numerical simulations³⁰ also suggest d_l for a spin-glass order may be close to three.

In this paper, we reinvestigate the possibility of an ordered phase at small but finite temperature T by a numerical DWRG,^{11,12} or defect energy scaling. The domain-wall or defect energy of the system is computed by using the Hamiltonian in the Coulomb gas (vortex) representation, which is more convenient for numerical work as it eliminates spin-wave contributions to the energy. Although the conventional DWRG method can handle only the scaling of the coupling constant $J(L)$ at scale L , which is proportional to the domain wall or defect energy, our method enables us to study the flows of both the coupling constant and the disorder strength, $A(L)$, at length scale L .⁸ We apply this to the case of general disorder strength, $0 \leq \alpha \leq 1$. The outline of the paper is as follows. In Sec. II we discuss the DWRG method and also our strategy. In Sec. III, we explicitly perform the transformation of the 3D Hamiltonian of Eq. (1) from the phase to the Coulomb gas representation. Our numerical method is explained in Sec. IV. Finally we discuss our numerical results in Sec. V and give a brief discussion of some of the effects of weak disorder, $\alpha < 1$, and of finite screening of vortex-vortex interactions.

II. STRATEGY

The general idea behind a DWRG is to compute, analytically or numerically, the energy $\Delta E(L)$ of a domain wall in a system of linear size L and fit this to a finite-size scaling form

$$\Delta E(L) \sim L^\theta, \quad (2)$$

where θ is a stiffness exponent, whose sign is of fundamental importance. If $\theta < 0$, $\Delta E(L)$ vanishes in the thermodynamic limit. The energy of the domain wall or defect excitation vanishes which implies that, for $T > 0$, the probability of the defect $P_L \sim e^{-\Delta E(L)/kT} \rightarrow 1$ as $L \rightarrow \infty$. This in turn implies that the density of such defects is finite when $T > 0$ and there will be no resistance to an infinitesimal applied force and the system has no order. This is analogous to the vanishing of the shear modulus in a liquid, the superfluid density in a superfluid or superconductor, and the spin stiffness constant in an isotropic magnet when $T > T_c$. On the other hand, if $\theta > 0$, such defects will have zero probability when $L = \infty$ and the system will have finite stiffness and will be ordered at sufficiently small $T > 0$.

In a uniform system without disorder, the definition of the energy of a domain wall of size L , $\Delta E(L)$, is intuitively obvious. The first step is to find the ground-state (GS) energy of a system of size L , which requires applying boundary conditions (BC) which are compatible with the GS configuration. For a ferromagnet, this is straightforward to implement as the GS configuration is known to be one with all spins parallel, and periodic BC are compatible with this. To

impose a spin domain wall perpendicular to the $\hat{\mathbf{x}}$ direction, one simply changes the BC to antiperiodic along $\hat{\mathbf{x}}$ and periodic in the other $d-1$ directions. Then it immediately follows that

$$\Delta E(L) = E_{ap}(L) - E_p(L) \sim L^{d-n}, \quad (3)$$

where $n=1$ for an Ising model and $n=2$ for a system with a continuous symmetry such as XY and Heisenberg models.

One would like to use the same strategy for *random* systems described by Eq. (1), as suggested by Anderson for Ising spin glasses.³⁴ However, it is not so clear how to proceed because, for a particular sample (realization of disorder), neither the GS configuration nor compatible BC are known so computing the defect energy $\Delta E(L)$ is problematical. Assuming $\Delta E(L)$ can be calculated, the stiffness exponent is defined by the scaling ansatz

$$\langle \Delta E(L) \rangle \sim L^\theta, \quad (4)$$

where $\langle \dots \rangle$ denotes an average over realizations of disorder. To our knowledge, it is not known how to calculate *analytically* either the GS energy $E_0(L)$ or the energy $E_p(L)$ of the system containing a defect *relative to this GS* which means that one must proceed numerically. A number of conceptual and technical difficulties are apparent. The first, and most important, is the technical problem of computing the energy difference $\Delta E(L)$ between the energies of the system subject to two different BC. We ultimately want the disorder averaged defect energy $\langle \Delta E(L) \rangle = \langle E_a(L) - E_b(L) \rangle$ where $E_a(L)$ is the lowest energy of a particular sample subject to BC denoted by a and $E_b(L)$ with BC denoted by b . We need the individual energies $E_a(L)$ and $E_b(L)$ essentially *exactly* because the uncertainty in $\langle \Delta E(L) \rangle$ must be kept as small as possible. Also, to our knowledge, there is no proof that the scaling ansatz of Eq. (4) is a correct description and, even if it is, the only thing we can be sure of is $\theta \leq (d-2)/2$. All results are based on fitting data to the scaling form of Eq. (4) so one is attempting *both* to verify the scaling ansatz and to estimate a numerical value of θ . For any conclusion to be believable, the data must have both very small errors and fit Eq. (4) extremely well. The first requirement of highly accurate data is the most important as the estimate of θ depends on this. Assuming that $E_a(L)$ and $E_b(L)$ can be determined exactly for each sample, then $\Delta E(L)$ is also known exactly for each sample and the errors in $\langle \Delta E(L) \rangle$ are $O(N^{-1/2}L^{d-1})$ where N is the number of samples of size L in d dimensions. If the energy minima $E_{a,b}(L)$ are not found exactly, a crude estimate of the errors in $\Delta E(L)$ is $O(N^{-1/2}L^d)$ but this is certainly too low as failure of the algorithm to find the true minima because of being trapped in a metastable state of energy $E > E_0$ will cause systematic errors of unknown magnitude. Empirically, we find that this can readily cause errors larger than $\langle \Delta E(L) \rangle$ which makes the data point useless. This is most likely to happen for large L because the CPU time required grows uncontrollably, as do the errors, so the large L data becomes unreliable. This technical difficulty limits the accessible sizes L to small values as one must keep errors in individual data points small.

We are forced to conclude that the accessible sizes L are limited by the necessity of finding essentially exact global energy minima of each of a number N of samples subject to certain, yet to be defined, BC. To our knowledge, there is no algorithm applicable to the systems of interest which will find exact minima in polynomial time, such as the branch and cut algorithm³⁵ for the 2D Ising spin glass or numerically exact combinatorial optimization algorithms³⁶ for gauge and vortex glass models in the infinite screening limit, so we have to live with the fact that our problem is NP complete and the required CPU time explodes as L increases. We use simulated annealing^{37,38} to estimate the lowest energies, which seems considerably more efficient than simple quenching to $T=0$, but we are unable to go beyond $L=7$ in 3D and $L=10$ in 2D. We wish to extract the stiffness exponent θ from the scaling ansatz of Eq. (4) with a single power law and this makes sense only if the errors on individual data points are very small and the fit to the assumed scaling form is extremely good. In our opinion, the only sensible strategy is to obtain very accurate estimates of $\langle \Delta E(L) \rangle$ for the limited sizes L which are feasible for the computer power available.

In the phase representation of Eq. (1), the configuration space to be searched for the global energy minima $E_{a,b}(L)$ is rather large as the phases $\theta_i \in (0, 2\pi]$ are continuous variables. Searching this space in finite time is not feasible as most of the allowed configurations of the θ_i are not even local energy minima. It is well known that random XY models with a Hamiltonian of Eq. (1) can be written in a Coulomb gas (CG) or vortex representation via a duality transformation^{40,41,33} which leaves the partition function invariant. This expresses the Hamiltonian in terms of charge or vortex configurations which are *already* local energy minima.^{42,43} Thus, a reformulation of the Hamiltonian of Eq. (1) as a CG performs a partial minimization. A further minimization of the CG Hamiltonian corresponds to searching the much smaller space of local minima. Reformulating the problem of Eq. (1) including the BC in CG language is clearly a worthwhile exercise as it dramatically reduces the number of configurations over which we have to minimize, despite introducing long-range Coulomb interactions between vortices. The transformation is carried out in Sec. III for the model of Eq. (1) in 3D.

The final problem is to define what is meant by a domain wall and the BC needed to induce a wall in a finite system of size L in d dimensions. We imagine the system of Eq. (1) on a torus in 2D or a hypertorus in 3D, which corresponds to imposing periodic BC in the phases $\theta_{\mathbf{i}+L\hat{\mathbf{e}}_\mu} = \theta_{\mathbf{i}}$ where $\hat{\mathbf{e}}_\mu$ is a unit vector in the direction $\mu=x, y, \dots, d$ and $\mathbf{i}=(i_x, \dots, i_d)$ with $i_\mu=(1, 2, \dots, L)$. The phases at corresponding sites (i, j) on opposite faces are coupled by some interaction $\tilde{V}(\theta_i, \theta_j, A_{ij})$ which may be regarded as defining the BC. In principle, the GS is obtained by minimizing the energy with respect to the L^d bulk variables θ_i and all forms of \tilde{V} . This program is beyond our ability and we restrict ourselves to those \tilde{V} which induce a spin or chiral defect, which are related to the continuous and reflection symme-

tries respectively. To impose a spin defect, we choose $\tilde{V} = V(\theta_i - \theta_j - A_{ij})$. The plaquettes between the opposite faces are indistinguishable from the others and play no special role. We therefore keep fixed the frustrations $f_{\mathbf{r}} = \sum_{\square \mathbf{r}} A_{ij}/2\pi$ where the sum is over the bonds in a clockwise direction of the elementary plaquette centered at \mathbf{r} . We still have the freedom to add Δ_μ to every bond in the μ direction between opposite faces which imposes a global twist Δ_μ in the phase around a loop circling the hypertorus in the direction μ . This is equivalent to a gauge transformation $A_{ij} \rightarrow A_{ij} + \Delta_\mu/L$ on every bond ij in the direction μ . The lowest energy, $E_0(\Delta_\mu)$, is 2π periodic in Δ_μ with a minimum at some Δ_μ^0 which depends on the sample. To induce a spin domain wall normal to \mathbf{x} , one changes the twists from their best twist (BT) values $\Delta_\mu^0 \rightarrow \Delta_\mu^0 + \pi\delta_{\mu,x}$. The minimum energy subject to this constraint gives $E_{sD}(L)$, the energy of the system of size L containing an extra spin defect. Note that $E_{sD} \geq E_0$ for every sample but E_0 is not necessarily the absolute minimum as some other functional form of \tilde{V} may give a lower energy. However, even if E_0 is not the true GS energy but is the energy of a state with some excitation from the GS, this method of inducing a spin defect ensures that any excitation in the BT configuration will also be present in the state with an extra spin domain wall so that $\Delta E_s^{BT}(L) \equiv E_{sD}(L) - E_0(L) \geq 0$ is not affected by these. It is convenient, but not necessary, to define the spin defect energy by a twist of π from the BT value Δ_x^0 . This choice yields the maximum defect energy $\Delta E(L)$. Any other choice $0 < \epsilon \leq \pi$ yields the same spin stiffness exponent θ_s^{BT} defined by

$$\langle \Delta E_s^{BT}(L, \epsilon) \rangle = \mathcal{A}(\epsilon) L^{\theta_s^{BT}}. \quad (5)$$

The size ϵ of the twist from the BT value Δ_x^0 affects only the amplitude $\mathcal{A}(\epsilon)$ which is a maximum at $\epsilon = \pi$.

A chiral domain wall is induced by imposing reflective BC,²⁸ which means that corresponding sites (ij) on opposite faces are connected by interactions $\tilde{V} = V(\theta_i + \theta_j - A_{ij})$ which is equivalent to a reflection of the spins about some axis. In principle, one follows the procedure for a spin domain wall to obtain the chiral defect energy $\Delta E_c = E_{cD} - E_0$ where E_{cD} is the minimum energy of the system with these modified interactions \tilde{V} connecting opposite faces. However, there is no reason to expect $E_{cD} > E_0$ as the BC defining E_0 may trap a chiral defect in some samples and, in such cases, the modified interactions \tilde{V} will cancel the chiral defect to give $E_{cD} < E_0$. This phenomenon has been observed previously in numerical simulations of the XY spin glass.^{28,30} We therefore define the chiral defect energy as $\Delta E_c^{BT}(L) \equiv |E_{cD}(L) - E_0(L)|$ and the chiral stiffness exponent θ_c^{BT} by a finite-size scaling ansatz analogous to Eq. (5),

$$\langle \Delta E_c^{BT}(L) \rangle \sim L^{\theta_c^{BT}}. \quad (6)$$

Note that the Hamiltonian of Eq. (1) is truly invariant under reflection $\theta_i \rightarrow -\theta_i$ in the XY spin-glass case when $A_{ij} = 0, \pi$ as $A_{ij} = \pm \pi$ are equivalent. The Hamiltonian with uniform distribution of A_{ij} such as a gauge glass is not truly

invariant under reflection which would also require $A_{ij} \rightarrow -A_{ij}$ for the Hamiltonian to be invariant as it lacks the reflection symmetry.

However, in an XY spin glass there are two possible types of order, spin glass order and a chiral glass order each with their own stiffness exponent of Eq. (5) and Eq. (6). Recently, an important prediction was made that $\theta_s = \theta_c < 0$ for an XY spin glass in dimension $d < d_l$ where d_l is the lower critical dimension.³³ Although not rigorous, the arguments are very plausible and supported by analytic calculations on simple one-dimensional systems in which $\theta_s = \theta_c$ exactly.^{31,32} There is a notable lack of analytic results in this field with which to test numerical simulations and to our knowledge this is the only one existing at present. We have checked our numerical method in $d = 2 < d_l$ and get agreement with the analytic prediction that $\theta_s = \theta_c = -0.37 \pm 0.015$ to within numerical uncertainty.⁴⁴ Assuming the conjecture³³ is correct, this agreement gives some confidence in our definition of domain-wall energies as discussed above and in our numerical method in $d = 3$ using the CG representation. There is no analogous equality of the stiffness exponents in an XY spin glass for $d = 3 > d_l$ so we do not attempt to estimate θ_c in 3D but concentrate on the spin stiffness exponent θ_s . Also, at present, we are unable to derive an expression for a CG Hamiltonian with reflective boundary conditions in $d = 3$. All previous work on the XY gauge glass^{5,6,8,9} and on the XY spin glass²⁸⁻³⁰ using the $T = 0$ DWRG method have used different definitions for domain wall energies. Minimization with respect to the global twists Δ_μ is omitted, the lowest energy with $\Delta_\mu = 0$ is called E_p , and the lowest energy with $\Delta_\mu = \pi$ is called E_{ap} . Neither of these BC is compatible with the GS configuration, as both must induce some excitation from E_0 . Nevertheless, the spin defect energy is defined by $\Delta E_s^{RT} \equiv |E_{ap} - E_p|$ and the spin stiffness exponent θ_s^{RT} by

$$\langle \Delta E_s^{RT}(L) \rangle \sim L^{\theta_s^{RT}}. \quad (7)$$

We call this a random twist (RT) measurement since, for a particular sample, the twists $\Delta_\mu = 0, \pi$ are two arbitrary random choices relative to the best twist Δ_μ^0 , which is the twist which yields the lowest energy. In a uniform ferromagnet, $\Delta_\mu^0 = 0$ which is realized by periodic BC and $\Delta_\mu^0 + \pi$ by antiperiodic BC. For a particular realization of randomness, Δ_μ^0 is the analog of periodic BC in a uniform ferromagnet.

Recently, the 2D Ising spin glass has been studied by a closely related defect energy scaling method.^{45,46} At first sight, this is a much simpler system than the XY spin and gauge glasses of this work but it turns out that the 2D Ising spin glass is a very subtle system whose properties depend on the form of the distribution of couplings. In fact, the most recent study⁴⁶ computes exact domain wall energies ΔE for systems with $L \leq 480$ by means of a minimum-weight perfect matching algorithm and finds that $\theta = -0.27$ for a Gaussian distribution of coupling strengths but $\theta = 0$ for a $\pm J$ distribution, implying that 2D is the lower critical dimension and $T_c = 0$.^{47,48} These results are similar to the earlier study⁴⁵ which was restricted to $L \leq 24$ with a cluster Monte Carlo technique and is less accurate.

III. TRANSFORMATION TO COULOMB GAS REPRESENTATION

In this section, we discuss the CG representation of the Hamiltonian of Eq. (1) including all finite-size contributions. This representation parametrizes the energy in terms of the topological excitations on a torus in 2D and a hypertorus in 3D and includes global excitations which wind around the whole hypertorus. These latter excitations are very important for a finite system and are vital for finite-size scaling considerations when one is limited to small system sizes L . Also, every allowed configuration of topological excitations is a local energy minimum,^{42,43} since spin-wave excitations decouple from the vortex excitations, which allows us to obtain a more accurate estimate of energy minima than using the phase representation of Eq. (1) with the limited CPU time available. The transformation of the two-dimensional XY model to the CG representation including boundary contributions has been discussed in detail in earlier works.^{33,40,41} In this section, we use the method of Ney-Nifle and Hilhorst³³ to transform to the CG representation in 3D.

We first replace the potential $V(\phi)$ in Eq. (1) by a piecewise parabolic potential which is equivalent to a Villain³⁹ potential at $T = 0$. The partition function for a $L \times L \times L$ gauge glass model in 3D is

$$Z = \int_{-\pi}^{+\pi} \prod_i d\theta_i \sum_{\{n_{ij}\}} \exp \left[-\beta J \sum_{\langle ij \rangle} (\theta_{ij} - A_{ij})^2 \right], \quad (8)$$

where $\theta_{ij} \equiv \theta_i - \theta_j - 2\pi n_{ij}$ and where $n_{ij} = -n_{ji}$ are integers on the bond $\langle ij \rangle$. By choosing one phase, θ_0 , as a reference, the partition function can be written as

$$\begin{aligned} Z = & \int_{-\pi}^{\pi} d\theta_0 \int_{-\infty}^{\infty} \prod_{\langle ij \rangle} d\theta_{ij} \exp \left[-\beta J (\theta_{ij} - A_{ij})^2 \right] \prod_{\mathbf{r}} \\ & \times \delta \left(\sum_{\square \mathbf{r}_{xy}} \theta_{ij} \text{mod} 2\pi \right) \delta \left(\sum_{\square \mathbf{r}_{yz}} \theta_{ij} \text{mod} 2\pi \right) \prod_{\mathbf{r}} \\ & \times \delta \left(\sum_{\square \mathbf{r}_{zx}} \theta_{ij} \text{mod} 2\pi \right) \delta \left(\sum_{\square \mathbf{x}} \theta_{ij} \text{mod} 2\pi \right) \prod_{\mathbf{r}} \\ & \times \delta \left(\sum_{\square \mathbf{y}} \theta_{ij} \text{mod} 2\pi \right) \delta \left(\sum_{\square \mathbf{z}} \theta_{ij} \text{mod} 2\pi \right). \end{aligned} \quad (9)$$

Here, \mathbf{r} is the coordinate of the center of an elementary cube of the original lattice which corresponds to the coordinate of a dual lattice site. \mathbf{r}_{xy} is the coordinate of the center of the elementary plaquette in the xy plane and similarly for \mathbf{r}_{yz} and \mathbf{r}_{zx} . Note that for a 2D system in the xy plane, \mathbf{r}_{xy} are the dual lattice sites \mathbf{r} . Since each cube has six faces (plaquettes), each of which is shared by two adjacent cubes, to each dual lattice site \mathbf{r} we assign three independent plaquettes with centers at \mathbf{r}_{xy} , \mathbf{r}_{yz} , and \mathbf{r}_{zx} as shown in Figs. 1 and 2. $\sum_{\square \mathbf{r}_{xy}} \theta_{ij}$ is the circulation of θ_{ij} round the plaquette in the xy plane of the cube at \mathbf{r} ,

$$\sum_{\square \mathbf{r}_{xy}} \theta_{ij} \equiv [\theta_{ij_x} + \theta_{j_x i'} - \theta_{j_y i'} - \theta_{ij_y}], \quad (10)$$




FIG. 1. An elementary cube of the original lattice with vertices at i, j_μ . The dual lattice site \mathbf{r} is at the center of the cube and the faces associated with \mathbf{r} have centers at \mathbf{r}_{xy} , \mathbf{r}_{yz} and \mathbf{r}_{zx} . The bond variables A_{ij_μ} , relabeled as A_r^μ , are associated with the cube at \mathbf{r} .

and $\sum_{\square_x} \theta_{ij}$ is the circulation of θ_{ij} along an arbitrary loop in the x direction around the hypertorus,

$$\sum_{\square_x} \theta_{ij} \equiv \sum_{i_x=1}^L \theta_{(i_x, i_y, i_z), (i_x+1, i_y, i_z)}, \quad (11)$$

where i_μ with $\mu=x, y, z$ is the coordinate of the original lattice site and i_y and i_z are fixed. Other summations are defined similarly. Note that one needs to consider only one global loop on the hypertorus in each direction. Circulations around other global loops can be expressed in terms of circulations around any three chosen global loops and around elementary plaquettes. It is clear from the definition of θ_{ij} and the periodic boundary condition imposed on the θ_i that these circulations are integer multiples of 2π . Since the delta functions can be rewritten as follows,




FIG. 2. Graphical explanation of our symbols. \mathbf{r} denotes the center of the cube and light solid lines join original lattice sites.




FIG. 3. The $L \times L \times L$ system is represented by a cube. The thick lines are our choices of the three global loops around the whole system.

$$\delta\left(\sum_{\square_{xy}} \theta_{ij} \bmod 2\pi\right) = \frac{1}{2\pi} \sum_{n_r^z=-\infty}^{\infty} \exp\left[in_r^z \sum_{\square_{xy}} \theta_{ij}\right],$$

$$\delta\left(\sum_{\square_x} \theta_{ij} \bmod 2\pi\right) = \frac{1}{2\pi} \sum_{n_x=-\infty}^{\infty} \exp\left[in_x \sum_{\square_x} \theta_{ij}\right],$$

the partition function now becomes

$$\begin{aligned} Z &= Z_0 \sum_{\mathbf{n}} \sum_{\{\mathbf{n}_r\}} \int_{-\infty}^{\infty} \prod_{\langle ij \rangle} d\theta_{ij} \exp[-\beta J (\theta_{ij} - A_{ij})^2] \\ &\times \exp\left[\sum_{\mathbf{r}} \left(in_r^x \sum_{\square_{yz}} \theta_{ij} + in_r^y \sum_{\square_{zx}} \theta_{ij} + in_r^z \sum_{\square_{xy}} \theta_{ij}\right)\right] \\ &\times \exp\left[in_x \sum_{\square_x} \theta_{ij} + in_y \sum_{\square_y} \theta_{ij} + in_z \sum_{\square_z} \theta_{ij}\right], \quad (12) \end{aligned}$$

where n_r^μ and n_μ with $\mu=x, y, z$ are integers, and $\sum_{\mathbf{n}} \equiv \prod_{\mu} \sum_{n_\mu}$ and $\sum_{\{\mathbf{n}_r\}} \equiv \prod_{\mu} \sum_{n_r^\mu}$. The sum $\sum_{\mathbf{r}}$ is over the dual lattice sites \mathbf{r} at the centers of the elementary cubes and Z_0 is an unimportant constant. To perform the integrations over $\{\theta_{ij}\}$, we choose the three global loops around the hypertorus as shown in Fig. 3. To deal with these global loops mathematically, we introduce the following quantities,

$$\delta_{\mathbf{r}}^x \equiv \begin{cases} 1 & \text{if } \mathbf{r} = (x, 1, 1) \text{ with } x = 1, 2, \dots, L, \\ 0 & \text{otherwise,} \end{cases} \quad (13)$$

similarly for $\delta_{\mathbf{r}}^y$ and $\delta_{\mathbf{r}}^z$. For example, the cubes at $\mathbf{r} = (x, 1, 1)$ have a part of the global loop in the x direction.

The definition of A_r^μ associated with the cube at \mathbf{r} is also shown in Fig. 1. As the plaquettes, one can assign three independent A_r^μ to each cube. After relabeling A_{ij} by A_r^μ , performing the integrations in Eq. (12) over θ_{ij} , the partition function becomes

$$\begin{aligned} Z &= Z_0 \sum_{\mathbf{n}, \mathbf{n}_r} \exp\left[-\frac{1}{4\beta J} \sum_{\mathbf{r}} \sum_{\alpha} \{(\nabla \times \mathbf{n}_r)_\alpha + \delta_{\mathbf{r}}^\alpha n_\alpha\}^2\right] \\ &\times \exp\left[-i \sum_{\mathbf{r}} \sum_{\alpha} \{(\nabla \times \mathbf{n}_r)_\alpha + \delta_{\mathbf{r}}^\alpha n_\alpha\} A_r^\alpha\right]. \quad (14) \end{aligned}$$

We use the following notation for the discrete derivative,

$$(\nabla \times \mathbf{n}_r)_x \equiv (n_r^y - n_{r-\hat{z}}^y) - (n_r^z - n_{r-\hat{y}}^z) \quad (15)$$

and similarly for other components of $\nabla \times \mathbf{n}_r$. Note that, when $\mathbf{r} = (1, y, z)$, then $\mathbf{r} - \hat{\mathbf{x}} = (L, y, z)$ due to periodic boundary conditions, and similarly for $\mathbf{r} - \hat{\mathbf{y}}$ and $\mathbf{r} - \hat{\mathbf{z}}$. Applying the Poisson-summation formula to n_r^μ ,

$$\sum_{n=-\infty}^{\infty} f(n) = \sum_{q=-\infty}^{\infty} \int_{-\infty}^{\infty} du e^{2\pi i q u} f(u), \quad (16)$$

the partition function of Eq. (14) becomes

$$\begin{aligned} Z = Z_0 \sum_{\mathbf{n}} \sum_{\{\mathbf{q}_r\}} \int_{-\infty}^{\infty} \prod_{\mathbf{r}} d\mathbf{u}_r \exp \left[2\pi i \sum_{\mathbf{r}} \mathbf{q}_r \cdot \mathbf{u}_r \right] \\ \times \exp \left[-\frac{1}{4\beta J} \sum_{\mathbf{r}} \sum_{\alpha=x,y,z} \{(\nabla \times \mathbf{u}_r)_\alpha + \delta_r^\alpha n_\alpha\}^2 \right] \\ \times \exp \left[-i \sum_{\mathbf{r}} \sum_{\alpha=x,y,z} \{(\nabla \times \mathbf{u}_r)_\alpha + \delta_r^\alpha n_\alpha\} A_r^\alpha \right]. \quad (17) \end{aligned}$$

It is convenient to change integration variables to v_r^μ to make Eq. (17) more symmetric,

$$\begin{aligned} v_r^x &\equiv u_r^x - \frac{y}{2L} n_z + \frac{z}{2L} n_y, \\ v_r^y &\equiv u_r^y - \frac{z}{2L} n_x + \frac{x}{2L} n_z, \\ v_r^z &\equiv u_r^z - \frac{x}{2L} n_y + \frac{y}{2L} n_x, \end{aligned} \quad (18)$$

where $\mathbf{r} = (x, y, z)$. The partition function of Eq. (17) now becomes

$$\begin{aligned} Z = Z_0 \sum_{\mathbf{n}} \sum_{\{\mathbf{q}_r\}} \int_{-\infty}^{\infty} \prod_{\mathbf{r}} dv_r \exp \left[2\pi i \sum_{\mathbf{r}} \mathbf{q}_r \cdot \mathbf{v}_r \right] \\ \times \exp \left[\frac{\pi i}{L} \sum_{\mathbf{r}} \mathbf{q}_r \cdot (\mathbf{r} \times \mathbf{n}) \right] \\ \times \exp \left[-\frac{1}{4\beta J} \sum_{\mathbf{r}} \sum_{\alpha=x,y,z} \left\{ (\nabla \times \mathbf{v}_r)_\alpha + \frac{n_\alpha}{L} \right\}^2 \right] \\ \times \exp \left[-i \sum_{\mathbf{r}} \sum_{\alpha=x,y,z} \left\{ (\nabla \times \mathbf{v}_r)_\alpha + \frac{n_\alpha}{L} \right\} A_r^\alpha \right]. \quad (19) \end{aligned}$$

The terms linear in n_μ in the exponent of Eq. (19) vanish due to the periodic boundary condition $\sum_{\mathbf{r}} (v_r^x - v_{r-\hat{y}}^x) = 0$. Equation (19) can be simplified by introducing frustration variables f_r^μ at site \mathbf{r} ,

$$f_r^x \equiv -\frac{1}{2\pi} \sum_{\square_{r,yz}} A_{ij} = \frac{1}{2\pi} (A_r^y - A_{r+\hat{z}}^y + A_{r+\hat{y}}^z - A_r^z), \quad (20)$$

from which f_r^y and f_r^z are obtained by cyclic permutation of xyz . After some algebra, the partition function of Eq. (19) becomes a form suitable for integration over v_r^μ ,

$$\begin{aligned} Z = Z_0 \sum_{\mathbf{n}} \sum_{\{\mathbf{q}_r\}} \int_{-\infty}^{\infty} \prod_{\mathbf{r}} dv_r \exp \left[\frac{2\pi i}{2L} \{\mathbf{q}_r \cdot (\mathbf{r} \times \mathbf{n})\} \right] \\ \times \exp \left[-\frac{1}{4\beta J} \sum_{\mathbf{r}} \sum_{\alpha=x,y,z} \{(\nabla \times \mathbf{v}_r)_\alpha\}^2 \right] \\ \times \exp \left[2\pi i \sum_{\mathbf{r}} (\mathbf{q}_r - \mathbf{f}_r) \cdot \mathbf{v}_r \right]. \quad (21) \end{aligned}$$

To evaluate the integrals over v_r^μ in Eq. (21) it is convenient to take the Fourier transform

$$\mathbf{v}_r = L^{-\frac{3}{2}} \sum_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}} \tilde{\mathbf{v}}(\mathbf{k}), \quad (22)$$

where $\mathbf{k} = (k_x, k_y, k_z)$ and $k_i = 2\pi/L m_i$ with $m_i = 0, 1, \dots, L-1$. We also decompose $\tilde{\mathbf{v}}(\mathbf{k})$ into longitudinal, $\tilde{v}_L(\mathbf{k})$, and transverse, $\tilde{v}_T(\mathbf{k})$ components where

$$\tilde{v}_L^\alpha(\mathbf{k}) = \sum_{\beta} \frac{(1 - e^{ik_\alpha})(1 - e^{-ik_\beta})}{\lambda_{\mathbf{k}}} \tilde{v}^\beta(\mathbf{k}) \equiv \sum_{\beta} L_{\alpha\beta} \tilde{v}^\beta(\mathbf{k}),$$

$$\tilde{v}_T^\alpha(\mathbf{k}) \equiv \sum_{\beta} (\delta_{\alpha\beta} - L_{\alpha\beta}) \tilde{v}^\beta(\mathbf{k}) \equiv \sum_{\beta} T_{\alpha\beta} \tilde{v}^\beta(\mathbf{k}),$$

$$\lambda_{\mathbf{k}} = 6 - 2 \cos k_x - 2 \cos k_y - 2 \cos k_z \quad (23)$$

define the longitudinal and transverse projection operators $L_{\alpha\beta}(\mathbf{k})$ and $T_{\alpha\beta}(\mathbf{k})$. In Fourier space, the integration over \mathbf{v}_r in Eq. (21) is straightforward,

$$\begin{aligned} \int \prod_{\mathbf{k}} d\tilde{v}_L(\mathbf{k}) d\tilde{v}_T(\mathbf{k}) \exp \left[-\frac{1}{4\beta J} \sum_{\mathbf{k}} \lambda_{\mathbf{k}} \left| \tilde{v}_T(\mathbf{k}) \right|^2 \right] \\ \times \exp \left[2\pi i \sum_{\mathbf{k}} \{ \tilde{\mathbf{p}}_T(\mathbf{k}) \cdot \tilde{\mathbf{v}}_T(\mathbf{k}) + \tilde{\mathbf{p}}_L(\mathbf{k}) \cdot \tilde{\mathbf{v}}_L(\mathbf{k}) \} \right], \quad (24) \end{aligned}$$

where $\tilde{\mathbf{p}}(\mathbf{k}) = \mathbf{q}(\mathbf{k}) - \mathbf{f}(\mathbf{k})$. Integration over $\tilde{v}_L(\mathbf{k})$ gives $\tilde{\mathbf{p}}_L(\mathbf{k}) = 0$. In real space, this is the condition whereby the discrete divergence of the vorticity (charge) \mathbf{q}_r at any dual lattice site \mathbf{r} obeys $\nabla \cdot \mathbf{q}_r = 0$ since $\nabla \cdot \mathbf{f}_r = 0$. The integration over the transverse components $\tilde{v}_T(\mathbf{k})$ are simple Gaussian integrals and are easily performed. Integration over $\tilde{v}_L(0)$ and $\tilde{v}_T(0)$ yields the neutrality condition

$$\tilde{\mathbf{p}}(0) = \sum_{\mathbf{r}} \mathbf{p}_r = 0. \quad (25)$$

The final step in this rather technical derivation of the Hamiltonian in the CG representation is to apply the Poisson-summation formula of Eq. (16) to eliminate the n_μ in favor of global vortices or charges $q_{\mu 1}$. The Gaussian integrations yield the partition function

$$Z = Z_0 \sum_{\{\mathbf{q}_r\}} \sum_{q_\mu} \exp[-\beta H(\mathbf{q}_r, \mathbf{f}_r, q_{\mu 1}, f_{\mu 1})]. \quad (26)$$

The Hamiltonian H is identified as

$$\begin{aligned} H = & (2\pi)^2 J \sum_{\mathbf{r}, \mathbf{r}'} (\mathbf{q}_r - \mathbf{f}_r) \cdot (\mathbf{q}_{r'} - \mathbf{f}_{r'}) G(\mathbf{r} - \mathbf{r}') \\ & + \frac{J}{2L} \left\{ \frac{\pi}{L} \sum_{\mathbf{r}} (z p_{\mathbf{r}}^y - y p_{\mathbf{r}}^z) + Q_x \right\}^2 + \frac{J}{2L} \left\{ \frac{\pi}{L} \sum_{\mathbf{r}} (x p_{\mathbf{r}}^z \right. \\ & \left. - z p_{\mathbf{r}}^x) + Q_y \right\}^2 + \frac{J}{2L} \left\{ \frac{\pi}{L} \sum_{\mathbf{r}} (y p_{\mathbf{r}}^x - x p_{\mathbf{r}}^y) + Q_z \right\}^2, \quad (27) \end{aligned}$$

where $G(\mathbf{r}) = L^{-3} \sum_{\mathbf{k} \neq 0} [\exp(i\mathbf{k} \cdot \mathbf{r}) - 1] / \lambda_{\mathbf{k}}$ is the lattice Green's function on a simple cubic lattice in 3D with periodic BC and the quantities Q_μ are

$$Q_x = \pi \sum_{\mathbf{r}} (z p_{\mathbf{r}}^y \delta_{y,1} - y p_{\mathbf{r}}^z \delta_{z,1}) + 2\pi L(q_{x1} - f_{x1}), \quad (28)$$

with Q_y and Q_z obtained from Q_x by cyclic permutations of xyz . In Eq. (28), f_{x1} is the circulation of A_{ij} along the chosen global loop around the hypertorus in the x direction,

$$2\pi f_{x1} = \sum_{x=1}^L A_{\mathbf{r}=(x,1,1)}^x \quad (29)$$

and similarly for f_{y1} and f_{z1} . The integers $q_{\mu 1}$ are interpreted as circulations of the phase around the three independent global loops encircling the hypertorus. Equations (27) and (28) are the main results of this section. These expressions give the energy of a system of size L on a hypertorus. To find the minimum energy E_0 the Hamiltonian $H = H(\mathbf{q}_r, \mathbf{f}_r, q_{\mu 1}, f_{\mu 1})$ is minimized with respect to the bulk integer valued vector charges \mathbf{q}_r , the integer valued global winding numbers $q_{\mu 1}$, and the global frustrations $f_{\mu 1}$. In the case of the XY spin glass the $q_{\mu 1} - f_{\mu 1}$ of Eq. (28) are restricted to be integer or half integer while for the gauge glass this can have any real value. Note that minimizing with respect to q_{x1} and f_{x1} is exactly minimizing with respect to the twist Δ_x and the best twist Δ_μ^0 corresponds to $f_{\mu 1}^0$, the value of $f_{\mu 1}$ at the energy minimum. Adding a global twist Δ_μ to the phases is exactly equivalent to changing the global frustration from its original value $f_{\mu 1} \rightarrow f_{\mu 1} + \Delta_\mu / 2\pi$. As discussed in Sec. II, a spin domain wall is induced by $\Delta_x^0 \rightarrow \Delta_x^0 + \pi$ which, in the CG representation, is $f_{x1}^0 \rightarrow f_{x1}^0 + 1/2$. Note that the frustrations \mathbf{f}_r are given in terms of the random A_{ij} and are kept fixed during the minimization.

To confront our numerical results with the only existing analytic prediction³³ we need expressions for both spin and chiral domain wall energies in dimension $d = 2 < d_l$. The CG representation of the XY spin and gauge glasses including all finite size corrections has been known for some time in 2D,^{41,33} and, for the sake of completeness, we quote the necessary results below. To study the spin stiffness, we join corresponding sites on opposite faces by $\tilde{V} = V(\theta_i - \theta_j - A_{ij})$ and the CG Hamiltonian is

$$H = (2\pi)^2 J \sum_{\mathbf{r}, \mathbf{r}'} (q_r - f_r) G(\mathbf{r} - \mathbf{r}') (q_{r'} - f_{r'}) + \frac{J}{L^2} (\sigma_x^2 + \sigma_y^2), \quad (30)$$

where

$$\begin{aligned} \sigma_x = & -2\pi \left[L(q_{x1} - f_{x1}) + \sum_{\mathbf{r}} (q_r - f_r) y \right], \\ \sigma_y = & -2\pi \left[L(q_{y1} - f_{y1}) - \sum_{\mathbf{r}} (q_r - f_r) x \right], \\ G(\mathbf{r}) = & \frac{1}{L^2} \sum_{\mathbf{k} \neq 0} \frac{e^{i\mathbf{k} \cdot \mathbf{r}} - 1}{4 - 2 \cos k_x - 2 \cos k_y}. \quad (31) \end{aligned}$$

Here, as in 3D, $\mathbf{r} = (x, y)$ represents the coordinates of the dual lattice sites and $G(\mathbf{r})$ is the lattice Green's function with periodic BC on the θ_i . From Eq. (8), we see that the difference of a factor 2 in the prefactor of Eq. (30) from other works is in the coupling constant J . Also note that the Hamiltonian of Eq. (30) describes the XY spin glass when the frustrations \mathbf{f}_r and the global frustrations $f_{\mu 1}$ are restricted to $(0, 1/2)$. In the gauge glass, they can have any real value. As in 3D, the topological charges q_r are integers as are $q_{\mu 1}$ because of the periodic BC in the θ_i .

The last piece of information we need is the CG representation of the Hamiltonian of the XY spin glass in 2D with a chiral domain wall imposed. This is to be found in the paper of Ney-Nifle and Hilhorst.³³ A single chiral domain wall is induced by joining opposite faces by interactions $\tilde{V} = V(\theta_i + \theta_j - A_{ij})$ which is equivalent to imposing reflective BC.^{28,33} In turn, this is equivalent to doubling the size in (say) the x direction to a $2L \times L$ lattice in which the extra half is a charge conjugated image of the other. This system has two chiral domain walls with the Hamiltonian³³

$$H_R = 2\pi^2 J \sum_{\mathbf{r}, \mathbf{r}'} (q_r - f_r) \tilde{G}(\mathbf{r} - \mathbf{r}') (q_{r'} - f_{r'}), \quad (32)$$

where $\tilde{G}(\mathbf{r})$ is the Green's function for a $2L \times L$ square lattice with *periodic* BC and also with $q_{\mathbf{r}+L\hat{x}} = -q_{\mathbf{r}}$ and $f_{\mathbf{r}+L\hat{x}} = -f_{\mathbf{r}}$. Note that the sign reversal of the frustrations \mathbf{f}_r is not necessary for the spin glass because $f_r = \pm 1/2$ are equivalent.

The form of the energy of Eq. (32) is used in the simulations to estimate θ_c , the chiral stiffness exponent, as it is intuitively more transparent and more convenient than the corresponding expression with a single chiral domain wall.³³ Unfortunately, we have been unable to derive the analogous expression in 3D to Eq. (32) so we have no independent estimate of θ_c (Ref. 30) for the 3D XY spin glass.

IV. NUMERICAL METHOD

A. Minimization algorithm

In Sec. II we argued that it is necessary to find the energies $E_0(L)$ and $E_D(L)$ essentially exactly for every sample to control the errors in the *small* domain wall energy

$\langle \Delta E(L) \rangle$. To our knowledge, for the systems of interest no algorithm exists which can locate the global energy minima in polynomial time. We are left with two methods: (i) repeated simple quenches from an arbitrary initial configuration to $T=0$ followed by a downward slide to the nearest local minimum and (ii) simulated annealing³⁷ which is considerably more efficient.³⁸ By this we mean that, for the same CPU time, simulated annealing finds a lower energy than simple quenching.

We start with the system in some randomly chosen configuration and quench to some T_0 , determined by trial and error, for each size L . Then we do a Monte Carlo (MC) sweep through the system by inserting charges $q=1$ and $q'=-1$ on an arbitrary pair of nearest-neighbor sites of the dual lattice in 2D and accepting or rejecting the move according to conventional MC rules. This has the effect of inserting new charges, annihilating charges, or moving a charge by one lattice spacing while maintaining charge neutrality $\sum q_r=0$. In 3D, the elementary excitation is a loop of charge around an elementary square with vertices at dual lattice sites. This maintains $\nabla \cdot \mathbf{q}_r=0$. The closed loop of charge can lie in any of the three orthogonal planes of the cubic lattice and the charge can circulate around the loop in either direction, making six possibilities with the center of the loop at a fixed but randomly selected position. The temperature T is then reduced to $T_1=\alpha T_0$ with $\alpha<1$ whose value is again determined by trial and error and the procedure iterated a large number, N , of times to reach a lowest temperature $T_N=\alpha^N T_0 \approx 0$. Of course, the system may be trapped in a deep metastable well with barriers too high for the MC passes to overcome so the whole annealing sequence is repeated M times from different random initial configurations and the lowest energy out of all the NM trials is recorded. Again, this does not guarantee that the global minimum energy is found but this method does have a few checks built in. At the crudest level, the best twist condition $E_{sD}(L) \geq E_0(L)$ must be obeyed for each sample since $E_0(L)$ is, by construction, the lowest energy of the system subject to the BC given by the interactions $\tilde{V}=V(\theta_i-\theta_j - A_{ij})$ across opposite faces. $E_{sD}(L)$ is obtained by $f_{x1}^0 \rightarrow f_{x1}^0 + 1/2$ where $f_{\mu 1}^0$ is the value of $f_{\mu 1}$ which makes the boundary terms of Eqs. (27) and (30) vanish, corresponding to the best twist Δ_μ^0 . It is clear that $E_{sD}(L)$ is the energy of the system containing an extra spin domain wall compared to the system with energy $E_0(L)$. As discussed earlier, $E_0(L)$ is not necessarily the true GS energy as the system may contain some chiral domain walls. However, by construction, $E_{sD}(L)$ is the energy of the system with the *same* chiral defects and an extra spin defect.

If any sample violates the BT condition, clearly the annealing is not sufficient and one can either increase the number MN of annealing attempts or just discard that sample. Increasing MN involves a significant increase in CPU time particularly for the larger sizes L so the choice depends on the time available. However, even if the BT condition is satisfied for every sample, there is no guarantee that the lowest energies found are *true* global minima. To improve the chances that true minima are achieved, where possible we

did two independent simulations on identical samples with different pseudorandom number sequences and, if the same energy minima are found in both simulations, this is defined to be the true minimum. This procedure is very expensive in CPU time and our resources did not permit this last check to be performed for every sample, particularly for the largest systems. This check was done for at least a few randomly selected samples, except for $L=7$ in the 3D gauge glass. Averaging over disorder was performed over as many samples as possible with the aim of making the uncertainty in the mean domain-wall energy $\langle \Delta E(L) \rangle$ less than 3% which requires averaging over at least 10^3 samples. The first set of checks are done on every sample which makes it fairly probable that exact minima $E_0(L)$ and $E_D(L)$ are obtained. We can assume with some confidence that the uncertainty in $\langle \Delta E(L) \rangle$ is purely statistical and $O(N^{1/2})$. Averaging over about 10^3 samples leads to an acceptable uncertainty of about 3% in $\langle \Delta E(L) \rangle$.

B. XY spin glass in 2D

This particular system with a Hamiltonian of Eq. (1) in 2D is not particularly interesting in the sense that there exists no finite temperature transition. It has been known for a long time that $d_l > 2$ and both spin and chiral stiffness exponents are negative. However, the situation has been controversial due to the possible existence of different stiffness or correlation length exponents for spin- and chiral glass order in 2D. Previous estimates of the spin and chiral stiffness exponents are summarized by $\theta_s \approx 2\theta_c \approx -0.78$ based on extensive numerical simulations such as DWRG and finite temperature MC simulations.^{28-30,49,50} However, Ney-Nifle and Hilhorst³³ made a nonrigorous but very plausible conjecture based on analytic considerations that for dimension $d \leq d_l$, $\theta_s = \theta_c \leq 0$. This is supported by exact analytic results on simple models of (i) an XY spin glass on a ladder lattice³¹ where the common correlation length exponent $\nu = |\theta|^{-1} = 0.5263 \dots$ and (ii) the XY spin glass on a tube lattice³² with $\nu = 0.5564 \dots$. The key disagreement between numerical estimates and analytic theory is in the conjectured equality of θ_s and θ_c . If one makes the hypothesis that earlier estimates are in error, then the most likely reason is that the simulations are computing the wrong quantity when estimating one or both of the stiffness exponents. It is extremely unlikely that all the simulations are in error for simple technical reasons since all find the same values of θ_s and θ_c , but $\theta_s \neq \theta_c$.

To test this hypothesis, we have performed simulations of the 2D XY spin glass in the CG representation using Eq. (30) to estimate the spin stiffness exponent θ_s using both BT and RT measurements with the results $\theta_s^{BT} = -0.37 \pm 0.015$ and $\theta_s^{RT} = -0.76 \pm 0.015$.⁴⁴ These numbers were obtained from sizes $L=4, 5, 6, 7, 8$, and 10 averaging over 2560 samples for $L \leq 8$ and 1152 samples for $L=10$. As expected, the value of θ_s^{RT} agrees with all previous estimates,^{28,30} all of which use the RT measurement in some form. Both the BT and the RT data fit the scaling ansatz of Eq. (4) equally well and some other information is needed to decide which value of θ_s , if either, is correct. Both cannot be correct as both are




FIG. 4. Top to bottom: L dependence of ΔE_s^{BT} , ΔE_s^{RT} , ΔE_c^{RT} , and ΔE_c^{BT} for the 2D XY spin glass.

supposed to measure the same quantity. The necessary information is in the chiral stiffness exponent θ_c which we measure by simulating Eq. (32) on a $2L \times L$ system. Again, both BT and RT measurements were made with the same range of L and the same number of samples as for θ_s with the results $\theta_c^{BT} = -0.37 \pm 0.010$ and $\theta_c^{RT} = -0.37 \pm 0.015$.⁴⁴ At first sight it is surprising that both measurements give the same value for θ_c to within numerical uncertainty while the values of θ_s^{RT} and θ_c^{BT} differ by a factor of 2. Note that the boundary terms σ_α of Eq. (30) which contain the twist parameter $f_{\mu 1}$ vanish in the BT condition and such boundary terms are absent from Eq. (32). Thus, any measurement with a CG representation in which boundary contributions are absent is automatically a BT measurement. Since $\theta_c = \theta_s^{BT} \approx -0.37$ and $\theta_c \neq \theta_s^{RT} \approx -0.76$, assuming that the conjecture is correct, we conclude that the BT measurement yields a reasonable estimate of the true θ_s while the commonly used RT measurement yielding $\theta_s^{RT} \neq \theta_c$ is not an appropriate method for the small values of L accessible at present. Our simulation results are shown in Fig. 4. We do not understand what, if anything, θ_s^{RT} means despite the apparent excellent fit of $\langle \Delta E_s^{RT}(L) \rangle$ to the scaling ansatz, since this is the energy difference of the system subject to two random choices of BC and we can see no reason why it should scale as L^{θ_s} over decades of L .

V. RESULTS

Our investigation of the 2D XY spin glass establishes that the BT method of measuring domain wall energies can reproduce the one and only available analytic prediction³³ which is relevant for our purposes. This gives some encouragement to venture into areas where no such analytic guide exists. In this section, we report some new results on the spin stiffness exponent θ_s for the XY spin glass in 3D (Sec. V A), and the gauge glass in both 2D and 3D (Sec. V B). We also perform simulations on systems with varying strengths of disorder (Sec. V C) where we study the renormalization

group flows for both the coupling constants $J(L)$ and disorder strength $A(L)$ with an increasing length scale L . All our results are numerical and the exponent $\theta_s = \theta_s^{BT}$ is estimated by fitting estimates of $\langle \Delta E_s^{BT}(L) \rangle$ to the scaling ansatz of Eq. (4). The system sizes L are very small as they are severely constrained by the necessity of controlling the errors in both $E_0(L)$ and $E_D(L)$, which are estimated by independent simulations. However, the fit to the scaling ansatz is very good despite the very few data points for all cases we have studied and is comparable with the same procedure carried out on the 2D Ising ferromagnet. In this case, simulations for sizes $L=2-5$ are sufficient to reproduce the exact $\theta_s = 1$ to high accuracy.

A. XY spin glass in 3D

This system has been somewhat controversial for some years and is not yet settled. It has been believed that $d_l \geq 4$ for spin-glass order, and $d_l < 3$ for chiral glass order and that, in 2D and 3D, spin and chiral variables decouple and order separately.^{27-30,49,50} This allows for the widely accepted scenario that in 3D, spin-glass order sets in at $T_{SG}=0$ whereas chiral glass order sets in at $T_{CG}>0$. This scenario is based on MC simulations at finite T (Refs. 29 and 51) and on $T=0$ defect energy scaling^{28,29} using the RT method. Attempts to show rigorously that $T_{SG}=0$ in 3D (Ref. 52) fail if reflection symmetry is broken^{29,53,54} at finite T . The first cracks in this widely accepted scenario appeared recently when Maucourt and Gempel³⁰ published the results of a large scale defect energy scaling study of the 3D XY spin-glass model of Eq. (1). They used the RT method, as all previous defect energy scaling studies have done, with sizes $L \leq 12$ in 2D and $L \leq 8$ in 3D. Although the fit to the scaling ansatz is not good due to strong crossover effects and large uncertainties in the large L data which was used to estimate θ_s^{RT} , it is clear that both $\langle \Delta E_s^{RT}(L) \rangle$ and $\langle \Delta E_c^{RT}(L) \rangle$ increase with L which implies that there is both spin- and chiral glass order at sufficiently small $T>0$. However, as argued above a valid numerical method must yield $\theta_s = \theta_c$ in 2D while they obtain $\theta_s \approx 2\theta_c \approx -0.78$,³⁰ in agreement with other estimates.^{28,29} Finite T simulations seem to suffer from severe equilibration difficulties^{29,51} which makes any conclusions from them also suspect.

In view of the lack of reliable results for the 3D XY spin glass, we have done some preliminary simulations on very small systems with $L=2-6$ in the CG representation with Eq. (27) where $f_{\mu 1}=0,1/2$ in Eq. (28). Following the method outlined in Sec. IV B we estimate $\theta_s^{BT} = +0.10 \pm 0.03$ while the data for $\langle \Delta E_s^{RT}(L) \rangle$ clearly decreases with L for $L=2-4$, with roughly the same slope as found by Kawamura²⁹ for the same range of L . Our value $\theta_s = 0.10 \pm 0.03$ (Ref. 44) is to be compared with the recent estimate $\theta_s^{RT} = +0.052 \pm 0.03$ for $L=5-7$.³⁰ Although these values are close numerically and are equal to within error bars, we are more inclined to believe in the former number as we consider θ_s^{BT} to give the true spin stiffness exponent and we speculate that further points will also lie on the scaling ansatz and will significantly reduce the 30% uncertainty. The




FIG. 5. L dependence of ΔE_s^{BT} and ΔE_s^{RT} in 3D. The error in the $L=6$ point is due to the rather few samples. The solid line is power-law fit and the dotted line is a guide to the eye.

results are shown in Fig. 5. Unfortunately, we have no estimate for the chiral stiffness exponent θ_c^{BT} for the 3D XY spin glass as we have been unable to derive the 3D analog of Eq. (32) to estimate $\Delta E_c^{BT}(L)$ and $\Delta E_c^{RT}(L)$. However, because the numerical values of θ_c^{BT} and θ_c^{RT} are the same in 2D and equal to θ_s as required, we are reasonably safe in assuming that existing estimates^{29,30} of θ_c in 3D are fairly accurate. Even though our spin stiffness exponent $\theta_s^{BT} \approx +0.1$ has a rather large uncertainty, it suggests that the lower critical dimension for spin-glass order is $d_l < 3$, as for chiral order. This is to be expected from analytic arguments.³³

B. Gauge glass in 2D and 3D

We have also performed simulations on the gauge glass in the CG representation using Eq. (30) in 2D and Eq. (27) in 3D. The only differences to the spin glass are in the values of $f_{\mu 1}$ and f_r which can have any value in the interval $[-1/2, 1/2)$. The frustrations $f_r = -\sum_{\square} A_{ij}/2\pi$ are correlated random variables, since the A_{ij} are the independent random variables. Similarly, in 3D, the frustrations \mathbf{f}_r of Eq. (20) are correlated random variables with each component in the interval $[-1/2, 1/2)$. The three global frustrations $f_{\mu 1}$ can take any value in the same interval. In both spin and gauge glasses, the vorticities q_r^μ and $q_{\mu 1}$ are integers. We have not computed θ_c in either 2D or 3D for the gauge glass, mainly because we have been unable to obtain the appropriate expression for the Hamiltonian in the CG representation with reflective BC in 3D and we are unable to understand what information such a simulation would yield. The major reason for doing such a simulation in the 2D spin-glass case was to confirm that our procedure is a useful numerical method to estimate the actual spin stiffness exponent θ_s for both spin and gauge glasses in 2D and 3D. To compare with earlier work^{5,6,8,9} we have also estimated θ_s^{RT} by keeping $f_{\mu 1}$ fixed at some random value during the minimization, exactly as for the spin glass. The results of the simulations⁵⁵ in 2D for




FIG. 6. Size L dependence of domain wall energy in 2D. Both RT and BT measurements are shown. Solid lines are power law fits. Error bars are not shown if smaller than symbol size.

system sizes $L=2-6, 8$, and 10 are shown in Fig. 6 for $\langle \Delta E_s^{BT}(L) \rangle$ and $\langle \Delta E_s^{RT}(L) \rangle$. Averages were performed over about 10^3 samples for each size L . In this case, all the checks discussed in Sec. IV A were performed so we assume the errors are purely statistical. Both fit well to the scaling ansatz of Eq. (4) with very similar errors with the values $\theta_s^{BT} = -0.36 \pm 0.010$ and $\theta_s^{RT} = -0.45 \pm 0.015$. The latter value is consistent with all earlier estimates of θ_s ,^{5,6,9} which is not a surprise as all these were done using the RT method in some form. However, as argued in Sec. IV B, this is not an accurate estimate of θ_s so this number is *not* the $T=0$ spin stiffness exponent. A more accurate estimate of this is $\theta_s^{BT} = -0.36 \pm 0.010$ which is significantly larger than θ_s^{RT} , so that the 2D gauge glass has a longer correlation length $\xi(T) \sim T^{-1/|\theta_s|}$ than previously thought.

We have also obtained some estimates of θ_s in 3D by performing simulations⁵⁵ of the gauge glass in the CG representation using Eq. (27) with the distribution of frustrations \mathbf{f}_r appropriate for this case as determined by taking the A_{ij} uniformly distributed in $(-\pi, +\pi]$. The system sizes are $2 \leq L \leq 7$ with disorder averaging over 10^3 samples for $L \leq 5$, 300 for $L=6$, and 60 for the largest size $L=7$. The uncertainty in $\langle \Delta E_s^{BT}(L) \rangle$ for $L=7$ is very large, but this data point is included to check that it is consistent with the behavior deduced from the more reliable data of the smaller sizes. The results are shown in Fig. 7 for $\langle \Delta E_s(L) \rangle$ for the unscreened gauge glass using both BT and RT measurements and for a gauge glass in 3D with screened interactions, $\lambda < \infty$. The BT data fit the scaling form of Eq. (4) very well for sizes $L \leq 6$ with exponent $\theta_s^{BT} = +0.31 \pm 0.010$. If the $L=7$ point is also included in the fit, we obtain $\theta_s^{BT} = +0.30 \pm 0.015$. These errors in θ_s come from a naive least squares fit of the data to a straight line and should not be taken too seriously. The $L=7$ data is suspect because four samples violated the BT condition $\Delta E_s^{BT}(L) \geq 0$ out of a total of only 64, despite running a highly vectorized code for a few thousand CPU hours on a Cray J90. Time did not permit any




FIG. 7. L dependence of domain-wall energy in 3D. Bottom curve is RT measurement for unscreened interaction. All others are BT measurements. Topmost curve is unscreened case $L=2-7$. Other curves are screened interactions with λ decreasing from top to bottom. Solid line is a power-law fit and dotted lines are guides for the eye.

further checks for attaining the global energy minimum for $L=7$. We cannot be sure that the remaining 60 samples which did not violate the BT condition reached their global energy minima nor that the energies $E_{s,D}$ are determined sufficiently accurately. The error bar on the $L=7$ point in Fig. 7 assumes that the uncertainty in $\langle \Delta E_s^{BT}(L=7) \rangle$ is purely statistical and the true uncertainty is probably *much* larger. At least an order-of-magnitude more CPU time is needed for sufficient annealing to reach the true minima and to perform the additional simulations with different random number sequences to check that the minimization algorithm is successful. This is just for a single batch of 64 samples and to reduce the uncertainty to 3%, yet another order of magnitude of CPU time would be needed to average over 10^3 samples. This is totally out of reach with the computing resources available to us. What data we have is entirely consistent with the scaling form of Eq. (4) with $\theta_s \approx +0.30$ with no sign of any deviation from this form.

The behavior of $\langle \Delta E_s^{RT}(L) \rangle$ is also shown in Fig. 7 for sizes $L \leq 6$ which is very much like the data obtained by earlier simulations. This clearly does not fit the scaling form of Eq. (4) for these small values of L , but if one insists on extracting a value of θ_s^{RT} from the data, one obtains consistency with previous estimates⁶ for the spin stiffness exponent $\theta_s^{RT} \approx +0.05 \pm 0.05$. As can be seen from Fig. 7, this estimate has no meaning as the data clearly does not obey the scaling ansatz. In fact, as noticed by Maucourt and Gempel,⁹ $\langle \Delta E_s^{RT}(L) \rangle$ seems to start increasing with L for $L > 5$ but, as we argue earlier, this may, or may not, eventually scale as $\langle \Delta E_s^{BT}(L) \rangle$ for sufficiently large L . Speculation along these lines is fruitless until computers which are many orders-of-magnitude faster become available or until an analytic solution is found. We conclude that, for the *unscreened* gauge glass in 3D, the spin stiffness exponent $\theta_s = +0.31 \pm 0.010$,

which is considerably larger than earlier estimates and indicates that the lower critical dimension $d_l < 3$. This is consistent with finite T MC results^{4,5,13,14} for the gauge glass in 3D which indicate that $T_c = \mathcal{O}(J)$. This value of T_c is very difficult to reconcile with the suggestion that $d_l \approx 3$ from previous DWRG studies^{5,6,8,9} as this implies a very small value of T_c/J . A very recent Monte Carlo study¹⁵ of the 2D and 3D unscreened gauge glass obtains values of θ_s which are very close to those of this work.

The prediction that $T_c = 0$ for the 2D gauge glass is still somewhat controversial. It is safe to say that all $T=0$ defect energy scaling studies agree that $\theta_s < 0$ in 2D although there is some disagreement on the actual value of θ_s . Finite but low-temperature Monte Carlo simulations^{56,57} claim that $T_c > 0$ for the 2D gauge glass. This result is contested by some very recent large scale simulations by Katzgraber and Young¹⁵ at much lower T in 2D. The data of this study scales best with $\theta_s = -0.39 \pm 0.03$ and $T_c = 0$ and is inconsistent with $T_c > 0$ but is consistent with the results of this work.

We have also studied the effects of screened vortex-vortex interactions on the spin domain-wall energy using the BT measurement in 3D. Screening of the Coulomb interaction of vortices is implemented by adding a mass term λ^{-2} to the denominator of the Green's function,⁵⁸

$$G(\mathbf{r}) = \frac{1}{L^3} \sum_{\mathbf{k} \neq 0} \frac{e^{i\mathbf{k} \cdot \mathbf{r}} - 1}{6 - 2\cos k_x - 2\cos k_y - 2\cos k_z + \lambda^{-2}}. \quad (33)$$

The results are also shown in Fig. 7. We average over 10^3 samples for $L=2-4$ and 250 for $L=5$ for several values of the screening length λ . Screening is clearly a relevant perturbation when λ is finite and $\theta_s^{BT} < 0$ but our small sizes do not permit an estimate of the value of θ_s^{BT} . For large screening lengths, $\langle \Delta E_s^{BT}(L) \rangle$ seems to scale the same as for the unscreened case but we expect that $\langle \Delta E_s^{BT}(L) \rangle$ will decrease as L^{θ_s} with $\theta_s < 0$ at length scales which are beyond our computing power for any $\lambda < \infty$. These results are consistent with those of Bokil and Young⁵⁸ who studied the question of screening using a RT measurement and with Rieger *et al.*⁵⁹ for very strong screening.

C. Varying disorder strength in 2D and 3D

We have also performed simulations with various strengths of disorder in the CG representation using Eq. (30) for the 2D case and Eq. (27) for 3D where the random bond variables A_{ij} are independently uniformly distributed in the range $[-\alpha\pi, \alpha\pi]$ with $0 \leq \alpha \leq 1$ so that $\langle A_{ij} \rangle = 0$ and $\langle |A_{ij}| \rangle = \alpha\pi/2$. Physical realizations of this model are, e.g., an XY magnet with random Dzyaloshinski-Moriya interactions¹⁸ and Josephson-junction arrays with positional disorder^{19,20} where both the effective coupling constant $J(L)$ and the effective disorder strength $A(L)$ at length scale L play a role. Studies in 2D (Refs. 18–20) suggest that weak disorder ($\alpha \geq 0$) does not affect the existence of an ordered phase at intermediate temperature but there is a reentrant transition to a disordered phase at low temperature. However, recent analytic^{22–25} and numerical⁸ studies show there




FIG. 8. RG flow in 2D. The flows are from right to left for $\alpha \geq 0.4$ and from left to right for $\alpha \leq 0.35$.

is an ordered phase for $T < T_c(\alpha)$ with $T_c(\alpha) > 0$ for $0 \leq \alpha < \alpha_c$. To study how the disorder strength behaves as the length scale L varies, one must, if possible, identify the scaled disorder strength $A(L)$ with some measurable quantity. An identification has been proposed in a recent numerical study⁸ where the effective disorder strength is defined as $2\pi A(L) \equiv \langle |\Delta^0(L)| \rangle$ with $A(1) = \alpha/4$ so that one can follow the flows of both $J(L)$ and $A(L)$ with an increasing length scale L . With this definition, $0 \leq A(L) \leq 1/4$. $\Delta^0(L)$ is the global phase twist minimizing the energy of a system of size L for a particular realization of disorder. For two phases with energy $E_{12} = V(\theta_1 - \theta_2 - A_{12})$, the minimum is at $\theta_1 - \theta_2 = A_{12}$ which is satisfied by applying a “global” phase twist of A_{12} . Hence, this follows the definition of $A(L)$ as a measure of disorder at scale L .

Since the numerical study⁸ used the Hamiltonian of Eq. (1) is the phase representation, we reinvestigate this model in the CG representation. The simulations were performed for $L=2-6, 8$, and 10 in 2D, $L=2-5$ in 3D and averaged over at least 10^3 samples. The results are shown in Fig. 8 in 2D and in Fig. 9 in 3D. We are interested in the stable fixed-point values at $L \rightarrow \infty$ J^* and A^* as these determine the nature of the phases. In 2D, weak disorder ($\alpha < \alpha_c \approx 0.37$) seems to be marginal and the system seems to iterate toward a glass phase with quasi-long-range order characterized by (J^*, A^*) where the fixed point values J^*, A^* are finite and depend on α . This is consistent with recent analytic studies.²²⁻²⁵ On the other hand, systems with strong disorder, $\alpha > \alpha_c$, seem to flow to a disordered fixed point $(J^*, A^*) = (0, 1/4)$ which corresponds to a nonsuperconducting glass.

In 3D, for $\alpha < \alpha_c \approx 0.57$, the system flows to a strong-coupling limit $J^* = \infty$. The disorder strength $A(L)$ appears to flow to a finite fixed point value A^* which depends on α . However this is not conclusive from our simulations, since only very small values $L=2-6$ are used. This can be interpreted as the zero field version of a Bragg glass phase. For large disorder, $\alpha > \alpha_c$, $[J(L), A(L)]$ seem to iterate to their maximum values of $(\infty, 1/4)$ corresponding to the gauge




FIG. 9. RG flow in 3D. The flows are from left to right for all α .

glass fixed point. It seems that, in the absence of screening, $\lambda \rightarrow \infty$, there are two glassy superconducting phases at $T=0$ which, in an applied magnetic field, correspond to a Bragg glass⁶⁰ for $\alpha < \alpha_c$ and to a vortex glass⁶¹ for $\alpha > \alpha_c$.

VI. DISCUSSION AND SUMMARY

In this paper, we reinvestigate the possibility of an ordered phase at small but finite temperature T by a numerical domain wall renormalization group method in a disordered XY model in 2D and 3D described by the Hamiltonian of Eq. (1) in the Coulomb gas representation. For the $\pm J$ XY spin glass in 3D, our simulations yield the spin glass stiffness exponent $\theta_s^{BT} \approx +0.10$ which suggests its lower critical dimension is $d_l < 3$. This value of θ_s^{BT} is very different from existing estimates of the chiral glass stiffness exponent in 3D $\theta_c \approx +0.47$ (Ref. 29) and $\theta_c \approx +0.56 \pm 0.18$.³⁰ The difference between θ_s and θ_c seems to support the decoupling of two degrees of freedom in 3D. For the gauge glass, we estimate the stiffness exponent $\theta_s^{BT} = -0.36 \pm 0.01$ in 2D and $\theta_s^{BT} = +0.31 \pm 0.01$ in 3D, which are considerably larger than all earlier estimates. The latter value is consistent with $T_c/J \sim \mathcal{O}(1)$ from finite temperature MC studies^{4,5,13,14} and also strongly suggests $d_l < 3$. The results for the XY spin glass in 3D are consistent with spin-glass order at $T > 0$ which is in contradiction with all other studies²⁷⁻²⁹ except one.³⁰ We also note that there are studies of the 3D Heisenberg spin glass⁶² which indicate that the behavior is similar to that of the XY spin glass found in Sec. V A except that they find a surprisingly large $\theta_s \approx 0.8$. The existence of a low- T state with spin-glass and chiral order is also found by studying nonequilibrium relaxation of the corresponding susceptibilities.⁶³

We also studied the effects of varying the disorder strength. In 2D, our simulations imply that weak disorder is marginal²²⁻²⁵ and a system with strong disorder flows to a disordered fixed point. There is no sign of a reentrant transition in our simulations. In 3D, weak disorder has little effect and the system flows to an ordered phase which is the zero-

field analog of a Bragg glass.⁶⁰ For strong disorder, the system seems to flow to a gauge glass fixed point. The disagreement between the stiffness exponent θ_s^{BT} and previous estimates is because these measure θ_s^{RT} whose meaning is less clear. The quantity $\Delta E^{RT}(L)$ seems more likely to suffer from large corrections to scaling as seen in Fig. 7, especially for the small system sizes L which are possible to simulate at present. However, we conjecture that both measurements would coincide if *much* larger values of L could be reached. Since our simulations are also limited to very small sizes L , it is not possible to draw any definite conclusions from them and more studies are needed to settle these problems in random systems more satisfactorily.

One interesting conjecture we can make concerns the Bragg and vortex glass states in disordered superconductors in an applied magnetic field. Recently, Giardinà *et al.*⁶⁴ studied the model of Eq. (1) in the strong screening limit $\lambda \rightarrow 0$ in 3D and found that two phases exist at $T=0$ in the presence of an applied external field. The low field, the small disorder phase has a well-ordered (by eye) vortex line lattice as a ground state with a stiffness exponent $\theta_s = +1.0$, implying this is a *superconducting* state. By contrast, the high-field, large disorder ground state is a disordered entangled vortex configuration, again by inspection, with $\theta_s \approx -1.0$,^{59,64} which implies this is a *nonsuperconducting* state. We conjecture that the low-field state is a Bragg glass and the high-field state is a disordered entangled vortex liquid. In this limit, the evidence is strongly in favor of a direct, disorder, or field-driven transition from a superconducting Bragg glass to a normal nonsuperconducting phase. This scenario seems to be favored by recent experiments.

In the absence of screening of the vortex-vortex interactions, the picture which results from this work is somewhat different, although the studies here are all done in zero applied field. One may argue that increasing the disorder is equivalent to increasing the field at fixed disorder. At low field or low disorder, the ground state is a *superconducting* Bragg glass with $\theta \approx +1.0$, exactly as with screening. Without screening, the main difference is that the high-field, large disorder phase is a true superconducting vortex *glass* with stiffness exponent $\theta \approx +0.30$, as proposed by Fisher *et al.*⁶¹ We tentatively conjecture that a true superconducting vortex glass phase does not exist in 3D *except* in the absence of screening ($\lambda \rightarrow \infty$). Unfortunately, this prediction about the existence of a superconducting phase in the presence of large disorder seems to contradict two recent studies.^{65,66} These

are both finite T Monte Carlo simulations of a 3D XY model with uniform A_{ij} given by the applied magnetic field and the disorder by random couplings J_{ij} . In these simulations, superconductivity is probed by the helicity modulus and, although these studies differ in several respects, they agree on one vital point which is that, in the absence of screening, a superconducting phase does *not* exist for strongly disordered J_{ij} in a uniform applied field. If our conjecture is correct that the models considered in this work are not affected by the application of a uniform magnetic field, then $\theta > 0$ which seems to imply a finite helicity modulus and superconductivity. The natural consequence of this argument is that, in the absence of screening, a strongly disordered system in a field will be superconducting which disagrees with recent simulations.^{65,66}

Clearly, further work is needed to resolve this contradiction which could be due to several possibilities. These include the fact, among others, that the models used may not be equivalent because of the different types of disorder imposed, the presence or absence of an external magnetic field, and the extreme difficulty of the simulations causing a finite helicity modulus to be missed. One difference which appears to be unimportant is that this work is at $T=0$ while the simulations^{65,66} are at $T>0$. This is apparent in the recent work of Katzgraber and Young¹⁵ who find essentially the same exponent $\theta > 0$ at $T>0$ for the 3D gauge glass. There is a very recent simulation⁶⁷ on precisely this point using the CG representation of the unscreened 3D system in a large applied field with random pinning of the vortex cores. This supports the existence of a low- T superconducting *vortex glass* phase at large disorder in a field in the absence of screening, as conjectured here. Our understanding of the experimental consequences for real systems with *mesoscopic* penetration depths $\lambda \sim \mathcal{O}(10^3)$ Å is lacking. Much work remains to be done to clarify experimental and theoretical situations. A challenge to simulations is, in a single model, to demonstrate unambiguously that only a Bragg glass exists in the presence of screening and at weak disorder while, without screening, a distinct vortex glass phase also exists at large disorder.

ACKNOWLEDGMENTS

Computations were performed at the Theoretical Physics Computing Facility at Brown University. J.M.K. thanks A. Vallat and B. Grossmann for many discussions about spin glasses, best twists, etc.

¹M. P. A. Fisher, Phys. Rev. Lett. **62**, 1415 (1989).
²G. Blatter, M. V. Feigel'man, V. J. Geshkenbein, A. I. Larkin, and V. M. Vinokur, Rev. Mod. Phys. **66**, 1125 (1994).
³T. Nattermann and S. Scheidl, Adv. Phys. **49**, 607 (2000).
⁴D. A. Huse and H. S. Seung, Phys. Rev. B **42**, 1059 (1990).
⁵J. D. Reger, T. A. Tokuyasu, A. P. Young, and M. P. A. Fisher, Phys. Rev. B **44**, 7147 (1991).
⁶M. J. P. Gingras, Phys. Rev. B **45**, 7547 (1992).

⁷M. P. A. Fisher, T. A. Tokuyasu, and A. P. Young, Phys. Rev. Lett. **66**, 2931 (1991).
⁸J. M. Kosterlitz and M. V. Simkin, Phys. Rev. Lett. **79**, 1098 (1997).
⁹J. Maucourt and D. R. Grempel, Phys. Rev. B **58**, 2654 (1998).
¹⁰C. Dekker, P. J. M. Wöltgens, R. H. Koch, B. W. Hussey, and A. Gupta, Phys. Rev. Lett. **69**, 2717 (1992).
¹¹J. R. Banavar and M. Cieplak, Phys. Rev. Lett. **48**, 832 (1982).

- ¹²W. L. McMillan, Phys. Rev. B **29**, 4026 (1983).
- ¹³C. Wengel and A. P. Young, Phys. Rev. B **56**, 5918 (1997).
- ¹⁴T. Olson and A. P. Young, Phys. Rev. B **61**, 12467 (2000).
- ¹⁵H. G. Katzgraber and A. P. Young, Phys. Rev. B **64**, 104426 (2001); cond-mat/0205206, Phys. Rev. B (to be published).
- ¹⁶R. H. Koch, V. Foglietti, W. J. Gallagher, G. Koren, A. Gupta, and M. P. A. Fisher, Phys. Rev. Lett. **63**, 1511 (1989).
- ¹⁷C. Dekker, W. Eidelloth, and R. H. Koch, Phys. Rev. Lett. **68**, 3347 (1992).
- ¹⁸M. Rubinstein, B. Shraiman, and D. R. Nelson, Phys. Rev. B **27**, 1800 (1983).
- ¹⁹E. Granato and J. M. Kosterlitz, Phys. Rev. B **33**, 6533 (1986).
- ²⁰E. Granato and J. M. Kosterlitz, Phys. Rev. Lett. **62**, 823 (1989).
- ²¹Y. Ozeki and H. Nishimori, J. Phys. A **26**, 3399 (1993).
- ²²T. Natterman, S. Scheidl, S. E. Korshunov, and M. S. Li, J. Phys. **I 5**, 555 (1995).
- ²³S. E. Korshunov and T. Natterman, Phys. Rev. B **53**, 2746 (1996).
- ²⁴S. Scheidl, Phys. Rev. B **55**, 457 (1997).
- ²⁵M.-C. Cha and H. A. Fertig, Phys. Rev. Lett. **74**, 4867 (1995).
- ²⁶J. Villain, J. Phys. C **10**, 4793 (1977); *ibid.* **11**, 745 (1978).
- ²⁷H. Kawamura and M. Tanemura, Phys. Rev. B **36**, 7177 (1987).
- ²⁸H. Kawamura and M. Tanemura, J. Phys. Soc. Jpn. **60**, 608 (1991).
- ²⁹H. Kawamura, Phys. Rev. B **51**, 12 398 (1995).
- ³⁰J. Maucourt and D. R. Grempel, Phys. Rev. Lett. **80**, 774 (1998).
- ³¹M. Ney-Nifle, H. J. Hilhorst, and M. A. Moore, Phys. Rev. B **48**, 10 254 (1993).
- ³²M. J. Thill, M. Ney-Nifle, and H. J. Hilhorst, J. Phys. A **28**, 4825 (1995).
- ³³M. Ney-Nifle and H. J. Hilhorst, Phys. Rev. B **51**, 8357 (1995).
- ³⁴P. W. Anderson, J. Less-Common Met. **62**, 291 (1978); P. W. Anderson and C. M. Pond, Phys. Rev. Lett. **40**, 903 (1979).
- ³⁵H. Rieger, L. Santen, U. Blasum, M. Diehl, and M. Jünger, J. Phys. A **29**, 3939 (1996); N. Kawashima and H. Rieger, Europhys. Lett. **39**, 85 (1997).
- ³⁶H. Rieger, in *Frustrated Systems: Ground State Properties Via Combinatorial Optimization*, Lecture Notes in Physics, Vol. 501 (Springer-Verlag, Heidelberg, 1998); R. Ahuja, T. Magnanti and J. Orlin, *Network Flows* (Prentice-Hall, Englewood Cliffs, NJ, 1993).
- ³⁷S. Kirkpatrick, C. D. Gellat, and M. P. Vecchi, Science **220**, 671 (1983).
- ³⁸M. V. Simkin, Phys. Rev. B **55**, 11 405 (1997).
- ³⁹J. Villain, J. Phys. (France) **36**, 581 (1975).
- ⁴⁰J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, Phys. Rev. B **16**, 1217 (1977).
- ⁴¹A. Vallat and H. Beck, Phys. Rev. B **50**, 4015 (1994).
- ⁴²J. M. Kosterlitz and D. J. Thouless, J. Phys. C **6**, 1181 (1973).
- ⁴³J. M. Kosterlitz, J. Phys. C **7**, 1046 (1974).
- ⁴⁴J. M. Kosterlitz and N. Akino, Phys. Rev. Lett. **82**, 4094 (1999).
- ⁴⁵F. Matsubara, T. Shirakura, and M. Shiomi, Phys. Rev. B **58**, 11 821 (1998).
- ⁴⁶A. K. Hartmann and A. P. Young, Phys. Rev. B **64**, 180404 (2001).
- ⁴⁷L. Saul and M. Kardar, Phys. Rev. E **48**, 3221 (1993).
- ⁴⁸J. Houdayer, Eur. Phys. J. B **22**, 479 (2001).
- ⁴⁹H. S. Bokil and A. P. Young, J. Phys. A **29**, L89 (1996).
- ⁵⁰P. Ray and M. A. Moore, Phys. Rev. B **45**, 5361 (1992).
- ⁵¹S. Jain and A. P. Young, J. Phys. C **19**, 3913 (1986).
- ⁵²H. Nishimori and Y. Ozeki, J. Phys. Soc. Jpn. **59**, 289 (1990).
- ⁵³M. Schwartz and A. P. Young, Europhys. Lett. **15**, 209 (1991).
- ⁵⁴Y. Ozeki and H. Nishimori, Phys. Rev. B **46**, 2879 (1992).
- ⁵⁵J. M. Kosterlitz and N. Akino, Phys. Rev. Lett. **81**, 4672 (1998).
- ⁵⁶M. Y. Choi and S. Y. Park, Phys. Rev. B **60**, 4070 (1999).
- ⁵⁷B. J. Kim, Phys. Rev. B **62**, 644 (2000).
- ⁵⁸H. S. Bokil and A. P. Young, Phys. Rev. Lett. **74**, 3021 (1995).
- ⁵⁹J. Kisker and H. Rieger, Phys. Rev. B **58**, R8873 (1998); F. O. Pfeiffer and H. Rieger, *ibid.* **60**, 6304 (1999).
- ⁶⁰T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **72**, 1530 (1994); Phys. Rev. B **52**, 1242 (1995).
- ⁶¹M. P. A. Fisher, Phys. Rev. Lett. **62**, 1415 (1989); D. S. Fisher, M. P. A. Fisher, and D. A. Huse, Phys. Rev. B **43**, 130 (1991).
- ⁶²S. Endoh, F. Matsubara, and T. Shirakura, J. Phys. Soc. Jpn. **70**, 1543 (2001).
- ⁶³T. Nakamura and S. Endoh, cond-mat/0110017 (unpublished).
- ⁶⁴C. Giardinà, N. V. Priezjev, and J. M. Kosterlitz, cond-mat/0202487 (unpublished).
- ⁶⁵Y. Nonomura and X. Hu, Phys. Rev. Lett. **86**, 5140 (2001).
- ⁶⁶P. Olsson and S. Teitel, Phys. Rev. Lett. **87**, 137001 (2001).
- ⁶⁷A. Vestergren, J. Lidmar, and M. Wallin, Phys. Rev. Lett. **88**, 117004 (2002).