

Mott transitions in correlated electron systems with orbital degrees of freedomY. Ōno,¹ M. Potthoff,^{2,*} and R. Bulla³¹*Department of Physics, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8602, Japan*²*Lehrstuhl Festkörpertheorie, Institut für Physik, Humboldt-Universität zu Berlin, D-10115 Berlin, Germany*³*Theoretische Physik III, Elektronische Korrelationen und Magnetismus, Institut für Physik, Universität Augsburg, D-86135 Augsburg, Germany*

(Received 18 November 2002; published 30 January 2003)

Mott metal-insulator transitions in an M -fold orbitally degenerate Hubbard model are studied by means of a generalization of the linearized dynamical mean-field theory. The method allows for an efficient and reliable determination of the critical interaction U_c for any integer filling n and different M at zero temperature. For half filling a linear dependence of U_c on M is found. Inclusion of the (full) Hund's rule exchange J results in a strong reduction of U_c . The transition turns out to change qualitatively from continuous for $J=0$ to discontinuous for any finite J . This result is confirmed using the exact-diagonalization method for $M=2$.

DOI: 10.1103/PhysRevB.67.035119

PACS number(s): 71.10.Fd, 71.27.+a, 71.30.+h

I. INTRODUCTION

The Mott metal-insulator transition (MIT) has been the subject of numerous experimental and theoretical investigations over the past decades.^{1–3} As a transition from a paramagnetic metal to a paramagnetic insulator that is driven by the competition between the conduction electrons' kinetic energy and their mutual Coulomb repulsion, the Mott MIT represents a prime example for a quantum phase transition. Realizations of the Mott transition can be found in different $3d$ transition-metal oxides^{1,2} as well as in alkali-doped fullerenes,^{4,5} for example. Its description is a challenging task for many-body theory.

The minimal model to describe the Mott transition is the Hubbard model⁶ which in its simplest form contains the local Coulomb interaction between the electrons in a conduction band formed by a *single* orbital only. The investigation of such a single-band model might be justified in some cases, in particular, as an effective model for low-energy properties. A consistent description of the experimental observations for the above-mentioned systems, however, certainly requires the use of more realistic models which include orbital degrees of freedom, possible crystal-field splittings, etc.

Theoretical advances in the past decade—mainly due to the development of the dynamical mean-field theory^{7,8} (DMFT)—have led to an increased understanding of multi-orbital Hubbard models. Recent investigations have either concentrated on fundamental questions such as the nature of the Mott MIT as a function of orbital degeneracy^{9–16} or, within the so-called local-density approximation (LDA) +DMFT approach,¹⁷ on a realistic description of the transition.

Nevertheless, there are still no reliable DMFT results for the zero-temperature ($T=0$) Mott transition in general multiband models: Numerical methods to solve the DMFT equations are either restricted to fairly large temperatures [quantum Monte Carlo (QMC) approach] or have not yet been extended to the multiband case [numerical renormalization-group (NRG) method]. Additionally, due to the sign problem within the QMC approach only a simplified exchange part of the Coulomb interaction can be taken into

account.^{13,18} The exact-diagonalization (ED) method is restricted to rather small system sizes, and data are available for $M \leq 2$ orbitals only.¹⁶ The M dependence of the critical interaction strength U_c for different integer fillings n is particularly interesting. Detailed results for U_c as a function of M and n are available from the Gutzwiller approximation and from a slave-boson approach.^{9,11} Numerical DMFT-QMC calculations have been performed for $M \leq 3$.^{12,13} Remarkably, in the limit of large orbital degeneracy $M \rightarrow \infty$ an analytical treatment of the DMFT becomes possible for the MIT.¹⁵ A scaling $U_c = U_{c,2} \propto M$ for the actual transition is found while $U_{c,1} \propto \sqrt{M}$ is obtained for the critical interaction where the insulating solution breaks down.¹⁵ This is consistent with the linear dependence for large M found in Refs. 9 and 11 and with the square-root dependence reported in Ref. 5. However, there is still a need for a DMFT method which works at $T=0$ and which allows for an efficient and reliable determination of U_c for arbitrary n and M and for a model including the full exchange interaction.

Here we present and apply a proper multiband generalization of the so-called linearized DMFT (L-DMFT) (Ref. 19) which is a simple but rather successful technique that originally was developed for the critical regime of the Mott MIT in the single-band model. Within the L-DMFT the lattice problem is mapped onto an Anderson impurity model with a single bath site only by considering a simplified self-consistency condition just at the critical point. The approach can be considered to be the simplest nontrivial variant of the more systematic projective self-consistent method (PSCM) by Moeller *et al.*²⁰ It allows for extremely fast numerical calculations or even analytical results to estimate the critical parameters. Extensions have been considered for the (Mott-Hubbard or charge-transfer) MITs in a d - p model,²¹ for a quantum-critical point in the periodic Anderson model,²² and for the $M=2$ degenerate Hubbard model at half filling and $J=0$.¹⁶ Furthermore, a general two-site DMFT (Ref. 23) has been developed which reduces to the L-DMFT at the MIT but is not restricted to the critical point. An application to multiband systems is likewise possible but has not yet been considered.

As concerns effective single-band models, the linearized

DMFT has been tested extensively by comparing with numerical results from the full DMFT. For the standard Hubbard model^{19,24} but also for different thin-film and semi-infinite surface geometries²⁵ as well as for the d - p model²⁶ a remarkable agreement has been found: Detailed trends of U_c as a function of electronic model parameters and as a function of parameters characterizing the lattice geometry are predicted reliably. For quantitative estimates errors of the order of a few percent have to be tolerated.

As is demonstrated here, a straightforward extension of the method to multiband systems is possible. We expect the L-DMFT to be a fast but reliable tool in this case as well since the effective impurity model includes the complete atomic part and the self-consistently determined bath with the complete degeneracy (although restricted to a single site only). Furthermore, the reliability of the L-DMFT for the multiband case is checked by comparing with different fully numerical DMFT techniques whenever results are available. All DMFT calculations discussed in this paper were performed on a Bethe lattice with $t^2 = 1$, corresponding to bandwidth of 4.

II. MODEL AND LINEARIZED DMFT

We consider a multiband Hubbard-type model $H = H_0 + H_1$ including an intraorbital hopping

$$H_0 = \sum_{ij\alpha\sigma} t_{ij} c_{i\alpha\sigma}^\dagger c_{j\alpha\sigma} \quad (1)$$

and the direct and the exchange part of the on-site Coulomb interaction:

$$H_1 = \frac{1}{2} \sum_{i\alpha\alpha'\sigma\sigma'} U_{\alpha\alpha'} c_{i\alpha\sigma}^\dagger c_{i\alpha'\sigma'}^\dagger c_{i\alpha'\sigma'} c_{i\alpha\sigma} + \frac{1}{2} \sum_{i\alpha\alpha'\sigma\sigma'} J_{\alpha\alpha'} c_{i\alpha\sigma}^\dagger c_{i\alpha'\sigma'}^\dagger c_{i\alpha\sigma} c_{i\alpha'\sigma'}, \quad (2)$$

using standard notations. i is a site index, and $\sigma = \uparrow, \downarrow$ refers to the spin direction. The different orbitals labeled by $\alpha = 1, \dots, M$ are considered to be equivalent, and M is the orbital degeneracy. Exploiting atomic symmetries, we have $U_{\alpha\alpha'} = (U + 2J)\delta_{\alpha\alpha'} + U(1 - \delta_{\alpha\alpha'})$ and $J_{\alpha\alpha'} = J(1 - \delta_{\alpha\alpha'})$ as usual (see, e.g., Ref. 27). Opposed to calculations using the QMC method (see, e.g., Refs. 13 and 18), H_1 includes the full exchange part and thus preserves rotational invariance.

The DMFT assumes the self-energy to be local. Furthermore, the self-energy as well as the Green function are diagonal with respect to the orbital index α for the model considered here. In case of equivalent orbitals and in the absence of any spontaneous symmetry breaking, we thus have $\Sigma_{ij\alpha\beta}(\omega) = \delta_{ij}\delta_{\alpha\beta}\Sigma(\omega)$ and likewise for the local Green function: $G_{ii\alpha\beta}(\omega) = \delta_{\alpha\beta}G(\omega)$. Within the framework of the DMFT, the model H is mapped onto an impurity model $H' = H'_0 + H'_1$ where H'_1 is the local part of the interaction (2) at a distinguished site i_0 and

$$H'_0 = \sum_{\alpha\sigma} t_0 c_{i_0\alpha\sigma}^\dagger c_{i_0\alpha\sigma} + \sum_{\alpha\sigma, k=2}^{n_s} \epsilon_k a_{k\alpha\sigma}^\dagger a_{k\alpha\sigma} + \sum_{\alpha\sigma, k=2}^{n_s} V_k (c_{i_0\alpha\sigma}^\dagger a_{k\alpha\sigma} + \text{H.c.}) \quad (3)$$

with $t_0 = t_{ii}$. The self-energy $\Sigma(\omega)$ of H' is identified with the self-energy of the lattice model H and yields, via the lattice Dyson equation, the on-site Green function $G(\omega)$. The latter determines the one-particle parameters of H' or, equivalently, the hybridization function $\Delta(\omega) = \sum_k V_k^2 / (\omega + \mu - \epsilon_k)$ via the DMFT self-consistency condition $\Delta(\omega) = \omega + \mu - t_0 - \Sigma(\omega) - 1/G(\omega)$.

Within the full DMFT an infinite number of bath degrees of freedom $n_s \rightarrow \infty$ are necessary to fulfill the self-consistency equation. This implies the need for further approximations to treat H' . Contrary, a simple two-site model is considered here. For $n_s = 2$ the model (3) can be solved exactly; however, a simplified self-consistency condition must be tolerated.

Our approach rests on the assumption that similar to the single-band model, the MIT is characterized by a vanishing weight of a low-energy quasiparticle resonance. Close to the MIT it is then plausible to neglect any internal structure of the quasiparticle peak and to assume that there is no influence of high-energy features on the low-energy part of the excitation spectrum. The quasiparticle peak is then approximated by a single pole at the Fermi energy ($\epsilon_c \equiv \epsilon_{k=2} = \mu$) which reproduces itself in the DMFT self-consistency cycle. One can proceed as for the single-band model¹⁹ and is finally left with the following simple algebraic self-consistency equation which determines the critical parameters ($V \equiv V_{k=2}$):

$$V^2 = z M_2^{(0)}. \quad (4)$$

Here $M_2^{(0)}$ is the second moment of the noninteracting density of states $M_2^{(0)} = \int dx x^2 \rho^{(0)}(x)$. Formally, Eq. (4) is the same as that in the single-band model.¹⁹ However, the quasiparticle weight z has to be determined from the degenerate model (3). We calculate $z = 1/[1 - \Sigma'(0)]$ from the spectral representation of $G(\omega)$ by summing the residua $\propto V^2$ in the limit $V \rightarrow 0$. The residua are obtained by standard (Brillouin-Wigner) degenerate perturbation theory up to the order V^2 . This yields an expression $z = z(V, \epsilon_c, t_0, U, J) = V^2 F(\epsilon_c - t_0, U, J) / M_2^{(0)} + \mathcal{O}(V^4)$. Using $\epsilon_c = \mu$ in the limit $V \rightarrow 0$ and setting $t_0 = t_{ii} = 0$ to fix the energy to zero, we obtain from Eq. (4) $F(\mu, U, J) = 1$ as a condition for the MIT. This condition is different for different orbital degeneracy M and integer filling $n = 1, \dots, 2M - 1$.

III. RESULTS AND DISCUSSION

For an integer filling n and for any U larger than a critical interaction $U_c = U_c(n, M)$ there are two critical values for the chemical potential $\mu_{\pm}(U)$ which can be obtained by solving $F(\mu, U) = 1$ for μ (the case $J = 0$ is considered first). For $U > U_c(n, M)$ and $\mu_-(U) < \mu < \mu_+(U)$ the system is a Mott insulator with integer filling n . On the other hand, for μ




FIG. 1. $\Delta\mu(U)$ for half filling $n=M$ and different M as obtained from Eq. (7). Results from the full DMFT using ED, NRG, and the PSCM (Refs. 14 and 20) are shown for comparison.

$\langle\mu_-(U)$ or $\mu_+(U)$, a noninteger filling is realized, and the system becomes metallic. The chemical potential is a continuous function of the filling n' in the vicinity of n for any $U < U_c(n, M)$ but $\mu(n')$ is discontinuous at $n' = n$ with a jump from $\mu_-(U)$ to $\mu_+(U)$ for $U > U_c(n, M)$. U_c is determined by minimization of the function $U(\mu)$ which is obtained by solving $F(\mu, U) = 1$ for U .

For half filling $n=M$ and arbitrary M we succeeded with a completely analytical calculation for $F(\mu, U)$:

$$F(\mu, U)/M_2^{(0)} = \frac{1}{2} \left[\frac{M+1}{MU-\mu} + \frac{M}{\mu-(M-1)U} \right]^2 + \frac{1}{2} \left[\frac{M}{MU-\mu} + \frac{M+1}{\mu-(M-1)U} \right]^2. \quad (5)$$

For $U < U_c$ and half filling $n=M$ in the symmetric model, the chemical potential is fixed to $\mu = (M - \frac{1}{2})U$ due to particle-hole symmetry. Then we find $F = M_2^{(0)}(4M + 2)^2/U^2$ which results in the critical interaction

$$U_c(n=M, M) = (4M+2)\sqrt{M_2^{(0)}}. \quad (6)$$

By solving $F(\mu, U) = 1$ with Eq. (5) for $U > U_c$ we obtain the jump in the chemical potential

$$\Delta\mu \equiv \mu_+ - \mu_- = U \sqrt{B - \sqrt{B^2 - 1 + \frac{U_c^2}{U^2}}}, \quad (7)$$

where $B \equiv 1 + 1/[2(2M+1)^2(U/U_c)^2]$. The result for $\Delta\mu(U)$ is shown in Fig. 1 for $M=1, \dots, 5$. Here and hereafter we set $M_2^{(0)} = 1$. For fixed M and $U \rightarrow \infty$, the jump in the chemical potential increases linearly with U . For $U \rightarrow U_c$, $\Delta\mu(U)$ shows a square-root behavior.

Qualitatively and even quantitatively, this agrees well with results from numerical methods to solve the full DMFT equations. In particular, we compare with results from ED calculations which we have performed with $n_s=6$ for $M=2$ (circles) and with data available for $T=0$ from the projective self-consistent method (crosses).¹⁴ The nondegenerate case $M=1$ has already been discussed in Ref. 24. For $M=2$ and $n_s=6$ the error due to finite-size effects in the ED is




FIG. 2. Critical interaction U_c for different fillings $n = 1, 2, \dots, 2M-1$ and different M as a function of n/M . Symbols: L-DMFT results. Lines: simple fit function $U_c(n, M) = U_c(M, M) - c(1 - (n/M))^2$ and $c = -0.217 + 2.671M$.

almost negligible. One can state that the agreement of the L-DMFT results with the ED and PSCM data is equally good for both $M=1$ and $M=2$.

The generalized L-DMFT predicts $U_c(n=M, M)$ to depend linearly on M , see Eq. (6). This agrees with previous results from the Gutzwiller method,⁹ slave-boson calculations,¹¹ the noncrossing approximation for low temperature,²⁸ and the projective technique for large M .¹⁵ This finding can easily be understood by looking at the weight factors in the Lehmann representation of the spectral density of the two-site Anderson model. Expanding into the different configurations, one finds that the ground state is composed of the order of M singlet states in the limit $V \rightarrow 0$ and that there are of the order of M excited states coupling to the ground state. Hence, there are of the order of M^2 processes to be considered in the V perturbation theory which are contributing to the spectral weight near $\omega=0$. This implies $z \propto M^2 V^2 / U^2$ and eventually results in $U_c \propto M$.

In Fig. 2 the dependence of U_c on both n and M is shown for all integer fillings and $M \leq 5$. We find U_c to be reduced away from half filling, consistent with previous results.^{5,9,12-14} The fit shows an almost perfect quadratic dependence of U_c on the filling.

Let us now discuss the influence of the Hund's rule coupling J on the Mott transition at $T=0$. Contrary to the QMC method, the full exchange part of the on-site Coulomb interaction can easily be included in the L-DMFT calculation. A quantitative influence, i.e., a reduction of U_c due to J has already been observed in Refs. 5 and 13. This behavior is also evident from our L-DMFT results, as shown below.

The situation is, however, more complicated: It turns out that a finite J leads to a *qualitative* change of the nature of the $T=0$ Mott transition. This is illustrated in Fig. 3 which shows the U dependence of the quasiparticle weight z for various values of J and $n=M=2$. Note that z is plotted not only close to the transition but also for U far below U_c which means that the self-consistency relation (4) is used for any U . This is reasonable as long as there is a clear separation between the low-energy (quasiparticle) and the high-energy




FIG. 3. U dependence of the quasiparticle weight z for $n=M=2$ and different J . Dashed lines correspond to metastable solutions; vertical lines indicate the actual Mott transition at $U=U_c$.

scale (Hubbard bands). Although the approximation becomes questionable for higher z and particularly in the limit $z \rightarrow 1$, the trend of $z(U)$ looks rather plausible in whole range.

In the critical regime where z goes to zero continuously, we find $z(U)$ to decrease with decreasing U which seems to be unphysical. In fact, this behavior belongs to a metastable state (dashed line) while for the true ground state (solid line) $z(U)$ is always decreasing with increasing U . The two metallic states with finite $z > 0$ but different slope dz/dU are coexisting in a finite range of interaction strengths, namely, for U larger than $U_{c,2}$ (given by $z=0$) and for U smaller than another critical value $U_{c,3} > U_{c,2}$ where they merge.²⁹

The actual critical interaction strength U_c where the transition to the insulator takes place must be determined by comparing the respective energies with the energy of the insulator. This is shown in Fig. 4 for $J=0.1$. For any $J > 0$ it is found that $U_{c,2} < U_c < U_{c,3}$. At U_c the energies of the metallic and insulating states cross as functions of U . Consequently, the transition at $T=0$ is discontinuous with a jump in physical quantities such as the double occupancy $\langle n_{\uparrow}n_{\downarrow} \rangle = dE/dU$. Equivalently, one can look at the local magnetic moment at the impurity site $\langle S^2 \rangle$ (see Fig. 5). The local mo-




FIG. 4. Energy of the two metallic solutions relative to the energy of the insulator for $J=0.1$. At $U_{c,2}$ (filled circle) the metastable metallic state continuously merges with the insulating state. At $U_{c,3}$ (cross) the two metallic states merge. The discontinuous phase transition takes place at U_c (open circle).




FIG. 5. Local moment at the impurity site $\langle S^2 \rangle$ as a function of U for different J . Dashed lines: metastable solutions; vertical lines: actual Mott transition for $U=U_c$.

ment increases with U , typical for a strongly correlated normal metal. At U_c it jumps to $\langle S^2 \rangle = 2$ for any finite J . Similarly, the quasiparticle weight z decreases with increasing U and drops to zero at U_c from a finite value $z > 0$. In Fig. 3 this is indicated by solid vertical lines.

The phase diagram in the (U, J) plane is shown in Fig. 6. The critical interaction for the actual transition is plotted together with $U_{c,2}$ and $U_{c,3}$. U_c is found to be very close to $U_{c,3}$ for all J values. Note that already a moderate value of $J \approx 0.3$ leads to a drastic reduction of U_c from $U_c = 10$ to $U_c \approx 5$.

Interestingly, similar results have been obtained beforehand by a rather different method: Applying a generalized Gutzwiller approximation to a degenerate two-band model, Bünemann *et al.*¹⁰ found the Mott transition to be discontinuous for any finite $J > 0$ and to be continuous for $J=0$ only. While the Gutzwiller method (for $J=0$) significantly overestimates the critical interaction strength as compared to the L-DMFT, the reduction of U_c by a finite J is stronger. Common to both methods, however, is the qualitative change of the Mott transition for a finite J .

We have additionally checked the nature of the Mott transition by a fully numerical evaluation of the DMFT equations using the ED method with $n_s=6$ sites for $n=M=2$. The results for $J=0$ and $J=0.1$ are shown in Fig. 7. The critical interaction U_c is found to be somewhat smaller when




FIG. 6. (U, J) phase diagram for $n=M=2$. U_c : critical interaction for the MIT. $U_{c,3}$: maximum U up to which a metallic solution is found. $U_{c,2}$: critical U where $z \rightarrow 0$ [$U_{c,2} = 14/3 - 3J$ for $J > 0$ and $M_2^{(0)} = 1$].




FIG. 7. DMFT-ED results ($n_s=6$) for the U dependence of the quasiparticle weight (left scale) and the local moment (right scale). $J=0$ and $J=0.1$ for $n=M=2$.

compared to the L-DMFT result as shown in Fig. 6. More importantly, however, the ED calculations show the same qualitative trend: The slope of $z(U)$ approaches $-\infty$ at a critical value $U=U_{c,3}$. This clearly indicates a second metastable metallic solution and a discontinuous transition and thus corroborates our results discussed above. Figure 7 also shows the U dependence of the local magnetic moment where the effect is even more pronounced. Note that in the insulator $\langle S^2 \rangle$ is close but not equal to the prediction of the linearized DMFT $\langle S^2 \rangle=2$ (cf. Fig. 5). This is due to residual local fluctuations which are neglected by the L-DMFT for the insulating phase.

The limit $J \rightarrow 0$ which appears to be somewhat exceptional can be analyzed easily. From Fig. 3 it is obvious that the critical interaction $U_{c,2}$ at which $z \rightarrow 0$ in the metastable phase shows a discontinuous jump from a value $U_{c,2} \approx 5 < U_{c,3}$ for $J > 0$ to $U_{c,2} = 10 = U_{c,3}$ for $J = 0$. To find the physical reason for this discontinuity of $U_{c,2}$, one can again look at the Lehmann representation of the spectral density for $V \rightarrow 0$. The main difference between the case $J = 0$ and the case $J > 0$ consists of the fact that for $J > 0$ only those configurations with a triplet on the impurity site contribute to the (singlet) ground state in the limit $V \rightarrow 0$ (see Fig. 5). As compared with $J = 0$ this is about a factor of 2 configurations less. Thus, the discontinuous jump of $U_{c,2}$ is caused by a strong suppression of orbital fluctuations due to a finite J (a similar argument in a slightly different context has been put forward in Ref. 16). On the other hand, it should be stressed

that the actual critical interaction U_c remains continuous in the limit $J \rightarrow 0$ as $U_c \rightarrow U_{c,3}$. Note that to uncover this subtle distinction it has been necessary to extend the theory beyond the $z \rightarrow 0$ limit.

IV. CONCLUSION

To summarize, we have presented an extension of the linearized dynamical mean-field theory (L-DMFT) to multi-orbital Hubbard models including the *full* exchange term. This allows a reliable calculation of the critical interaction strength U_c for the zero-temperature Mott transition analytically for all M at half filling and numerically for all integer fillings $n \neq M$ and $M \leq 5$. We find a linear dependence of U_c on the number of orbitals. Remarkably, a qualitative change of the nature of the $T=0$ Mott transition is observed for any finite J : the transition becomes discontinuous with a finite jump of the quasiparticle weight at U_c close to $U_{c,3}$.

For finite temperatures one might speculate that a finite J increases the tendency for a first-order Mott transition. This could be of experimental relevance for the Mott transition in transition-metal oxides.

The L-DMFT for correlated electron systems with orbital degrees of freedom is a handy and trustworthy approach with large potential for future applications. One might consider, for instance, the multiband Hubbard model investigated in the context of the LDA+DMFT approach for the transition-metal oxide V_2O_3 .¹⁸ The L-DMFT can be easily extended to answer questions about the importance of the e_g^σ bands (neglected in Ref. 18) on the Mott transition, or the influence of the *full* exchange term. We have already observed a significant quantitative change of U_c when excluding the spin-dependent part from the exchange term in Eq. (2). This should have consequences for a realistic description of the MIT in transition-metal oxides which are worthy of study.

ACKNOWLEDGMENTS

We acknowledge the support of the Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture (Y.Ö.) and the Deutsche Forschungsgemeinschaft through the Sonderforschungsbereich 290 (M.P.) and 484 (R.B.).

*Present address: Institut für Theoretische Physik und Astrophysik, Universität Würzburg, Am Hubland, 97074 Würzburg, Germany.

¹N. F. Mott, Proc. Phys. Soc., London, Sect. A **62**, 416 (1949); *Metal-Insulator Transitions*, 2nd ed. (Taylor & Francis, London, 1990).

²M. Imada, A. Fujimori, and Y. Tokura, Rev. Mod. Phys. **70**, 1039 (1998).

³F. Gebhard, *The Mott Metal-Insulator Transition*, Springer Tracts in Modern Physics, Vol. 137 (Springer, Berlin, 1997).

⁴O. Gunnarsson, Rev. Mod. Phys. **69**, 575 (1997).

⁵E. Koch, O. Gunnarsson, and R. M. Martin, Phys. Rev. B **60**, 15 714 (1999).

⁶J. Hubbard, Proc. Phys. Soc., London, Sect. A **276**, 238 (1963); M. C. Gutzwiller, Phys. Rev. Lett. **10**, 59 (1963); J. Kanamori,

Prog. Theor. Phys. **30**, 275 (1963).

⁷W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989).

⁸A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).

⁹J. P. Lu, Phys. Rev. B **49**, 5687 (1994).

¹⁰J. Büneemann and W. Weber, Phys. Rev. B **55**, 4011 (1997); J. Büneemann, W. Weber, and F. Gebhard, *ibid.* **57**, 6896 (1998).

¹¹R. Frésard and G. Kotliar, Phys. Rev. B **56**, 12 909 (1997); R. Frésard and M. Lamboley, J. Low Temp. Phys. **126**, 1091 (2002).

¹²M. J. Rozenberg, Phys. Rev. B **55**, R4855 (1997).

¹³J. E. Han, M. Jarrell, and D. L. Cox, Phys. Rev. B **58**, 4199 (1998).

¹⁴G. Kotliar and H. Kajueter, Phys. Rev. B **54**, R14 211 (1996).

- ¹⁵S. Florens, A. Georges, G. Kotliar, and O. Parcollet, Phys. Rev. B **66**, 205102 (2002).
- ¹⁶A. Koga, Y. Imai, and N. Kawakami, Phys. Rev. B **66**, 165107 (2002).
- ¹⁷For a review, see K. Held, I. A. Nekrasov, G. Keller, V. Eyert, N. Blümer, A. K. McMahan, R. T. Scalettar, T. Pruschke, V. I. Anisimov, and D. Vollhardt, in *Quantum Simulations of Complex Many-Body Systems: From Theory to Algorithms*, edited by J. Grotendorst, D. Marx, and A. Muramatsu, NIC Series Vol. 10 (NIC Directors, Forschungszentrum Jülich, 2002), p. 175.
- ¹⁸K. Held, G. Keller, V. Eyert, D. Vollhardt, and V. I. Anisimov, Phys. Rev. Lett. **86**, 5345 (2001).
- ¹⁹R. Bulla and M. Potthoff, Eur. Phys. J. B **13**, 257 (2000).
- ²⁰G. Moeller, Q. Si, G. Kotliar, M. Rozenberg, and D. S. Fisher, Phys. Rev. Lett. **74**, 2082 (1995); see also G. Moeller, Ph.D. thesis, Rutgers University, 1994.
- ²¹Y. Ōno, R. Bulla, and A. C. Hewson, Eur. Phys. J. B **19**, 375 (2001).
- ²²K. Held and R. Bulla, Eur. Phys. J. B **17**, 7 (2000).
- ²³M. Potthoff, Phys. Rev. B **64**, 165114 (2001).
- ²⁴Y. Ōno, R. Bulla, A. C. Hewson, and M. Potthoff, Eur. Phys. J. B **22**, 283 (2001).
- ²⁵M. Potthoff and W. Nolting, Eur. Phys. J. B **8**, 555 (1999); Phys. Rev. B **60**, 7834 (1999).
- ²⁶Y. Ohashi and Y. Ōno, J. Phys. Soc. Jpn. **70**, 2989 (2001); *ibid.* Suppl. **71**, 217 (2002).
- ²⁷A. M. Oleś and G. Stollhoff, Phys. Rev. B **29**, 314 (1984).
- ²⁸M. Zöflf, Ph.D. thesis, University of Regensburg, 2001.
- ²⁹Taking into account additionally the insulating ($z=0$) solution for $U > U_{c,1}$ ($U_{c,1} \leq U_{c,2}$), this amounts to *three* coexisting solutions of the mean-field equations in the above interval.