

## Solving gapped Hamiltonians locally

M. B. Hastings\*

Center for Nonlinear Studies and Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

(Received 24 August 2005; revised manuscript received 7 December 2005; published 22 February 2006)

We show that any short-range Hamiltonian with a gap between the ground and excited states can be written as a sum of local operators, such that the ground state is an approximate eigenvector of each operator separately. We then show that the ground state of any such Hamiltonian is close to a generalized matrix product state. The range of the given operators needed to obtain a good approximation to the ground state is proportional to the square of the logarithm of the system size times a characteristic “factorization length.” Applications to many-body quantum simulation are discussed. We also consider density matrices of systems at non zero temperature.

DOI: [10.1103/PhysRevB.73.085115](https://doi.org/10.1103/PhysRevB.73.085115)

PACS number(s): 75.10.Jm, 05.10.Cc

### I. INTRODUCTION

The application of numerical renormalization group to quantum systems is a natural idea with a long history. Despite Wilson’s success with the Kondo model,<sup>1</sup> other early attempts based on keeping low-lying eigenstates in each block were less successful.<sup>2</sup> The basic idea of these methods is to break a system into subsystems, solve each of the subsystems, and then join the solutions together.

This leads to the following general question: how do you solve a system if you know the solution of its subsystems? Consider, for example, the following toy impurity problem: a single spin-1/2 impurity embedded in a band gap insulator. Suppose that the electrons in the insulator do not interact with each other, but only with the impurity spin. The problem of the electrons alone can readily be solved, even in an infinite system, by filling Bloch states up to the Fermi level, and one finds exponentially decaying correlations in this system. If the interaction with the impurity is strong, the impurity problem does not admit an analytic solution, but given the finite correlation length, one might approximately solve a finite periodic region around the impurity on a computer. How, though, can one write down a solution for the combined system? How can one “sew” the two solutions together?

In one-dimension, density matrix renormalization group (DMRG)<sup>3</sup> provides a means to do exactly this and has been extremely successful. The states that it finds are matrix product states.<sup>4–6</sup> There exist some promising higher dimensional generalization of these matrix product states.<sup>4,7</sup> Precise bounds on how well one can approximate a given quantum state by a matrix product state are still lacking, however.

Recently, there have been several advances in understanding the connection between a gap and the locality of correlation functions,<sup>8–12</sup> providing a firm analytical basis for the notion that a gap implies exponentially decaying correlations while a power law density of states implies correlations are bounded by an algebraic decay. In this paper, we will similarly use the existence of a gap to study this problem of sewing states together.

All of the known Hamiltonians that give matrix product states, such as the AKLT<sup>13</sup> Hamiltonian, have the property that the ground state is an eigenvector of each term in the

Hamiltonian separately. We refer to this as a local projective Hamiltonian. The first portion of this paper will construct a form of arbitrary gapped Hamiltonians such that the ground state is an *approximate* eigenvector of each term separately, writing the Hamiltonian as a sum of terms in Eq. (8). This will be a first step to building a matrix product form of the ground state. We construct such a form in this paper, but do not bound the number of states required in the matrix product construction. Such a bound will be given in a future work. While the proofs here will be to some extent constructive, they will assume that certain properties of the ground state are known, and thus they are not so useful in themselves for the problem of finding ground states.

Next, we briefly discuss applications to numerical simulation. Of course, one application is in analyzing existing algorithms, but we suggest the possibility of different algorithms based on the proofs in the first part. In this case, we will discuss how to find the needed properties of the ground state used in the proofs in the first part.

The last portion of the paper will consider systems at nonzero temperature. In this case we will show that the density matrix of the system can be written in a matrix product form, which provides a higher dimensional generalization of the one-dimensional matrix product form for density matrices.<sup>21</sup>

### II. APPROXIMATE LOCAL PROJECTIVE FORM OF THE HAMILTONIAN

Consider the AKLT Hamiltonian,<sup>13</sup>  $H = \sum H_i$ , with  $H_i = \vec{S}_i \cdot \vec{S}_{i+1} + (1/3)(\vec{S}_i \cdot \vec{S}_{i+1})^2$ . This Hamiltonian has an exact matrix product ground state. For a chain of  $N-1$  sites, suppose there are ground states labeled by an index  $\beta$ . Then, a chain of  $N$  sites is supposed to have ground states labeled by an index  $\alpha$ , with  $|\alpha\rangle = \sum_{\beta,s} A_{\alpha,\beta}(s) \beta \otimes s$ , where  $s$  denotes a complete set of states on the  $N$ th site (in this case, there are three such states), and  $A_{\alpha,\beta}(s)$  is the matrix defining the matrix product state. This then gives a wave function which sews the solutions of the two subsystems together. One way to find such wave functions is DMRG, while another is the variational matrix product method. For the AKLT chain, the ground state has  $\alpha=1, 2$  with

$$\sum_s A_{\alpha,\beta}(s)s = \begin{pmatrix} 0 & i\sqrt{2}+\rangle \\ i\sqrt{2}-\rangle & -0 \end{pmatrix}. \quad (1)$$

For more general Hamiltonians, the range of the indices  $\alpha, \beta, s$  may be larger and the matrix  $A_{\alpha,\beta}(s)$  may be different.

This ground state not only minimizes the Hamiltonian  $H$ , but also minimizes each  $H_i$  individually, and thus is an eigenvector of each  $H_i$ . This observation is fundamental to the work in this section. We take an arbitrary Hamiltonian and approximately rewrite it in an *approximate local projective form*, defined to be a form in which the Hamiltonian is a sum of local terms  $M_i$  such that the ground state is close (as defined below) to an eigenvector of each  $M_i$  separately. A Hamiltonian with such a form can truly be solved by solving subsystems separately. Break a chain of  $N$  sites up into two subchains of  $N-m$  and  $m$  sites. For each chain, find the eigenvectors of the Hamiltonian with the correct eigenvalue (the same eigenvalue as the ground state of the full system has for the Hamiltonian of the subchain). The ground state of the full chain will be a linear combination of outer products of the given states in each subchain. The matrix product method realizes this for  $m=1$ .

We consider an arbitrary Hamiltonian  $H$  which is assumed to have some number of degenerate ground states and then a gap  $\Delta E$  to the rest of the spectrum. That is, define  $\Psi_a\rangle$  to be an eigenstate state of  $H$  with energy  $E_a$ , and let  $\Psi_a\rangle$  for  $a=0\cdots n-1$  be  $n$  distinct ground states while for  $a\geq n$  we have  $E_a\geq\Delta E$ . We consider the case  $E_0=E_1=\cdots=E_{n-1}=0$ .<sup>14</sup> We assume that  $H$  obeys the finite range conditions.<sup>8,15</sup> That is, the Hamiltonian  $H$  can be written as a sum of terms  $H_i$  such that each  $H_i$  has a bounded operator norm,  $\|H_i\|\leq J$  for some  $J$  and such that each  $H_i$  acts only on sites within some interaction range  $R$  of site  $i$ ,<sup>16</sup> while there are at most  $S$  sites  $j$  within distance  $R$  of sites  $i$  for any  $i$ . Introduce some metric on the lattice  $d(i,j)$  as the distance between sites  $i$  and  $j$ , while  $d(O,j)$  is defined to be the distance between an operator  $O$  and a site  $j$ : that is, the minimum, over sites  $i$  on which operator  $O$  acts, of  $d(i,j)$ . Then, for any operator  $O_j$  which acts on a site  $j$  with  $d(i,j)>R$ , we have  $[H_i, O_j]=0$ .

We now construct the approximate local projective form, and in the next section construct the ground states. Following,<sup>9</sup> define

$$\tilde{H}_i^0 = \frac{\Delta E}{\sqrt{2\pi q}} \int_{-\infty}^{\infty} dt \tilde{H}_i(t), \quad (2)$$

where

$$\begin{aligned} \tilde{H}_i(t) &\equiv H_i(t) \exp[-(t\Delta E)^2/(2q)], \\ H_i(t) &= \exp(iHt) H_i \exp(-iHt), \end{aligned} \quad (3)$$

with  $q$  to be chosen later. That is,  $H_i(t)$  is defined following the usual Heisenberg evolution of operators, while  $\tilde{H}_i(t)$  is equal to  $H_i(t)$  multiplied by a Gaussian which cuts off the integral in Eq. (2) at times  $t$  of order  $\sqrt{q}/\Delta E$ . The notation of Ref. 9 for these operators  $\tilde{H}_i^0$  is chosen to indicate that we make an approximation (hence the tilde) to the zero frequency (hence the zero) part of the  $H_i$ .

Now, here is the key point of the paper. We claim that  $\tilde{H}_i^0$ , acting on a ground state, gives back another ground state up to some exponentially small difference. To see this, compute  $(\tilde{H}_i^0)_{ab} \equiv \langle \Psi_a, \tilde{H}_i^0 \Psi_b \rangle$ , the matrix element of  $\tilde{H}_i^0$  between states  $\langle \Psi_a$  and  $\Psi_b \rangle$ .<sup>17</sup> A direct computation gives

$$(\tilde{H}_i^0)_{ab} = (H_i)_{ab} \exp\left[-q \left(\frac{E_a - E_b}{\Delta E}\right)^2 / 2\right]. \quad (4)$$

Let

$$P_{\text{low}} = \sum_{0 \leq a \leq n} \Psi_a \langle \Psi_a \quad (5)$$

and

$$P_{\text{high}} = 1 - P_{\text{low}}. \quad (6)$$

Thus,  $P_{\text{low}}$  projects onto the space of ground states while  $P_{\text{high}}$  projects onto the remaining states. Then, from Eq. (4), the norm

$$\|P_{\text{high}} \tilde{H}_i^0 \Psi_a\rangle\| \leq \|H_i^0\| \exp(-q/2) \leq J \exp(-q/2), \quad (7)$$

as claimed.

One important fact is that

$$\sum_i \tilde{H}_i^0 = \sum_i H_i = H. \quad (8)$$

Also,  $H_i$  and  $\tilde{H}_i^0$  have the same matrix elements in the subspace of ground states:  $(\tilde{H}_i^0)_{ab} = (H_i)_{ab}$  if  $0 \leq a \leq n$  and  $0 \leq b \leq n$ . Finally,

$$\|H_i^0\| \leq \|H_i\| \leq J.$$

The  $\tilde{H}_i^0$  are *local* in that the commutator of  $H_i$  with any operator  $O_j$  which acts only on a site  $j$  is exponentially small in  $d(i,j)$ . This follows since Eq. (2) defines  $\tilde{H}_i^0$  as an integral of  $H_i(t)$  over times  $t$ ; the Gaussian in Eq. (3) cuts this integral off for a sufficiently long time while for short time  $H_i(t)$  is local. The precise statement shown in the Appendix is that, for any operator  $O_j$  which acts only on a site  $j$ ,

$$\|[\tilde{H}_i^0, O_j]\| \leq J \|O_j\| \{g(c_1 l, l) + 2 \exp[-(c_1 l \Delta E)^2/(2q)]\}, \quad (9)$$

where  $l=d(H_i, j)$  and the function  $g(c_1 l, l)$  is an exponentially decaying function of  $l/\xi_C$  for some microscopic length scale  $\xi_C$  of order the interaction range  $R$ . The constant  $c_1$  is a characteristic inverse velocity of propagation in the system; the existence of a finite velocity of propagation, as discussed in the Appendix, is essential in showing that  $H_i(t)$  is local for short time. Equation (9) implies that  $\tilde{H}_i^0$  is local in that it has a small commutator with operators which are far enough from  $i$ .

We can further define  $M_i$  to be an approximation to  $\tilde{H}_i^0$  which is truly finite range:  $M_i$  will exactly commute with  $O_j$  if  $d(i, j)$  is greater than a certain range  $l_{\text{proj}}$ . To do this, define

$$M_i = (\Delta E / \sqrt{2\pi q}) \int dt \exp[-(t\Delta E)^2/2] H_i^{\text{trunc}}(t), \quad (10)$$

where

$$H_i^{\text{trunc}}(t) = \exp(iH_{\text{loc}}t) H_i \exp(-iH_{\text{loc}}t), \quad (11)$$

$$H_{\text{loc}} = \sum_{j, d(i,j) \leq l_{\text{proj}} - R} H_j. \quad (12)$$

Thus,  $H_{\text{loc}}$  is the sum of terms  $H_j$  with  $d(i,j)$  less than  $l_{\text{proj}} - R$ , so that  $H_i^{\text{trunc}}(t)$  only acts on sites within distance  $l_{\text{proj}}$  of  $i$ . Thus, the procedure to define the  $M_i$  is very simple: one uses the definition Eq. (3), but as one evolves  $H_i(t)$  one drops terms which involve sites more than  $l_{\text{proj}}$  from  $i$ .

In the Appendix, we show that

$$\begin{aligned} \|\tilde{H}_i^0 - M_i\| &\leq J\{N(l_{\text{proj}})g(c_1 l_{\text{proj}}, l_{\text{proj}}) \\ &+ 2 \exp[-(c_1 l_{\text{proj}} \Delta E)^2/(2q)]\}, \end{aligned} \quad (13)$$

where  $N(l_{\text{proj}})$  is defined to be the number of sites  $j$  with  $l_{\text{proj}} - R < d(i,j) \leq l_{\text{proj}} + R$ . Note also that  $\|M_i\| \leq J$ .

We now pick  $q$ . For a given range  $l_{\text{proj}}$  of the  $M_i$ , we want to minimize  $|P_{\text{high}} M_i \Psi_a\rangle|$ , so that the ground states are approximate eigenstates of the  $M_i$ . By a triangle inequality,

$$\begin{aligned} |P_{\text{high}} M_i \Psi_a\rangle| &\leq |P_{\text{high}} \tilde{H}_i^0 \Psi_a\rangle| + \|\tilde{H}_i^0 - M_i\| \leq J\{\exp(-q/2) \\ &+ N(l_{\text{proj}})g(c_1 l_{\text{proj}}, l_{\text{proj}}) \\ &+ 2 \exp[-(c_1 l_{\text{proj}} \Delta E)^2/(2q)]\}. \end{aligned}$$

To get the best bound, we pick  $q = c_1 l_{\text{proj}} \Delta E$ . Then,

$$|P_{\text{high}} M_i \Psi_a\rangle| \leq \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})], \quad (14)$$

where  $\mathcal{O}$  denotes a quantity of order  $\exp(-l_{\text{proj}}/l_{\text{fac}})$ , with  $l_{\text{fac}}$  being the characteristic *factorization length*. The length  $l_{\text{fac}}$  is equal to the minimum of  $(c_1 \Delta E)^{-1}$  and  $\xi_C$ , and thus for small  $\Delta E$ ,  $l_{\text{fac}} = (c_1 \Delta E)^{-1}$ .

With the given  $q$ , the bound in Eq. (13) becomes

$$\|\tilde{H}_i^0 - M_i\| \leq J[N(l_{\text{proj}})g(c_1 l_{\text{proj}}, l_{\text{proj}}) + 2 \exp(-c_1 l_{\text{proj}} \Delta E/2)].$$

This difference is exponentially small in  $l_{\text{proj}}/l_{\text{fac}}$ , so that difference between the ground state energy per site of  $H = \sum_i H_i$  and that of the Hamiltonian  $M = \sum_i M_i$  is exponentially small in  $l_{\text{proj}}/l_{\text{fac}}$ . Defining  $N$  to be the number of sites  $i$  in the system, if  $N\|\tilde{H}_i^0 - M_i\|$  is less than of order  $\Delta E$ , then the ground state of  $M$  has a nonvanishing projection onto the ground state of  $H$ . This requires an  $l_{\text{proj}}$  which is of order  $\ln(N)$ .

We claim that these  $M_i$  realize the approximate local projective form,

$$H \approx M = \sum_i M_i. \quad (15)$$

We start with the simplest case of only one ground state,  $n = 1$ . Then, Eq. (14) implies that the ground state  $\Psi_0\rangle$  is close to an eigenvector of each  $\tilde{M}_i$ . That is,

$$|M_i \Psi_0\rangle - \langle M_i \Psi_0\rangle| \leq \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})],$$

where  $\langle \dots \rangle$  denotes the ground state expectation value. By picking  $l_{\text{proj}}$  large, we can make this difference as small as desired.

The  $M_i$  give the desired approximate local projective form for the case of a unique ground state. We will show in the next section how to construct matrix product states that are approximate ground states of this Hamiltonian. However, we first consider the case of multiple degenerate ground states.

If there are multiple ground states with a gap to the rest of the spectrum, then the situation is slightly more complicated.

Equation (7) implies that the  $\tilde{H}_i^0$  acting on ground states gives states which are close to ground states, but no longer necessarily implies that the ground states are eigenstates of the  $\tilde{H}_i^0$ . Instead, it depends to some extent on what basis we choose for the ground states. As a simple example, consider the Majumdar-Ghosh Hamiltonian for a one-dimensional spin-1/2 chain:  $H = \sum_i H_i$  with  $H_i = J \sum_i [\vec{S}_i \cdot \vec{S}_{i+1} + (1/2) \vec{S}_i \cdot \vec{S}_{i+2}]$ . This Hamiltonian has two exact ground states; in one state sites  $i=1, 2$  are in a singlet, sites  $i=3, 4$  are in a singlet, and so on, while in the other state sites  $i=2, 3$  are in a singlet, sites  $i=4, 5$  are in a singlet, and so on. Denote the first state by  $\Psi_{\text{even}}\rangle$  and the second state by  $\Psi_{\text{odd}}\rangle$ . Now, the states  $\Psi_{\text{even}}\rangle$  and  $\Psi_{\text{odd}}\rangle$  break translational symmetry; the expectation value  $\langle \Psi_{\text{even}}, H_i \Psi_{\text{odd}} \rangle$  is an alternating function of  $i$ . However, in an infinite system the expectation value  $\langle \Psi_{\text{odd}}, H_i \Psi_{\text{odd}} \rangle$  vanishes. Thus, in the subspace formed by the two vectors  $\Psi_{\text{even, odd}}\rangle$ , the  $H_i$  are diagonal and therefore the  $\tilde{H}_i^0$  are also diagonal in this subspace, so with the given  $M_i$ , there is no problem in this basis of ground states: the states  $\Psi_{\text{even, odd}}\rangle$  are approximate eigenstates of the  $M_i$  for large  $q$ . Of course, as is well known for the Majumdar-Ghosh chain, if we were to pick  $H_i = (J/2)[\vec{S}_i \cdot \vec{S}_{i+1} + \vec{S}_i \cdot \vec{S}_{i-1} + (1/2)\vec{S}_{i-1} \cdot \vec{S}_{i+1}]$ , then the states  $\Psi_{\text{even, odd}}\rangle$  would be exact eigenstates of the  $H_i$ , but let us suppose that we do not know that this form of the Hamiltonian is available.

Suppose instead we choose to form ground states which are eigenvectors of the translation operator by  $\Psi_S\rangle = \Psi_{\text{even}}\rangle + \Psi_{\text{odd}}\rangle$  and  $\Psi_A\rangle = \Psi_{\text{even}}\rangle - \Psi_{\text{odd}}\rangle$ . Then, the  $H_i$  are *not* diagonal in this subspace, and the states  $\Psi_{S,A}\rangle$  are not approximate eigenvectors of the  $\tilde{H}_i^0$ , no matter how large  $q$  is. One way to get around this is to go to an enlarged unit cell of two sites, setting  $H = \sum_j H'_j$  where  $H'_j = H_{2j} + H_{2j+1}$  and then the states  $\Psi_{S,A}\rangle$  are approximate eigenvectors of  $M'_j$ . However, the simplest solution is to use the states  $\Psi_{\text{even, odd}}\rangle$  instead of  $\Psi_{S,A}\rangle$ .

Thus, the important question is: can we simultaneously diagonalize all of the  $M_i$  in the subspace formed by the ground states  $\Psi_a\rangle$  for  $0 \leq a \leq n$ ? If so, then we can ensure that, by the appropriate choice of basis for the ground states, each ground state is an approximate eigenvector of each of the  $M_i$  and we will have

$$|(M_i - \langle \Psi_a, M_i \Psi_a \rangle) \Psi_a\rangle| \leq \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})], \quad (16)$$

If the states  $\Psi_a\rangle$  in this basis break translational symmetry, then the expectation value of  $M_i$  in a state  $\Psi_a\rangle$  may depend

on  $i$  even if  $H$  is translationally invariant. In order to simultaneously diagonalize the  $M_i$  in the subspace of ground states, we need the  $M_i$  to commute in this subspace.

The conditions for the  $M_i$  to commute in this subspace are discussed in an Appendix. We will show that, except for a few artificial examples, the  $M_i$  approximately commute. In particular, if  $H$  is translationally invariant, we will show that the commutator of the  $M_i$  in this subspace vanishes exponentially in the system size, and thus we can pick  $|\Psi_a\rangle$  such that Eq. (16) holds. For  $H$  which are not translationally invariant, we show that most of the  $M_i$  commute in this subspace. More precisely, for any basis of the ground states, define  $o(i)$  to be the operator norm of the off-diagonal part of  $M_i$  in the low energy sector in that basis (the off-diagonal part of  $M_i$  is a matrix which has zeros on the diagonal, but whose off-diagonal elements are the same as  $M_i$ ). We will show in the Appendix how to choose a basis in which we can bound  $\sum_i o(i)$  by Eq. (B3). This implies that we can pick the  $|\Psi_a\rangle$  such that

$$|M_i|\Psi_a\rangle - \langle\Psi_a|M_i|\Psi_a\rangle|\Psi_a\rangle| \leq \mathcal{J}\mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})] + o(i),$$

with the sum of  $o(i)$  bounded.

### III. MATRIX PRODUCT STATE FOR THE LOCAL PROJECTION HAMILTONIAN

We pick a given ground state  $|\Psi_a\rangle$  and approximate it by a matrix product state, with an appropriately bounded error. For simplicity, we consider the case in which we can pick the  $|\Psi_a\rangle$  such that Eq. (16) holds. This includes, as discussed above, all translationally invariant systems, as well as many translationally invariant systems, those without local zero energy excitations as discussed in the Appendix. We review previous work on matrix product or valence bond states, and then provide various constructions of the matrix product state in the given case. The value of  $l_{\text{proj}}$  required to obtain a good approximation to the ground state will be seen to grow as a power of the logarithm of the system size.

#### A. Matrix product and valence bond states

The one-dimensional matrix product state, as discussed above, takes a chain of  $N-1$  sites with a set of ground states  $\beta$ , and constructs the ground states of an  $N$  site chain by  $|\alpha\rangle = \sum_{\beta,s} A_{\alpha,\beta}(s)s \times \beta$ , where  $\beta$  are ground states of an  $N-1$  site chain. Matrix product ground states in one dimension can always be written as valence bond states.<sup>4</sup> Construct an enlarged Hilbert space on each site, labeling states on site  $i$  by two indices  $\alpha_i, \beta_i$ . A wave function is constructed in the enlarged space, such that this wave function  $\psi(\alpha_1, \beta_1, \alpha_2, \beta_2, \dots)$  is a product of wave functions  $\psi(\beta_1, \alpha_2)\psi(\beta_2, \alpha_3), \dots$ . Finally, a map is written on each site from the original Hilbert space  $s$  to the enlarged Hilbert space  $\alpha_i\beta_i$ , and the wave function on the original Hilbert space is defined to be the wave function of the mapped state on the enlarged Hilbert space. To make this concrete, suppose the matrix product construction gives  $|\alpha\rangle = \sum_{\beta,s} A_{\alpha,\beta}(s)s \times \beta$ . Then, set  $\psi(\beta_i, \alpha_{i+1}) = \delta(\beta_i, \alpha_{i+1})$ . Let  $F$  map state  $s_i$  on site  $i$  onto  $\sum_{\alpha_i\beta_i} A_{\alpha_i\beta_i}(s_i)\alpha_i\beta_i$ . Alternately,

this map  $F$  can be viewed as a projection from the space of states  $\alpha_i\beta_i$  onto the states  $s_i$ .<sup>18</sup> Then, the matrix product wave function is given by  $\psi[F(s_1, s_2, \dots)]$ .

In the AKLT case,  $\alpha_i$  labels one of the two spin-1/2's and  $\beta_i$  labels the other one. The product wave function is a product of singlet pairs, while the map  $F$  projects the two spin-1/2's onto a spin-1. The matrix product state for the AKLT chain can be written also as a valence bond state. Each site has a spin-1, which may be represented by two spin-1/2 spins. One spin-1/2 is in a singlet with a spin-1/2 on the next site to the right, and one is in a singlet with a spin-1/2 on the next site to the left. This state is then projected onto the spin-1 state of the two spin-1/2 spins on each site.

Matrix product states and valence bond states are equivalent. However, the discussion above was confined to pure states (wave functions) on finite systems. In Ref. 4, matrix product and valence bond states were also constructed for mixed states (density matrices) and again shown to be equivalent.

Thus far we have discussed systems in one dimension. A higher dimensional system can always be viewed as a one-dimensional system as follows. For a  $d$ -dimensional system in which each site is labeled by  $d$  coordinates, all of the sites with a particular value of one coordinate can be grouped into one super site, leaving a one-dimensional chain. This method is very limited in practice; for a system of linear size  $L$ , the size of the Hilbert space on a single super site is exponential in  $L^{d-1}$  and thus the range of the indices  $\alpha, \beta$  is also exponentially large. This method amounts to studying a one-dimensional ladder system.

Valence bond states are often regarded as a more appropriate way for constructing states in more than one dimension. To construct such a state,<sup>4,7</sup> on an arbitrary lattice in any number of dimensions, for each site  $i$  one constructs one  $k$ -dimensional auxiliary Hilbert space per bond, where the bond connects site  $i$  to site  $j$ . A wave function is defined in the enlarged Hilbert space, which is a product of wave functions on each bond, where the wave function on each bond is a function of states at the "ends" of the bond (in one dimension, these are the indices  $\beta_i, \alpha_{i+1}$ ). For each site, a map is defined from the original Hilbert space to the product of the auxiliary Hilbert spaces on that site. In this paper, we restrict to methods based on super sites for higher dimensions, while a future publication will provide valence bond constructions in this case.<sup>19</sup>

Manipulating these higher dimensional valence bond states is difficult. After Ref. 4, most higher dimensional work involved special examples where the Hamiltonian was exactly equal to a sum of projection operators,<sup>20</sup> so that the ground state is exactly a higher dimensional generalization of the AKLT ground state. However, in an important advance,<sup>7</sup> valence bond states were suggested as a good ansatz for arbitrary Hamiltonians, with a numerical technique being used to compute the state. We now provide a construction of the matrix product or valence bond states.

#### B. Construction of matrix product state

We first give the matrix product construction in one dimension. The approach discussed above in one dimension is




FIG. 1. Illustration of iterative and hierarchical procedures. (a) Iterative procedure. Filled circles represent sites, ovals surrounding sites representing grouping of sites into a super site. The first stage groups sites 1 and 2. The second groups the combined site with site 3, and so on. (b) Hierarchical procedure. The first stage groups pairs of sites, the second stages groups four sites into a single site, and so on.

an iterative procedure: from the ground states of an  $N-1$  site chain, we construct those of an  $N$  site chain. This procedure is shown schematically in Fig. 1(a). We first find the set of allowed states on the first two sites, then the set of allowed states on the first three sites, and so on. Our procedure will instead be a “hierarchical” procedure, as shown in Fig. 1(b). We will find a set of allowed states on pairs of sites; we then find sets of allowed states on groups of four sites, and so on. In practice in DMRG, the iterative approach works better, but we find the hierarchical approach gives better bounds here.

Consider a one-dimensional system with given  $l_{\text{proj}}$ . First group the sites into super sites with sites  $1 \leq i \leq 2l_{\text{proj}}$  grouped into one super site, sites  $2l_{\text{proj}} + 1 \leq i \leq 4l_{\text{proj}} + 1$  into another super site, and so on, grouping  $2l_{\text{proj}}$  sites into each super site. Suppose each site has an  $m$ -dimensional Hilbert space. The dimension of the space of states on the sites  $1 \leq i \leq 2l_{\text{proj}}$  is at most  $m^{2l_{\text{proj}}}$ . From now on, we refer to these super sites simply as “sites;” with this grouping, the operators  $M_i$  act only on pairs of neighboring sites, with no longer range interaction or three site interactions. We let  $M_{k,k+1}$  be the sum of the operators  $M_i$  which act on sites  $k$  and  $k+1$ . Label the super sites of the system by  $i=1, \dots, N$ . The system is periodic, so that site  $N+1$  is identical to site  $i$ ; similarly,  $d(N, 1)=1, d(N, 2)=2$ , and so on.

Pick a given  $\Psi_a$ . Let  $\alpha_k^1$  label a complete basis of states  $\alpha_k^1$  on site  $k$ . From Eq. (16),

$$|(M_{k,k+1} - \langle \Psi_a, M_{k,k+1} \Psi_a \rangle) \Psi_a| \leq J l_{\text{proj}} \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})].$$

Let  $P_k^1$  project onto the eigenvectors of  $M_{k,k+1}$  with eigenvalues  $\lambda$  such that  $|\lambda - \langle \Psi_a, M_{k,k+1} \Psi_a \rangle| \leq x$ , for some  $x$  to be chosen later. Let  $\alpha_k^2$  label these different eigenvectors  $\alpha_k^2$ . Then,

$$\begin{aligned} x \langle \Psi_a, (1 - P_k^1) \Psi_a \rangle &\leq |(M_{k,k+1} - \langle \Psi_a, M_{k,k+1} \Psi_a \rangle) \Psi_a| \\ &\leq J l_{\text{proj}} \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})]. \end{aligned}$$

Then, choose  $x = J l_{\text{proj}} \mathcal{O}[\exp(-l_{\text{proj}}/[\log_2(N)l_{\text{fac}}])]$ , so that

$$\langle \Psi_a, P_k^1 \Psi_a \rangle \geq 1 - b_1, \quad (17)$$

where  $b_1 = \exp[-[1 - 1/\log_2(N)](l_{\text{proj}}/l_{\text{fac}})]$ . This implies that  $M_{k,k+1}$  has at least one eigenvalue  $\lambda$ , such that

$$|\lambda - \langle \Psi_a, M_{k,k+1} \Psi_a \rangle| \leq J l_{\text{proj}} \mathcal{O}[\exp\{-l_{\text{proj}}/[\log_2(N)l_{\text{fac}}]\}].$$

Therefore, there is at least one state  $\alpha_k^2$  for each site  $k$ .

For each pair of sites  $k, k+1$  surrounded by an oval on the second line of Fig. 1(b), we calculate the  $P_k^1$  of the preceding paragraph. Then, for each group of four sites,  $k, k+1, k+2, k+3$  surrounded by any oval on the third line of Fig. 1(b), we let  $P_k^2$  project onto the eigenvectors of  $P_k^1 P_{k+2}^1 M_{k+1, k+2} P_{k+2}^1 P_k^1$  in the space of states  $\alpha_k^2 \otimes \alpha_{k+2}^2$ , such that the eigenvector has eigenvalue  $\lambda$  such that

$$\begin{aligned} |\lambda - \langle \Psi_a, M_{k+1, k+2} \Psi_a \rangle| &\leq J l_{\text{proj}} \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})] \\ &\quad + 2b_1^{1/\log_2(N)}. \end{aligned}$$

Let  $\alpha_k^3$  label the resulting eigenvectors  $\alpha_k^3$ . Define  $\Psi_a^1 = P_k^1 P_{k+2}^1 \Psi_a / \sqrt{\langle \Psi_a, P_k^1 P_{k+2}^1 \Psi_a \rangle}$ . This vector is normalized to unit norm. Then,  $|\Psi_a^1 - \Psi_a| \leq 2b_1$ . Thus,

$$\begin{aligned} |(M_{k+1, k+2} - \langle \Psi_a^1, M_{k+1, k+2} \Psi_a^1 \rangle) \Psi_a^1| \\ \leq J l_{\text{proj}} \mathcal{O}[\exp(-l_{\text{proj}}/l_{\text{fac}})] + 2b_1 J l_{\text{proj}}. \end{aligned}$$

Therefore,  $\langle \Psi_a^1, P_k^2 \Psi_a^1 \rangle \geq 1 - b_2$ , where  $b_2 = [\exp(-l_{\text{proj}}/l_{\text{fac}}) + 2b_1]^{1-1/\log_2(N)}$ .

Proceeding in this fashion, we find that  $\langle \Psi_a^{m-1}, P_k^m \Psi_a^{m-1} \rangle \geq 1 - b_m$ , where  $b_m = [\exp(-l_{\text{proj}}/l_{\text{fac}}) + 2b_{m-1}]^{1-1/\log_2(N)}$ . There are  $h \equiv \log_2(N)$  levels of this construction. Thus, after the last step,  $b_h$  is bounded by a quantity of order  $N \exp[-(l_{\text{proj}}/l_{\text{fac}}) \times (1 - 1/h)^h] \sim N \exp(-l_{\text{proj}}/el_{\text{fac}})$ . Choose  $l_{\text{proj}}$  such that  $l_{\text{proj}}/l_{\text{fac}} \geq \mathcal{O}[\ln(J/\Delta E) \ln(N)^2]$ . Then, there is at least one state  $\alpha^h$ , and one may show that

$$\begin{aligned} \left\langle \alpha^h, \sum_i M_i \alpha^h \right\rangle &\leq \left\langle \Psi_a, \sum_i M_i \Psi_a \right\rangle \\ &\quad + NJ \mathcal{O}[\exp\{-l_{\text{proj}}/[\log_2(N)l_{\text{fac}}]\}]. \end{aligned} \quad (18)$$

Thus, the energy of this state  $\alpha^h$  will be within  $\Delta E$  of the ground state energy.

Thus, this procedure yields a matrix product state close to the ground state in energy. At the same time, this procedure does not yield too many states. All of the states  $\alpha^h$  are within energy of order  $\Delta E$  of the ground state. We can choose  $l_{\text{proj}}$  so that they lie within energy  $\Delta E/2$  of the ground state; it can then be shown that the number of distinct  $\alpha^h$  is at most  $2n$ , where  $n$  is the number of ground states. By choosing the bound on the expectation value of the energy even smaller, the ground state may be approximated to arbitrary accuracy.

This procedure can be extended to systems in more than one dimension. One possibility is to treat a higher dimensional system as a one-dimensional system by grouping all the sites with a particular value of a coordinate into a super site as described in the section reviewing the matrix product and valence bond constructions. Another possibility is to group  $2^d$  sites together at each stage of the hierarchy. In

either case, it is still only necessary to take an  $l_{\text{proj}}$  which grows as a power of  $\ln(N)$  to get a good approximation to the ground state.

#### IV. QUANTUM SIMULATION

One application to quantum simulation of matrix product or valence bond states in higher dimensions is variational,<sup>7</sup> and the results here may be useful in analyzing such algorithms. However, another application of these results to quantum simulation involves using the construction of the preceding sections. The results in this section are illustrative, and will be worked out for more practical examples in a future presentation.<sup>19</sup> The goal here is to show, for at least some simple examples, that in principle one can actually use computations on finite systems to write down wave functions for much larger, or even infinite, systems, and to provide a variational technique that gives a *lower* bound on the energy.

To do this, we propose to calculate the matrices  $M_i$ , determine the correct eigenvalue of each  $M_i$ , and then use this to determine the ground state wave function for a large system. At this point, there is a very natural objection. The procedure requires that we find the correct eigenvalue of the  $M_i$ . If we can do this, then we know the ground state energy of the system (up to an exponentially small error). If  $l_{\text{proj}}$  is sufficiently large to obtain approximately correct eigenvalues for the  $M_i$ , then why not just do an exact diagonalization of the system on a system of size  $l_{\text{proj}}$  and compute the ground state energy that way? One answer is that the ground state energy is not the only interesting aspect of the system. Correlation functions are much more important, and it is often a difficult task to determine the long-range order from the quantum numbers (such as momentum and spin) of the low-lying states found in exact diagonalization. The procedure we outline of sewing together solutions will provide a way of taking a solution on a finite size system and extending it to a wave function for a system of much larger, or even infinite, size. This wave function can then be used to compute long-distance behavior of correlation functions. Also, as we will see below, in some cases this yields better energy estimates than exact diagonalization of the same size system.

We now discuss the first step, computing the  $M_i$ . One way is to directly calculate the  $M_i$ , using Eq. (10) and using Eq. (11) to define  $H_i^{\text{trunc}}(t)$ . One way to compute  $H_i^{\text{trunc}}(t)$  is via a series expansion  $H_i^{\text{trunc}}(t) = H_i + it[H^{\text{loc}}, H_i] + \dots$ . Another way is to exactly diagonalize a finite system of size  $l_{\text{proj}}$ , compute the matrix elements of  $H_i$  between the eigenvalues, and use Eq. (4) to get matrix elements of  $\tilde{H}_i^0$ . In both cases it is necessary to choose  $q$  to minimize the difference  $\|\tilde{H}_i^0 - M_i\|$  given by Eq. (13) and also the norm  $|P_{\text{high}}\tilde{H}_i^0\Psi_a\rangle$  of Eq. (7), giving a  $q$  of order  $c_1 l_{\text{proj}} \Delta E$ .

Given that the ground state is an approximate eigenstate of the  $M_i$  computed in this way, the question arises: what is the eigenvalue? That is, what is the approximate expectation value of  $\langle H_i \rangle$  in the ground state? In some cases, finding the correct eigenvalue is an extremely difficult problem. For example, consider an Ising spin-glass Hamiltonian,  $H = \sum_i H_i$  with  $H_i = \sum_j J_{ij} S_i^z S_j^z$ , where  $J_{ij}$  is some set of random couplings

between nearby spins. This is a purely classical problem, since all of the  $H_i$  commute, and any state in which each spin has a definite value of  $S^z$  is an eigenvector of every  $H_i$ . However, to find the ground state is clearly a difficult task. In this highly disordered system, each  $H_i$  has a different expectation value and one must find the correct eigenvalue for each one. For ordered systems the task is much easier. If the ground state does not break translational symmetry then each  $M_i$  has the same ground state expectation value. If there is a symmetry breaking ground state with an enlarged unit cell, there are still only a discrete number of different expectation values for the  $M_i$ . For example, in the Majumdar-Ghosh chain, the ground states are invariant under translation by two sites, and there are two different ground state expectation values for the  $M_i$ .

We have performed some simple numerical experiments on systems of free particles on a lattice with different gapped band structures. In this case there is no symmetry breaking, and if  $H$  is translationally invariant, then the expectation values of the  $M_i$  in the ground state are also translationally invariant. The lowest eigenvalue of  $\tilde{H}_i^0$  for given  $q$  always provides a lower bound to the energy as in Eq. (19). However, we have found that as  $q$  is increased, the lowest eigenvalue increased and converges rapidly to the ground state energy per site. There are several other eigenvalues very close in energy to the lowest, and then a gap to the rest, so in this case at least the identification of the correct eigenvalue is easy and this technique in fact provides a way to compute the energy per site. An upper bound to the energy per site as well as a macroscopic wave function can be obtained by using one of the matrix product constructions above. The most important question, of course, is how well this procedure can be extended to complicated interacting systems.

Although the main point of the present paper is the formal construction of the  $M_i$ , we finally discuss here two brief attempts to apply these techniques to interacting systems. First, the techniques here provide a variational lower bound to the energy of the system. Consider a translationally invariant Hamiltonian,  $H = \sum_i H_i$ , with all the  $H_i$  equal. Then, the ground state energy per site,  $E_0/N$  is at least equal to the smallest eigenvalue of  $H_i$ . Further,

$$E_0/N \geq \lambda_{\min}(H_i + ia_1[H_i, H] + a_2[[H_i, H], H] + \dots), \quad (19)$$

where  $\lambda_{\min}(O)$  is equal to the smallest eigenvalue of  $O$ , and  $a_1, a_2, \dots$  are arbitrary constants. The operators  $\tilde{H}_i^0$  are given by a particular choice of the constants  $a_k$  that can be obtained from Eq. (2). Other choices are possible, and varying over the constants will provide a variational lower bound for the energy. Thus, this provides an interesting complementary approach to other quantum simulation techniques, since almost all other techniques provide either approximate estimates or variational *upper* bounds.

We have studied how this bound is approached on a spin-1 Heisenberg chain. The ground state energy of this chain is known very accurately from DMRG. The present method (19) here is not intended to compare to DMRG, but rather we compare to exact diagonalization. In essence, this

method provides another type of boundary condition, instead of the usual periodic or antiperiodic boundary condition, with the advantage that in this case we know rigorously how the ground state energy compares to the energy from this procedure. We have some freedom to pick the constants  $a_1, a_2, \dots$ . We also have freedom to choose  $H_i$  to be an operator on a pair of neighboring sites, on three neighboring sites, or in general on any supercell of  $m$  sites. Since  $H$  is real symmetry in this case, we pick  $a_k=0$  for  $k$  odd. If  $m=2$ , at the most trivial level of  $a_k=0$  for all  $k$ , we need to diagonalize a Hamiltonian with  $n=2$  sites. For  $a_2 \neq 0$ ,  $a_k=0$  for  $k>2$ , we need to diagonalize a Hamiltonian with  $n=6$  sites, and so on. If we instead pick  $m=3$  we need to diagonalize a Hamiltonian with  $n=3$  sites if all  $a_k=0$ , a Hamiltonian with  $n=7$  sites if  $a_2 \neq 0$ ,  $a_k=0$  for  $k>2$ , and a Hamiltonian with  $n=11$  sites if  $a_2 \neq 0$ ,  $a_4 \neq 0$ ,  $a_k=0$  for  $k>2$ .

We have chosen to take  $m$  odd for the following reason. This technique provides a lower bound for the energy of the Hamiltonian, both for the infinite system and for the particular  $n$ -site Hamiltonian. However, we know for the spin-1 chain that on even size systems the energy obtained is already less than the ground state energy. Thus, the most efficient results will be obtained for odd size systems, and hence we pick an odd  $m$ .

For  $m=3$  and  $a_k=0$  for all  $k$  we find  $E_0/N \geq -1.5$ , which is equal to one-half the energy of an open three site chain. For  $m=3$  and  $a_2 \neq 0$ , with all higher  $a_k=0$ , we find  $E_0/N \geq -1.42569$  after picking  $a_2 = -0.075$ . This requires diagonalizing a seven site system; the most naive estimate that can be obtained from a seven site system, taking  $m=7$  gives a worse bound of  $E_0/N \geq -1.43909$ . For  $m=5$ ,  $a_2 = -0.075$  and all higher  $a_k=0$ , we find  $E_0/N \geq -1.4156$ . Slight improvements on these bounds can be found by better optimization of  $a_2$ . This last estimate requires diagonalizing a nine site system. One we move to an 11 site system, we have the option of either considering  $m=7$ ,  $a_2 \neq 0$ , or  $m=5$ ,  $a_2 \neq 0$ ,  $a_4 \neq 0$ , with all higher  $a_k$  vanishing.

For purposes of computing the energy, then, this technique offers some slight improvements over exact diagonalization. Using exact diagonalization with periodic boundary conditions in this particular case, even size systems offer lower bounds, while odd size systems offer upper bounds, while we find a lower bound in every case. The estimate from a nine site system using this method, for example, is better than that found using exact diagonalization of an eight site system, where the energy is estimated to be  $-1.417$ , but not as good as that found by diagonalizing a 10 site system, where one finds  $-1.4094$ , while for a seven site system the estimate is better than that found from diagonalizing a six site system, where one finds  $-1.44$ . Further this method offers the only way to obtain rigorous lower bounds. This may become especially important in studying frustrated systems. On a frustrated system with spiral order, for example, one has no foreknowledge that exact diagonalization of a periodic chain of a given length will provide a lower or upper estimate on the energy. The energy estimates of this unfrustrated chain obtained from exact diagonalization of odd size systems are poor compared to those obtained by exact diagonalization of even size systems; on a frustrated chain one has no notion of which sizes will yield accurate estimates of the energy.

In one dimension on unfrustrated systems, then, this technique does not give much improvement on the energy, as one can obtain a better estimate by exact diagonalization of a system of one site more. However, in higher dimensions, even on an unfrustrated system, this technique may become more useful again. Suppose we have an unfrustrated system of size  $L$ -by- $L$  in higher dimensions, and suppose it follows the pattern found here, that the most accurate estimate of the energy from exact diagonalization is found from periodic systems with even  $L$  where one obtains a lower bound. The present technique offers the possibility of obtaining accurate estimates of the energy from a system of size  $L-1$  instead of size  $L$ , which means studying a system with  $2L-1$  fewer sites.

We know that the ground state wave function of the full system has a bounded projection onto states other than those with close to the given eigenvalue of  $H_i + a_2[[H_i, H], H] + \dots$ . Indeed, for the particular choice of Eq. (2), the projection onto such states is provably exponentially small in the size of the system considered. Thus, one can follow a procedure of breaking a chain or lattice into blocks, building the  $M_i$  for each block and diagonalizing it in each block, restricting to the states with the given eigenvalues in each block, and then studying the behavior of the full Hamiltonian in this reduced space of states. As a first test of this algorithm, we take  $m=3$ . If we simply take the project onto the states with lowest eigenvalues of the three site Hamiltonian, all  $a_k=0$ , this provides a poor approximation to the eigenstates of larger systems. For example, on a seven site chain, with the three sites taken in the middle of the chain, there are  $3^5 = 243$  states such that the three site Hamiltonian has an energy per bond given by  $E/2 = -1.5$ . The Hamiltonian of the seven site chain with open boundary conditions has a lowest energy state with energy per bond equal to  $-1.43909$ , but if we project onto the given 243 states above, we only achieve an energy per bond of  $-1.36496$ . However, if we take  $a_2 = -0.075$  with  $a_k=0$  for  $k>2$ , there are 243 eigenvalues of the Hamiltonian with  $E/2 \leq -1.34$ , and then the 244th state has eigenvalue  $-0.945706$ . Projecting the Hamiltonian of the seven site chain with open boundary onto these 243 states we find a energy per bond of  $-1.43331$ . Thus, we have accurately selected the needed states, having projected from 2187 states to 243 states, or from 393 states with  $S_z=0$  to 51 states with  $S_z=0$ .

A further test breaking a 14 site periodic spin-1 chain into seven site blocks and taking  $m=3$  and  $a_2 = -0.075$  to project onto states in each block showed that the lowest energy wave function in this subspace had energy per site equal to  $-1.39545$ , compared to the exact result of  $-1.40394$  for this size. If instead of joining subblocks we had added sites to a subblock one at a time, and used the present method of projecting onto the states in each subblock, we would arrive at an algorithm similar to DMRG. However, the goal is not to compare to DMRG in one dimension, but rather to present an algorithm that can be extended to higher dimensions, which DMRG cannot.

A second technique, which may also offer the possibility of improved accuracy on the energies compared to exact diagonalization is based on the idea that the operators  $\tilde{H}_i^0$  are

defined by the choice of constants  $a_k=0$  for  $k$  odd, and  $a_k = (-1)^{k/2}[(k-1)!!/k!](q/\Delta E^2)^k$  for  $k$  even. We then have a series expansion for  $\tilde{H}_i^0$  in powers of  $q$ . We can then perturbatively expand the eigenvalues of  $\tilde{H}_i^0$  in powers of  $q$  and extrapolate to  $q=\infty$  from a finite number of terms. This is a speculative approach that is currently being studied.

## V. NONZERO TEMPERATURE

We now turn to systems at nonzero temperature. In this case the temperature enables us to construct an approximate matrix product form for the density matrix, regardless of whether or not there is a gap. The unnormalized density matrix for the system is equal to  $\rho = \exp(-\beta H)$ .

We will construct an approximate matrix product form,  $\rho(\beta, l_{\text{proj}})$ , so that

$$\rho(\beta) \approx \rho(\beta, l_{\text{proj}}) = \sum_{\{\alpha_i\}} \rho_1(\alpha_1) \rho_2(\alpha_2) \cdots F_1(\{\alpha_j\}) F_2(\{\alpha_j\}) \cdots, \quad (20)$$

where for each site  $i$  we assign an index  $\alpha_i$  defined below, and sum over all values of that index. The operator  $\rho_i(\alpha_i)$  acts only on site  $i$ , and the functions  $F_i$  obey a finite range constraint: each  $F_i$  depends only on the  $\alpha_j$  for  $d(i, j) \leq l_{\text{proj}}$  for some  $l_{\text{proj}}$  defined below. The error between  $\rho(\beta)$  and  $\rho(\beta, l_{\text{proj}})$  will be exponentially small in  $l_{\text{proj}}$ , while the range of the indices  $\alpha_i$  will depend on  $l_{\text{proj}}$ . Specifically, we bound the error by showing that for any operator  $O$ ,  $Z^{-1} |\text{Tr}[O\rho(\beta)] - \text{Tr}[O\rho(\beta, l_{\text{proj}})]| \leq c \|O\|$ , for some constant  $c$ , where we define  $Z = \text{Tr}[\rho(\beta)]$ .

Before defining  $\rho(\beta, l_{\text{proj}})$ , we recall the Trotter-Suzuki<sup>22</sup> decomposition of the path integral. In this case, we write  $\rho(\beta) \approx \rho_n$ , where  $\rho_n = [\prod_i \exp(-\beta H_i/n)]^n$ , where the product ranges over all  $i$  in some given sequence; since the different  $H_i$  do not commute, the result depends on the particular sequence chosen. We claim that each of the  $\rho_n$  can be written exactly in a matrix product form as in the right-hand side of Eq. (20). A given operator  $\exp(-\beta H_i/n)$  acts on sites within a distance  $R$  of site  $i$ , and can be written as  $\exp(-\beta H_i/n) = \sum_{\{\alpha_{i,j}\}} F_i(\{\alpha_{i,j}\}) \prod_{j, d(i,j) \leq R} O_j(\alpha_{i,j})$ , where the operator  $O_j$  acts only on site  $j$  and the range of values of index  $\alpha_{i,j}$  is exponentially large in  $S$ , the number of sites within distance  $R$  of site  $i$ . Here,  $F_i(\{\alpha_{i,j}\})$  is some function of the  $S$  different indices  $\alpha_{i,j}$  with the given  $i$ . The operator  $\rho_n$  is a finite product of these operators  $\exp(-\beta H_i/n)$ . Each term in this product can be written in matrix product form. For each site  $i$ , the operator  $\exp(-\beta H_i/n)$  appears  $n$  times in the given product, and on the  $m$ th time it appears, we use a set of indices  $\alpha_{i,j}^m$  to provide the matrix product form as above. For each site,  $j$ , we have  $n$  different indices  $\alpha_{i,j}^m$  for each site  $i$  within distance  $R$  of site  $j$ , and thus at most  $nS$  different indices for each site  $j$ . Grouping all such indices for a given site  $j$  into one index  $\alpha_j$ , and defining  $F_i(\{\alpha\}) = \prod_{m=1}^n F_i(\{\alpha_{i,j}^m\})$ , we arrive at the matrix product form for  $\rho_n$ .

Thus, the Trotter decomposition gives an approximate matrix form, but the error in this approximation compared to the exact result is not very good. In contrast, the stochastic

series expansion<sup>23</sup> provides a much better way of approximating the desired  $\rho(\beta)$  with much smaller error, but it is difficult to write the stochastic series expansion result in a matrix product form. Below, we will propose a matrix product form with a bound on error comparable to that in the stochastic series expansion.

## A. Percolation transition at high temperature

The exponential  $\exp(-\beta H)$  can be expanded as a power series  $1 - \beta H + \cdots$ . We will show that at sufficiently high temperatures,  $\beta^{-1} \sim J$ , there is a ‘‘percolation transition’’ in this exponential, as we now describe. Any given term in the power series expansion is a product of  $H_i$  for different sites  $i$ . For each term, we define a set of ‘‘active bonds’’ and ‘‘clusters’’ as follows: for each  $H_i$  which appears in the given term, we connect by active bonds all sites acted on by  $H_i$ , so that the length of the active bonds is at most  $2R$ . We define a cluster to be a set of sites, all connected to each other by active bonds, and not connected to any other sites outside the cluster by active bonds. Then, for  $\beta \leq \beta_0$ , where  $\beta_0$  is specified below, define  $\rho(\beta, l_{\text{proj}})$  to include only the terms in the power series such that no two sites  $i, j$ , with  $d(i, j) > l_{\text{proj}}$ , are in the same cluster. Thus, each term in  $\rho(\beta, l_{\text{proj}})$  is a product of operators, each operator acting on the sites within a given cluster, such that each cluster has a diameter at most  $l_{\text{proj}}$ . See Fig. 2. For  $\beta \leq \beta_0$ , at temperatures above the percolation transition, we will be able to bound the difference between  $\rho(\beta)$  and  $\rho(\beta, l_{\text{proj}})$ .

Let  $C$  be some set of sites  $i$ . Define  $H_C = \sum_{i \in C} H_i$ . Define  $B(C)$  to be equal to the set of all sites  $j$  such that  $j \neq i$  for any  $i \in C$  and such that  $H_j$  acts on a site  $i$  for some site  $i \in C$ . Then,  $H_{B(C)} = \sum_{i \in B(C)} H_i$ . Then, if a given term in the power series expansion for  $\exp(-\beta H)$  includes  $H_i$  for every  $i \in C$ , but does not include  $H_j$  for all  $j \in B(C)$ , then this term in the power series expansion has a cluster which includes exactly the sites acted on by  $H_i$  for all  $i \in C$ .

For any operators  $O_1, O_2, \dots$ ,

$$Z^{-1} \text{Tr}[O O_1(i\tau_1) O_2(i\tau_2) \cdots \exp(-\beta H)] \leq \|O\| \|O_1\| \|O_2\| \cdots,$$

where  $O(i\tau) = \exp(-H\tau) O \exp(H\tau)$  and  $0 \leq \tau_1 \leq \tau_2 \leq \cdots \leq \beta$ . Thus, if  $C$  has  $n_C$  sites and  $B(C)$  has  $n_{B(C)}$  sites, we find that for any operator  $O$ ,

$$Z^{-1} \text{Tr}\{O \exp[-\beta(H - H_C - H_{B(C)})]\} \leq \|O\| \exp[(n_C + n_{B(C)})J\beta], \quad (21)$$

as can be seen by using a power series expansion of

$$\begin{aligned} & Z^{-1} \text{Tr}\{O \exp[-\beta(H - H_C - H_{B(C)})]\} \\ &= Z^{-1} \text{Tr} \left[ O T \exp \left( \int_0^\beta d\tau [H_C(i\tau) + H_{B(C)}(i\tau)] \right) \right. \\ & \quad \left. \times \exp(-\beta H) \right], \end{aligned}$$

where  $T$  gives the  $\tau$ -ordered exponential.

We now define  $\rho(\beta, C)$  to be equal to the sum of all terms in the power series expansion of  $\exp(-\beta H)$  that include  $H_i$




FIG. 2. Example of a set of active bonds shown as solid lines; dashed lines represent bonds which are not active while circles represent sites. The Hamiltonian is a sum of terms which act on pairs of neighboring sites, so that the bonds connect only neighboring sites. Using a Manhattan metric for the lattice, the set of active bonds shown here is a term in  $\rho(\beta, l_{\text{proj}}=2)$ . There are four distinct clusters, three with diameter two and one with diameter one.

for all  $i \in C$ , but do not include  $H_j$  for any  $j \in B(C)$ . For any  $O$ ,

$$\begin{aligned} \text{Tr}[O\rho(\beta, C)] &\leq [\exp(J\beta) - 1]^{n_C} \\ &\quad \times \exp(\beta\|H_C + H_{B(C)}\|)\text{Tr}[O \exp(-\beta H)], \end{aligned}$$

as can be seen by a power series expansion. Using  $\|H_C + H_{B(C)}\| \leq (n_C + n_{B(C)})J \leq SN_C$  gives

$$Z^{-1} \text{Tr}[O\rho(\beta, C)] \leq \|O\| [\exp(J\beta) - 1]^{n_C} \exp(n_C S J \beta). \quad (22)$$

Then,  $\rho(\beta) - \rho(\beta, l_{\text{proj}})$  is a sum over terms which include a sequence of active bonds connecting any two sites  $i, j$  separated by a distance at least  $l_{\text{proj}}$ . This difference  $\rho(\beta) - \rho(\beta, l_{\text{proj}}) = -\sum_{m=1}^{\infty} (-1)^m \rho_m$ , where  $\rho_m$  is equal to the sum of all terms in the power series expansion of  $\rho$  which include at least  $m$  clusters with diameter greater than  $l_{\text{proj}}$ . Using Eq. (22),  $Z^{-1} \text{Tr}(O\rho_1) \leq \sum_{\text{clusters}} y^{n_C} \|O\|$ , where the sum ranges over all different clusters on the lattice which connect two sites separated by distance greater than  $l_{\text{proj}}$  and  $n_C$  is the number of sites in the cluster and where  $y = [\exp(J\beta) - 1] \exp(SJ\beta)$ . Similarly,  $Z^{-1} \text{Tr}(O\rho_m) \leq (1/m!) \times (\sum_{\text{clusters}} y^{n_C})^m \|O\|$ .

We now show the ‘‘percolation transition.’’ The number of clusters of  $n_C$  sites is bounded on a regular lattice by  $Nx^{n_C}$  for some constant  $x$ , as is known for lattice animals,<sup>24</sup> where  $N$  is the number of sites on the lattice. Thus for sufficiently small  $\beta$ ,

$$\sum_{m=1}^{\infty} (1/m!) \left( \sum_{\text{clusters}} y^{n_C} \right)^m \leq \exp \left( N \sum_{n=l_{\text{proj}}+1}^{\infty} (xy)^n \right) - 1.$$

For fixed  $J, S$ , we can make  $y$  as close to zero as desired by taking sufficiently small  $\beta$ . Thus, there exists some small enough  $\beta_0$  such that for  $\beta \leq \beta_0$ ,

$$\sum_{m=1}^{\infty} (1/m!) \left( \sum_{\text{clusters}} y^{n_C} \right)^m \leq \exp[N \exp(-l_{\text{proj}}/\xi_{\text{perc}})] - 1,$$

where  $\xi_{\text{perc}}$  is of order the interaction range  $R$ . Then,

$$\begin{aligned} Z^{-1} |\text{Tr}[O\rho(\beta)] - \text{Tr}[O\rho(\beta, l_{\text{proj}})]| \\ \leq \{ \exp[N \exp(-l_{\text{proj}}/\xi_{\text{perc}})] - 1 \|O\|. \end{aligned} \quad (23)$$

### B. Matrix product form at high temperature

We now show that  $\rho(\beta, l_{\text{proj}})$  realizes the matrix product form Eq. (20) for  $\beta \leq \beta_0$ . Each term in  $\rho(\beta, l_{\text{proj}})$  is a product of terms acting on clusters of sites. We now specify the indices  $\alpha_i$ . First, if site  $i$  is in a cluster, the index  $\alpha_i$  indicates all the sites in that cluster. There is some redundancy in this description: if two sites  $i$  and  $j$  are in the same cluster, then the indices  $\alpha_i$  and  $\alpha_j$  specify the same set of sites. Each  $\alpha_i$  will also specify some additional information. The sum of all terms which include a given cluster is some operator acting on the sites in the cluster; this operator can be decomposed into a sum of products of operators that act on individual sites in the cluster. The indices  $\alpha_i$  will also keep track of the terms in this sum. The number of different values that each index  $\alpha_i$  can assume is exponentially large in  $l_{\text{proj}}^d$  in this construction. However, this is much smaller than the number of different values that would be needed to describe a general operator  $\rho$ : in general, one might need a range of values which is exponentially large in the system size. Thus,  $\rho(\beta, l_{\text{proj}})$  can be exactly written in a matrix product form, as on the right-hand side of Eq. (20).

Further, we have shown that  $\rho(\beta, l_{\text{proj}})$  is a good approximation to  $\rho(\beta)$ . This can be expressed in terms of the trace norm. For any operator  $O$ , the trace norm  $|O|$  is equal to  $\text{Tr}(\sqrt{O^\dagger O})$ , where  $\text{Tr}(\cdots)$  denotes the trace over all states and the unique positive square-root is taken. For a Hermitian operator, such as  $\rho(\beta)$  or  $\rho(\beta, l_{\text{proj}})$ ,  $|O|$  is equal to the sum of the absolute values of the eigenvalues. For a positive definite Hermitian operator, such as  $\rho(\beta)$ , the trace norm is equal to the trace. Use Eq. (23), valid for arbitrary  $O$ , and choose  $O$  so that in a basis of eigenvectors of  $\rho(\beta) - \rho(\beta, l_{\text{proj}})$ ,  $O$  is a diagonal matrix with each diagonal entry equal to  $\pm 1$ , the sign being chosen the same as the sign of the corresponding eigenvalue. Then,

$$|\rho(\beta) - \rho(\beta, l_{\text{proj}})| \leq \{\exp[N \exp(-l_{\text{proj}}/\xi_{\text{perc}})] - 1\} |\rho(\beta)|. \quad (24)$$

By taking an  $l_{\text{proj}}$  that is of order  $\ln(N)\xi_{\text{perc}}$ , we can obtain a small error  $|\rho(\beta) - \rho(\beta, l_{\text{proj}})|$ .

We now obtain a matrix product form at arbitrary temperature  $\beta$  that guarantees positive definiteness of  $\rho(\beta, l_{\text{proj}})$ . For  $\beta > \beta_0$ , set

$$\rho(\beta, l_{\text{proj}}) = \rho(\beta/n, l_{\text{proj}})^n, \quad (25)$$

where  $n$  is the smallest even integer such that  $\beta/n \leq \beta_0$ . By picking  $n$  even we guarantee that  $\rho(\beta, l_{\text{proj}})$  is positive definite. Equation (25) realizes a matrix product form (20): for each site  $i$  there are  $n$  different indices  $\alpha_i$ , one from each of the  $\rho(\beta/n, l_{\text{proj}})$ . We group these into one single index, and call the new index  $\alpha_i$ , giving Eq. (20). The number of different values that the new  $\alpha_i$  can assume is exponentially large in  $(\beta/\beta_0)^{l_{\text{proj}}}$ . We can bound the error at arbitrary temperature as follows.

For each  $\rho(\beta/n, l_{\text{proj}})$  we define clusters. We can show, as above, that

$$Z^{-1} \text{Tr}\{O\rho(k\beta/n)\rho(\beta/n, C)\rho[(n-k-1)\beta/n]\} \leq \|O\|y^{nc},$$

for any  $0 \leq k \leq n-1$ , where  $y = [\exp(J\beta/n) - 1]\exp(SJ\beta/n)$ .

Define  $\rho_m$  to be the sum over all terms in which there are at least  $m$  clusters with diameter greater than  $l_{\text{proj}}$ . Then,

$$Z^{-1} \text{Tr}(O\rho_m) \leq (1/m!) \left( \sum_{\text{clusters}} ny^{nc} \right)^m \|O\|.$$

Following the same arguments as before

$$\begin{aligned} Z^{-1} |\text{Tr}[O\rho(\beta)] - \text{Tr}[O\rho(\beta, l_{\text{proj}})]| \\ \leq \{\exp[(\beta/\beta_0)N \exp(-l_{\text{proj}}/\xi_{\text{perc}})] - 1\} \|O\| \end{aligned} \quad (26)$$

and so we need an  $l_{\text{proj}}$  that grows logarithmically in the system size to obtain a good approximation. At the value of  $l_{\text{proj}}$  that gives a good approximation,  $l_{\text{proj}} \sim \sum \xi_{\text{perc}} \ln(N)$ , the number of different values that each index  $\alpha$  can assume is of order

$$\exp\{\mathcal{O}[\xi_{\text{perc}}^d \ln(N)^d](\beta/\beta_0)\}. \quad (27)$$

The final matrix product form turns a quantum statistical mechanics problem into a classical statistical mechanics problem, as tracing over the  $\rho_i$  gives a probability distribution for the  $\alpha_i$ . However, the resulting classical problem may suffer from a sign problem.

## DISCUSSION

We have shown how to construct an approximate local projective form of short range, gapped Hamiltonians, and used this to show that the ground states are close to matrix product states. The main goal of the present paper is at a formal level; the techniques developed in this paper provide a way to, in principle at least, extend calculations on small systems to wave functions on much larger system sizes. It is worth comparing to Ref. 25, which showed how to write a very general class of Hamiltonians as a sum of projection

operators. The projection operators there were two-by-two matrices which makes it much easier to find the ground state. Here, the projection operators are large matrices, and the task of constructing the ground state from these operators is tricky. However, the important advance here is that the projection operators are local. This strongly constrains the ground state and leads to the matrix product for the ground state.

In addition to the formal interest, this work may find practical use in numerical simulation, as, at least in simple cases, it is possible to directly calculate the local projective form. Further, we have provided (19), a variational lower bound on energy.

We finally return to the impurity problem raised in the introduction. The following procedure will generate a good wave function for the whole system. First, numerically compute the  $M_i$  for some region around the impurity. Next, compute the appropriate  $M_i$  for the system outside that region; this can be done analytically since that part of the system is noninteracting. Then, determine the appropriate eigenvalues for the  $M_i$  and get a basis of states  $\alpha_{\text{impurity}} \otimes \alpha_{\text{insulator}}$ . Finally, use the  $M_i$  that connect the two regions to determine a wave function in this basis of states. This procedure can be followed even if there are many impurities embedded in the system and the resulting wave function can be used as a starting point for further improvement.

## ACKNOWLEDGMENT

This work was supported by DOE Contract No. W-7405-ENG-36.

## APPENDIX A: LOCALITY

The locality of the  $\tilde{H}_i^0$  relies on the finite group velocity result.<sup>8,15,26</sup> Given the finite-range conditions on the Hamiltonian above, one can bound the commutator  $\|[A(t), B(0)]\|$ , where  $A(t) = \exp(i\mathcal{H}t)A \exp(-i\mathcal{H}t)$ , and show that this commutator is exponentially small for times  $t$  less than  $c_1 l$  where  $l$  is the distance between  $A$  and  $B$  and  $c_1$  is some characteristic inverse velocity. The bound is that  $\|[A(t), B(0)]\| \leq \|A\| \|B\| \sum_j g[t, d(A, j)]$ , where the sum ranges over sites  $j$  which appear in operator  $B$  and where the function  $g$  has the property that for  $|t| \leq c_1 l$ ,  $g(c_1 l, l)$  is exponentially decaying in  $l$  for large  $l$  with decay length  $\xi_C$ . Also, for  $t < t'$ ,  $g(t, l) \leq (t/t')g(t', l)$ .

Consider  $[\tilde{H}_i^0, O_j]$  for some  $O_j$  which acts only on site  $j$ . This equals  $(\Delta E/\sqrt{2\pi q}) \int_{-\infty}^{\infty} dt [H_i(t), O_j] \exp[-(t\Delta E)^2/(2q)]$ . Applying a triangle inequality to the integral we have

$$\begin{aligned} \|[\tilde{H}_i^0, O_j]\| &\leq (\Delta E/\sqrt{2\pi q}) \int_{-\infty}^{\infty} dt \| [H_i(t), O_j] \| \\ &\quad \times \exp[-(t\Delta E)^2/(2q)]. \end{aligned}$$

Let  $l = d(H_i, j)$  be the distance between  $H_i$  and site  $j$ . We split the integral over times  $t$  into a sum of one integral over  $|t| \leq c_1 l$  and one integral over  $|t| \geq c_1 l$ . For  $|t| \leq c_1 l$ , we use the

finite group velocity bound to bound  $\| [H_i(t), O_j] \|$  while for  $|t| \geq c_1 l$  we use  $\| [H_i(t), O_j] \| \leq 2J \| O_j \|$ . Thus

$$\begin{aligned} \| [\tilde{H}_i^0, O_j] \| &\leq J \| O_j \| \left( \frac{\Delta E}{\sqrt{2\pi q}} \int_{|t| \leq c_1 l} g(c_1 l, l) \exp[-(t\Delta E)^2/(2q)] \right. \\ &\quad \left. + 2 \frac{\Delta E}{\sqrt{2\pi q}} \int_{|t| \geq c_1 l} \exp[-(t\Delta E)^2/(2q)] \right) \\ &\leq J \| O_j \| \{ g(c_1 l, l) + 2 \exp[-(c_1 l \Delta E)^2/(2q)] \}, \quad (\text{A1}) \end{aligned}$$

giving the claimed bound on the commutator (9).

We now bound the difference  $\| \tilde{H}_i^0 - M_i \|$ . We have

$$\begin{aligned} \| \tilde{H}_i^0 - M_i \| &\leq (\Delta E / \sqrt{2\pi q}) \\ &\quad \times \int dt \exp[-(t\Delta E)^2/(2q)] \| H_i(t) - H_i^{\text{trunc}}(t) \|. \quad (\text{A2}) \end{aligned}$$

The difference between  $H_i(t)$  and  $H_i^{\text{trunc}}(t)$  is due to the different Hamiltonians used to define the time evolution.

We now bound the difference between  $H_i(t)$  and  $H_i^{\text{trunc}}(t)$ . This result will also be needed in the nonzero temperature calculation. We can replace Eq. (12) by  $H_{\text{loc}} = \sum_{j, d(i,j) \leq l_{\text{proj}} - R} H_j + \sum_{j, d(i,j) > l_{\text{proj}} + R} H_j$ ; by adding the terms  $H_j$  with  $d(i, j) > l_{\text{proj}} + R$  we do not change  $H_i^{\text{trunc}}(t)$ . Then,

$$\| H_i(t) - H_i^{\text{trunc}}(t) \| \leq \sum_{j, l_{\text{proj}} - R < d(i,j) \leq l_{\text{proj}} + R} \int_0^t dt' \| [H_j, H_i(t')] \|.$$

Using the finite group velocity bound we have

$$\begin{aligned} \| H_i(t) - H_i^{\text{trunc}}(t) \| &\leq \sum_{j, l_{\text{proj}} - R < d(i,j) \leq l_{\text{proj}} + R} J^2 S \\ &\quad \times \int_0^t dt' g(t', l_{\text{proj}} - 2R), \end{aligned}$$

where  $l_{\text{proj}} - 2R$  is the minimum distance between the operators  $H_j$  and  $H_i$  and  $S$  is defined to be the number of sites acted on by the operator  $H_j$ . However, the integral  $J S \int_0^t dt' g(t', l_{\text{proj}} - 2R)$  is bounded by  $g(t, l_{\text{proj}})$ .<sup>8</sup> Thus,

$$\begin{aligned} \| H_i(t) - H_i^{\text{trunc}}(t) \| &\leq \sum_{j, l_{\text{proj}} - R < d(i,j) \leq l_{\text{proj}} + R} J g(t, l_{\text{proj}}) \\ &\equiv N(l_{\text{proj}}) J g(t, l_{\text{proj}}), \quad (\text{A3}) \end{aligned}$$

where  $N(l_{\text{proj}})$  is defined to be the number the number of sites  $j$  with  $l_{\text{proj}} - R < d(i, j) \leq l_{\text{proj}} + R$ .

Equation (A3) can be expressed in a more general form. For any operator  $O_i$ , we can define  $O_i^{\text{trunc}}(t)$  by  $O_i^{\text{trunc}} = \exp(iH_{\text{loc}} t) O_i \exp(-iH_{\text{loc}} t)$ , with  $H_{\text{loc}} = \sum_{j, d(i,j) \leq l_{\text{trunc}} - R} H_j$ . Then,  $O_i^{\text{trunc}}(t)$  only involves sites within distance  $l_{\text{trunc}}$  of  $i$  (assuming that  $O_i$  originally involved only sites within that distance). Then,

$$\begin{aligned} \| O_i(t) - O_i^{\text{trunc}}(t) \| &\leq \sum_{j, l_{\text{trunc}} - R < d(i,j) \leq l_{\text{trunc}} + R} \| O \| g(t, l_{\text{trunc}}) \\ &\equiv N(l_{\text{trunc}}) \| O \| g(t, l_{\text{trunc}}), \quad (\text{A4}) \end{aligned}$$

where  $N(l_{\text{trunc}})$  is defined to be the number the number of sites  $j$  with  $l_{\text{trunc}} - R < d(i, j) \leq l_{\text{trunc}} + R$ .

In Eq. (A2), for  $|t| < c_1 l_{\text{proj}}$  we can use the bound on  $g$  to bound the difference  $\| H_i(t) - H_i^{\text{trunc}}(t) \|$ , while for  $|t| > c_1 l_{\text{proj}}$  we use  $\| H_i(t) - H_i^{\text{trunc}}(t) \| \leq 2J$ . Thus,

$$\begin{aligned} \| \tilde{H}_i^0 - M_i \| &\leq J \{ N(l_{\text{proj}}) g(c_1 l_{\text{proj}}, l_{\text{proj}}) \\ &\quad + 2 \exp[-(c_1 l_{\text{proj}} \Delta E)^2/(2q)] \}, \quad (\text{A5}) \end{aligned}$$

giving Eq. (13).

## APPENDIX B: COMMUTATOR IN LOW ENERGY SECTOR

Here we consider sufficient conditions for the  $M_i$  to commute in the low energy sector. We consider the case that  $n$  is uniformly bounded above, and  $\Delta E$  is uniformly bounded below, independent of system size. The important result is that for most systems of interest, including all translationally invariant systems, the commutator of the  $M_i$  is exponentially small in the system size; for a few examples certain of the  $M_i$  do not commute, but even in that case the lack of commutation only poses problems in a finite region of the system (in this case, as we will see, there are local zero energy degrees of freedom).

The operators  $M_i$  are local, in that each acts only within a finite range  $l_{\text{proj}}$  of site  $i$ . Consider then the operators  $\tilde{M}_i^0$ , for some  $q$ . This  $q$  is not necessarily the  $q$  of Eq. (2). Using the locality results, we have

$$\| [\tilde{M}_i^0, M_j] \| \leq J^2 \{ N_M g(c_1 l, l) + 2 \exp[-(c_1 l \Delta E)^2/(2q)] \}, \quad (\text{B1})$$

where  $l = d(i, j) - 2l_{\text{proj}}$  is the smallest distance between sites acted on by  $M_i$  and  $M_j$  and  $N_M$  is the total number of sites acted on by  $M_j$ .

Now,  $P_{\text{low}} \tilde{M}_i^0 P_{\text{low}} = P_{\text{low}} M_i P_{\text{low}}$ . Therefore,

$$[P_{\text{low}} \tilde{M}_i^0 P_{\text{low}}, P_{\text{low}} M_j P_{\text{low}}] = [P_{\text{low}} M_i P_{\text{low}}, P_{\text{low}} M_j P_{\text{low}}].$$

The second commutator is the low energy commutator that we wish to evaluate. We can bound the first commutator by a triangle inequality,

$$\begin{aligned} \| [P_{\text{low}} \tilde{M}_i^0 P_{\text{low}}, P_{\text{low}} M_j P_{\text{low}}] \| &\leq \| [\tilde{M}_i^0, M_j] \| + (\| P_{\text{high}} \tilde{M}_i^0 P_{\text{low}} \| \\ &\quad + \| P_{\text{low}} \tilde{M}_i^0 P_{\text{high}} \|) \| M_j \|. \end{aligned}$$

Following Eq. (7),  $\| P_{\text{high}} \tilde{M}_i^0 P_{\text{low}} \| \leq J \exp(-q/2)$  and also  $\| P_{\text{low}} \tilde{M}_i^0 P_{\text{high}} \| \leq J \exp(-q/2)$ . Combining this with Eq. (B1),

$$\begin{aligned} \| [P_{\text{low}} M_i P_{\text{low}}, P_{\text{low}} M_j P_{\text{low}}] \| &\leq J^2 \{ N_M g(c_1 l, l) \\ &\quad + 2 \exp[-(c_1 l \Delta E)^2/(2q)] \\ &\quad + 2 \exp[-q/2] \}. \end{aligned}$$

Pick  $q=c_1 l \Delta E$ , we find that the commutator of

$$\begin{aligned} \|[P_{\text{low}} M_i P_{\text{low}}, P_{\text{low}} M_j P_{\text{low}}]\| &\leq J^2 [N_M g(c_1 l, l) \\ &+ 4 \exp(-c_1 l \Delta E/2)]. \end{aligned} \quad (\text{B2})$$

Thus, the commutator of  $M_i$  with  $M_j$  in the low energy sector is exponentially small in  $d(i, j)$ .

For a translationally invariant,  $d$ -dimensional, system on a lattice, this suffices to show that the commutator of  $M_i$  with  $M_j$  in the low energy sector is exponentially small in the system size. We can introduce coordinates for each site,  $(\vec{x}, y)$ . The  $d$ -dimensional vector  $\vec{x}$  labels the particular unit cell to which the site belongs, while the coordinate  $y$  indicates the particular site within that cell. The Hamiltonian is invariant under translation so we can pick a basis in which the  $n$  ground states are eigenvectors of the translation operators. In this basis, the matrix elements of the  $M$  are related by  $(M_{\vec{x}, y})_{ab} = (M_{\vec{x}+\vec{x}', y})_{ab} \exp[i\vec{x}' \cdot (\vec{k}_a - \vec{k}_b)]$ , where  $\vec{k}_a$  is the momentum of ground state  $a$ , for  $0 \leq a \leq n-1$ .

In the simplest case (a case which applies to every system of which we are aware), the  $\vec{k}_a$  are vectors of rational multiples  $p/q$  of  $2\pi$ , such that  $p, q$  are integers with some denominator  $q$  which is independent of system size for large enough systems. That is,  $\vec{k}_a = 2\pi(p_a^1/q_a^1, \dots, p_a^d/q_a^d)$ . Then, it is possible to identify a supercell, such that all of the  $n$  ground states are unchanged by translation by a supercell. The size of this supercell in a given direction  $m$  is equal to the least common multiple of the  $n$  different  $q_a^m$ . For example, in the Majumdar-Ghosh chain, the two lowest states have momentum 0 and  $\pi$ , so that in one case  $p=0, q=1$ , and in the other  $p=1, q=2$ . Then, the ground states are unchanged under translation by two sites. For a system with momenta  $0, \pi/3, 2\pi/3$  for the ground states, the ground states would be unchanged under translation by three sites. Then, consider any commutator  $[M_{\vec{x}_1, y_1}, M_{\vec{x}_2, y_2}]$ . This equal  $[M_{\vec{x}_1, y_1}, M_{\vec{x}_2+\vec{x}', y_2}]$  where  $\vec{x}'$  translates  $\vec{x}_2$  by some number of supercells. By choosing  $\vec{x}'$  correctly, we can make the distance between  $(\vec{x}_1, y_1)$  and  $(\vec{x}_2+\vec{x}', y_2)$  of order the linear size of the system. Then, we can use the result that the commutator vanishes exponentially in the spacing between the operators to show that the original commutator  $[M_{\vec{x}_1, y_1}, M_{\vec{x}_2, y_2}]$  is exponentially small in the system size, as desired.<sup>27</sup>

A slightly more complicated, but very artificial, case is that in which the  $p, q$  depend on system size in such a way that the size of the supercell is equal to the linear size of the system. It is not clear that such a thing can actually happen in a system with a finite number  $n$  of low energy states. Even in

this case it is possible to show the commutator of any two  $M_i, M_j$  in the low energy sector is exponentially small in the system size, by using the smallness of the commutator  $[M_i, M_j]$  for large  $d(i, j)$  and expanding the commutator in intermediate states. However, the proof is sufficiently artificial that we do not give it here.

What if a system does not have translational invariance? In this case, the  $M_i$  need not commute in the low energy sector. Consider the following example system, a one-dimensional system of spin-1/2 on each site  $i$ . For  $i \neq 0, 1, 2$ , we have  $H_i = S_i^z$ , while  $H_0 = S_0^x + S_0^y$ ,  $H_1 = -S_0^x - S_1^z$ , and  $H_2 = -S_0^y + S_2^z$ . This is a complicated way of writing the Hamiltonian  $H = \sum_{i \neq 0} S_i^z$ , since all the terms acting on site 0 cancel between  $H_0, H_1$ , and  $H_2$ . This Hamiltonian has a doubly degenerate ground state, with all spins pointing down, except for spin-0 which can point in either direction. For large enough  $q, l_{\text{proj}}$ , the commutator of the  $M_i$  in the low energy sector is close to that of the  $H_i$  in the low energy sector. However, clearly  $H_0$  and  $H_1$  do not commute in the low energy sector, since  $[S_0^y, S_0^x] \neq 0$ . Similarly,  $H_1$  and  $H_2$  do not commute, and  $H_0$  and  $H_2$  do not commute. Still  $H_0$  commutes in the low energy sector with  $H_i$  for  $i \geq 2$ .

In this case, the lack of commutation is localized near site 0. The two ground states differ only locally, on site 0, and only the  $H_i$  for  $i=0, 1, 2$  fail to commute, while the others are diagonal in the low energy subspace.

We now show how to choose a basis of the ground states to bound  $\sum_i o(i)$ . Consider a system with  $n=2$ . Then, find the  $j$  which maximizes  $\delta\lambda_j$ , where  $\delta\lambda_j$  is the difference between the two eigenvalues of  $M_j$  in the low energy sector, and work in a basis which diagonalizes  $M_j$  in the low energy sector. Then,  $o(i) \leq \|[P_{\text{low}} M_i P_{\text{low}}, P_{\text{low}} M_j P_{\text{low}}]\| / (\delta\lambda_j)$ . From Eq. (B2), the commutator is an exponentially decaying function of  $d(i, j)$ , and summing over all  $i$  we find that

$$\sum_i o(i) \leq \mathcal{O}(J^2/c_1 l \Delta E \delta\lambda_j), \quad (\text{B3})$$

independent of system size.

For a system with  $n > 2$ , we can proceed similarly; we first find the  $M_j$  with the maximum difference between its largest and smallest eigenvalues, and use this to bound the off-diagonal matrix elements of other  $M_i$  between the corresponding eigenvectors. We then find an  $M_k$  which has two different eigenvalues, with different eigenvectors from  $M_j$ , and show that off-diagonal matrix elements between those two states are exponentially small in  $d(i, k)$ . We proceed like this until we have bounded all off-diagonal matrix elements.

\*Electronic address: hastings@lanl.gov

<sup>1</sup>K. Wilson, Rev. Mod. Phys. **47**, 773 (1975).

<sup>2</sup>J. W. Bray and S. T. Chui, Phys. Rev. B **19**, 4876 (1979); C. Y. Pan and X. Chen, *ibid.* **36**, 8600 (1987).

<sup>3</sup>S. R. White, Phys. Rev. Lett. **69**, 2863 (1992).

<sup>4</sup>M. Fannes, B. Nachtergaele, and R. F. Werner, Commun. Math. Phys. **144**, 443 (1992).

<sup>5</sup>S. Ostlund and S. Rommer, Phys. Rev. Lett. **75**, 3537 (1997).

<sup>6</sup>J. Dukelsky, M. A. Martin-Delgado, T. Nishino, and G. Sierra, Europhys. Lett. **43**, 457 (1998).

- <sup>7</sup>F. Verstraete and J. I. Cirac, cond-mat/0407066 (unpublished).
- <sup>8</sup>M. B. Hastings, Phys. Rev. B **69**, 104431 (2004).
- <sup>9</sup>M. B. Hastings, Phys. Rev. Lett. **93**, 140402 (2004).
- <sup>10</sup>M. B. Hastings, Phys. Rev. Lett. **93**, 126402 (2004).
- <sup>11</sup>M. B. Hastings, Europhys. Lett. **70**, 824 (2005).
- <sup>12</sup>M. B. Hastings and X.-G. Wen, Phys. Rev. B **72**, 045141 (2005).
- <sup>13</sup>I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987); Commun. Math. Phys. **115**, 477 (1988).
- <sup>14</sup>The case of almost degenerate low energy states, with a splitting which is very small, say exponentially small in the system size, can be handled with only very minor changes to the calculations in this paper.
- <sup>15</sup>E. Lieb and D. Robinson, Commun. Math. Phys. **28**, 251 (1972).
- <sup>16</sup>One can also consider interactions which decay exponentially in space.
- <sup>17</sup>We use the notation  $\langle \Psi_1, \Psi_2 \rangle$ , for the inner product, rather than  $\langle \Psi_1 | \Psi_2 \rangle$ , to avoid confusion with the use of  $|\cdot\rangle$ , to denote a norm.
- <sup>18</sup>F. Verstraete, D. Porras, and J. I. Cirac, Phys. Rev. Lett. **93**, 227205 (2004).
- <sup>19</sup>M. B. Hastings (unpublished).
- <sup>20</sup>T. Kennedy, E. H. Lieb, and H. Tasaki, J. Stat. Phys. **53**, 383 (1988); H. Niggeman, A. Klumper, and J. Zittartz, Z. Phys. B: Condens. Matter **104**, 103 (1997); H. Niggemann, A. Klumper, and J. Zittartz, Eur. Phys. J. B **13**, 15 (2000).
- <sup>21</sup>F. Verstraete, J. J. Garcia-Ripoll, and J. I. Cirac, Phys. Rev. Lett. **93**, 207204 (2004).
- <sup>22</sup>M. Suzuki, Prog. Theor. Phys. **56**, 1454 (1976).
- <sup>23</sup>A. W. Sandvik and J. Kurkijarvi, Phys. Rev. B **43**, 5950 (1991).
- <sup>24</sup>D. A. Klarner, Can. J. Math. **19**, 851 (1967).
- <sup>25</sup>C. Castelnovo, C. Chamon, C. Mudry, and P. Pujol, cond-mat/0502068 (unpublished).
- <sup>26</sup>B. Nachtergaele and R. Sims, math-ph/0506030 (unpublished).
- <sup>27</sup>This argument depends on translation invariance. In general, other forms of self-similarity could be used, as in M. B. Hastings and T. Koma, math-ph/0507008 (in press).