

Scaling behavior of entanglement in two- and three-dimensional free-fermion systems

Weifei Li,¹ Letian Ding,¹ Rong Yu,¹ Tommaso Roscilde,^{1,2} and Stephan Haas¹¹Department of Physics and Astronomy, University of Southern California, Los Angeles, California 90089, USA²Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-strasse 1, 85748 Garching, Germany

(Received 16 February 2006; revised manuscript received 12 June 2006; published 17 August 2006)

Exactly solving a spinless fermionic system in two and three dimensions, we investigate the scaling behavior of the block entropy in critical and noncritical phases. The scaling of the block entropy crucially depends on the nature of the excitation spectrum of the system and on the topology of the Fermi surface. Noticeably, in the critical phases the scaling violates the area law and acquires a logarithmic correction *only* when a well-defined Fermi surface exists in the system. When the area law is violated, we accurately verify a conjecture for the prefactor of the logarithmic correction, proposed by D. Gioev and I. Klich [Phys. Rev. Lett. **96**, 100503 (2006)].

DOI: 10.1103/PhysRevB.74.073103

PACS number(s): 05.30.-d, 03.65.Ud, 73.43.Nq

The nature of many-body entanglement in various solid-state models has been the focus of recent interest. The motivation for this effort is twofold. On the one side, these systems are of interest for the purpose of quantum information processing and quantum computation.¹ At a fundamental level, the study of entanglement represents a *purely quantum* way of understanding and characterizing quantum phases and quantum phase transitions in many-body physics.²⁻⁵

A striking feature of entangled states $|\Psi\rangle$ is that a *local* accurate description of such states is impossible; namely, each subsystem A of the total system U can have a *finite* entropy, quantified as the von Neumann entropy $S_A = -\text{Tr} \rho_A \log_2 \rho_A$ of its reduced density matrix $\rho_A = \text{Tr}_{U \setminus A} |\Psi\rangle\langle\Psi|$, whereas the total system clearly has zero entropy. The *entropy of entanglement* S_A of the subsystem is a reliable estimate of the entanglement between the subsystem A and the rest, $U \setminus A$. Assuming that the system U corresponds to the whole universe in d dimensions, a fundamental question concerns the scaling behavior of the entropy of entanglement S_L of a hypercubic subsystem L^d (hereafter denoted as a *block*) with its size L . Indeed, such scaling probes directly the spatial range of entanglement: when the block size exceeds the characteristic length over which two sites are entangled, the block entropy should become subadditive and scale at most as the area of the block boundaries, following a so-called *area law* $S_L \sim L^{d-1}$. A crucial question is then if and how the scaling of the block entropy changes when the nature of the quantum many-body state evolves in a critical way by passing through a quantum phase transition and how the characteristic spatial extent of entanglement relates to the correlation length of the system.

This question has been extensively addressed in the case of one-dimensional spin systems,⁵⁻⁸ in chains of harmonic oscillators,^{9,10} and in related conformal field theories (CFT's).^{11,12} Here it is found unambiguously that in states with exponentially decaying (connected) correlators S_L follows the area law $S_L \sim L^0$, saturating to a finite value, whereas for critical states, displaying power-law decaying correlations, a *logarithmic correction* to the area is always present: $S_L = [(c + \bar{c})/6] \log_2 L$, where c is the central charge of the related CFT. The asymptotic value of the block entropy is found to diverge logarithmically with the correlation length, $S_\infty \sim \log_2(\xi)$, which clearly establishes the relation-

ship between entanglement and correlations. The above picture holds true only in presence of *short-range* interactions; on the contrary, in presence of long-range interactions the divergence of the correlation length can be still accompanied by the area law.^{10,13}

In higher spatial dimensions fewer results are available and the general relationship between the block entropy scaling and the correlation properties of the quantum state is still unclear even for short-range interactions. In free-boson systems, it has been generally proven that the area law is satisfied for noncritical systems.^{10,14} For free-fermion systems, on the other hand, it has been proven^{15,16} that critical systems with short-range hoppings and a finite Fermi surface exhibit a logarithmic correction to the area law

$$S_L = C/3(\log_2 L)L^{d-1}. \quad (1)$$

In this paper, we investigate a general quadratic fermionic Hamiltonian both in $d=2$ and $d=3$. Upon tuning the Hamiltonian parameters, this model has distinct critical phases *with* and *without* a finite Fermi surface, as well as noncritical phases. The scaling behavior of the block entropy is accurately obtained through exact diagonalization. In noncritical states we find that the area law indeed holds, and we confirm that logarithmic corrections to such law are present in critical states with a finite Fermi surface, as found in Refs. 15 and 16. The prefactor C of the L dependence of S_L in Eq. (1) is found to be very accurately predicted by a formula based on the Widom conjecture.¹⁶ On the other hand, for critical states with a Fermi surface of zero measure, we find that the corrections to the area law are either *absent* or *sublogarithmic*. This means that the relationship between entanglement and correlations in higher-dimensional systems is different than in $d=1$ and that a crucial role is played by the geometry of the Fermi surface or, alternatively, by the density of states at the ground-state energy.

We consider a bilinear spinless fermionic system on a d -dimensional hypercubic lattice with hopping and pairing between nearest-neighbor lattice sites

$$H = \sum_{\langle ij \rangle} [c_i^\dagger c_j - \gamma(c_i^\dagger c_j^\dagger + c_j c_i)] - \sum_i 2\lambda c_i^\dagger c_i. \quad (2)$$

λ is the chemical potential, while γ is the pairing potential. The sum of $\sum_{\langle ij \rangle}$ extends over nearest-neighbor pairs. The




FIG. 1. (Color online) Phase diagram of the model, Eq. (2), for the case $d=2$. The roman numbers for the various phases are explained in the text. Representative contour plots of the dispersion relation Λ_k are also shown. There the black areas corresponds to $\Lambda_k=0$ and the white areas to the top of the band.

above Hamiltonian is a $d > 1$ generalization of the one-dimensional (1D) spinless fermionic Hamiltonian which is obtained by Jordan-Wigner transformation of the XY model in a transverse field.¹⁷ Although in $d > 1$ the exact relationship to the XY model is lost, we can imagine the above Hamiltonian to represent the effective fermionic degrees of freedom of an interacting system with quantum-critical phases.

A more insightful expression for the Hamiltonian of Eq. (2) is obtained upon Fourier transformation to momentum space:

$$H = \sum_k [-2t_k c_k^\dagger c_k + i\Delta_k (c_k^\dagger c_{-k}^\dagger + c_{-k} c_k)],$$

$$t_k = \lambda - \sum_{\alpha=1}^d \cos k_\alpha, \quad \Delta_k = \gamma \sum_{\alpha=1}^d \sin k_\alpha. \quad (3)$$

The pairing potential in k space, Δ_k , clearly reveals a p -wave symmetry.

This Hamiltonian can be diagonalized exactly by a Bogoliubov transformation to give

$$H = \sum_k \Lambda_k f_k^\dagger f_k, \quad \Lambda_k = 2\sqrt{t_k^2 + \Delta_k^2}. \quad (4)$$

Depending on the parameters γ and λ , this system has a rich phase diagram, including metallic, insulating, and (p -wave) superconducting regimes, as shown in Fig. 1. The different phases are certainly distinguished by the different decay of the correlation function, which tells apart the critical from the noncritical phases. Nonetheless, a classification which turns out to be relevant for the study of entanglement is based on the features of the gapless excitation manifold $\Lambda_k=0$. Such a manifold can be characterized by the *density of states* at the ground-state energy $g(0)$ and by the so-called

codimension^{18,19} \bar{d} , defined as the dimension of k space minus the dimension of the $\Lambda_k=0$ manifold. We notice that the existence of a finite Fermi surface at zero energy implies that $\bar{d}=1$ and $g(0) > 0$, while in the absence of a finite Fermi surface we have $\bar{d} \geq 2$ and $g(0) > 0$ or $g(0)=0$ depending on the dispersion relation Λ_k around its nodes.

According to this classification, we can distinguish three phases.

Phase I: $\{0 \leq \lambda \leq d, \gamma=0\}$ and $\{\lambda=0, \gamma>0\}$ if $d=2$. For $\gamma=0$, Eq. (2) reduces to a simple tight-binding model, which is in a metallic state with a 2D-fold symmetric Fermi surface as far as $\lambda \leq d$. In $d=2$, for $\lambda=0$ the system is still a metal with a well-defined Fermi surface, which is simply $k_x = k_y = \pm \pi$, and whose symmetry is lowered by the presence of the γ term in the Hamiltonian. In this phase, $g(0) > 0$ and $\bar{d}=1$ everywhere except at the point $\{\lambda=d, \gamma=0\}$ where $\bar{d}=2$.

Phase II: $\{0 < \lambda \leq d, \gamma > 0\}$ and $\{\lambda=0, \gamma > 0\}$ if $d=3$. Away from the boundary lines of this phase, the system is in a p -wave superconducting state, with a finite pairing amplitude $\langle c_k^\dagger c_{-k}^\dagger \rangle \neq 0$. Such a pairing amplitude vanishes at the boundaries of this region. The dispersion relation Λ_k has point nodes in $d=2$ and line nodes in $d=3$. Everywhere in this phase $g(0)=0$ and $\bar{d}=2$.

Phase III: $\{\lambda > d\}$. In this phase the system is in an insulating state with a gap in the excitation spectrum. Here $g(0)=0$ and $\bar{d}=d$.

This shows that, in terms of the spectral properties, the above system has *two distinct critical phases* (I and II), which are both gapless, and a *noncritical phase* (III). Numerical evaluation of the correlation function through integration over the first Brillouin zone (FBZ),

$$\langle c_i^\dagger c_j \rangle = \int_{\text{FBZ}} \frac{d^d k}{(2\pi)^d} \frac{t_k}{2\Lambda_k} e^{ik \cdot (i-j)}, \quad (5)$$

shows an expected power-law decay in the critical phases and an exponential decay in the noncritical one.

We then proceed to the evaluation of the block entropy of entanglement. The ground state of Eq. (2) is known to be a Gaussian state, whose density matrix can be expressed as the exponential of a quadratic fermion operator.^{20,21} To obtain the reduced density matrix of a L^d subsystem, Grassman algebra is needed.²² Using a Bogoliubov transformation, the reduced density matrix ρ_L can then be written as

$$\rho_L = A \exp\left(-\sum_{l=1}^L \varepsilon_l d_l^\dagger d_l\right), \quad (6)$$

where d_l^\dagger, d_l are the new Fermi operators after the transformation and A is a normalization constant to ensure $\text{Tr}(\rho) = 1$. The single-particle eigenvalues ε_l can be obtained from $\langle c_i^\dagger c_j \rangle$ and $\langle c_i^\dagger c_j^\dagger \rangle$ by the following formula:²¹

$$\begin{aligned} & \left(C - F - \frac{I}{2}\right) \left(C + F - \frac{I}{2}\right) \\ &= \frac{1}{4} P \text{diag} \left\{ \tanh^2\left(\frac{\varepsilon_1}{2}\right), \tanh^2\left(\frac{\varepsilon_2}{2}\right), \dots, \tanh^2\left(\frac{\varepsilon_L}{2}\right) \right\} P^{-1}, \end{aligned} \quad (7)$$

where $C_{i,j} = \langle c_i^\dagger c_j \rangle$ and $F_{i,j} = \langle c_i^\dagger c_j^\dagger \rangle$; P is the orthogonal matrix that diagonalizes the left side of the above equation. The Block entropy can then be calculated in terms of ε_l as

$$S_L = \sum_{l=1}^L \left\{ \ln[1 + \exp(-\varepsilon_l)] + \frac{\varepsilon_l}{\exp(\varepsilon_l) + 1} \right\}. \quad (8)$$

In $d=1$ the above formulas reproduce the scaling of the block entropy as observed in the XY model in a transverse field.^{5,6} In $d=2$ the phase diagram is richer, and we need to consider the various phases one by one. We begin with the critical metallic phase (I), $\gamma=0, 0 \leq \lambda < d$. For this case a logarithmic correction to the area law, $S_L = [C(\lambda)/3](\log_2 L)L^{d-1}$, is observed for all values of λ , as shown in Fig. 2. This is in full agreement with the results of Refs. 15 and 16, which predict this behavior in presence of a finite Fermi surface. More specifically, Ref. 16 also supplies us with an explicit prediction for the λ dependence of $C(\lambda)$, based on the Widom's conjecture,²³ in the form

$$C(\lambda) = \frac{1}{4(2\pi)^{d-1}} \int_{\partial\Omega} \int_{\partial\Gamma(\lambda)} |n_x \cdot n_p| dS_x dS_p, \quad (9)$$

where Ω is the volume of the block normalized to 1, $\Gamma(\lambda)$ is the volume enclosed by the Fermi surface, and the integration is carried over the surface of both domains. A numerical fit of the calculated asymptotic behavior of S_L through the formula $S_L = \frac{C}{3} L^{d-1} \log_2(L) + BL^{d-1} + AL^{d-2} + \dots$ provides us with the exact result for the $C(\lambda)$ prefactor. In Fig. 3 the prediction of Ref. 16, Eq. (9), for the case $\{0 \leq \lambda \leq d, \gamma=0\}$ is compared to our numerical results both for $d=2$ and $d=3$. The agreement is clearly striking. Moreover, for




FIG. 2. Scaling of the block entropy S_L in $d=2$ for $\gamma=0$ (left panel) and $\lambda=0$ (right panel). The solid lines correspond to fits according to the formula $S_L = \frac{C}{3} L \log_2(L) + BL + A$.

$\{\lambda=0, \gamma>0\}$ in $d=2$ the formula (9) predicts $C=1$, which is also accurately verified by our data as shown in Fig. 2. This proves that the formula (9) is essentially providing a complete analytic form for the leading behavior of the block-entropy scaling in arbitrary dimensions for systems with a finite Fermi surface.

We then turn to the other two phases II and III. Two scans through these phases, at fixed $\gamma=1$ and at fixed $\lambda=1$, are shown in Fig. 4. We observe that logarithmic corrections are *absent* in both and only sublogarithmic corrections are possible. For $\{\lambda \geq d, \gamma=0\}$ $S_L=0$ identically and the state is not entangled. While the area law is expected to hold in the noncritical phase III, it is surprising to observe it enforced also in the critical phase II, which has a diverging correlation




FIG. 3. λ dependence of the C coefficient in Eq. (1) in $d=2$ and $d=3$. The values extracted from fits to our numerical data are compared with the predictions of Ref. 16. In $d=2$, the exact form of $C(\lambda)$ can be obtained, which is equal to $\frac{2}{\pi} \cos^{-1}(\lambda-1)$.




FIG. 4. (Color online) Scaling of the block entropy S_L in $d=2$ for $\gamma=1$ (left panel) and $\lambda=1$ (right panel).

length. This clearly reveals that the connection between block-entropy scaling and correlation properties is not straightforward in $d>1$.

Our results for the entanglement behavior, codimension, density of states, and correlation properties are summarized in Table I. A crucial difference between the two critical phases I and II is the codimension \bar{d} and the density of states at the ground-state energy. We have $\bar{d}=1$ and $g(0)>0$ in phase I, which shows logarithmic corrections to the area law, whereas $\bar{d}=2$ and $g(0)=0$ in phase II, in which the area law is verified up to sublogarithmic corrections. The special case $\{\lambda=0, \gamma>0\}$ makes this difference stand out as we move from $d=2$ to $d=3$. In this parameter region, for $d=2$ we have $\bar{d}=1$ and $g(0)>0$, while for $d=3$ we have $\bar{d}=2$ and $g(0)=0$. $S(L)\sim L\log(L)$ is observed for the former case,

TABLE I. Summary of the entanglement scaling properties, codimension, density of states, and decay of correlations in the three phases of the model, Eq. (2), in $d=2,3$.

	S_L	\bar{d}	$g(0)$	$\langle c_i^\dagger c_j \rangle$
Phase I	$\sim(\log_2 L)L^{d-1}$	1	>0	Power-law decay
Phase II	$\sim L^{d-1}$	2	0	Power-law decay
Phase III	$\sim L^{d-1}$	d	0	Exp. decay

while $S(L)\sim L^2$ for the latter, which does not show a logarithmic correction.

It is therefore tempting to conjecture that a codimension $\bar{d}=1$ or, alternatively, a finite density of states at the ground-state energy $g(0)>0$ is a *sufficient condition* for critical phases in $d>1$ to show violations of the area law. For the fermionic system under consideration, $\bar{d}=1$ requires the existence of a finite Fermi surface, which is the basic assumption of the proof of area-law violation in Refs. 15 and 16. This conjecture would generalize the results for $d=1$, where the codimension can only take the value $\bar{d}=1$, and only critical phases with $g(0)>0$ have been explored in the literature.

Further investigations in systems with $d>1$ are clearly needed to confirm this picture and to clarify whether more severe violations of the area law are possible in the presence of infinitely degenerate ground states or in systems with a fractal Fermi surface.¹⁵

Note added. During the completion of this manuscript we became aware of Ref. 24 whose results are in full agreement with the ones reported in our work.

We thank A. Cassidy, P. Sengupta, I. Grigorenko, and M. Wolf for useful discussions. T.R. acknowledges support of the European Union through the SCALA project. This work was supported by the Petroleum Research Foundation, Grant ACS PRF No. 41757 and the Department of Energy, Grant No. DE-FG02-05ER46240.

¹G. Burkard, in *Handbook of Theoretical and Computational Nanotechnology*, edited by M. Rieth and W. Schommers (American Scientific, 2006); cond-mat/0409626 (unpublished), and references therein.

²A. Osterloh, L. Amico, G. Falci, and R. Fazio, *Nature (London)* **416**, 608 (2002).

³T. J. Osborne and M. A. Nielsen, *Phys. Rev. A* **66**, 032110 (2002).

⁴F. Verstraete, M. Popp, and J. I. Cirac, *Phys. Rev. Lett.* **92**, 027901 (2004).

⁵G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, *Phys. Rev. Lett.* **90**, 227902 (2003).

⁶J. I. Latorre, E. Rico, and G. Vidal, *Quantum Inf. Comput.* **4**, 48 (2004).

⁷G. Refael and J. E. Moore, *Phys. Rev. Lett.* **93**, 260602 (2004).

⁸N. Laflorencie, *Phys. Rev. B* **72**, 140408(R) (2005).

⁹S. O. Skrøvseth, *Phys. Rev. A* **72**, 062305 (2005).

¹⁰M. Cramer, J. Eisert, M. B. Plenio, and J. Dreissig, *Phys. Rev. A* **73**, 012309 (2006).

¹¹P. Calabrese and J. Cardy, *J. Stat. Mech.: Theory Exp.* 2004, P06002.

¹²V. E. Korepin, *Phys. Rev. Lett.* **92**, 096402 (2004).

¹³W. Dür, L. Hartmann, M. Hein, M. Lewenstein, and H.-J. Briegel, *Phys. Rev. Lett.* **94**, 097203 (2005).

¹⁴M. B. Plenio, J. Eisert, J. Dreissig, and M. Cramer, *Phys. Rev. Lett.* **94**, 060503 (2005).

¹⁵M. M. Wolf, *Phys. Rev. Lett.* **96**, 010404 (2006).

¹⁶D. Gioev and I. Klich, *Phys. Rev. Lett.* **96**, 100503 (2006).

¹⁷E. Lieb, T. Schultz, and D. Mattis, *Ann. Phys. (N.Y.)* **16**, 407 (1961).

¹⁸G. E. Volovik, cond-mat/0505089 (unpublished).

¹⁹G. E. Volovik, *The Universe in a Helium Droplet* (Clarendon Press, Oxford (2003).

²⁰M. Gaudin, *Nucl. Phys.* **15**, 89 (1960).

²¹I. Peschel, *J. Phys. A* **36**, L205 (2003).

²²M. C. Chung and I. Peschel, *Phys. Rev. B* **64**, 064412 (2001).

²³H. Widom, *Toeplitz centennial (Tel Aviv, 1981), Operator Theory: Advances Applications Vol. 4* (Birkhuser, Basel-Boston, MA, 1982).

²⁴T. Barthel, M. C. Chung, and U. Schollwoeck, cond-mat/0602077 (unpublished).