

Nonequilibrium magnetotransport through a quantum dot: An interpolative perturbative approach

A. A. Aligia

Comisión Nacional de Energía Atómica, Centro Atómico Bariloche and Instituto Balseiro, 8400 Bariloche, Argentina

(Received 12 June 2006; revised manuscript received 28 July 2006; published 27 October 2006)

We study the differential conductance, spectral density, and magnetization for a quantum dot coupled to two conducting leads as a function of bias voltage V_{ds} , magnetic field B , and temperature T . The system is modeled with the Anderson model solved using a spin-dependent interpolative perturbative approximation in the Coulomb repulsion U that conserves the current. For large enough magnetic field, the differential conductance as a function of bias voltage shows split peaks. This splitting is larger than the corresponding splitting in the spectral density of states $\rho_d(\omega)$, in agreement with experiment.

DOI: [10.1103/PhysRevB.74.155125](https://doi.org/10.1103/PhysRevB.74.155125)

PACS number(s): 72.15.Qm, 73.63.Kv, 85.75.-d

I. INTRODUCTION

Electron transport through nanoscale quantum dots (QDs) has been a subject of great interest in the last years. A QD consists of a confined droplet of electrons and it has been predicted that the system behaves as a single-electron transistor described by the Anderson model.^{1,2} Experiments with one QD in the linear response regime^{3–6} displayed clearly the Kondo physics in this almost ideal “one-impurity” system and confirmed the predictions based on the Anderson model.^{1,2,7} The unitary limit of ideal transmittance has been reached.⁶ Calculations using the accurate Wilson’s numerical renormalization group (NRG) have shown a good agreement with these experiments.⁸ The effects of a magnetic field and the possibility to use transport through a QD or a site coupled to a QD as a spin filter have also been addressed theoretically,^{9–12} including NRG results.¹⁰

In contrast to the linear response regime, the experimental situation in which a finite bias voltage between drain and source V_{ds} is applied,^{13–18} i.e., the nonequilibrium case, is much harder to treat theoretically. The most accurate techniques used to study the impurity Anderson and Kondo models at equilibrium¹⁹ cannot be easily extended to the nonequilibrium situation. The NRG has not been implemented out of equilibrium. A formalism to generalize exact Bethe ansatz results for the nonequilibrium case has been proposed recently,²⁰ but its application to the Kondo model requires further elaboration. Some results based on integrability but not exact have been presented.²¹ Also a Kondo model has been solved exactly for particular parameters.²² The non-crossing approximation (NCA), which has been successfully used to calculate thermodynamic properties of generalized Anderson models above or near the Kondo temperature T_K ,^{23–25} has also been used for nonequilibrium situations.^{26–28} Unfortunately, at low temperatures this method fails to fulfill Fermi liquid relations in the equilibrium case,²⁹ and the spectral density presents a spurious peak at the Fermi level in the presence of a magnetic field B .²⁷ A likely cause of this peak is given by Moore and Wen, who study the splitting of the Kondo resonance with B from the Bethe-ansatz solution at equilibrium.³⁰ Slave-boson mean-field approximations have been used successfully in the equilibrium case,^{12,31–36} but they rely on the minimization of the free energy, which is not at a minimum in the nonequilibrium

case. Nevertheless, these approximations have been used for problems out of equilibrium.^{37,38} A relatively simple but accurate approach based on perturbation theory is the local moment approach,^{39–41} but its extension to the nonequilibrium case is not trivial because it uses Fermi liquid relations valid at equilibrium. For the Kondo model, a perturbative approach in the exchange constant has been used, which is valid when either V_{ds} or B are much larger than T_K .⁴²

Another approximation for the Anderson model is perturbation theory in $U/(\pi\Delta)$, where U is the Coulomb repulsion and Δ is the resonant level width.^{43–46} It can be extended naturally to the nonequilibrium case using the Keldysh formalism.^{47–50} In the equilibrium case, the second order approximation has been used for several nanoscopic systems.^{12,35,51–54} A shortcoming of this method is that it cannot describe the exponential decrease of T_K in the Kondo regime as U is increased. One way to avoid this problem is to use renormalized perturbation theory (RPT).^{55,56} Another method to extend the validity of the approximation to larger values of U is to use an interpolative perturbative approach (IPA),^{57–60} which corrects the second-order result in order to reproduce exactly the atomic limit $U/\Delta \rightarrow +\infty$. The IPA has been shown to describe well the conductance through a QD for $U \leq 8\Delta$.⁵² The results agree with those obtained more recently using the finite temperature density matrix renormalization group method.⁶¹ In addition, comparison of the spin dependent IPA^{12,53} with exact diagonalization in finite systems shows very good agreement for $U=6.25\Delta$.⁵³ Note that in some experimental situations $\Delta \sim 0.15$ meV, $U \sim 0.6$ meV ($U/\Delta=4$),^{3,5} while in recent nonequilibrium experiments $2\Delta \sim 0.33$ meV, $U \sim 1.2$ meV ($U/\Delta=7.3$).¹⁶ The extension of perturbation theory in U^2 to the nonequilibrium case has been considered by Hershfield, Davies, and Wilkins.⁶² They found that for finite bias voltage V_{ds} , the current is conserved only in the symmetric Anderson model, in which the on-site energy at the dot $E_d = -U/2 + (\mu_L + \mu_R)/2$, where μ_L (μ_R) is the chemical potential at the left (right) lead. Recent use of perturbation theory in $U/(\pi\Delta)$ for $V_{ds} \neq 0$ was limited to the symmetric case, for calculations of the current noise,⁶³ and other properties including terms of third and fourth order in U ,^{64–67} and in the particular in presence of magnetic field B ,^{65,66} motivated by recent experiments.^{14–16} However, even in the symmetric case, the *spin*

current is not conserved by the approximation when $V_{ds} \neq 0$. In the Kondo regime, an exact expression for the differential conductance has been derived using RPT,⁶⁶ but it is limited to V_{ds} small in comparison with T_K .

There is some controversy concerning the effect of V_{ds} on the Kondo peak in the spectral density for $B=0$. Calculations based on the NCA^{26–28} and on the equation of motion method^{68,69} predict a splitting of the peak. However, this method has limitations in the Kondo regime, particularly in the particle-hole symmetric case, for which the Green's function is independent of temperature.⁷⁰ Instead, second-order perturbation theory in U predicts a fading of the peak with increasing V_{ds} without splitting. It has been suggested that terms of order U^3 and U^4 might cause the peak to split.^{64,65} However, a recent study with an improvement of the higher order corrections, which reproduces correctly the equilibrium expressions, indicates that the spectral density retains the qualitative features of the second-order result.⁶⁷ Physically, one might expect that if the dot is hybridized with left and right leads at chemical potentials μ_L and μ_R with matrix elements V_L and V_R , respectively, the effective distribution function fluctuates between that corresponding to μ_L and μ_R (at least if $V_L, V_R \gtrsim |\mu_L - \mu_R|$) and no split Kondo peak at the Fermi level is expected.⁶⁷ However, as stated above, different approximations lead to different results. The experimental situation is also controversial: a fading of the central peak in the differential conductance with increasing bias voltage was reported for a carbon nanotube quantum dot,¹⁷ while a splitting of the Kondo resonance in the spectral density was observed in a three-terminal quantum ring.¹⁸ Since the diameter of the ring is of the order of 500 nm, it might be possible that to model the system as one effective site connected to conducting leads is not a good approximation and the space dependence of the energy distribution function plays an essential role.⁷¹

In this work, we generalize the spin dependent IPA, based on perturbation theory in U up to second order, to the non-equilibrium case. The problem of the conservation of the current for each spin is solved by a simple trick. The partition of the Hamiltonian $H=H_0+H'$ into an unperturbed part H_0 and a perturbation H' is in principle arbitrary. In the past, for example, H_0 has been chosen so that the Friedel-Langreth sum rule⁷² is satisfied at zero temperature.^{52,60} Here we also use the spin dependent version of this rule^{12,55} to calculate the spectral density at equilibrium. For finite bias voltage V_{ds} , we determine H_0 asking that the current is conserved for each spin projection. The formalism is applied to calculate the differential conductance dI/dV_{ds} , spectral density, and magnetization as a function of bias voltage and magnetic field B for various temperatures. Our results are compared with experiment and previous calculations. Recent measurements^{14–16} of dI/dV_{ds} report results which seem to disagree with accurate calculations^{73,74} of the spectral density at equilibrium for $B \neq 0$. However, Hewson *et al.* pointed out that these experiments should not be interpreted in terms of the equilibrium Green's functions.⁶⁶ In spite of shortcomings of the self-consistent procedure at small temperatures, when both V_{ds} and B are small but different from zero, important conclusions can be drawn from our results.

The paper is organized as follows. In Sec. II the model is described, the nonequilibrium perturbation formalism is

briefly reviewed, useful expressions are derived, and the generalization of the IPA to the nonequilibrium situation is presented. Section III contains the results of the application of the method to magnetotransport, spectral density, and magnetization. Section IV contains a summary and a brief discussion. Some details of the calculations were moved to the Appendix.

II. MODEL AND APPROXIMATIONS

A. Hamiltonian

We consider a QD interacting with two conducting leads, one at the left and one at the right, with chemical potentials μ_L and μ_R , respectively, with $\mu_L - \mu_R = eV_{ds}$. As usual, the QD is modeled by one effective site with one relevant orbital with an important on-site Coulomb repulsion U and an on-site energy E_d controlled by the gate voltage V_g . The Hamiltonian is that of an Anderson model, which is split into a noninteracting part H_0 and a perturbation H' as

$$H = H_0 + H',$$

$$H_0 = \sum_{k\alpha\sigma} \varepsilon_{k\alpha} c_{k\alpha\sigma}^\dagger c_{k\alpha\sigma} + \sum_{\sigma} \varepsilon_{eff}^{\sigma} n_{d\sigma} + \sum_{k\alpha\sigma} (V_{k\alpha} c_{k\alpha\sigma}^\dagger d_{\sigma} + \text{H.c.}),$$

$$H' = (E_d - \varepsilon_{eff}^{\sigma}) \sum_{\sigma} n_{d\sigma} - B(n_{d\uparrow} - n_{d\downarrow}) + U n_{d\uparrow} n_{d\downarrow}. \quad (1)$$

Here $\alpha=L,R$ refers to the left and right leads. The operator $c_{k\alpha\sigma}^\dagger$ creates an electron in the state with wave vector k and spin σ at the lead α . Similarly d_{σ}^\dagger creates an electron with spin σ at the QD. The number operator $n_{d\sigma} = d_{\sigma}^\dagger d_{\sigma}$, and B is the effect of an applied magnetic field on the on-site energy at the QD for each spin. The first term of H_0 describes the leads and the third its hybridization with the QD. H' contains the dot on-site energy, Zeeman splitting, and Coulomb repulsion, respectively. The two constants $\varepsilon_{eff}^{\sigma}$ are arbitrary since they cancel in H_0+H' . However, in our perturbation treatment, the results depend on them and they are determined self-consistently as described in Sec. II E.

B. Nonequilibrium perturbation theory

We use the notation of Lifshitz and Pitaevskii.⁴⁸ There are four different Green's functions. For the electrons at the dot, they can be chosen as

$$G_{d\sigma}^r(t, t') = -i\theta(t-t') \langle d_{\sigma}(t) d_{\sigma}^{\dagger}(t') + d_{\sigma}^{\dagger}(t') d_{\sigma}(t) \rangle,$$

$$G_{d\sigma}^a(t, t') = i\theta(t'-t) \langle d_{\sigma}(t) d_{\sigma}^{\dagger}(t') + d_{\sigma}^{\dagger}(t') d_{\sigma}(t) \rangle,$$

$$G_{d\sigma}^{<}(t, t') = i \langle d_{\sigma}^{\dagger}(t') d_{\sigma}(t) \rangle,$$

$$G_{d\sigma}^{>}(t, t') = -i \langle d_{\sigma}(t) d_{\sigma}^{\dagger}(t') \rangle. \quad (2)$$

The first two are the retarded and advanced Green's functions, already present in the equilibrium case. Similar Green's functions can be constructed involving conduction electrons. Since obviously $G^a - G^r = G^{<} - G^{>}$, only three

Green's functions are independent, and in the following we use this relation to eliminate $G^>$. Also, three independent self-energy functions Σ^r , Σ^a , and $\Sigma^<$ can be defined, which allow us to write a matrix Dyson's equation⁴⁸

$$\mathbf{G} = \mathbf{g} + \mathbf{g}\Sigma\mathbf{G}, \quad (3)$$

where

$$\mathbf{G} = \begin{pmatrix} 0 & G^a \\ G^r & 2G^< + G^r - G^a \end{pmatrix}, \quad (4)$$

\mathbf{g} has a similar expression in terms of the corresponding Green's functions for the unperturbed Hamiltonian H_0 , and

$$\Sigma = \begin{pmatrix} \Sigma^r - \Sigma^a - 2\Sigma^< & \Sigma^r \\ \Sigma^a & 0 \end{pmatrix}. \quad (5)$$

In general, the products in Eq. (3) have to be understood as convolutions in time and space. However, in this work, in which we consider the stationary case and perturbation theory in U up to second order,⁶⁷ these products become just ordinary products involving the Fourier transforms in $t-t'$ of the Green's functions for the d electrons $G_{d\sigma}^{\nu}(\omega)$ and the corresponding self-energies $\Sigma_{\sigma}^{\nu}(\omega)$. In addition, $G_{d\sigma}^a(\omega) = [G_{d\sigma}^r(\omega)]^*$ and $\Sigma_{\sigma}^a(\omega) = [\Sigma_{\sigma}^r(\omega)]^*$, where the asterisk denotes complex conjugation. Therefore, our task reduces to find suitable approximations for $\Sigma_{\sigma}^r(\omega)$ and $\Sigma_{\sigma}^<(\omega)$.

After doing the matrix product in Eq. (3), from the entries (1,2) and (2,1) one obtains

$$[G_{d\sigma}^r(\omega)]^{-1} = [g_{d\sigma}^r(\omega)]^{-1} - \Sigma_{\sigma}^r(\omega), \quad (6)$$

and its complex conjugate. After some algebra and using previous equations, the remaining equation [from the entry (2,2)] can be written in the form

$$G_{d\sigma}^<(\omega) = |G_{d\sigma}^r(\omega)|^2 \left(\frac{g_{d\sigma}^<(\omega)}{|g_{d\sigma}^r(\omega)|^2} - \Sigma_{\sigma}^<(\omega) \right). \quad (7)$$

The noninteracting Green's functions can be obtained easily from the equations of motion⁵⁰ using H_0 :

$$[g_{d\sigma}^r(\omega)]^{-1} = \omega - \varepsilon_{eff}^{\sigma} - \sum_{\alpha} [E_{\alpha}(\omega) - i\Delta_{\alpha}(\omega)],$$

$$g_{d\sigma}^<(\omega) = 2i|g_{d\sigma}^r(\omega)|^2 \sum_{\alpha} \Delta_{\alpha}(\omega) f(\omega - \mu_{\alpha}), \quad (8)$$

where $f(\omega)$ is the Fermi function and E_{α} , Δ_{α} for leads $\alpha = L, R$, are the real and imaginary parts of the sum

$$\sum_k \frac{|V_{k\alpha}|^2}{\omega - \varepsilon_{k\alpha} + i\eta} = E_{\alpha}(\omega) - i\Delta_{\alpha}(\omega), \quad (9)$$

with η a positive infinitesimal.

We approximate the retarded self-energy as

$$\Sigma_{\sigma}^r(\omega) = E_d^{\sigma} - \varepsilon_{eff}^{\sigma} + U\langle n_{d\bar{\sigma}} \rangle + \Sigma_{\sigma int}^{r2}(\omega), \quad (10)$$

where $E_d^{\uparrow} = E_d - B$, $E_d^{\downarrow} = E_d + B$, and $\bar{\sigma}$ means $-\sigma$. The last term is an interpolative expression based on the correction of order U^2 , as described in Sec. II D. The remaining terms would correspond to the first order correction in H' if $\langle n_{d\bar{\sigma}} \rangle$ were

evaluated with H_0 . However, $\langle n_{d\bar{\sigma}} \rangle$ is evaluated self-consistently using

$$\langle n_{d\sigma} \rangle = \frac{-i}{2\pi} \int d\omega G_{d\sigma}^<(\omega), \quad (11)$$

and this is equivalent to a partial summation of an infinite series of diagrams.⁶⁵

The correction for $\Sigma_{\sigma}^<(\omega)$ of first order in H' vanishes. We approximate $\Sigma_{\sigma}^<(\omega)$ by an expression $\Sigma_{\sigma int}^<(\omega)$ based on the second order term $\Sigma_{\sigma}^{<2}(\omega)$, also described in Sec. II D.

The diagrams for the corrections to the self-energy of second order in U are drawn in Ref. 48. In terms of the independent unperturbed Green's functions the expressions are

$$\begin{aligned} \Sigma_{\uparrow}^{r2}(\omega) = U^2 \int \frac{d\omega_1}{2\pi} \int \frac{d\omega_2}{2\pi} & [g_{\uparrow}^r(\omega_1)g_{\downarrow}^r(\omega_2)g_{\downarrow}^<(\omega_1 + \omega_2 - \omega) \\ & + g_{\uparrow}^r(\omega_1)g_{\downarrow}^<(\omega_2)g_{\downarrow}^<(\omega_1 + \omega_2 - \omega) \\ & + g_{\uparrow}^<(\omega_1)g_{\downarrow}^r(\omega_2)g_{\downarrow}^<(\omega_1 + \omega_2 - \omega) \\ & + g_{\uparrow}^<(\omega_1)g_{\downarrow}^<(\omega_2)g_{\uparrow}^a(\omega_1 + \omega_2 - \omega)], \end{aligned} \quad (12)$$

$$\begin{aligned} \Sigma_{\uparrow}^{<2}(\omega) = -U^2 \int \frac{d\omega_1}{2\pi} \int \frac{d\omega_2}{2\pi} \\ \times g_{\uparrow}^<(\omega_1)g_{\downarrow}^<(\omega_2)g_{\downarrow}^>(\omega_1 + \omega_2 - \omega), \end{aligned} \quad (13)$$

and the same interchanging spin up and down, with $g_{\sigma}^>(\omega) = g_{\sigma}^<(\omega) + 2 \text{Im} g_{\sigma}^r(\omega)$. It can be shown that these expressions are equivalent to those given by Hershfield *et al.*⁶² These integrals which have the form of convolutions in frequency can be calculated conveniently using fast Fourier transform to the time representation.^{46,75} For the case of constant $E_{\alpha}(\omega)$ and $\Delta_{\alpha}(\omega)$, it is more convenient to evaluate one of the integrals analytically, as shown in the Appendix.

Although it is not convenient for the numerical evaluation, it is interesting to express the above self-energy corrections in terms of the unperturbed spectral density of states at the dot $\rho_{d\sigma}^0(\omega) = -\text{Im} g_{d\sigma}^r(\omega) / \pi$. In terms of this density

$$g_{d\sigma}^r(\omega) = \int d\varepsilon \frac{\rho_{d\sigma}^0(\varepsilon)}{\omega - \varepsilon + i\eta},$$

$$g_{d\sigma}^<(\omega) = 2i\pi\rho_{d\sigma}^0(\omega)\tilde{f}(\omega), \quad (14)$$

where $\tilde{f}(\omega)$ is a weighted distribution function of the Fermi functions at the two leads:

$$\tilde{f}(\omega) = \frac{\sum_{\alpha} \Delta_{\alpha}(\omega) f(\omega - \mu_{\alpha})}{\sum_{\alpha} \Delta_{\alpha}(\omega)}. \quad (15)$$

Replacing Eqs. (14) in Eqs. (12) and (13) we obtain after some algebra

$$\begin{aligned} \Sigma_{\uparrow}^{r2}(\omega) = U^2 \int d\varepsilon_1 d\varepsilon_2 d\varepsilon_3 \frac{\rho_{d\uparrow}^0(\varepsilon_1)\rho_{d\downarrow}^0(\varepsilon_2)\rho_{d\downarrow}^0(\varepsilon_3)}{\omega + \varepsilon_3 - \varepsilon_1 - \varepsilon_2 + i\eta} \\ \times [1 - \tilde{f}(\varepsilon_2)]\tilde{f}(\varepsilon_3) + \tilde{f}(\varepsilon_1)\tilde{f}(\varepsilon_2)[1 - \tilde{f}(\varepsilon_3)], \end{aligned} \quad (16)$$

$$\begin{aligned} \Sigma_{\uparrow}^{\leq 2}(\omega) = & -2i\pi U^2 \int d\epsilon_1 d\epsilon_2 \rho_{d\uparrow}^0(\epsilon_1) \rho_{d\downarrow}^0(\epsilon_2) \rho_{d\downarrow}^0(\epsilon_1 + \epsilon_2 - \omega) \\ & \times \tilde{f}(\epsilon_1) \tilde{f}(\epsilon_2) [1 - \tilde{f}(\epsilon_1 + \epsilon_2 - \omega)]. \end{aligned} \quad (17)$$

At equilibrium $\mu_L = \mu_R$. Then $\tilde{f}(\omega)$ becomes the usual distribution function and $\Sigma_{\sigma}^{r2}(\omega)$ coincides with the result of ordinary perturbation theory. Equations (16) and (17) are used in Sec. II D to construct the corresponding interpolating expressions $\Sigma_{\sigma\text{int}}^{r2}(\omega)$ and $\Sigma_{\sigma\text{int}}^{\leq}(\omega)$.

C. The current

Following Meir and Wingreen,⁷⁶ the current with spin σ flowing between the left lead and the dot is

$$j_{L\sigma} = \frac{2ie}{h} \int d\omega \Delta_L(\omega) [2if(\omega - \mu_L) \text{Im} G_{d\sigma}^r(\omega) + G_{d\sigma}^{\leq}(\omega)]. \quad (18)$$

Similarly, the current with spin σ flowing between the dot and the right lead is

$$j_{R\sigma} = -\frac{2ie}{h} \int d\omega \Delta_R(\omega) [2if(\omega - \mu_R) \text{Im} G_{d\sigma}^r(\omega) + G_{d\sigma}^{\leq}(\omega)]. \quad (19)$$

Of course, since the current is conserved one should have

$$j_{L\sigma} = j_{R\sigma}, \quad (20)$$

but perturbation theory does not satisfy this equality for generic values of the parameters.⁶² In this work we determine $\varepsilon_{\text{eff}}^{\sigma}$ imposing Eqs. (20).

D. The interpolative perturbative approach (IPA)

Extending previous ideas,⁵⁷⁻⁶⁰ we want to replace the second-order contributions to the self-energies $\Sigma_{\sigma}^{r2}(\omega)$ and $\Sigma_{\sigma}^{\leq 2}(\omega)$, by other ones $\Sigma_{\sigma\text{int}}^{r2}(\omega)$ and $\Sigma_{\sigma\text{int}}^{\leq}(\omega)$, which coincide with the previous ones to order U^2 for small U , but also reproduce the high frequency and atomic limits. For the sake of clarity in the exposition we choose $\sigma = \uparrow$. The final results are also valid interchanging spin up and down.

We propose⁶⁰

$$\Sigma_{\uparrow\text{int}}^{r2}(\omega) = \frac{A_1 \Sigma_{\uparrow}^{r2}(\omega)}{1 - A_2 \Sigma_{\uparrow}^{r2}(\omega)}, \quad (21)$$

where A_1 is determined so that $\Sigma_{\uparrow\text{int}}^{r2}(\omega)$ reproduces the leading behavior at high frequencies, and afterwards A_2 is determined to reproduce the exact result in the atomic limit $\Delta_{\alpha}(\omega)/U \rightarrow 0$.

Up to order $1/\omega$, the exact self-energy is determined by the first and second moments of the spectral density of d states $\rho_{d\sigma}(\omega)$, which can be evaluated independently of $\rho_{d\sigma}(\omega)$ using particular commutators.⁷⁷ Proceeding as in the equilibrium case,⁶⁰ allowing dependence on spin,^{12,53} and using Eqs. (6) and (10), one obtains that the correct leading behavior of the retarded second-order self-energy is

$$\Sigma_{\uparrow\text{int}}^{r2}(\omega) = \langle n_{d\downarrow} \rangle (1 - \langle n_{d\downarrow} \rangle) U^2 \omega^{-1} + O(\omega^{-2}). \quad (22)$$

For $\omega \rightarrow \infty$, the integrals over ϵ_i in Eq. (16) decouple and then

$$\Sigma_{\uparrow}^{r2}(\omega) = \langle n_{d\downarrow}^0 \rangle (1 - \langle n_{d\downarrow}^0 \rangle) U^2 \omega^{-1} + O(\omega^{-2}), \quad (23)$$

where $\langle n_{d\sigma}^0 \rangle = (-i/2\pi) \int d\omega g_{d\sigma}^{\leq}(\omega)$ is the expectation value of $n_{d\sigma}$ calculated using the noninteracting lesser Green's function [see Eq. (14)]. From the above equations, one sees that choosing

$$A_1 = \langle n_{d\downarrow} \rangle (1 - \langle n_{d\downarrow} \rangle) / [\langle n_{d\downarrow}^0 \rangle (1 - \langle n_{d\downarrow}^0 \rangle)], \quad (24)$$

the moments of $\rho_{d\sigma}(\omega)$ up to the second one are reproduced exactly.

In the atomic limit, it can be easily verified using equations of motion⁵⁰ that the exact retarded Green's function is

$$G_{d\uparrow}^{r,\text{at}}(\omega) = \frac{1 - \langle n_{d\downarrow} \rangle}{\omega - E_d^{\uparrow} + i\eta} + \frac{\langle n_{d\downarrow} \rangle}{\omega - E_d^{\uparrow} - U + i\eta}. \quad (25)$$

From Eqs. (6) and (10), this implies

$$\Sigma_{\uparrow\text{int}}^{r2,\text{at}}(\omega) = \frac{\langle n_{d\downarrow} \rangle (1 - \langle n_{d\downarrow} \rangle) U^2}{\omega - E_d^{\uparrow} - (1 - \langle n_{d\downarrow} \rangle) U + i\eta}. \quad (26)$$

The usual contribution in U^2 to the retarded self-energy can be calculated easily in the atomic limit using Eq. (16) because the noninteracting spectral densities $\rho_{d\sigma}^0(\omega)$ become delta functions. The result can be written in the form

$$\Sigma_{\uparrow}^{r2,\text{at}}(\omega) = \frac{\langle n_{d\downarrow}^0 \rangle (1 - \langle n_{d\downarrow}^0 \rangle) U^2}{\omega - \varepsilon_{\text{eff}}^{\uparrow} + i\eta}. \quad (27)$$

From Eqs. (21), (24), (26), and (27), the coefficient A_2 is determined. The final result is

$$\begin{aligned} \Sigma_{\uparrow\text{int}}^{r2}(\omega) &= \frac{\langle n_{d\downarrow} \rangle (1 - \langle n_{d\downarrow} \rangle) \Sigma_{\uparrow}^{r2}(\omega)}{\langle n_{d\downarrow}^0 \rangle (1 - \langle n_{d\downarrow}^0 \rangle) + [\varepsilon_{\text{eff}}^{\uparrow} - E_d^{\uparrow} - (1 - \langle n_{d\downarrow} \rangle) U] [\Sigma_{\uparrow}^{r2}(\omega)/U^2]}. \end{aligned} \quad (28)$$

This result has the same form as in the spin dependent situation in equilibrium.^{12,53} However, the expectation values are calculated with distribution functions out of equilibrium.

From the equations of motion in the atomic limit (see Sec. III of Ref. 50), one obtains that the lesser Green's function for $V_{k\alpha} = 0$ is the sum of two delta functions $G_{d\uparrow}^{\leq,\text{at}}(\omega) = ia\delta(\omega - E_d^{\uparrow}) + ib\delta(\omega - E_d^{\uparrow} - U)$, where a and b are two real unknown constants. Using Eq. (25), we can write $G_{d\uparrow}^{\leq,\text{at}}(\omega) = -2iy(\omega) \text{Im} G_{d\uparrow}^{r,\text{at}}(\omega)$, where the function $y(\omega)$ is unknown. Electron-hole symmetry in the symmetric Anderson model imposes that $y(-\omega) = 1 - y(\omega)$. At equilibrium, $\mu_L = \mu_R = \mu$ and it can be shown from their definitions [Eqs. (2)]⁷⁶ that $y(\omega) = f(\omega - \mu)$. Out of equilibrium, $y(\omega)$ is ill-defined at $V_{k\alpha} = 0$, since the dot is disconnected to the leads, while even for small nonzero $V_{k\alpha}$ the problem is not trivial. In the general case, we assume that $y(\omega) = \tilde{f}(\omega)$, the average distribution function defined in Eq. (15). This has the correct limit at equilibrium or when the limit $V_{k\alpha} \rightarrow 0$ is taken for one of the leads first and then for the other. Using this choice, Eqs.

(6)–(8) imply for the lesser self-energy in the atomic limit

$$\Sigma_{\uparrow\text{int}}^{<,at}(\omega) = 2i\tilde{f}(\omega)\text{Im}\Sigma_{\uparrow\text{int}}^{r2,at}(\omega). \quad (29)$$

On the other hand, using Eqs. (8), (17), and (27), we can write for the second order correction in the same limit

$$\Sigma_{\uparrow}^{2<,at}(\omega) = 2i\tilde{f}(\omega)\text{Im}\Sigma_{\uparrow}^{r2,at}(\omega). \quad (30)$$

From these equations, and taking into account that $\Sigma_{\uparrow\text{int}}^{r2}(\omega)$ coincides with $\Sigma_{\uparrow}^{r2}(\omega)$ for small U and with $\Sigma_{\uparrow}^{r2,at}(\omega)$, the simplest interpolative expression for $\Sigma_{\uparrow}^{<}(\omega)$ is

$$\Sigma_{\uparrow\text{int}}^{<}(\omega) = \frac{\text{Im}\Sigma_{\uparrow\text{int}}^{r2}(\omega)}{\text{Im}\Sigma_{\uparrow}^{r2}(\omega)}\Sigma_{\uparrow}^{2<}(\omega), \quad (31)$$

when $\text{Im}\Sigma_{\uparrow}^{r2}(\omega) \neq 0$, and zero otherwise. Note that if $\text{Im}\Sigma_{\uparrow}^{r2}(\omega) = 0$, also $\Sigma_{\uparrow}^{2<}(\omega) = 0$, as can be seen from Eqs. (16) and (17).

E. The self-consistency problem

Equation (28) replaced in Eq. (10) and the same interchanging \uparrow and \downarrow define the retarded self-energies. Similarly Eq. (31) and the same for spin down are the lesser self-energies used. The expressions for $\Sigma_{\sigma}^{r2}(\omega)$ and $\Sigma_{\sigma}^{2<}(\omega)$ for constant Δ_{α} are given in the Appendix. The self-energies replaced in Eqs. (6)–(8) give the Green's functions for the interacting system that depend in general on four unknown quantities: $\varepsilon_{\text{eff}}^{\sigma}$ and $\langle n_{d\sigma} \rangle$. They are determined by the self-consistent solution of Eqs. (11) and (20). The resulting values of $\varepsilon_{\text{eff}}^{\sigma}$ do not have any particular physical meaning.

In the present work, we present results for the symmetric Anderson model, $E_d = -U/2$, $\mu_L = -\mu_R = eV_{ds}/2$, $\Delta_L = \Delta_R$ independent of energy, in the presence of a magnetic field B . In this case, due to electron-hole symmetry $\varepsilon_{\text{eff}}^{\sigma} = -\varepsilon_{\text{eff}}^{\bar{\sigma}}$ and $j_{L\sigma} = j_{R\bar{\sigma}}$. This reduces the problem to two self-consistent equations with two unknowns.

Unfortunately, for small temperature T , small nonzero V_{ds} and small nonzero B , there were instabilities in the numerical algorithm used. Also in some regions no solution could be found, while in other cases the solution jumped to another value. Therefore our results should be regarded as complementary to those obtained using renormalized perturbation theory (RPT), which correspond to $T=0$ and small V_{ds} .⁶⁶ We have tried an alternative approach, relaxing the condition of conservation of current and fixing $\varepsilon_{\text{eff}}^{\sigma}$ either at the value that satisfies the Friedel-Langreth sum rule (see below) or $\varepsilon_{\text{eff}}^{\sigma} = 0$. In the former case, the solution of Eqs. (11) for $\langle n_{d\sigma} \rangle$ was lost for very small V_{ds} , while in the second, the peak in the differential conductance dI/dV_{ds} was already split for very small values of B (of the order of $T_K/30$) in contradiction with the results of RPT. Therefore these alternatives were abandoned.

In general, we have started from the solution for large V_{ds} (with small reasonable $|\varepsilon_{\text{eff}}^{\sigma}|$), which was easy to find, and used it as an initial guess for lower V_{ds} . The procedure was repeated until reaching a small nonzero value of V_{ds} or some instability.

For $V_{ds}=0$, the problem is ill-posed since the current is conserved for any $\varepsilon_{\text{eff}}^{\sigma}$. Also the first derivative of $j_{L\sigma} - j_{R\sigma}$




FIG. 1. (a) Spectral density at the quantum dot as a function of energy for $V_{ds}=0$ and (b) differential conductance as a function of applied voltage, for $T=0$, and different values of B . We remind the reader that $T_K=0.45$.

with respect to V_{ds} vanishes and the second is very small for any $\varepsilon_{\text{eff}}^{\sigma}$. For $V_{ds}=T=0$, we have calculated the spectral density replacing the condition $j_{L\sigma}=j_{R\sigma}$ by the Friedel-Langreth sum rule⁷² for nonzero magnetic field, which for constant $\Delta=\Delta_R(\omega)+\Delta_L(\omega)$ can be written in the form^{12,55}

$$\pi\Delta\rho_{d\sigma}(0) = \frac{\Delta^2}{\Delta^2 + [\varepsilon_{\text{eff}}^{\sigma} + \Sigma_{\sigma}^r(0)]^2} = \sin^2(\pi\langle n_{d\sigma} \rangle). \quad (32)$$

III. NUMERICAL RESULTS

In this section we take $\Delta_R(\omega)=\Delta_L(\omega)$ independent of energy and $\Delta=\Delta_R+\Delta_L=1$ as the unit of energy. The origin of energy ($\omega=0$) is set at $(\mu_L+\mu_R)/2$. We have chosen $U=5$ for the numerical analysis because we want U large enough so that the system is at the Kondo regime, but for $U>6.5$, the magnetic susceptibility for $V_{ds}=T=0$ decreases with increasing U , indicating the failure of the IPA for a quantitative analysis at small non-zero magnetic field. These parameters lead to a Kondo temperature $T_K=0.45$ as determined from the width of the spectral density.

To represent the spectral density for $V_{ds}=T=0$, we have determined self-consistently the occupation numbers $\langle n_{d\sigma} \rangle$ and effective energies $\varepsilon_{\text{eff}}^{\sigma}$ using the Friedel-Langreth sum rule, while to calculate the conductivity at finite V_{ds} we have replaced this rule by the condition of conservation of current as described above.

In Fig. 1 we represent the differential conductance dI/dV_{ds} as a function of the applied voltage between drain and source V_{ds} for zero temperature, and compare it with the total spectral density of states $\rho_d(\omega)=\rho_{d\uparrow}(\omega)+\rho_{d\downarrow}(\omega)$ for $V_{ds}=0$ and $\omega=eV_{ds}$. The spectral density at $B=0$ shows a central peak corresponding to the Kondo peak with a half width at half maximum $T_K=0.45$ and two shoulders at $\omega=E_d=-2.5$ and $\omega=E_d+U=2.5$, which tend to separate with




FIG. 2. Spectral density as a function of energy for $T=0$, different values of the applied voltage $V=eV_{ds}$, and (a) $B=0$, (b) $B=1$. Note that $T_K=0.45$.

increasing magnetic field. The central peak shows a splitting which is tiny for $B=0.2$ (not shown) and increases with applied magnetic field. The splitting of the peak is larger than the Zeeman splitting $2B$ by a factor ~ 1.5 for $B=0.3$ and increases to slightly below 2 (~ 1.8) for $B=1.5$. This behavior agrees with previous studies of the spectral density under an applied magnetic field for large enough U .^{10,26,30,40,56,65}

The differential conductance dI/dV_{ds} was obtained by numerical differentiation of the total current $I=j_{L\uparrow}+j_{L\downarrow}=j_{R\uparrow}+j_{R\downarrow}$. For $0 < B < 0.5$ and $0 < V_{ds} < 1.5$, we could not find a self-consistent solution of the system of equations, and therefore we are not able to describe how the peak in the differential conductance splits with applied magnetic field at $T=0$. As described below, the situation improves with applied temperature. For $B=0$, the peak in dI/dV_{ds} is narrower than that of the spectral density with a half width at half maximum of 0.383. In addition the shoulders are displaced from the central peak. In general, the split peaks in dI/dV_{ds} are displaced to higher $|eV_{ds}|$ when compared with the same peaks in $\rho_d(\omega)$. Because of the lack of results for small non-vanishing magnetic field and voltage, our calculations for $T=0$ are not enough to establish if this tendency persists for small splittings.

In Fig. 2, we show the evolution of the spectral density with applied voltage. For $B=0$, the Kondo peak decreases slowly without splitting. For $V_{ds}=3$ (not shown), $\rho_d(\omega)$ is flat near $\omega=0$. Further increase in V_{ds} leads to only very small changes in $\rho_d(\omega)$. For large magnetic field, the changes in $\rho_d(\omega)$ with V_{ds} are less dramatic. For large V_{ds} the spectral density for different applied magnetic fields looks similar, but somewhat flatter and broader for larger B .

In Fig. 3 we show the magnetization $m=\langle n_{d\uparrow} \rangle - \langle n_{d\downarrow} \rangle$ as a function of applied voltage between drain and source for different values of the magnetic field and temperature. Because of our problems with the self-consistent solution for $T=0$, the values of $B > T_K=0.45$ taken at zero temperature




FIG. 3. Magnetization as a function of the applied voltage for $T=0$ (except dashed dot line for which $T=2T_K/3=0.3$) and several values of B .

are large enough so that the Kondo effect is at least partially destroyed. In this case, an increase in V_{ds} leads to a decrease in the magnetization which is expected as a consequence of the flattening of the spectral density of states discussed above. These curves for $T=0$ and comparatively large values of B show an initial slow decrease with increasing applied voltage, an inflection point for $eV_{ds} \sim 4B$, and an asymptotic approach to $m=0$ for large V_{ds} . Instead for small B (see the curve for $B=0.2$ and $T=3$), the magnetization first increases and then decreases with applied voltage. This can be interpreted in terms of a destruction of the Kondo singlet state for $V_{ds} \lesssim 2T_K$, which moves the system towards the local moment regime, followed by the same effects of flattening of the spectral density present for higher B .

For temperatures $T \sim T_K/2$, we have obtained some results in the problematic region for self-consistency ($0 < B < T_K$ and $0 < V_{ds} \lesssim 2T_K$). Some of them are shown in Fig. 4. However, still the value of B at which the splitting of the central peak takes place could not be identified precisely and abrupt changes in the solution take place, for example, for $B=T=0.3$ and $B=T=0.2$. In spite of this, these results and those for $T=0.5$ shown below suggest a transition from one maxima to two maxima in the differential conductance dI/dV_{ds} as a function of applied voltage for $B_c \sim 0.3$




FIG. 4. Spectral density a function of energy for $V_{ds}=0.1$ and (b) differential conductance as a function of applied voltage for $T=2T_K/3=0.3$, and different values of B .




FIG. 5. Spectral density as a function of energy for $T=2T_K/3=0.3$, different values of the applied voltage $V=eV_{ds}$, and (a) $B=0.2$, (b) $B=0.4$. The thin line corresponds to the contribution with spin up $\rho_{d\uparrow}(\omega)$ for $V=0.1$.

$=(2/3)T_K$. This value is consistent with that obtained by Hewson and co-workers⁶⁶ $B_c=0.584T_K$. Comparison of all split maxima of dI/dV_{ds} with the corresponding ones of the spectral density shows again that all the former lie at values of $|eV_{ds}|$ larger than the corresponding values of $|\omega|$ for the maxima of $\rho_d(\omega)$. This provides a natural explanation of the disagreement of the splitting of the peaks observed in experiments of the differential conductance in the presence of a magnetic field,^{14–16} when compared with accurate results of the spectral density.⁷³ The observed splitting for sufficiently large field was found to be higher than the corresponding one of the spectral density.

In Fig. 5, we show the evolution of $\rho_d(\omega)$ with applied voltage for $T=0.3$. For $B=0.2$ and small V_{ds} , the spectral density displays only one maximum at $\omega=0$, although its components $\rho_{d\uparrow}(\omega)$ and $\rho_{d\downarrow}(\omega)$ have maxima at $\omega=-\omega_M$ and $\omega=\omega_M$, respectively, with $\omega_M>0$, as shown in Fig. 5(a) for $eV_{ds}=0.1$. We remind the reader that due to our choice of parameters, $\rho_{d\downarrow}(\omega)=\rho_{d\uparrow}(-\omega)$. As in the case of zero temperature (Fig. 2), the changes driven by the applied voltage on $\rho_{d\sigma}(\omega)$ are smaller for higher magnetic fields.

In Fig. 6 we show dI/dV_{ds} as a function of V_{ds} for $T=0.5$, and the spectral density $\rho_d(\omega)$ for $eV_{ds}=0.1$. As before and to facilitate comparison, the abscissa of $\rho_d(\omega)$ is chosen in such a way that ω in Fig. 6(a) is the same as eV_{ds} in Fig. 6(b). The scales are the same as in Fig. 4. For this temperature, there were no instabilities in the algorithm for self-consistent equations and we could follow the splitting of the central Kondo peak. The transition from one maxima to two in the differential conductance takes place slightly below $B=0.3$. With increasing temperature, both the spectral density and the differential conductance become flatter, and the changes are more noticeable for small values of the magnetic field. Again, the structure of dI/dV_{ds} is broader than that of $\rho_d(\omega)$ and the maxima are more separated from $\omega=0$.

IV. SUMMARY AND DISCUSSION

We have generalized the spin-dependent interpolative perturbative approximation (IPA), based on second-order pertur-




FIG. 6. Same as Fig. 4 for $T=0.5 \sim T_K=0.45$.

bation theory in the Coulomb repulsion U , to the nonequilibrium case. This permits one to extend the validity of the results of nonequilibrium second order perturbation theory in $U/(\pi\Delta)$ to higher values of U . The effective unperturbed energy at the dot has been determined self-consistently asking that the current for each spin is conserved. This allows us to correct the shortcoming of ordinary perturbation theory in $U/(\pi\Delta)$ that the spin current is not conserved, except for the symmetric Anderson model without an applied magnetic field.

We have applied the method to study the effects of a magnetic field B on the differential conductance dI/dV_{ds} , starting from the symmetric Anderson model in which the on-site energy $E_d=-U/2$, setting the origin of energies at the average of the chemical potentials, and the resonant level width for both leads are equal ($\Delta_L=\Delta_R$) and independent of energy. The self-consistent procedure fails for temperatures below the Kondo temperature T_K , when $0<B<T_K$ and $0<V_{ds}\leq 2T_K$.

The procedure can in principle be applied to the general case (without starting from the symmetric case). Then, one has to solve four self-consistent equations for the effective occupations up and down, and the corresponding effective unperturbed levels. Similar problems were solved for the equilibrium case in the presence of a magnetic field¹² and to calculate persistent currents in a ring with an odd number N of electrons.⁵³ In the latter case, however, the procedure failed for large N . In the present case, one might expect more numerical difficulties after our experience with the simpler case treated here.

In spite of the above-mentioned failure, our results indicate that the critical magnetic field B_c for the splitting of the Kondo resonance in dI/dV_{ds} as a function of V_{ds} is near and slightly below $(2/3)T_K$. This is consistent with the results of renormalized perturbation theory⁶⁶ which give $B_c=0.584T_K$.

In the region of split peaks, the splitting is larger than the corresponding one of the spectral density of states $\rho_d(\omega)$ as a function of ω . This provides a natural explanation of why recent experiments of the differential conductance in the presence of a large magnetic field give a splitting larger than

two times the Zeeman splitting, as expected for the spectral density of states of states at equilibrium.^{14–16} Instead, previous results using slave bosons obtained a reduction of the peak splitting.³⁸

The effect of increasing bias voltage on the spectral density $\rho_d(\omega)$ is just an overall flattening of the energy dependence, without introducing new peaks. This is in agreement with recent calculations in the symmetric case⁶⁷ including terms of third and fourth order in U .

For small applied magnetic field, so that the system is in the Kondo regime, the magnetization as a function of the applied bias voltage V_{ds} first increases because due to the destruction of the Kondo singlet state, the system enters in the localized moment regime. For $V_{ds} \gtrsim 2T_K$, or $B > T_K$, the effect of flattening of the spectral density dominates and the magnetization decreases with increasing V_{ds} .

ACKNOWLEDGMENTS

I thank M. J. Sánchez for many helpful discussions. I am partially supported by CONICET. This work was sponsored by PICT 03-13829 of ANPCyT and PIP 5254 of CONICET.

APPENDIX: SELF-ENERGIES FOR CONSTANT Δ_α

Usually for problems in QDs, the density of states of the wide band of conduction electrons at the leads, as well as the hybridization with the dot $V_{k\alpha}$, can be taken as constant within the narrow energy range of T_K or other energy scales in the problem.⁵² In this case E_α and Δ_α become independent of energy and one of the integrals in Eqs. (12) and (13) can be done analytically. E_α merely renormalizes E_d .

Decomposing $g_{d\sigma}^<(\omega)$ as well as products of unperturbed Green functions as sums of simple fractions with single poles times Fermi functions, and decomposing products of Fermi functions using

$$f(x)f(y) = \frac{1}{\exp[(x+y)/T] - 1} [f(-x) - f(y)], \quad (\text{A1})$$

the first integral in frequency is decomposed into a sum of terms of the form

$$I_0 = \int d\omega f(\omega) \frac{1}{\omega - a \pm ib}, \quad (\text{A2})$$

with a and b real. This integral can be evaluated using the digamma function:⁷⁸

$$I_0 - \int d\omega f(\omega) \frac{1}{\omega \pm i\eta} = \psi\left(\frac{1}{2} + \frac{b \pm ia}{2\pi T}\right) - \psi\left(\frac{1}{2}\right). \quad (\text{A3})$$

After a lengthy algebra, the corrections of order U^2 to the self energies become

$$\begin{aligned} \Sigma_\uparrow^{r2}(\omega) &= \left(\frac{U}{2\pi}\right)^2 \int d\nu [g_{d\uparrow}^r(\omega - \nu)I_1(-\nu) + g_{d\uparrow}^<(\omega - \nu)I_2(\nu)], \\ \Sigma_\uparrow^{<2}(\omega) &= -\left(\frac{U}{2\pi}\right)^2 \int d\nu g_{d\uparrow}^<(\omega - \nu)I_1(\nu), \end{aligned} \quad (\text{A4})$$

and the same interchanging spins up and down, with

$$\begin{aligned} I_1(\nu) &= \frac{2}{\Delta^2} \text{Re} \left\{ \frac{2i\Delta}{\nu(\nu + 2i\Delta)} \sum_\alpha \{ (A_\alpha + B_{\bar{\alpha}}) [\varphi_{2\alpha}^*(-\nu) - \varphi_{1\alpha}] \right. \\ &\quad \left. + (A_\alpha + B_\alpha) [\varphi_{2\alpha}(\nu) - \varphi_{1\alpha}^*] \right\}, \end{aligned}$$

$$I_2(\nu) = -\sum_\alpha \frac{2i\Delta_\alpha}{\nu(\nu + 2i\Delta)} [\varphi_{1\alpha} + \varphi_{1\alpha}^* - \varphi_{2\alpha}(\nu) - \varphi_{2\alpha}^*(-\nu)], \quad (\text{A5})$$

where $\Delta = \Delta_L + \Delta_R$, $\bar{\alpha} = R(L)$ if $\alpha = L(R)$, and $\varphi_{1\alpha}$ and the functions $\varphi_{2\alpha}(\nu)$, $A_\alpha(\nu)$, and $B_\alpha(\nu)$ are

$$\begin{aligned} \varphi_{1\alpha} &= \psi\left(\frac{1}{2} + \frac{\Delta + i(\varepsilon_{eff}^\downarrow - \mu_\alpha)}{2\pi T}\right), \\ \varphi_{2\alpha}(\nu) &= \psi\left(\frac{1}{2} + \frac{\Delta + i(\varepsilon_{eff}^\downarrow - \mu_\alpha - \nu)}{2\pi T}\right), \\ A_\alpha(\nu) &= \frac{\Delta_\alpha^2}{\exp(\nu/T) - 1}, \end{aligned}$$

$$B_\alpha(\nu) = \frac{\Delta_L \Delta_R}{\exp[(\nu + \mu_\alpha - \mu_{\bar{\alpha}})/T] - 1}. \quad (\text{A6})$$

In equilibrium, when $\mu_L = \mu_R$, the expression for $\Sigma_\uparrow^{r2}(\omega)$ can be easily shown to coincide with that given by Horvatić and Zlatić.⁴⁴

¹L. I. Glazman and M. E. Raikh, JETP Lett. **47**, 452 (1988).

²T. K. Ng and P. A. Lee, Phys. Rev. Lett. **61**, 1768 (1988).

³D. Goldhaber-Gordon, H. Shtrikman, D. Mahalu, D. Abusch-Magder, U. Meirav, and M. A. Kastner, Nature (London) **391**, 156 (1998).

⁴S. M. Cronenwet, T. H. Oosterkamp, and L. P. Kouwenhoven, Science **281**, 540 (1998).

⁵D. Goldhaber-Gordon, J. Göres, M. A. Kastner, H. Shtrikman, D. Mahalu, and U. Meirav, Phys. Rev. Lett. **81**, 5225 (1998).

⁶W. G. van der Wiel, S. de Franceschi, T. Fujisawa, J. M. Elzerman, S. Tarucha, and L. P. Kouwenhoven, Science **289**, 2105 (2000).

⁷T. A. Costi, A. C. Hewson, and V. Zlatić, J. Phys.: Condens. Matter **6**, 2519 (1994).

⁸W. Izumida, O. Sakai, and S. Suzuki, J. Phys. Soc. Jpn. **70**, 1045 (2001).

⁹P. Recher, E. V. Sukhorukov, and D. Loss, Phys. Rev. Lett. **85**, 1962 (2000).

¹⁰T. A. Costi, Phys. Rev. B **64**, 241310(R) (2001).

¹¹M. E. Torio, K. Hallberg, S. Flach, A. E. Miroshnichenko, and M.

- Titov, *Eur. Phys. J. B* **37**, 399 (2004).
- ¹²A. A. Aligia and L. A. Salguero, *Phys. Rev. B* **70**, 075307 (2004); **71**, 169903(E) (2005).
- ¹³S. De Franceschi, R. Hanson, W. G. van der Wiel, J. M. Elzerman, J. J. Wijkema, T. Fujisawa, S. Tarucha, and L. P. Kouwenhoven, *Phys. Rev. Lett.* **89**, 156801 (2002).
- ¹⁴A. Kogan, S. Amasha, D. Goldhaber-Gordon, G. Granger, M. A. Kastner, and H. Shtrikman, *Phys. Rev. Lett.* **93**, 166602 (2004).
- ¹⁵D. M. Zumbühl, C. M. Marcus, M. P. Hanson, and A. C. Gossard, *Phys. Rev. Lett.* **93**, 256801 (2004).
- ¹⁶S. Amasha, I. J. Gelfand, M. A. Kastner, and A. Kogan, *cond-mat/0411485* (to be published).
- ¹⁷J. Nygard, W. F. Koehl, N. Mason, L. Di Carlo, and C. M. Marcus, *cond-mat/0410467* (to be published).
- ¹⁸R. Leturcq, L. Schmid, K. Ensslin, Y. Meir, D. C. Driscoll, and A. C. Gossard, *Phys. Rev. Lett.* **95**, 126603 (2005).
- ¹⁹A. C. Hewson, in *The Kondo Problem to Heavy Fermions* (Cambridge University Press, Cambridge, England, 1993).
- ²⁰P. Mehta and N. Andrei, *cond-mat/0508026* (to be published).
- ²¹R. M. Konik, H. Saleur, and A. Ludwig, *Phys. Rev. B* **66**, 125304 (2002).
- ²²A. Schiller and S. Hershfield, *Phys. Rev. B* **51**, 12896(R) (1995).
- ²³N. E. Bickers, D. L. Cox, and J. W. Wilkins, *Phys. Rev. B* **36**, 2036 (1987).
- ²⁴R. Monnier, L. Degiorgi, and B. Delley, *Phys. Rev. B* **41**, 573 (1990); Yu Zhou, S. P. Bowen, D. D. Koelling, and R. Monnier, *ibid.* **43**, 11071 (1991).
- ²⁵N. O. Moreno, A. Lobos, A. A. Aligia, E. D. Bauer, S. Bobev, V. Fritsch, J. L. Sarrao, P. G. Pagliuso, J. D. Thompson, C. D. Batista, and Z. Fisk, *Phys. Rev. B* **71**, 165107 (2005).
- ²⁶Y. Meir, N. S. Wingreen, and P. A. Lee, *Phys. Rev. Lett.* **70**, 2601 (1993).
- ²⁷N. S. Wingreen and Y. Meir, *Phys. Rev. B* **49**, 11040 (1994).
- ²⁸A. Rosch, J. Kroha, and P. Wölfle, *Phys. Rev. Lett.* **87**, 156802 (2001).
- ²⁹E. Müller-Hartmann, *Z. Phys. B: Condens. Matter* **57**, 281 (1984).
- ³⁰J. E. Moore and X-G Wen, *Phys. Rev. Lett.* **85**, 1722 (2000).
- ³¹K. Kang, S. Y. Cho, J. J. Kim, and S. C. Shin, *Phys. Rev. B* **63**, 113304 (2001).
- ³²H. Hu, G-M. Zhang, and L. Yu, *Phys. Rev. Lett.* **86**, 5558 (2001).
- ³³A. A. Aligia, K. Hallberg, B. Normand, and A. P. Kampf, *Phys. Rev. Lett.* **93**, 076801 (2004).
- ³⁴C.-Y. Lin, A. H. Castro Neto, and B. A. Jones, *Phys. Rev. B* **71**, 035417 (2005).
- ³⁵A. A. Aligia and A. M. Lobos, *J. Phys.: Condens. Matter* **17**, S1095 (2005), and references therein.
- ³⁶A. M. Lobos and A. A. Aligia, *Phys. Rev. B* **74**, 165417 (2006).
- ³⁷R. Aguado and D. C. Langreth, *Phys. Rev. Lett.* **85**, 1946 (2000).
- ³⁸B. Dong and X. L. Lei, *Phys. Rev. B* **63**, 235306 (2001).
- ³⁹D. E. Logan, M. P. Eastwood, and M. A. Tusch, *J. Phys.: Condens. Matter* **10**, 2673 (1998); M. T. Glossop and D. E. Logan, *ibid.* **14**, 6737 (2002).
- ⁴⁰D. E. Logan and N. L. Dickens, *J. Phys.: Condens. Matter* **13**, 9713 (2001).
- ⁴¹N. S. Vidhyadhiraja and D. E. Logan, *Eur. Phys. J. B* **39**, 313 (2004).
- ⁴²A. Rosch, J. Paaske, J. Kroha, and P. Wölfle, *Phys. Rev. Lett.* **90**, 076804 (2003).
- ⁴³K. Yosida and K. Yamada, *Prog. Theor. Phys.* **46**, 244 (1970); **53**, 1286 (1975); K. Yamada, *ibid.* **53**, 970 (1975).
- ⁴⁴B. Horvatić and V. Zlatić, *Phys. Status Solidi B* **99**, 251 (1980).
- ⁴⁵B. Horvatić and V. Zlatić, *Phys. Rev. B* **30**, 6717 (1984); B. Horvatić, D. Šokčević, and V. Zlatić, *ibid.* **36**, 675 (1987).
- ⁴⁶V. Zlatić, B. Horvatić, B. Dolički, S. Grabowski, P. Entel, and K. D. Schotte, *Phys. Rev. B* **63**, 035104 (2000).
- ⁴⁷L. V. Keldysh, *Sov. Phys. JETP* **20**, 1018 (1965).
- ⁴⁸E. M. Lifshitz and A. L. Pitaevskii, *Physical Kinetics* (Pergamon, Oxford, 1981).
- ⁴⁹G. D. Mahan, *Many Particle Physics* (Kluwer/Plenum, New York, 2000).
- ⁵⁰C. Niu, D. L. Lin, and T-H. Lin, *J. Phys.: Condens. Matter* **11**, 1511 (1999).
- ⁵¹A. A. Aligia, *Phys. Rev. B* **64**, 121102(R) (2001); A. Lobos and A. A. Aligia, *ibid.* **68**, 035411 (2003).
- ⁵²A. A. Aligia and C. R. Proetto, *Phys. Rev. B* **65**, 165305 (2002).
- ⁵³A. A. Aligia, *Phys. Rev. B* **66**, 165303 (2002).
- ⁵⁴A. Oguri, *Phys. Rev. B* **63**, 115305 (2001), and references therein.
- ⁵⁵A. C. Hewson, *Phys. Rev. Lett.* **70**, 4007 (1993).
- ⁵⁶A. C. Hewson, *cond-mat/0509296* (to be published).
- ⁵⁷A. Martín-Rodero, M. Baldo, F. Flores, and R. Pucci, *Solid State Commun.* **44**, 911 (1982).
- ⁵⁸A. Martín-Rodero, E. Louis, F. Flores, and C. Tejedor, *Phys. Rev. B* **33**, 1814 (1986).
- ⁵⁹A. L. Yeyati, A. Martín-Rodero, and F. Flores, *Phys. Rev. Lett.* **71**, 2991 (1993).
- ⁶⁰H. Kajueter and G. Kotliar, *Phys. Rev. Lett.* **77**, 131 (1996).
- ⁶¹I. Maruyama, N. Shibata, and K. Ueda, *J. Phys. Soc. Jpn.* **73**, 3239 (2004).
- ⁶²S. Hershfield, J. H. Davies, and J. W. Wilkins, *Phys. Rev. B* **46**, 7046 (1992).
- ⁶³M. Hamasaki, *Phys. Rev. B* **69**, 115313 (2004).
- ⁶⁴T. Fujii and K. Ueda, *Phys. Rev. B* **68**, 155310 (2003).
- ⁶⁵T. Fujii and K. Ueda, *J. Phys. Soc. Jpn.* **74**, 127 (2005).
- ⁶⁶A. C. Hewson, J. Bauer, and A. Oguri, *J. Phys.: Condens. Matter* **17**, 5413 (2005).
- ⁶⁷M. Hamasaki, *cond-mat/0506752* (to be published).
- ⁶⁸R. Świrkowicz, J. Barnaś, and M. Wilczyński, *Phys. Rev. B* **68**, 195318 (2003).
- ⁶⁹R. C. Monreal and F. Flores, *Phys. Rev. B* **72**, 195105 (2005).
- ⁷⁰V. Kashcheyevs, A. Aharony, and O. Entin-Wohlman, *Phys. Rev. B* **73**, 125338 (2006).
- ⁷¹H. Pothier, S. Guéron, N. O. Birge, D. Esteve, and M. H. Devoret, *Phys. Rev. Lett.* **79**, 3490 (1997).
- ⁷²D. C. Langreth, *Phys. Rev.* **150**, 516 (1966).
- ⁷³T. A. Costi, *Phys. Rev. Lett.* **85**, 1504 (2000).
- ⁷⁴W. Hofstetter, *Phys. Rev. Lett.* **85**, 1508 (2000).
- ⁷⁵H. Schweitzer and G. Czycholl, *Z. Phys. B: Condens. Matter* **83**, 93 (1991).
- ⁷⁶Y. Meir and N. S. Wingreen, *Phys. Rev. Lett.* **68**, 2512 (1992); A-P. Jauho, N. S. Wingreen, and Y. Meir, *Phys. Rev. B* **50**, 5528 (1994).
- ⁷⁷W. Nolting and W. Borgiel, *Phys. Rev. B* **39**, 6962 (1989).
- ⁷⁸M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1972).