

Kondo and Dicke effect in quantum dots side coupled to a quantum wire

Pedro A. Orellana,¹ Gustavo A. Lara,² and Enrique V. Anda³

¹*Departamento de Física, Universidad Católica del Norte, Casilla 1280, Antofagasta, Chile*

²*Departamento de Física, Universidad de Antofagasta, Casilla 170, Antofagasta, Chile*

³*Departamento de Física, P. U. Católica do Rio de Janeiro, C.P. 38071-970, Rio de Janeiro, RJ, Brazil*

(Received 9 June 2006; revised manuscript received 5 September 2006; published 22 November 2006)

Electron tunneling through quantum dots side coupled to a quantum wire, in equilibrium and nonequilibrium Kondo regime, is studied. The mean-field finite- U slave-boson formalism is used to obtain the solution of the problem. We have found that the transmission spectrum shows a structure with two antiresonances localized at the renormalized energies of the quantum dots. The density of states (DOS) of the system shows that when the Kondo correlations are dominant there are two Kondo regimes with its own Kondo temperature. The above behavior of the DOS can be explained by quantum interference in the transmission through the two different resonance states of the quantum dots coupled to common leads. This result is analogous to the Dicke effect in optics. We investigate the many-body Kondo states as a function of the parameters of the system.

DOI: [10.1103/PhysRevB.74.193315](https://doi.org/10.1103/PhysRevB.74.193315)

PACS number(s): 73.21.La, 73.63.Kv, 72.10.Fk, 85.35.Be

I. INTRODUCTION

The Kondo effect in quantum dots (QDs) has been extensively studied.¹⁻³ The QDs allow studying systematically the quantum-coherence many-body Kondo state, due to the possibility of continuous tuning the relevant parameters governing the properties of this state, in equilibrium and nonequilibrium situations. Recently, Kondo effect has been studied in side attach quantum dot⁴ and parallel quantum dots.^{5,6} Recent electron transport experiments showed that Kondo and Fano resonances occur simultaneously.⁷ Multiple scattering of traveling electronic waves on a localized magnetic state are crucial for the formation of both resonances. The condition for the Fano resonance is the existence of two scattering channels: a discrete level and a broad continuum band.⁸

An alternative configuration consists of two single QDs side attached to a perfect quantum wire (QW). This structure is reminiscent of the cross-bar-shaped quantum wave guides.⁹ In this case, the QDs act as scattering centers in close analogy with the traditional Kondo effect.¹⁰ This configuration was previously studied theoretically by Stefański and Tamura *et al.*¹¹ and experimentally by Sasaki *et al.*¹²

In this work we study the transport properties of two single quantum dots side coupled to a quantum wire in the Kondo regime. We use the finite- U slave boson mean-field approach, which was initially developed by Kotliar and Ruckenstein¹³ and used later by Dong and Lei to study the transport through coupled double quantum dots connected to leads.¹⁴ This approach enforces the correspondence between the impurity fermions and the auxiliary bosons to a mean-field level to release the $U=\infty$ restriction. In quantum dots, this approach allows to treat the dot-lead coupling nonperturbatively for an arbitrary strength of the Coulomb interaction U .¹⁴ We have found that the transmission spectrum shows a structure with two antiresonances localized at the renormalized energies of the quantum dots. The DOS of the system shows that when the Kondo correlations are dominant there are two Kondo regimes each with its own Kondo temperature. The above behavior of the DOS can be explained by quantum interference in the transmission through the two different resonance states of the quantum dots coupled to

common lead. This phenomenon is in analogy to the Dicke effect in quantum optics, that takes place in the spontaneous emission of two closely lying atoms radiating a photon into the same environment.¹⁵ In the electronic case, however, the decay rates (level broadening) are produced by the indirect coupling of the up-down QDs, giving rise to a fast (*superradiant*) and a slow (*subradiant*) mode. Recently, Brandes reviewed the Dicke effect in mesoscopic systems.¹⁶

II. MODEL

Let us consider two single quantum dot (2QD) side coupled to a perfect quantum wire (QW) (see Fig. 1). We adopt the two-impurities Anderson Hamiltonian. Each dot has a single level energy ε_l (with $l=1,2$), and a intradot Coulomb repulsion U . The two side attached quantum dots are coupled to the QW with coupling t_0 . The QW sites have local energies $\varepsilon_{wi,\sigma}=0$ and a hopping parameter t .

The corresponding Hamiltonian model is

$$H_0 = -t \sum_{i,\sigma} (c_{i,\sigma}^\dagger c_{i+1,\sigma} + \text{H.c.}) + \sum_{l=1,2,\sigma} \left[-t_{l,\sigma} (c_{0,\sigma}^\dagger f_{l,\sigma} + \text{H.c.}) + \left(\varepsilon_{l,\sigma} + \frac{U}{2} \hat{n}_{l,-\sigma} \right) \hat{n}_{l,\sigma} \right] \quad (1)$$

where $c_{i,\sigma}^\dagger$ ($c_{i,\sigma}$) is the creation (annihilation) operator of an electron with spin σ at the i th site of the quantum wire; $f_{l,\sigma}^\dagger$ ($f_{l,\sigma}$) is the creation (annihilation) operator of an electron with spin σ in the l th QD and $\hat{n}_{l,\sigma}$ is the corresponding number operator.

To find the solution of this correlated fermions system, we appeal to an analytical approach where, generalizing the infinite- U slave-boson approximation¹⁷ the Hilbert space is enlarged at each site, to contain in addition to the original fermions a set of four bosons¹³ represented by the creation (annihilation) operators e_l^\dagger (e_l), $p_{l,\sigma}^\dagger$ ($p_{l,\sigma}$), and d_l^\dagger (d_l) for the l th dot. They act as projectors onto empty, single occupied (with spin up and down) and doubly occupied electron states, respectively. Then, each creation (annihilation) operator of an electron with spin σ in the l th QD, is substituted by $f_{l,\sigma}^\dagger \tilde{Z}_{l,\sigma}^\dagger$ ($\tilde{Z}_{l,\sigma} f_{l,\sigma}$) where




FIG. 1. Scheme of side-coupled quantum dots attached laterally to a perfect quantum wire (QW). The QW is coupled to the left (L) and right (R) noninteracting leads.

$$\tilde{Z}_{l,\sigma} = \frac{e_l^\dagger p_{l,\sigma} + p_{l,-\sigma}^\dagger d_l}{\sqrt{1 - d_l^\dagger d_l - p_{l,\sigma}^\dagger p_{l,\sigma}} \sqrt{1 - e_l^\dagger e_l - p_{l,-\sigma}^\dagger p_{l,-\sigma}}}. \quad (2)$$

The denominator is chosen to reproduce the correct $U \rightarrow 0$ limit in the mean-field approximation without changing neither the eigenvalues nor the eigenvector.

The constraint, i.e., the completeness relation $\sum_{\sigma} p_{l,\sigma}^\dagger p_{l,\sigma} + b_l^\dagger b_l + d_l^\dagger d_l = 1$ and the condition among fermions and bosons $n_{l,\sigma} - p_{l,\sigma}^\dagger p_{l,\sigma} - d_l^\dagger d_l = 0$, will be incorporated with Lagrange multipliers $\lambda_l^{(1)}$ and $\lambda_l^{(2)}$ into the Hamiltonian. Also in the mean-field approximation all the boson operators are replaced by their expectation value which can be chosen, without loss of generality, as real numbers.

The Hamiltonian in this new and enlarged Hilbert space, is, $\mathcal{H} = H_b + H_e$, where

$$H_b = \sum_{l=1,2} \lambda_l^{(1)} (p_{l,\uparrow}^2 + p_{l,\downarrow}^2 + e_l^2 + d_l^2 - 1) - \sum_{l=1,2,\sigma} \lambda_{l,\sigma}^{(2)} (p_{l,\sigma}^2 + d_l^2) + U \sum_{l=1,2} d_l^2, \quad (3)$$

depends explicitly only upon the boson expectation values and the Lagrange multipliers, and

$$H_e = -t \sum_{i,\sigma} (c_{i,\sigma}^\dagger c_{i+1,\sigma} + \text{H.c.}) + \sum_{l=1,2,\sigma} [-\tilde{t}_{l,\sigma} (c_{0,\sigma}^\dagger f_{l,\sigma} + \text{H.c.}) + \tilde{\varepsilon}_{l,\sigma} n_{l,\sigma}] \quad (4)$$

is a tight-binding Hamiltonian that depends implicitly on the boson expectation values through the parameters: $\tilde{\varepsilon}_{l,\sigma} = \varepsilon_{l,\sigma} + \lambda_{l,\sigma}^{(2)}$, $\tilde{t}_{l,\sigma} = t_0 \langle \tilde{Z}_{l,\sigma} \rangle$.

As we work at zero temperature, the boson operators expectation values and the Lagrange multipliers are determined by minimizing the energy $\langle \mathcal{H} \rangle$ with respect to these quanti-




FIG. 2. Transmission spectrum in equilibrium for $V_g = -3$, $\delta V = 0.1$ and various values of U .




FIG. 3. DOS for $V_g = -3$, $\delta V = 0.1$ (solid line), 0.5 (dashed line). The on site energy U , is (a) 1, (b) 3, (c) 5, and (d) 6.

ties. It is obtained in this way, a set of nonlinear equations for each quantum dot, relating the expectation values of the four bosonic operators, the three Lagrange multipliers, and the electronic expectation values,

$$p_{l,\sigma}^2 = \langle \hat{n}_{l,\sigma} \rangle - d_l^2, \quad (5a)$$

$$e_l^2 = 1 - \sum_s \langle \hat{n}_{l,s} \rangle + d_l^2, \quad (5b)$$

$$\lambda_l^{(1)} = \frac{t_0}{e_l} \sum_s \langle f_{l,s}^\dagger c_{0,s} \rangle \frac{\partial \langle \tilde{Z}_{l,s} \rangle}{\partial e_l}, \quad (5c)$$

$$\lambda_l^{(1)} - \lambda_{l,\sigma}^{(2)} = \frac{t_0}{p_{l,\sigma}} \sum_s \langle f_{l,s}^\dagger c_{0,s} \rangle \frac{\partial \langle \tilde{Z}_{l,s} \rangle}{\partial p_{l,\sigma}}, \quad (5d)$$

$$U + \lambda_l^{(1)} - \sum_s \lambda_{l,s}^{(2)} = \frac{t_0}{d_l} \sum_s \langle f_{l,s}^\dagger c_{0,s} \rangle \frac{\partial \langle \tilde{Z}_{l,s} \rangle}{\partial d_l}. \quad (5e)$$

where l is the dot index, s, σ are spin indexes, and $\langle \tilde{Z}_{l,s} \rangle$ satisfies,

$$\langle \tilde{Z}_{l,s} \rangle = \frac{p_{l,s} (e_l + d_l)}{\sqrt{(1 - d_l^2 - p_{l,s}^2)(1 - e_l^2 - p_{l,-s}^2)}}. \quad (6)$$

To obtain the electronic expectation values $\langle \dots \rangle$, the Hamiltonian, H_e , is diagonalized and their stationary states can be written as




FIG. 4. Current (solid line) and differential conductance (dashed line) for $V_g = -3$, on site energy, $U = 6$ for (a) $\delta V = 0.1$ and (b) $\delta V = 0.5$.




FIG. 5. Current (solid line) and differential conductance (dashed line) for $V_g = -3$, on site energy, $U = 6$ for (a) $\delta V = 0.1$ and (b) $\delta V = 0.5$.

$$|\psi_k\rangle = \sum_{j=-\infty}^{\infty} a_j^k |j\rangle + \sum_{l=1}^2 b_l^k |l\rangle, \quad (7)$$

where a_j^k and b_l^k are the probabilities amplitudes to find the electron at the site j and at the l th QD, respectively, with energy $\omega = -2t \cos k$. As we study the paramagnetic case the spin index is neglected.

The amplitudes a_j^k and b_l^k obey the following linear difference equations:

$$\omega a_j^k = -t(a_{j+1}^k + a_{j-1}^k), \quad j \neq 0, \quad (8a)$$

$$\omega a_0^k = -t(a_1^k + a_{-1}^k) - \tilde{t}_1 b_1^k - \tilde{t}_2 b_2^k, \quad (8b)$$

$$(\omega - \tilde{\varepsilon}_1) b_1^k = -\tilde{t}_1 a_0^k, \quad (8c)$$

$$(\omega - \tilde{\varepsilon}_2) b_2^k = -\tilde{t}_2 a_0^k. \quad (8d)$$

In order to study the solutions of Eqs. (8), we assume that the electrons are described by a unitary incident amplitude plane wave and reflection and transmission amplitudes r and τ , respectively. That is,

$$a_j^k = e^{ik \cdot j} + r e^{-ik \cdot j} \quad (k \cdot j < 0), \quad (9a)$$

$$a_j^k = \tau e^{ik \cdot j} \quad (k \cdot j > 0). \quad (9b)$$

Inserting Eqs. (9) into Eqs. (8), we get an inhomogeneous system of linear equations for τ , r , a_j^k , and b_l^k , leading to the following expression in equilibrium:

$$\tau = \frac{1}{1 + i \left(\frac{\tilde{\Gamma}_1}{\omega - \tilde{\varepsilon}_1} + \frac{\tilde{\Gamma}_2}{\omega - \tilde{\varepsilon}_2} \right)}, \quad (10)$$

where $\tilde{\Gamma}_l = \pi \tilde{t}_l^2 \rho_0(\omega)$ ($l=1,2$) is the renormalized coupling between each quantum dot and the leads of density of states $\rho_0(\omega)$. In spite of the apparent simplicity of the expression, it is necessary to remember that the quantity \tilde{t}_l implicitly depends on the expectation values of the boson operators also as fermion operators.

The transmission probability is given by $T = |\tau|^2$,

$$T(\omega) = \frac{1}{1 + \left(\frac{\tilde{\Gamma}_1}{\omega - \tilde{\varepsilon}_1} + \frac{\tilde{\Gamma}_2}{\omega - \tilde{\varepsilon}_2} \right)^2}. \quad (11)$$

From the amplitudes b_1^k and b_2^k we obtain the local density of states (LDOS) at the quantum dot l (with $l=1,2$). In equilibrium that is,

$$\rho_l(\omega) = \frac{1}{\pi \tilde{\Gamma}_l} \frac{\left(\frac{\tilde{\Gamma}_l}{\omega - \tilde{\varepsilon}_l} \right)^2}{1 + \left(\frac{\tilde{\Gamma}_1}{\omega - \tilde{\varepsilon}_1} + \frac{\tilde{\Gamma}_2}{\omega - \tilde{\varepsilon}_2} \right)^2}. \quad (12)$$

In the nonequilibrium case, we suppose a finite source-drain biased with a symmetric voltage drop. The incident electrons from the left-hand side (L), they are in equilibrium with thermodynamical potential $\mu_L = V/2$, and the incidents from the right-hand side (R), they are in equilibrium with thermodynamical potential $\mu_R = -V/2$.

Once the amplitudes $a_{j,\sigma}^k$ and $b_{j,\sigma}^k$ are known, the electronic expectation value is obtained from

$$\langle f_l^\dagger c_j \rangle = \frac{1}{2} \sum_{\alpha=L,R} \frac{1}{N} \sum_{k_\alpha} f(\epsilon_{k_\alpha} - \mu_\alpha) b_l^{k_\alpha} a_j^{k_\alpha}. \quad (13)$$

And the current is obtained from

$$J = 2 \frac{2e}{\hbar} t \sum_{\alpha, k_\alpha} f(\epsilon_{k_\alpha} - \mu_\alpha) \text{Im} \{ a_0^{k_\alpha} a_1^{k_\alpha} \}, \quad (14)$$

where $f(\epsilon_{k_\alpha} - \mu_\alpha)$ is the Fermi function for incident electrons from the α side.

III. RESULTS

We solve numerically the set of nonlinear equations and take typical values for the parameters that define the system, $t = 25\Gamma$, $t_0 = 5\Gamma$ where $\Gamma = \pi t_0^2 \rho_0(0)$ is taken to be the unit of energy.

We consider first the situation in equilibrium where the two dots local state energies are set by $\varepsilon_1 = V_g - \delta V$ and $\varepsilon_2 = V_g + \delta V$. We choose the value of $V_g = -3\Gamma$. From now on all energies are in units of Γ .

The transmission probability, T , is displayed in Fig. 2 for various values of U . The transmission probability always reaches zero at $\omega = \tilde{\varepsilon}_1$ and $\tilde{\varepsilon}_2$ and unitary value at $\omega = (\tilde{\varepsilon}_1 + \tilde{\varepsilon}_2)/2$. For small values of U the antiferromagnetic spin-spin correlation between the dots is dominant and the system does not possess a Kondo regime.¹⁸ Increasing U , a sharp feature develops close to the Fermi energy ($\omega = 0$), indicating the appearance of a Kondo resonance.

For U sufficiently large the transmission can be written approximately as the superposition of Fano and Breit-Wigner line shapes,

$$T(\omega) \approx \frac{(\epsilon + q)^2}{\epsilon^2 + 1} + \frac{\tilde{\Delta}^2}{\omega^2 + \tilde{\Delta}^2}, \quad (15)$$

where $\epsilon = \omega/2\tilde{\Gamma}$, $q = 0$, with $\tilde{\Delta} = \delta\tilde{V}^2/2\tilde{\Gamma}$.

The DOS gives us more details about the formation of the Kondo resonance. The DOS is displayed in Fig. 3. In the Kondo regime the DOS can be written as the superposition of the two Lorentzian. These results imply the existence of two Kondo temperature $T_{1K}=2\tilde{\Gamma}$ and $T_{2K}=\tilde{\Delta}=\delta\tilde{V}^2/\tilde{\Gamma}$, associated to each Kondo regime,

$$\rho(\omega) \approx \frac{1}{\pi} \frac{2\tilde{\Gamma}}{\omega^2 + 4\tilde{\Gamma}^2} + \frac{1}{\pi} \frac{\tilde{\Delta}}{\omega^2 + \tilde{\Delta}^2}. \quad (16)$$

The above behavior of the DOS is due to quantum interference taking place in the transmission through the two different discrete states (the two quantum dot levels) coupled to common leads. This phenomenon resembles the Dicke effect in optics, which takes place in the spontaneous emission of a pair of atoms radiating a photon with a wavelength much larger than the separation between them.¹⁵ The luminescence spectrum is characterized by a narrow and a broad peak, associated with long- and short-lived states, respectively. The former state, weakly coupled to the electromagnetic field, is called *subradiant*, and the latter, strongly coupled, *superradiant* state. In the present case this effect is due to the indirect coupling between up-down QDs through the QW. The states strongly coupled to the QW yield an effective width $2\tilde{\Gamma}$ while those weakly coupled to the QW give a Dicke state with width $\tilde{\Delta}$. A similar result was found for a parallel double quantum dot without electron-electron interaction.¹⁹

The current and the differential conductance dJ/dV are two significant and experimentally measured quantities, which have been calculated numerically at finite source-drain biases.

Figure 4 displays the characteristic J - V (solid line) and the differential conductance dJ/dV - V (dashed line) for two values of δV . For $\delta V=0.1\Gamma$ the current shows a pronounced plateau around zero bias while for $\delta V=0.5\Gamma$ the plateau is less defined. However in both cases the differential conductance shows an anomaly at zero bias.

Figure 5 shows details of the current and differential conductance around zero bias.

We can obtain the expressions for the current and the differential conductance by integrating over ω the transmission probability given in Eq. (15),

$$J \approx \frac{2e}{h} \left[V - 2\tilde{\Gamma} \arctan\left(\frac{V}{2\tilde{\Gamma}}\right) + \tilde{\Delta} \arctan\left(\frac{V}{\tilde{\Delta}}\right) \right],$$

$$\frac{dJ}{dV} \approx \frac{2e^2}{h} \left(1 - \frac{4\tilde{\Gamma}^2}{\left(\frac{V}{2}\right)^2 + 4\tilde{\Gamma}^2} + \frac{\tilde{\Delta}^2}{\left(\frac{V}{2}\right)^2 + \tilde{\Delta}^2} \right), \quad (17)$$

We identify each term of the above equation as follows. The first term on the right-hand side of Eqs. (17) is the contribution arising from an ideal unidimensional conductor. The second term comes from the Kondo-Fano state with temperature T_{1k} giving a quasiplateau for the current and almost zero differential conductance when $|V| \ll \tilde{\Gamma}$. The third term arises from the Kondo-Dicke state weakly coupled to the wire. It is responsible for an abrupt increase of the current and an amplification on the differential conductance around zero bias. Finally, for $|V| > \tilde{\Gamma}$, the Kondo effect disappears.

IV. SUMMARY

We have studied the transport through two single side-coupled quantum dots using the finite- U slave boson mean field approach at $T=0$. We have found that the transmission spectrum shows a structure with two antiresonances localized at the renormalized energies of the quantum dots. The DOS of the system shows that when the Kondo correlations are dominant there are two Kondo regimes each with its own Kondo temperature. The above behavior of the DOS is due to quantum interference in the transmission through the two different resonance states of the quantum dots coupled to common leads. This result is analogous to the Dicke effect in optics. These phenomena have been analyzed as a function of the relevant parameters of the system.

ACKNOWLEDGMENTS

Two of the authors (G.A.L. and P.A.O.) acknowledge financial support from Milenio ICM P02-054F. One of the authors (P.A.O.) also thanks FONDECYT (Grants Nos. 1060952 and 7020269), and one of the authors (G.A.L.) thanks U.A. (PEI-1305-04). One of the authors (E.V.A.) acknowledges support from the Brazilian agencies CNPq (CIAM project) and FAPERJ.

¹L. I. Glazman and M. É. Raikh, JETP Lett. **47**, 452 (1988); T. K. Ng and P. A. Lee, Phys. Rev. Lett. **61**, 1768 (1988).

²D. Goldhaber-Gordon *et al.*, Nature (London) **391**, 156 (1998); D. Goldhaber-Gordon *et al.*, Phys. Rev. Lett. **81**, 5225 (1998).

³S. M. Cronenwett *et al.*, Science **281**, 540 (1998).

⁴R. Franco *et al.*, Phys. Rev. B **67**, 155301 (2003); M. E. Torio *et al.*, *ibid.* **65**, 085302 (2002).

⁵Yoichi Tanaka and Norio Kawakami, Phys. Rev. B **72**, 085304 (2005).

⁶Rui Sakano and Norio Kawakami Phys. Rev. B **72**, 085303 (2005).

⁷J. Göres *et al.*, Phys. Rev. B **62**, 2188 (2000).

⁸U. Fano, Phys. Rev. **124**, 1866 (1961).

⁹P. Debray *et al.*, Phys. Rev. B **61**, 10950 (2000).

¹⁰K. Kang *et al.*, Phys. Rev. B **63**, 113304 (2001).

¹¹Piotr Stefański, Solid State Commun. **128**, 29 (2006); Hiroyuki Tamura and Leonid Glazman, Phys. Rev. B **72**, 121308(R) (2005).

¹²S. Sasaki *et al.*, Phys. Rev. B **73**, 161303(R) (2006).

¹³G. Kotliar and A. E. Ruckenstein, Phys. Rev. Lett. **57**, 1362 (1986), and references cited therein.

¹⁴B. Dong and X. L. Lei, Phys. Rev. B **63**, 235306 (2001).

¹⁵R. H. Dicke, Phys. Rev. **89**, 472 (1953).

¹⁶T. Brandes, Phys. Rep. **408**, 315 (2005).

¹⁷P. Coleman, Phys. Rev. B **29**, 3035 (1984).

¹⁸C. A. Busser *et al.*, Phys. Rev. B **62**, 9907 (2000).

¹⁹P. A. Orellana *et al.*, Phys. Rev. B **70**, 233315 (2005).