

Hyperfine-interaction-induced critical exponents in the quantum Hall effect

V. Kagalovsky¹ and I. Vagner^{2,3,*}

¹*Sami Shamoon College of Engineering, P.O. Box 45, Beer-Sheva, 84100, Israel*

²*RCQCE-Research Center for Quantum Communication Engineering, Holon Institute of Technology, 52 Golomb Street, Holon 58102, Israel*

³*Cavendish Laboratory, Cambridge University, Madingley Road, Cambridge CB3 0HE, United Kingdom*

(Received 29 November 2006; published 9 March 2007)

We study localization-delocalization transition in quantum Hall systems with a random field of nuclear spins acting on two-dimensional (2D) electron spins via hyperfine contact (Fermi) interaction. We use the Chalker-Coddington network model, which corresponds to the projection onto the lowest Landau level. The inhomogeneous nuclear polarization acts on the electrons as an additional confining potential and, therefore, introduces additional parameter p (the probability to find a polarized nucleus in the vicinity of a saddle point of random potential) responsible for the change from quantum to classical behavior. In this manner we obtain two critical exponents corresponding to quantum and classical percolations. We also study how the 2D extended state develops into the one-dimensional critical state.

DOI: [10.1103/PhysRevB.75.113304](https://doi.org/10.1103/PhysRevB.75.113304)

PACS number(s): 73.43.-f, 31.30.Gs, 73.20.Fz, 73.20.Jc

Celebrated quantum Hall effect (QHE) is realized in a two-dimensional (2D) electron gas subjected to a strong perpendicular magnetic field and a random potential.^{1,2} The uniqueness of this phenomenon is in the high precision of the plateaux in the Hall component and in the very rich physics in the interplateau transitions.³ Here we will study the influence of the nuclear-spin fields⁴ on the critical exponents in QHE.

The rich physics of the random potential in quantum Hall systems could be roughly divided into spin-independent and spin-dependent (spintronics) electron-scattering processes. Magnetic impurities perturb the QHE transport very strongly and will not be considered here. Recently, sharply growing attention was attracted to the physics of the hyperfine interactions in the QHE. It was suggested theoretically⁴ and observed experimentally^{5,6} that the underlying nuclear-spin structure can provide the microscopic information on the 2D electron wave functions and provide strong influence on the precision and other parameters of a QHE system.

The interaction between electron and nuclear spins in heterojunctions under QHE conditions is due, usually, to the *hyperfine* Fermi contact interaction.^{4,7} This interaction is represented by the Hamiltonian:

$$\hat{H}_{int} = -\gamma_n \hbar \vec{I}_i \cdot \vec{H}_e, \quad (1)$$

where γ_n is the nuclear gyromagnetic ratio, \vec{I}_i is the nuclear spin, and \vec{H}_e is the magnetic field on the nuclear site, produced by electron orbital and spin magnetic moments:

$$\vec{H}_e = -g\beta \sum_e \frac{8\pi}{3} \hat{s}_e \delta(\vec{r}_e - \vec{R}_i). \quad (2)$$

Here \vec{r}_e is the electron radius vector, \hat{s}_e is the electron-spin operator, $\beta = e\hbar/m_0c$ is the Bohr magneton, g is the electronic g -factor, and \vec{R}_i is the nucleus radius vector.

It follows from Eqs. (1) and (2) that once the nuclear spins are polarized, i.e., if $\langle \sum_i \vec{I}_i \rangle \neq 0$, the charge-carrier spins feel the effective, time-dependent hyperfine field B_{hf}

$= B_{hf}^0 \exp(-t/T_1)$ (T_1 is a nucleus relaxation time), which lifts the spin degeneracy even in the absence of external magnetic field. In GaAs/AlGaAs, one may achieve the spin splitting due to hyperfine field of the order of the Fermi energy.^{5,6} The inhomogeneous nuclear polarization acts on the electrons as additional (to the scalar potential of the impurities) confining potential $V_{hf} = -\mu_B B_{hf}$.⁸

The nuclear-spin polarization, once created, remains finite for macroscopically long times. Intensive experimental studies^{5,6} of this phenomenon in QHE systems have provided a more detailed knowledge on the hyperfine interaction between the nuclear and electron spins in heterojunctions and quantum wells. It was observed that the nuclear-spin relaxation time is rather long (up to 10^3 s) and the hyperfine field acting on the charge-carrier spins is extremely high, up to 10^4 G.⁵ The nuclear relaxation time depends strongly on the vicinity to the impurity and its sign.³ The presence of the impurity (long-range potential) provides the necessary energy conservation in the spin-exciton creation process leading to the nuclear-spin relaxation. We can, therefore, expect the following scenario: nuclear spins being polarized by some external field will then relax differently depending on whether they are close to maxima or minima of the scalar potential created by impurities. Therefore, they should strongly affect tunneling of electrons through saddle-point potential.

When random potential varies smoothly (its correlation length is much larger than the magnetic length as, e.g., in GaAs heterostructures), a semiclassical description becomes relevant: electrons move along the lines of constant potential. When two equipotential lines come close to each other (near a saddle point), tunneling is feasible. In this Brief Report, we investigate how this picture will be affected by strong nuclear polarization. We find that scaling of the localization length is modified [Eq. (8)], which is the main result of this work.

In the Chalker-Coddington (CC) network model,⁹ electrons move along unidirectional links forming closed loops in analogy to semiclassical motion on contours of constant




FIG. 1. Network model with missing nodes.

potential. Scattering between links is allowed at nodes in order to map tunneling through saddle-point potentials. Propagation along links yields a random phase ϕ ; thus, links are presented by diagonal matrices with elements in the form $\exp(i\phi)$. Transfer matrix for one node relates a pair of incoming and outgoing amplitudes on the left to a corresponding pair on the right; it has the form

$$\mathbf{T} = \begin{pmatrix} \sqrt{1 + \exp(-\pi\epsilon)} & \exp(-\pi\epsilon/2) \\ \exp(-\pi\epsilon/2) & \sqrt{1 + \exp(-\pi\epsilon)} \end{pmatrix}. \quad (3)$$

In order for a system to be invariant, on average, under 90° rotation the transmission and reflection at the next neighbor node are interchanged, i.e., the transfer matrix has the same form as in Eq. (3) with a parameter $\epsilon' = -\epsilon$.⁹ In order to obtain this relation, one simply interchanges Z_3 and Z_4 (see Fig. 1) and brings a transfer matrix to the form of Eq. (3). We therefore describe scattering at the nodes indicated in Fig. 1 by circles with transfer matrix $\mathbf{T}(\epsilon)$ and at the nodes indicated by boxes with $\mathbf{T}(-\epsilon)$.

The node parameter ϵ is the relative distance between the electron energy and the barrier height. It is related to the physical quantities describing the system

$$\epsilon \equiv \left[E - \left(n + \frac{1}{2} \right) E_2 - V_0 \right] / E_1, \quad (4)$$

where E_1 measures the ratio between saddle-point parameters and magnetic field, E_2 is the distance between Landau levels at strong magnetic fields, and V_0 is a reference point of a scalar potential (see Ref. 10 for details). It is easy to see that the most “quantum” case (equal probabilities to scatter to the left and to the right) is at $\epsilon=0$; in fact, numerical calculations⁹ show that there is an extended state at that energy. Numerical simulations on the network model are performed in the following way: one studies a system with fixed width M and periodic boundary conditions in the transverse direction. Multiplying transfer matrices for N slices and then diagonalizing the resulting total transfer matrix, it is possible to extract the smallest Lyapunov exponent λ [the eigenvalues of the transfer matrix are $\exp(\lambda N)$]. The localization length ξ_M is proportional to $1/\lambda$. Repeating calculations for differ-

ent system widths and different energies, it is possible to show that the localization length ξ_M satisfies a scaling relation

$$\frac{\xi_M}{M} = f\left(\frac{M}{\xi(\epsilon)}\right). \quad (5)$$

In the QHE, the thermodynamic localization length $\xi(\epsilon) \sim |\epsilon|^{-\nu}$ and $\nu=2.5 \pm 0.5$. This is the main result,⁹ and it is in good agreement with experimental data for spin-split resolved levels,¹¹ numerical simulations using other models,¹² and semiclassical argument^{13,14} that predicts $\nu=7/3$.

It is possible to model classical percolation using CC model as well. It was shown¹⁵ that when the relative height of the barriers fluctuate in the infinite range, the percolation becomes classical (no tunneling is allowed) and classical percolation exponent $\nu_{cl}=4/3$ is retrieved. On the other hand, when the fluctuations are finite, their width acts as irrelevant parameter^{16–18} and does not affect ν .

In the present work, we modify CC model in the following way. We expect that the presence of a polarized nucleus near a saddle point of the scalar potential will modify a tunneling parameter ϵ in Eq. (3) by changing V_0 to $V_0 \pm V_{hf}$. Moreover, we also expect that due to different relaxation rates (in the vicinity of impurities of different signs), the following scenario can be realized: nuclei situated near different types of saddle point (nodes of the model) will be polarized in opposite directions, breaking, therefore, the isotropy of the system. We model this situation by introducing a parameter $0 \leq p < 1$ describing the probability that there is a polarized nucleus near a particular saddle point. Due to the effect of high hyperfine fields described above, as a rough approximation, we can expect that the barrier becomes “infinite,” i.e., the transfer matrix at the node is now a unit matrix. On the model language, it means that the quasiparticle stays on the same horizontal link (see Fig. 1) and the isotropy of the model is therefore broken. Obviously, when $p=1$, a 2D system is broken into M one-dimensional chains, and, due to the fact that there is no backscattering, all states are extended independent of energy ϵ and system width M . We, therefore, expect the smallest Lyapunov exponent $\lambda=0$, in contradistinction to the “ordinary” 2D extended state, where λ is finite, and infinite thermodynamic localization length is recovered only after finite-size scaling. In this sense, $p=1$ case is close to a one-dimensional (1D) metal found for a dirty superconductor with broken time-reversal and spin-rotational symmetries.¹⁹

Before we present numerical results, let us discuss the possible form for the scaling of the renormalized localization length. Now, when we have “wiped out,” on average, a fraction p of the nodes, a quasiparticle should travel a larger distance [times $1/(1-p)$] in order to experience the same number of scattering events. Therefore, naively, one would expect that the effective system width is now $M(1-p)^{-1}$ and the scaling is

$$\frac{\xi_M}{M} = (1-p)^{-1} f\left(\frac{M}{\xi(\epsilon)}\right). \quad (6)$$




FIG. 2. Renormalized localization length at critical energy $\epsilon=0$ as function of the fraction of missing nodes p for different system widths. The solid line is the best fit $1.24(1-p)^{-1.3}$. The dashed line is the fit with “naive” exponent $\nu=1$.

On the other hand, we should take into account that the “missing” node actually does not allow the quasiparticle to propagate in the transverse direction (we have chosen the system in such a way that, if there is no scattering, the quasiparticle stays on the same *horizontal* link). Usually for CC model and its generalizations, the typical value of the renormalized localization length for the extended state is of the order of 1, meaning that in the extended state the quasiparticle is able to traverse the system of width M . Therefore, in the present situation, we could expect even larger value of ξ_M in the extended state, i.e., $(1-p)^{-\nu}$ dependence with $\nu > 1$.

In order to find both critical exponents, we start by studying a p dependence for $\epsilon=0$, corresponding to the development of a 2D extended state into a 1D extended state. The results for system widths $M=16, 32, 64$ are presented in Fig. 2, allowing the following fit:

$$\frac{\xi_M}{M} = (1-p)^{-1.3} f(0), \quad (7)$$

where $f(0)$ is the value of the renormalized localization length in the extended state $\epsilon=0$ for the standard CC model ($p=0$). This value for the critical exponent is suspiciously close to the classical percolation exponent $\nu_{cl}=4/3$. We also show visibly worse fit of the data with the “naive” critical exponent $\nu=1$.

We next use the value ν found in Fig. 2 and study numerically renormalized localization length for various $\epsilon \neq 0$ and $p < 1$. All our data collapse on one curve with abscissa in the form $M/\xi(\epsilon)$, where thermodynamic localization length




FIG. 3. Data collapse for all energies ϵ , system widths M , and all fractions $p \neq 1$ of missing nodes.

diverges as $\xi \sim \epsilon^{-\nu_q}$ with quantum percolation exponent $\nu_q \approx 2.5$. The results of the scaling are presented in Fig. 3.

We argue that one can understand the appearance of the classical percolation exponent in Eq. (7) by considering a quasiparticle on the standard CC model deeply into the localized regime. In this case, localization length ξ_M is M independent, meaning that a quasiparticle does not “feel” the boundaries of the system, and its thermodynamic localization length $\xi = \xi_M$. Therefore, a quasiparticle travels on the perimeter of the classical cluster of the typical size ξ . Then by increasing the fraction p of the missing nodes, we increase the size of the classical cluster, actually making it infinite as p approaches 1. Therefore, $(1-p)$ acts as energy in the classical percolation problem, explaining the value $1.3 \approx 4/3$.

Finally, all the numerical data we have obtained support the following scaling relation:

$$\frac{\xi_M}{M} = (1-p)^{-\nu_{cl}} f(M\epsilon^{\nu_q}), \quad (8)$$

We stress that this result produces both quantum and classical percolation exponents from the same data. To summarize, we have studied the influence of nuclear spins on the localization-delocalization transition in quantum Hall systems. We have found that the fraction p of polarized nuclei acts as a relevant parameter, leading to a different scaling relation for the localization length [Eq. (8)].

One of us (V. K.) appreciates valuable discussions with Ferdinand Evers, Alexander Mirlin, Yuval Gefen, Baruch Horowitz, and Yshai Avishai. I.V. is grateful to Mike Pepper, Cavendish, for hospitality during the summer of 2005, when part of this paper was accomplished. I.V. acknowledges the EuroMagNET of FP6, RII3-CT-2004-506-239.

*Deceased.

- ¹K. v. Klitzing, G. Dorda, and M. Pepper, *Phys. Rev. Lett.* **45**, 494 (1980); D. C. Tsui, H. L. Störmer, and A. C. Gossard, *ibid.* **48**, 1559 (1982); R. B. Laughlin, *ibid.* **50**, 1395 (1983).
- ²R. Prange and S. Girvin, *The Quantum Hall Effect* (Springer, Berlin, 1983); *Perspectives in Quantum Hall Effect*, edited by S. Das Sarma and A. P. Pinczuk (Wiley, New York, 1997).
- ³S. V. Iordansky, *Solid State Commun.* **43**, 1 (1982); S. Luryi and R. F. Kazarinov, *Phys. Rev. B* **27**, 1386 (1983); R. Joint, *J. Phys. C* **18**, L331 (1985); I. D. Vagner and M. Pepper, *Phys. Rev. B* **37**, 7147 (1988).
- ⁴I. D. Vagner and T. Maniv, *Phys. Rev. Lett.* **61**, 1400 (1988); *Physica B* **204**, 141 (1995); for a recent review see I. D. Vagner, *J. Sci. Eng.* 1565-4990 **1**, 152 (2004).
- ⁵A. Berg, M. Dobers, R. R. Gerhardt, and K. V. Klitzing, *Phys. Rev. Lett.* **64**, 2563 (1990); J. N. Smet, R. A. Deutschmann, F. Ertl, W. Wegscheider, G. Abstreiter, and K. von Klitzing, *Nature (London)* **415**, 281 (2002).
- ⁶Go Yusa, Koji Muraki, Key Takashima, Katsushi Nashimoto, and Yoshiro Kirayama, *Nature (London)* **430**, 439 (2004); G. Yusa, K. Hashimoto, K. Muraki, T. Saku, and Y. Hirayama, *Phys. Rev. B* **69**, 161302(R) (2004); M. Xiao, I. Martin, Eli Yablonovich, and H. W. Jiang, *Nature (London)* **430**, 435 (2004).
- ⁷A. Abragam, *The Principles of Nuclear Magnetism* (Clarendon, Oxford, 1961); C. P. Slichter, *Principles of Magnetic Resonance*, 3rd ed. (Springer Verlag, Berlin, 1991).
- ⁸Yu. V. Pershin, S. N. Shevchenko, I. D. Vagner, and P. Wyder, *Phys. Rev. B* **66**, 035303 (2002).
- ⁹J. T. Chalker and P. D. Coddington, *J. Phys. C* **21**, 2665 (1988).
- ¹⁰H. A. Fertig and B. I. Halperin, *Phys. Rev. B* **36**, 7969 (1987).
- ¹¹S. Koch, R. J. Haug, K. v. Klitzing, and K. Ploog, *Phys. Rev. Lett.* **67**, 883 (1991).
- ¹²B. Huckestein and B. Kramer, *Phys. Rev. Lett.* **64**, 1437 (1990).
- ¹³G. V. Milnikov and I. M. Sokolov, *JETP Lett.* **48**, 536 (1988).
- ¹⁴V. Kagalovsky, B. Horovitz, and Y. Avishai, *Europhys. Lett.* **31**, 425 (1995).
- ¹⁵D. H. Lee, Z. Wang, and S. Kivelson, *Phys. Rev. Lett.* **70**, 4130 (1993).
- ¹⁶B. Huckestein, *Phys. Rev. Lett.* **72**, 1080 (1994).
- ¹⁷J. T. Chalker and J. F. G. Eastmond (unpublished); J. F. G. Eastmond, Ph.D. thesis, Oxford University, 1992.
- ¹⁸V. Kagalovsky, Ph.D. thesis, Ben Gurion University, 1995.
- ¹⁹J. T. Chalker, N. Read, V. Kagalovsky, B. Horovitz, Y. Avishai, and A. W. W. Ludwig, *Phys. Rev. B* **65**, 012506 (2002).