

Andreev-Lifshitz supersolid hydrodynamics including the diffusive mode

Matthew R. Sears and Wayne M. Saslow*

Department of Physics, Texas A&M University, College Station, Texas 77843-4242, USA

(Received 20 July 2010; revised manuscript received 1 September 2010; published 19 October 2010)

We have reexamined the Andreev-Lifshitz theory of supersolids. This theory implicitly neglects uniform bulk processes that change the vacancy number and assumes an internal pressure P in addition to lattice stress λ_{ik} . Each of P and λ_{ik} takes up a part of an external, or applied, pressure P_a (necessary for solid ^4He). The theory gives four pairs of propagating elastic modes, of which one corresponds to a fourth-sound mode, and a single diffusive mode, which has not been analyzed previously. The diffusive mode has three distinct velocities, with the superfluid velocity much larger than the normal fluid velocity, which in turn is much larger than the lattice velocity. The mode structure depends on the relative values of certain kinetic coefficients and thermodynamic derivatives. We consider pressurization experiments in solid ^4He at low temperatures in light of this diffusion mode and a previous analysis of modes in a normal solid with no superfluid component.

DOI: [10.1103/PhysRevB.82.134523](https://doi.org/10.1103/PhysRevB.82.134523)

PACS number(s): 67.80.bd, 67.80.dj, 05.70.Ln

I. INTRODUCTION

Since the late 1960s there have been theoretical suggestions that solids might display flow behavior similar to what is found in superfluids.¹⁻⁴ For that reason there has been a great deal of interest in solid ^4He as a candidate to be a *supersolid*.⁵ The first experimental indication of superflow was the appearance of a nonclassical moment of inertia (NCRI), first observed by Chan's group, since confirmed by many other laboratories, and strongly linked to disorder.⁶⁻¹⁵ In addition, the shear modulus shows anomalous behavior,¹⁶ although not enough to explain the NCRI experiments.¹⁷ Non-NCRI superflow has been searched for but not observed.¹⁸ Evidence is growing that restricts the possible temperature range over which supersolidity can occur.¹⁹ Moreover, for NCRI experiments with rim velocity v at temperature T , the observed hysteresis in v - T space suggests multiple apparent phase transitions.^{10,20,21}

A recent experiment on a pancake-shaped sample, where a pressure change is applied to one side, finds an exponential decay with time of the pressure response on the opposite side.²² The response is slower at lower temperatures, rather than saturating as for a quantum transition, perhaps an indication that the system is *not* supersolid. We have recently studied the lattice diffusion mode of a normal solid (see Ref. 23), based on equations obtained by eliminating the superfluid velocity \vec{v}_s from the theory of Andreev and Lifshitz.¹ We obtained both the diffusion constant and the eigenmode structure (by which we mean the ratios of the deviations from equilibrium of various thermodynamic quantities) for a solid under an externally applied pressure P_a (necessary to solidify ^4He , even at $T=0$).

Whereas an ordinary solid has eight degrees of freedom,²³ the addition of \vec{v}_s (the gradient of a phase) gives a supersolid nine degrees of freedom. For a plane wave, where k_i is the wave vector with magnitude k , the degrees of freedom are given by two scalar thermodynamic quantities (which can be taken to be the mass density ρ and the entropy density s), the lattice vector u_i , the normal fluid velocity vector v_{ni} , and $v_s = k_i v_{si}/k$, where v_{si} is the superfluid velocity. The scalar quantity v_s has been defined because v_{si} is expressible as the

gradient of a phase ϕ . The total momentum density is thus given by $g_i = \rho_n v_{ni} + \rho_s v_{si}$, where ρ_n and ρ_s are the respective densities of the normal and superfluid components. (In principle, both ρ_n and ρ_s are tensors but calculations for hcp ^4He indicate that they are nearly isotropic.^{24,25}) The nine degrees of freedom imply there are nine normal modes. For a uniform infinite system these modes are four pairs of propagating elastic waves (previously studied for both zero^{1,26-28} and nonzero²⁹ P_a), with frequency $\omega \sim k$ and a diffusive mode, with $\omega \sim ik^2$ (whose structure has not been previously studied).

The present work studies this diffusive mode that occurs in a supersolid when \vec{v}_s is included. We employ a variation on the notation of Ref. 26, which gives a more explicit derivation of the equations of motion than does Ref. 1, and extends Ref. 1 to include nonlinear terms.³⁰ References 1 and 26 implicitly assume that uniform vacancy-number-changing bulk processes are negligible.

The Andreev-Lifshitz (AL) theory is remarkable in that it assigns an internal pressure P , in addition to lattice stress λ_{ik} , to a supersolid, in order to continuously go to the superfluid limit. Each of P and λ_{ik} take up part of P_a . Reference 23 finds for a solid, by thermodynamic considerations, the dependence of P on P_a . The consequences of distinct P , λ_{ik} , and P_a had not previously been considered. Reference 23 calculates the effect of P_a on the propagating elastic and diffusive modes of an ordinary solid. Reference 29 calculates the effect of P_a on the propagating elastic modes of a supersolid, as well as the efficiency with which a heater or a transducer generates these modes. The present work considers the effect of P_a on the diffusive mode of a supersolid.

As for the lattice diffusion mode for the normal solid, the diffusive mode for the supersolid is characterized not by the diffusion of a single thermodynamic variable but by specific amounts of each, determined by the eigenmode structure. A dissipative term in the equation of motion for the lattice displacement permits the lattice velocity to differ from \vec{v}_n . We find the relationships between the normal, superfluid, and lattice velocities in this mode. From the lattice velocity one can obtain the lattice displacement and lattice strain deviation. Because the mode is diffusive, the rate of change in the

momentum density, and thus the total stress deviation, are nearly zero, so lattice stress deviations must be canceled by an opposing pressure deviation P' , thus determining P' . Again because the mode is diffusive, the rate of change in \vec{v}_s , and thus the total chemical-potential deviation, are nearly zero, so the P' must be compensated by an opposing contribution due to a temperature deviation T' . This diffusion mode is therefore characterized by its diffusion constant and specific ratios of the normal and superfluid velocities, and the temperature and pressure, relative to the lattice velocity. In practice we use the entropy and mass densities rather than temperature and pressure. The theory permits vacancies to diffuse but there are no bulk sources or sinks for them.

Section II gives the AL supersolid theory in our notation. Section III derives the normal modes for the supersolid. Section IV provides a summary. Appendix estimates the sizes of several quantities relevant to the diffusive mode.

II. ANDREEV-LIFSHITZ SUPERSOLID

In what follows we employ the primary quantities energy density ϵ , lattice displacement u_i , and nonsymmetrized strain $w_{ik} = \partial_i u_k$.

A. Thermodynamics

The thermodynamic equations for a supersolid are given by

$$d\epsilon = Tds + \lambda_{ik} dw_{ik} + \mu d\rho + \vec{v}_n \cdot d\vec{g} + \vec{j}_s \cdot d\vec{v}_s, \quad (1)$$

$$\epsilon = -P + Ts + \lambda_{ik} w_{ik} + \mu\rho + \vec{v}_n \cdot \vec{g} + \vec{j}_s \cdot \vec{v}_s, \quad (2)$$

$$0 = -dP + sdT + w_{ik} d\lambda_{ik} + \rho d\mu + \vec{g} \cdot d\vec{v}_n + \vec{v}_s \cdot d\vec{j}_s. \quad (3)$$

Here λ_{ik} is an elastic tensor density (with the same units as pressure P), μ is the chemical potential (with units of velocity squared),

$$\vec{g} = \rho_n \vec{v}_n + \rho_s \vec{v}_s \quad (4)$$

is the momentum density, and

$$\vec{j}_s = \vec{g} - \rho \vec{v}_n = \rho_s (\vec{v}_s - \vec{v}_n) \quad (5)$$

is a momentum density defined so that $d\epsilon = \vec{g} \cdot d(\delta\vec{v})$ under a Galilean boost $\delta\vec{v}$. Since $(\vec{g}, \vec{v}_n, \vec{v}_s)$ are all vectors under Galilean boosts, we deduce that $\rho_n + \rho_s = \rho$.

We find it convenient to define

$$\vec{j}_n \equiv \rho \vec{v}_n \quad (6)$$

so that

$$\vec{g} = \vec{j}_n + \vec{j}_s. \quad (7)$$

Unlike \vec{j}_s , the quantity \vec{j}_n is a momentum density both in units and in its properties under Galilean boosts.

B. Dynamics

The linearized equations of motion relevant to obtaining the normal modes, considering only the independent variables s , u_i , ρ , v_{ni} , and v_{si} , are

$$\partial_t s + \partial_j f_j = \frac{R}{T} \quad (R \geq 0), \quad (8)$$

$$\partial_t u_i = U_i, \quad (9)$$

$$\partial_t \rho + \partial_i g_i = 0, \quad (10)$$

$$\partial_t g_i + \partial_k \Pi_{ik} = 0, \quad (11)$$

$$\partial_t v_{si} + \partial_i \theta = 0. \quad (12)$$

Here, the fluxes f_i , Π_{ik} , g_i , θ , and the ‘‘source’’ U_i are given by

$$f_i = s v_{ni} - \frac{\kappa_{ij}}{T} \partial_j T - \frac{\alpha_{ij}}{T} \partial_l \lambda_{lj}, \quad (13)$$

$$U_i = v_{ni} + \frac{\alpha_{ij}}{T} \partial_j T + \beta_{ij} \partial_l \lambda_{lj}, \quad (14)$$

$$\Pi_{ik} = (P \delta_{ik} - \lambda_{ki}) - \eta_{iklm} \partial_m v_{nl} - \zeta_{ik} \partial_{lj} s, \quad (15)$$

$$\theta = \mu - \zeta_{ik} \partial_k v_{ni} - \chi \partial_{kj} s, \quad (16)$$

$$g_i = \rho v_{ni} + j_{si}, \quad (17)$$

and we take $R \approx 0$, as it is second order in deviations. AL use both $\sigma_{ik} \approx -\Pi_{ik}$ (a notation we employ below) and $j_i = g_i$. The term in Eq. (14) proportional to β_{ij} allows the lattice velocity \dot{u}_i to differ from the velocity v_{ni} associated with mass flow.

Recall that a diffusion constant D is proportional to a characteristic velocity times a characteristic mean-free path, so it has units of m^2/s . In terms of a D , the dissipative coefficients have the following units: κ_{ij} has units of s times D ; α_{ij} has units of D ; β_{ij} has units of inverse pressure times D ; η_{iklm} has units of ρ times D ; ζ_{ik} has units of D ; and χ has units of inverse density times D .

III. NORMAL MODES IN A SUPERSOLID

As noted earlier, this system has nine variables: s , ρ , u_i , v_{ni} , and v_{si} . With deviations from equilibrium denoted by primes, we use the nine variables s' , ρ' , u'_i ,

$$g'_i \approx \rho_n v'_{ni} + \rho_s v'_{si}, \quad (18)$$

and

$$v'_s = \frac{k_i v'_{si}}{k}. \quad (19)$$

As noted above, there correspondingly are nine normal modes. For an infinite system we assume a disturbance of the form $\exp[i(\vec{k} \cdot \vec{r} - \omega t)]$, where the real wave vector \vec{k} is considered to be known but ω is unknown. For the disturbance to decay in time, $\text{Im}(\omega) < 0$. We find that six modes come in three degenerate pairs, with g'_i and u'_i strongly coupled, and correspond to ordinary elasticity. Two other modes also form a degenerate pair, corresponding to fourth sound, with the

superfluid component in motion and the normal component essentially at rest.^{29,31,32} The ninth and final modes are diffusive, with v'_n and v'_s in opposing directions, and nearly constant chemical potential and stress.

We consider the (off-diagonal) temperature-lattice transport coefficient $\alpha_{ij}=0$, and set to zero the thermal-expansion coefficient. We also neglect the viscosities η_{iklm} , ζ_{ik} , and χ , which to lowest order do not contribute to the modes.³³ We consider an isotropic solid, for which $\kappa_{ij}=\kappa\delta_{ij}$ and $\beta_{ij}=\beta\delta_{ij}$. Unless otherwise specified, thermodynamic derivatives with respect to ρ , s , or w_{ik} are taken with the other two variables held constant.

A. Elastic modes

The elastic modes are obtained by neglecting dissipative and nonlinear terms in Eqs. (8)–(12). Although the elastic modes of a supersolid had previously been found for $P_a=0$,^{1,26,27} Ref. 29 explicitly finds the elastic modes for nonzero P_a (recall that a $P_a \gtrsim 25$ bar is necessary to solidify ⁴He). A summary of the results and convenient notation are provided here.

For the isotropic case, we define

$$\frac{\partial P}{\partial w_{ik}} \equiv \frac{\partial P}{\partial w} \delta_{ik}, \quad (20)$$

$$\frac{\partial \lambda_{ik}}{\partial \rho} \equiv \frac{\partial \lambda}{\partial \rho} \delta_{ik}, \quad (21)$$

$$\frac{\partial \lambda}{\partial w} \equiv K + \frac{4}{3} \mu_V. \quad (22)$$

In this case the static value of the strain (dependent on the applied pressure) is isotropic,^{23,34}

$$w_{ik}^{(0)} = \frac{w_{ll}^{(0)}}{3} \delta_{ik} \approx -\frac{P_a}{3K} \delta_{ik}. \quad (23)$$

1. Longitudinal elastic modes

For $\vec{k} \cdot \vec{j}_n \neq 0 \neq \vec{k} \cdot \vec{j}_s$ and $\vec{k} \times \vec{v}_n = 0$, there are two degenerate pairs of solutions to the equations of motion, a pair that corresponds to first sound and a pair that corresponds to fourth sound. With

$$f_s \equiv \frac{\rho_s}{\rho}, \quad (24)$$

to first order in f_s , first sound frequencies are given by

$$\frac{\omega_1^2}{k^2} = c_1^2 + f_s \left[c_1^2 - 2\tilde{c}^2 + \frac{\tilde{c}^4}{c_1^2} + w_{ll}^{(0)} \frac{\partial \lambda}{\partial \rho} \left(\frac{\tilde{c}^2}{c_1^2} - 1 \right) \right] \quad (25)$$

and fourth sound frequencies are given by

$$\frac{\omega_4^2}{k^2} = f_s \left(c_0^2 - \frac{\tilde{c}^4}{c_1^2} - w_{ll}^{(0)} \frac{\partial \lambda}{\partial \rho} \frac{\tilde{c}^2}{c_1^2} \right). \quad (26)$$

Here, the velocities c_0 , c_1 , and \tilde{c} satisfy

$$c_0^2 \equiv \rho \frac{\partial \mu}{\partial \rho}, \quad (27)$$

$$c_1^2 \equiv \frac{\partial P}{\partial \rho} - \frac{\partial \lambda}{\partial \rho} + \frac{1}{\rho} \left(\frac{\partial \lambda}{\partial w} - \frac{\partial P}{\partial w} \right), \quad (28)$$

$$\tilde{c}^2 \equiv c_0^2 - \frac{\partial \lambda}{\partial \rho}. \quad (29)$$

If σ rather than s were held constant, c_0 would be the sound velocity in an ordinary (nonsuper)liquid, and c_1 would be the sound velocity in an ordinary solid.²³ Reference 29 shows that for $P_a \ll K$ we have $c_1^2 \gg \tilde{c}^2 \gg c_0^2$ and strain $w_{ll}^{(0)} \ll 1$. It is also convenient to define the “fluidlike” and “solidlike” velocities c_{lL} and c_{lS} , which satisfy²³

$$c_{lL}^2 \equiv \frac{\partial P}{\partial \rho} - \frac{\partial \lambda}{\partial \rho}, \quad c_{lS}^2 \equiv \frac{1}{\rho} \left(\frac{\partial \lambda}{\partial w} - \frac{\partial P}{\partial w} \right) \quad (30)$$

so that

$$c_1^2 = c_{lL}^2 + c_{lS}^2. \quad (31)$$

For an ordinary solid, the derivatives in Eq. (30) are taken at constant σ rather than s .

2. Transverse elastic modes

For $\vec{k} \cdot \vec{j}_n = 0 = \vec{k} \cdot \vec{j}_s$ and $\vec{k} \times \vec{v}_n \neq 0$, there are two degenerate pairs of elastic modes. They each have a frequency satisfying

$$\omega_t = k \sqrt{\frac{\mu_V}{\rho_n}}, \quad (32)$$

which is larger than the ordinary (nonsuper)solid transverse frequency by the factor $\sqrt{\rho/\rho_n}$. Such an effect, to our knowledge, has not been observed.

B. Diffusive mode

For the diffusive mode, we keep the dissipative terms in the equations of motion, Eqs. (8)–(12), so that $\dot{u}'_i \neq v'_{ni}$. With $w'_{jl} = ik_j u'_l$, rewriting Eqs. (8)–(12) in terms of the variables v'_{ni} , v'_{si} , ρ' , s' , and u'_i gives

$$\omega s' = k_i s v'_{ni} - ik^2 \frac{\kappa}{T} \left(\frac{\partial T}{\partial s} s' + \frac{\partial T}{\partial \rho} \rho' + \frac{\partial T}{\partial w_{jl}} ik_j u'_l \right), \quad (33)$$

$$\omega u'_i = i v'_{ni} - \beta k_k \left(\frac{\partial \lambda_{ki}}{\partial s} s' + \frac{\partial \lambda_{ki}}{\partial \rho} \rho' + \frac{\partial \lambda_{ki}}{\partial w_{jl}} ik_j u'_l \right), \quad (34)$$

$$\omega \rho' = k_i g'_i = k_i (\rho_n v'_{ni} + \rho_s v'_{si}), \quad (35)$$

$$\begin{aligned} \omega g'_i = & -k_k \sigma'_{ik} = k_k \left[\left(\frac{\partial P}{\partial s} \delta_{ik} - \frac{\partial \lambda_{ik}}{\partial s} \right) s' + \left(\frac{\partial P}{\partial \rho} \delta_{ik} - \frac{\partial \lambda_{ik}}{\partial \rho} \right) \rho' \right. \\ & \left. + \left(\frac{\partial P}{\partial w_{jl}} \delta_{ik} - \frac{\partial \lambda_{ik}}{\partial w_{jl}} \right) ik_j u'_l \right], \end{aligned} \quad (36)$$

$$\omega v'_{si} = k_i \mu' = k_i \left(\frac{\partial \mu}{\partial s} s' + \frac{\partial \mu}{\partial \rho} \rho' + \frac{\partial \mu}{\partial w_{jl}} i k_j u'_l \right). \quad (37)$$

Recall that we have neglected the viscosity as a higher order effect in k^2 as $k \rightarrow 0$. We assume that

$$\omega = -iD_D k^2, \quad (38)$$

where the diffusion constant $D_D > 0$ is to be determined.

At first sight this system promises to yield a quintic in ω , associated with the longitudinal modes. However, the assumption that there is a diffusive mode (whose consistency we must verify) permits us to reduce this to a single linear equation. In some sense a single diffusive mode is expected because we have already obtained four pairs of propagating modes. We detail our procedure because it both illuminates the physics and clarifies the mathematics.

(1) *Method of solution.* Since we take the long-wavelength limit, we neglect terms that are higher order in k . In the present analysis we are merely interested in an order of magnitude estimation so we drop subscripts. When later solving for the frequency and mode structure we use appropriate subscripts.

When written in terms of powers of k [using Eq. (38)], mass and momentum conservation [Eqs. (35) and (36)] imply that

$$k^2 \rho' \sim k g', \quad (39)$$

$$k^2 g' \sim k \tilde{\sigma}'. \quad (40)$$

Here we use $\tilde{\sigma}$ to distinguish a stress from σ , the entropy/mass. Combination of Eqs. (39) and (40) yields $k^2 \rho' \sim \tilde{\sigma}'$. Since expanding $\tilde{\sigma}'$ in terms of the other variables gives a term proportional to ρ' , for small k the term $k^2 \rho'$ is negligible, so $\tilde{\sigma}' \rightarrow 0$ as $k \rightarrow 0$. The diffusive mode therefore is characterized by a negligible stress deviation. Physically this means that the fluidlike stress deviation nearly cancels the solidlike stress deviation. When $\tilde{\sigma}'$ is expanded in terms of the other variables, the condition $\tilde{\sigma}' \approx 0$ provides a relationship between s' , ρ' , and ku' .

We now turn to the superfluid, Eq. (37), which gives

$$k^2 v'_s \sim k \mu'. \quad (41)$$

We now assume that $\mu' \rightarrow 0$ as $k \rightarrow 0$, to be verified below. When μ' is expanded in terms of the other variables, the condition $\mu' \approx 0$ provides a second relationship between s' , ρ' , and ku' . In the remaining equations, for s' and u' , we choose to eliminate ρ' and u' in favor of s' .

Neither of the equations for s' nor u' [Eqs. (33) and (34)] involves v'_s . Hence, on eliminating ρ' and u' in favor of s' , Eqs. (33) and (34) involve s' and v'_n , as well as the unknown ω . This leaves us with two linear equations for two unknowns: the ratio of v'_n to s' , and ω . Once these are determined, we use conservation of mass to relate the still unknown v'_s to v'_n and ρ' , both of which having been found in terms of s' . We find that at low temperatures $\omega \rho'$ can be neglected relative to kv'_n so that Eq. (35) gives $0 \approx g' = \rho_n v'_n + \rho_s v'_s$. This is not a result of an analysis in powers of k as $k \rightarrow 0$ but rather from relations between various thermodynamic quantities.

In what follows, several Maxwell relations from Eq. (1) are used,

$$\frac{\partial \mu}{\partial s} = \frac{\partial T}{\partial \rho}, \quad \frac{\partial \lambda_{ik}}{\partial s} = \frac{\partial T}{\partial w_{ik}}, \quad \frac{\partial \lambda_{ik}}{\partial \rho} = \frac{\partial \mu}{\partial w_{ik}}. \quad (42)$$

Further, Ref. 34 gives, for the elastic stress,

$$\lambda_{ik} = \left(\frac{\partial \lambda}{\partial w} - 2\mu_V \right) \delta_{ik} w_{ll} + \mu_V (w_{ik} + w_{ki}), \quad (43)$$

where $\partial \lambda / \partial w$ is defined in Eq. (22). Since, as in Eq. (23), the static strain is isotropic (i.e., $w_{ik}^{(0)} \sim \delta_{ik}$), Eq. (43) implies that the static elastic stress also is isotropic (i.e., $\lambda_{ik}^{(0)} \sim \delta_{ik}$). Thus $(\partial \lambda_{ik} / \partial \rho)_{w_{jl}}$ and $(\partial \lambda_{ik} / \partial s)_{w_{jl}}$ also are isotropic, which permits us to define

$$\frac{\partial T}{\partial w_{ik}} = \frac{\partial \lambda_{ik}}{\partial s} \equiv \frac{\partial \lambda}{\partial s} \delta_{ik}, \quad \frac{\partial \mu}{\partial w_{ik}} = \frac{\partial \lambda_{ik}}{\partial \rho} \equiv \frac{\partial \lambda}{\partial \rho} \delta_{ik}. \quad (44)$$

Note that Eq. (43) gives

$$\frac{\partial \lambda_{ik}}{\partial w_{jl}} = \left(\frac{\partial \lambda}{\partial w} - 2\mu_V \right) \delta_{ik} \delta_{jl} + \mu_V (\delta_{ij} \delta_{kl} + \delta_{kj} \delta_{il}). \quad (45)$$

(2) *Rewriting stress equation.* For the isotropic case, using Eqs. (20), (21), (30), (42), (44), and (45), Eq. (36) gives, for negligible total stress,

$$\begin{aligned} 0 \approx & \left(\frac{\partial P}{\partial s} - \frac{\partial \lambda}{\partial s} \right) k_i s' + \left(\frac{\partial P}{\partial \rho} - \frac{\partial \lambda}{\partial \rho} \right) k_i \rho' - \left(\frac{\partial \lambda}{\partial w} - \mu_V \right. \\ & \left. - \frac{\partial P}{\partial w} \right) i k_j k_l u'_l - \mu_V i k^2 u'_i \\ \approx & - \frac{\partial \tilde{\sigma}}{\partial s} k_i s' + c_{LL}^2 k_i \rho' - (\rho c_{IS}^2 - \mu_V) i k_j k_l u'_l - \mu_V i k^2 u'_i, \end{aligned} \quad (46)$$

where we define

$$\frac{\partial \tilde{\sigma}}{\partial s} \equiv \frac{\partial \lambda}{\partial s} - \frac{\partial P}{\partial s}. \quad (47)$$

Since each term of Eq. (46) except the last is proportional to k_i , we have that u'_i is proportional to k_i . Thus, $k_i k_j u'_l = k^2 u'_l$, and Eq. (46) becomes, on taking the dot product with k_i / k^2 and dropping the indices on $k_i u'_i$,

$$0 \approx - \frac{\partial \tilde{\sigma}}{\partial s} s' + c_{LL}^2 \rho' - \rho c_{IS}^2 i k u'. \quad (48)$$

Further, since $u'_i \sim k_i$, substitution of Eqs. (44) and (45) into Eq. (34) gives $v'_{ni} \sim k_i$. Then, since Eq. (37) gives $v'_{si} \sim k_i$, the diffusive mode is purely longitudinal ($v'_{si} \sim v'_{ni} \sim u'_i \sim k_i$), and we therefore drop indices for v'_s , v'_n , and u' dotted with k . Moreover, for $u'_i \sim k_i$, Eq. (45) gives

$$\frac{\partial \lambda_{ki}}{\partial w_{jl}} i k_k k_j u'_l = \frac{\partial \lambda}{\partial w} i k^2 u'_i. \quad (49)$$

(3) *Rewriting μ' equation.* Since we assume that $\mu' \approx 0$, we neglect the left-hand side (LHS) of Eq. (37); this yields

$$0 \approx \frac{\partial \mu}{\partial s} s' + \frac{\partial \mu}{\partial \rho} \rho' + \frac{\partial \mu}{\partial w_{jl}} ik_j u'_l. \quad (50)$$

Substitution from Eqs. (42) and (44) gives

$$0 \approx \frac{\partial T}{\partial \rho} s' + \frac{\partial \mu}{\partial \rho} \rho' + \frac{\partial \lambda}{\partial \rho} iku'. \quad (51)$$

(4) *Combining stress and μ' equations.* Solving Eqs. (48) and (51) for ρ' and u' gives

$$\rho' = \frac{Y_3 \rho s'}{Y_1 s}, \quad -iku' = \frac{Y_2 s'}{Y_1 s}, \quad (52)$$

where we introduce three quantities, each with units of velocity to the fourth power,

$$Y_1 \equiv \frac{\partial \lambda}{\partial \rho} c_{IL}^2 + c_0^2 c_{IS}^2, \quad (53)$$

$$Y_2 \equiv s \frac{\partial T}{\partial \rho} c_{IL}^2 + c_0^2 s \frac{\partial \bar{\sigma}}{\rho \partial s}, \quad (54)$$

$$Y_3 \equiv \frac{s \partial \lambda}{\rho \partial \rho} \frac{\partial \bar{\sigma}}{\partial s} - s \frac{\partial T}{\partial \rho} c_{IS}^2. \quad (55)$$

Here we employ Eq. (27). Equation (52) holds for any $\omega \sim k^2$.

Appendix uses the results of Ref. 23 to estimate the sizes of Y_1 , Y_2 , and Y_3 . With θ_D the Debye temperature, k_B the Boltzmann constant, and m_4 the atomic mass of ${}^4\text{He}$, we find

$$Y_1 \approx -\frac{2P_a^2}{\rho^2}, \quad Y_2 \approx -\frac{24\pi^4 T^3 k_B T P_a}{9 \theta_D^3 m_4 \rho},$$

$$Y_3 \approx -\frac{24\pi^4 T^3 k_B T K}{9 \theta_D^3 m_4 \rho}. \quad (56)$$

Note that Y_1 is independent of T . To evaluate these we take $\theta_D \approx 25$ K,³⁵ $m_4 \approx 6.7 \times 10^{-27}$ kg, $\rho \approx 2 \times 10^3$ kg/m³, $P_a \approx 30$ bar, and $K \approx 300$ bar.²⁹ Further, following evidence that a supersolid phase of ${}^4\text{He}$ can only exist at $T < 55$ mK,¹⁹ we take $T \approx 50$ mK. Then, Eq. (56) yields

$$Y_1 \approx -4.5 \times 10^6 \text{ m}^4/\text{s}^4, \quad Y_2 \approx -3.25 \times 10^{-1} \text{ m}^4/\text{s}^4,$$

$$Y_3 \approx -3.25 \text{ m}^4/\text{s}^4, \quad (57)$$

so that $Y_1 \gg Y_3 \gg Y_2$. This inequality applies for any $T < 55$ mK, and therefore applies at any temperature relevant to supersolid ${}^4\text{He}$ experiments subject to $P_a \ll K$.

(5) *Rewriting s' and u'_i equations.* Substituting Eqs. (44), (49), and (52) into Eq. (33), and into Eq. (34) multiplied by $-iksY_1/Y_2$, yields

$$\omega s' = ksv'_n - ik^2 \frac{\kappa}{T} \left(\frac{\partial T}{\partial s} + \frac{\rho Y_3}{sY_1} \frac{\partial T}{\partial \rho} - \frac{Y_2}{sY_1} \frac{\partial \lambda}{\partial s} \right) s', \quad (58)$$

$$\omega s' = k \frac{sY_1}{Y_2} v'_n + ik^2 \beta \left(\frac{sY_1}{Y_2} \frac{\partial \lambda}{\partial s} + \frac{\rho Y_3}{Y_2} \frac{\partial \lambda}{\partial \rho} - \frac{\partial \lambda}{\partial w} \right) s'. \quad (59)$$

We simplify Eqs. (58) and (59) by the following argument. We take $\partial \lambda / \partial s$ to have the same linear T dependence as $\partial P / \partial s$ in a harmonic solid or $\partial \lambda / \partial s \sim T$. Then, since $s \sim T^3$ and $Y_2, Y_3 \sim T^4$, all terms $\sim \kappa$ in Eq. (58) have the same temperature dependence, and the same is true for all terms $\sim \beta$ in Eq. (59). Thus, since $Y_1 \sim 10^6 \times Y_3$ and $Y_3 \sim 10 \times Y_2$, in the parentheses of Eqs. (58) and (59) the first term dominates. Thus Eqs. (58) and (59) approximately give, on rearranging,

$$\left(\omega + ik^2 \frac{\kappa}{T} \frac{\partial T}{\partial s} \right) s' = ksv'_n, \quad (60)$$

$$\left(\omega - ik^2 \beta \frac{sY_1}{Y_2} \frac{\partial \lambda}{\partial s} \right) s' = k \frac{sY_1}{Y_2} v'_n. \quad (61)$$

Subtracting Eq. (61) from Eq. (60) and dividing by ks yields

$$v'_n = ik \frac{s'}{s} \left(\frac{1}{T} \frac{\partial T}{\partial s} \right) \left(\kappa + \frac{Y_1}{Y_2} \beta s T \frac{\partial \lambda}{\partial T} \right), \quad (62)$$

which holds for any $\omega \sim k^2$. Here we use $(\partial \lambda / \partial s) / (\partial T / \partial s) = \partial \lambda / \partial T$, where ρ and w_{ik} are implicitly held constant for each derivative. Hence, Eqs. (52) and (62) show the ratios of (ρ' , ku' , and v'_n) to s' to be frequency independent. We now find the frequency of the diffusive mode using Eqs. (60) and (61). Mass conservation from (35) then relates v'_s and s' , thus yielding all variables in terms of s' .

1. Diffusive mode frequency

Crossmultiplication of Eqs. (60) and (61) yields

$$\omega + ik^2 \frac{\kappa}{T} \frac{\partial T}{\partial s} = \frac{Y_2}{Y_1} \omega - ik^2 \beta s \frac{\partial \lambda}{\partial s}. \quad (63)$$

The frequency of the diffusive mode thus is

$$\omega = -ik^2 \frac{\bar{\kappa}}{T} \frac{\partial T}{\partial s}, \quad (64)$$

where

$$\bar{\kappa} \equiv \left[\frac{\kappa + \beta s T \frac{\partial \lambda / \partial s}{\partial T / \partial s}}{1 - \frac{Y_2}{Y_1}} \right] \approx \kappa + \beta s T \frac{\partial \lambda}{\partial T}. \quad (65)$$

Here we use $Y_1 \gg Y_2$. Recall that $\partial \lambda / \partial T$ is taken at constant ρ and w_{ik} . The frequency thus has a part associated with thermal diffusion ($\sim \kappa$) and a part associated with lattice diffusion ($\sim \beta$).

Before finding the full mode structure, it is worth commenting on Eqs. (64) and (65). If $\beta s T (\partial \lambda / \partial T) \ll \kappa$, then we have $D_D \rightarrow (\kappa / T) (\partial T / \partial s)$, as for ordinary thermal diffusion. As noted above, however, μ is constant in the long-wavelength limit for the diffusive mode of the supersolid (to be verified below), unlike in the case of a fluid or ordinary solid. Thus, even if the frequency were precisely as for nor-

mal thermal diffusion, the mode structure (e.g., v'_n/v'_s , etc.) would nonetheless be different than for the usual case.

2. v'_n , v'_s , and \dot{u}' in the diffusive mode

Equation (35) gives

$$v'_s = \frac{\omega\rho'}{k\rho_s} - \frac{\rho_n v'_n}{\rho_s}. \quad (66)$$

By Eq. (52), the first term on the right-hand side (RHS) of Eq. (66) is given by

$$\frac{\omega\rho'}{k\rho_s} = \frac{\omega\rho s'}{k\rho_s s} \left[\frac{Y_3}{Y_1} \right]. \quad (67)$$

Further, using Eq. (65) and $Y_1 \gg Y_2$, Eq. (62) can be written as

$$\begin{aligned} v'_n &= ik \frac{\bar{\kappa}}{T} \frac{\partial T}{\partial s} \frac{s'}{s} \left[1 + \left(\frac{Y_1}{Y_2} - 1 \right) \frac{\beta s T}{\bar{\kappa}} \frac{\partial \lambda}{\partial T} \right] \\ &\approx - \frac{\omega s'}{k s} \left[1 + \frac{Y_1}{Y_2} \frac{\beta s T}{\bar{\kappa}} \frac{\partial \lambda}{\partial T} \right]. \end{aligned} \quad (68)$$

Thus the second term on the RHS of Eq. (66) is given by

$$- \frac{\rho_n v'_n}{\rho_s} \approx \frac{\omega \rho_n s'}{k \rho_s s} \left[1 + \frac{Y_1}{Y_2} \frac{\beta s T}{\bar{\kappa}} \frac{\partial \lambda}{\partial T} \right]. \quad (69)$$

Since experiments²² indicate that $\rho_n \geq 0.8\rho$, on using $Y_1 \gg Y_3$, Eqs. (67) and (69) give $-(\rho_n/\rho_s)v'_n \gg (\omega\rho'/k\rho_s)$. Equation (66) therefore becomes, on employing Eq. (62),

$$v'_s \approx - \frac{\rho_n}{\rho_s} v'_n = - ik \frac{\rho_n s'}{\rho_s s} \left(\frac{1}{T} \frac{\partial T}{\partial s} \right) \left(\kappa + \frac{Y_1}{Y_2} \beta s T \frac{\partial \lambda}{\partial T} \right) \quad (70)$$

or, equivalently, $g' = \rho_n v'_n + \rho_s v'_s \approx 0$. Thus the superfluid velocity is opposite the normal velocity, with a weighting given by ρ_n/ρ_s . Since $\rho_n/\rho_s \geq 4$ we approximately have $|v'_s| \gg |v'_n|$. Note that Eq. (70) explicitly relates v'_s to s' , thus completely specifying the eigenmode.

We now verify the assumption that $\mu' \rightarrow 0$ for $k \rightarrow 0$. We do so by showing the LHS of Eq. (37) to be negligible compared to any given term on the RHS [e.g., $k^2 v'_s \ll (\partial\mu/\partial s)s'$]. Counting powers of k , Eqs. (62) and (70) give $v'_s \sim v'_n \sim ks'$. Thus $k^2 v'_s \sim k^3 s' \ll (\partial\mu/\partial s)s'$, which shows the consistency of the assumption.

Furthermore, using Eq. (52) to write the lattice velocity \dot{u}' gives

$$\dot{u}' = -i\omega u' = \frac{Y_2 \omega s'}{Y_1 k s}. \quad (71)$$

Comparison to Eq. (68) yields

$$v'_n = -\dot{u}' \frac{Y_1}{Y_2} \left[1 + \frac{Y_1}{Y_2} \frac{\beta s T}{\bar{\kappa}} \frac{\partial \lambda}{\partial T} \right]_{\rho, w_{ik}}. \quad (72)$$

Since $Y_1 \gg Y_2$, unless $[\beta s T (\partial\lambda/\partial T)/\bar{\kappa}] \approx -Y_2/Y_1$ (an unlikely coincidence), we have $v'_n \approx -\dot{u}' (Y_1/Y_2) \gg |\dot{u}'|$. Then, by Eq. (70), for $\rho_n \gg \rho_s$, we have $|v'_s| \gg |v'_n| \gg |\dot{u}'|$. Since $v'_n \neq \dot{u}'$, mass motion is distinct from lattice motion.

IV. SUMMARY

We have reexamined the supersolid hydrodynamics of Andreev and Lifshitz, including the effects of nonzero applied pressure P_a . For $P_a \neq 0$, a solid responds with both lattice stress λ_{ik} and internal pressure P . The dependence of P and λ_{ik} on P_a is found in Ref. 23, and employed here to describe the eigenmodes. We first summarized the results for the four degenerate pairs of longitudinal and transverse elastic mode frequencies (including fourth sound); because we include P_a and the associated strain, the results differ somewhat from those of previous work. In addition, again including P_a and the associated strain, in the long-wavelength limit we have obtained the previously unstudied diffusive eigenmode.

The diffusive mode frequency, under certain conditions, is similar to the frequency of ordinary thermal diffusion. However, the mode involves no deviations in net stress or net chemical potential, so its properties differ from ordinary thermal diffusion. To produce zero net stress deviation, the solidlike elasticity component is canceled by the previously neglected fluidlike component associated with lattice defects. To produce zero net chemical potential deviation, the temperature and pressure deviations must be related. With zero net stress deviation we find that at low temperature there also is zero net momentum. With the normal fluid density dominating the superfluid density, this means that the superfluid velocity is much larger than the normal fluid velocity. Because the lattice displacement is coupled to the elastic strain with a large coefficient but the normal fluid velocity is coupled to the fluidlike strain (a pressure) with a small coefficient, zero net stress deviation implies that the normal fluid velocity is much greater than the lattice velocity. This is an unusual phenomenon since in the other modes the lattice velocity and normal fluid velocity are nearly equal.

A previous work studied the lattice diffusion mode for a normal solid having distinct velocities associated with momentum (e.g., the normal fluid velocity) and lattice elasticity.²³ The motivation was to consider that the time delay in the pressurization experiments of Ref. 22 might be due to that mode, under the assumption that the sample studied is not supersolid. Similar considerations can be made for the diffusive mode we have just studied, because both modes are diffusive in nature, and thus would show a dependence on the sample thickness d as d^2 . A study of this dependence would be of interest, to confirm that the effect observed in Ref. 22 is diffusive in nature.

APPENDIX: RELATIVE SIZES OF Y_1 , Y_2 , AND Y_3 FOR SMALL P_a

Using the results of Refs. 23 and 29 and estimating certain thermodynamic derivatives under the condition $P_a \ll K$ allows us to estimate the sizes of Y_1 , Y_2 , and Y_3 . Reference 23 gives for a normal solid, to lowest order in P_a/K ,

$$w_{ll}^{(0)} = - \frac{P_a}{K}, \quad (A1)$$

$$\frac{\partial \lambda}{\partial \rho} = -c_{IL}^2 = \frac{VP_a}{\rho K} \frac{\partial K}{\partial V} \Big|_{\sigma, w_{ik}, N}, \quad (\text{A2})$$

$$\frac{\partial P}{\partial \rho} = \frac{V^2 P_a^2}{2\rho K^2} \frac{\partial^2 K}{\partial V^2} \Big|_{\sigma, w_{ik}, N}, \quad (\text{A3})$$

$$\frac{\partial P}{\partial w} = -P_a \left(1 - \frac{V}{K} \frac{\partial K}{\partial V} \Big|_{\sigma, w_{ik}, N} \right), \quad (\text{A4})$$

$$c_1^2 = c_{IS}^2 = \frac{\partial \lambda}{\partial w} = \frac{K + \frac{4}{3}\mu_V}{\rho}. \quad (\text{A5})$$

Although Ref. 23 evaluates the derivatives of λ_{ik} and P at constant σ rather than s , at $T=0$ we have $\sigma \approx 0 \approx s$, so holding either quantity constant should give nearly equivalent results for supersolid ^4He . All derivatives of K here are taken at constant σ , w_{ik} , and N , so we now drop the subscripts. Reference 29 also finds

$$c_0^2 \approx \frac{VP_a^2}{\rho K^2} \left(\frac{V}{2} \frac{\partial^2 K}{\partial V^2} + \frac{\partial K}{\partial V} \right). \quad (\text{A6})$$

Constant w_{ik} constant is equivalent to constant density of lattice sites. Because K is a measure of the material stiffness, one expects K to increase as V decreases, for constant N and w_{ik} , i.e., $\partial K/\partial V < 0$. Then by Eqs. (A2) and (A4) we have $\partial \lambda/\partial \rho < 0$ and $\partial P/\partial w < 0$.

For the putative supersolid, we approximate $\partial \tilde{\sigma}/\partial s$ using Eqs. (3), (42), and (44),

$$\frac{\partial \tilde{\sigma}}{\partial s} = \frac{\partial \lambda}{\partial s} - \frac{\partial P}{\partial s} \approx \frac{\partial \lambda}{\partial s} - s \frac{\partial T}{\partial s} - \rho \frac{\partial T}{\partial \rho} - w_{ll}^{(0)} \frac{\partial \lambda}{\partial s}. \quad (\text{A7})$$

Recall that, unless otherwise specified, derivatives with respect to ρ , s , or w_{ik} are taken with the other two held constant. Equation (A1) shows that for $P_a \ll K$, we have $w_{ll}^{(0)} \ll 1$. Also,²³ as noted earlier, $\rho(\partial T/\partial \rho) \approx \gamma s(\partial T/\partial s)$, where $\gamma \approx 10$. Thus,

$$\frac{\partial \tilde{\sigma}}{\partial s} \approx \frac{\partial \lambda}{\partial s} - (1 + \gamma^{-1}) \rho \frac{\partial T}{\partial \rho}. \quad (\text{A8})$$

On neglecting μ_V , Eqs. (43) and (A1) give $\partial \lambda/\partial s \approx (\partial K/\partial s) w_{ll}^{(0)} \approx (P_a/K)(\partial K/\partial s)$.

Substitution of Eqs. (A2), (A5), (A6), and (A8) into Eqs. (53)–(55) gives, to lowest order in P_a/K ,

$$Y_1 = \frac{\partial \lambda}{\partial \rho} c_{IL}^2 + c_0^2 c_{IS}^2$$

$$\approx -\frac{V^2 P_a^2}{\rho^2 K^2} \left[\left(\frac{\partial K}{\partial V} \right)^2 + \left(K + \frac{4}{3}\mu_V \right) \left(\frac{1}{2} \frac{\partial^2 K}{\partial V^2} + \frac{1}{V} \frac{\partial K}{\partial V} \right) \right], \quad (\text{A9})$$

$$Y_2 = s \frac{\partial T}{\partial \rho} c_{IL}^2 + c_0^2 \frac{s}{\rho} \frac{\partial \tilde{\sigma}}{\partial s} \approx -s \frac{\partial T}{\partial \rho} \frac{VP_a}{\rho K} \frac{\partial K}{\partial V}, \quad (\text{A10})$$

$$Y_3 = \frac{s}{\rho} \frac{\partial \lambda}{\partial \rho} \frac{\partial \tilde{\sigma}}{\partial s} - s \frac{\partial T}{\partial \rho} c_{IS}^2 \approx -s \frac{\partial T}{\partial \rho} \frac{K + \frac{4}{3}\mu_V}{\rho}. \quad (\text{A11})$$

Note that all terms $\sim \partial \tilde{\sigma}/\partial s$ are higher order in P_a/K and therefore are neglected. Approximating K to be linear in V and neglecting μ_V , Eqs. (A9)–(A11) give

$$Y_1 \approx -\frac{2P_a^2}{\rho^2}, \quad Y_2 \approx -s \frac{\partial T}{\partial \rho} \frac{P_a}{\rho}, \quad Y_3 \approx -s \frac{\partial T}{\partial \rho} \frac{K}{\rho}. \quad (\text{A12})$$

For $P_a \ll K$ we have $Y_3 \gg Y_2$.

To approximate the relative magnitudes of Y_3 and Y_2 to Y_1 , we now find an explicit form for $s(\partial T/\partial \rho)_s$. At low temperatures phonon gas statistical mechanics gives

$$s = \frac{2\pi^2 k_B^4 T^3}{15\hbar^3 \bar{u}^3}, \quad (\text{A13})$$

where \bar{u} is an average sound velocity and k_B is the Boltzmann constant. Further,²⁹

$$\frac{\partial T}{\partial \rho} \approx \frac{T}{\bar{u}} \frac{\partial \bar{u}}{\partial \rho} \approx \frac{10T}{3\rho}. \quad (\text{A14})$$

Combining Eqs. (A13) and (A14) gives

$$s \frac{\partial T}{\partial \rho} \approx \frac{4\pi^2 k_B^4 T^4}{9\rho \hbar^3 \bar{u}^3}. \quad (\text{A15})$$

In terms of the Debye temperature $\theta_D \approx (6\pi^2 n_v)^{1/3} (\hbar \bar{u}/k_B)$, where n_v is the number density of vibrations (essentially one per lattice site),

$$s \frac{\partial T}{\partial \rho} \approx \frac{24\pi^4 k_B T^4}{9(\rho/n_v)\theta_D^3} \approx \frac{24\pi^4 T^3 k_B T}{9\theta_D^3 m_4}. \quad (\text{A16})$$

Here m_4 is the atomic mass of ^4He , and we have taken $m_4 n_v \approx \rho$. Equation (A16) substituted into Eq. (A12) gives Eq. (56).

*wasaslow@tamu.edu

¹A. F. Andreev and I. M. Lifshitz, Sov. Phys. JETP **29**, 1107 (1969).

²D. J. Thouless, Ann. Phys. (N.Y.) **52**, 403 (1969). This contains the remark that, for a lattice of bosons, vacancies could be “in

the lowest Bloch state with a finite probability, so the system would be ‘super’ but not ‘fluid.’”

³G. V. Chester, Phys. Rev. A **2**, 256 (1970).

⁴A. J. Leggett, Phys. Rev. Lett. **25**, 1543 (1970).

⁵For a recent review, see S. Balibar and F. Caupin, J. Phys.: Con-

- dens. Matter **20**, 173201 (2008).
- ⁶E. Kim and M. Chan, *Nature (London)* **427**, 225 (2004).
- ⁷E. Kim and M. Chan, *Science* **305**, 1941 (2004).
- ⁸A. S. C. Rittner and J. D. Reppy, *Phys. Rev. Lett.* **97**, 165301 (2006).
- ⁹M. Kondo, S. Takada, Y. Shibayama, and K. Shirahama, *J. Low Temp. Phys.* **148**, 695 (2007).
- ¹⁰Y. Aoki, J. C. Graves, and H. Kojima, *Phys. Rev. Lett.* **99**, 015301 (2007).
- ¹¹A. Penzev, Y. Yasuta, and M. Kubota, *J. Low Temp. Phys.* **148**, 677 (2007).
- ¹²A. S. C. Rittner and J. D. Reppy, *Phys. Rev. Lett.* **98**, 175302 (2007).
- ¹³A. S. C. Rittner and J. D. Reppy, *Phys. Rev. Lett.* **101**, 155301 (2008).
- ¹⁴X. Lin, A. C. Clark, and M. H. W. Chan, *Nature (London)* **449**, 1025 (2007).
- ¹⁵A. C. Clark, J. T. West, and M. H. W. Chan, *Phys. Rev. Lett.* **99**, 135302 (2007).
- ¹⁶J. Day and J. Beamish, *Nature (London)* **450**, 853 (2007).
- ¹⁷M. H. W. Chan, *Science* **319**, 1207 (2008).
- ¹⁸J. Day and J. Beamish, *Phys. Rev. Lett.* **96**, 105304 (2006).
- ¹⁹O. Syshchenko, J. Day, and J. Beamish, *Phys. Rev. Lett.* **104**, 195301 (2010).
- ²⁰B. Hunt, E. Pratt, V. Gadagkar, M. Yamashita, A. V. Balatsky, and J. C. Davis, *Science* **324**, 632 (2009).
- ²¹H. Choi, S. Kwon, D. Y. Kim, and E. Kim, *Nat. Phys.* **6**, 424 (2010).
- ²²S. C. Rittner and J. D. Reppy, *J. Phys.: Conf. Ser.* **150**, 032089 (2009).
- ²³M. R. Sears and W. M. Saslow, *Phys. Rev. B* (to be published).
- ²⁴W. M. Saslow and S. Jolad, *Phys. Rev. B* **73**, 092505 (2006).
- ²⁵D. E. Galli, L. Reatto, and W. M. Saslow, *Phys. Rev. B* **76**, 052503 (2007).
- ²⁶W. M. Saslow, *Phys. Rev. B* **15**, 173 (1977).
- ²⁷M. Liu, *Phys. Rev. B* **18**, 1165 (1978).
- ²⁸C.-D. Yoo and A. T. Dorsey, *Phys. Rev. B* **81**, 134518 (2010).
- ²⁹M. R. Sears and W. M. Saslow, *Phys. Rev. B* (to be published).
- ³⁰Reference 26 finds equations of motion identical to those of Ref. 1 but also includes nonlinear terms. However, Ref. 27 notes that Ref. 26 employs the non-Galilean $\vec{j}_s = \rho_s \vec{v}_s$ in place of the Galilean $\vec{j}_s = \rho_s (\vec{v}_s - \vec{v}_n)$. This does not affect the equations of motion until the normal modes are calculated.
- ³¹K. R. Atkins, *Phys. Rev.* **113**, 962 (1959).
- ³²I. Rudnick and K. A. Shapiro, *Phys. Rev. Lett.* **9**, 191 (1962).
- ³³The deviation of the momentum flux tensor Π'_{ik} [Eq. (15)] has a term from the viscosity $\sim \eta_{iklm} k_m v'_l$ and a term from the stress tensor $\sim \lambda'_{ki}$. Reference 34 gives $\lambda'_{ik} \sim ik_j u'_j$. For both the propagating modes and the diffusive mode, we find that $u'_l \sim v'_n / \omega$. Thus, for both the propagating modes ($\omega \sim k$) and the diffusive mode ($\omega \sim k^2$) the term in Π'_{ik} due to viscosity is, at the least, of order k relative to the term λ'_{ki} , and is therefore neglected in the long-wavelength limit.
- ³⁴L. D. Landau and E. M. Lifshitz, *Theory of Elasticity*, 3rd ed. (Pergamon, Oxford, 1986).
- ³⁵R. H. Crepeau, O. Heybey, D. M. Lee, and S. A. Strauss, *Phys. Rev. A* **3**, 1162 (1971).