

Coulomb interaction in graphene: Relaxation rates and transportM. Schütt,¹ P. M. Ostrovsky,^{1,2} I. V. Gornyi,^{1,3} and A. D. Mirlin^{1,4,5}¹*Institut für Nanotechnologie, Karlsruhe Institute of Technology, D-76021 Karlsruhe, Germany*²*L. D. Landau Institute for Theoretical Physics RAS, 119334 Moscow, Russia*³*A. F. Ioffe Physico-Technical Institute, 194021 St. Petersburg, Russia*⁴*Institut für Theorie der kondensierten Materie, Karlsruhe Institute of Technology, D-76128 Karlsruhe, Germany*⁵*Petersburg Nuclear Physics Institute, 188350 St. Petersburg, Russia*

(Received 23 November 2010; revised manuscript received 24 February 2011; published 25 April 2011)

We analyze the inelastic electron-electron scattering in undoped graphene within the Keldysh diagrammatic approach. We demonstrate that finite temperature strongly affects the screening properties of graphene, which, in turn, influences the inelastic-scattering rates as compared to the zero-temperature case. Focusing on the clean regime, we calculate the quantum scattering rate which is relevant for dephasing interference processes. We identify a hierarchy of regimes arising due to the interplay of a plasmon enhancement of the scattering and finite-temperature screening of the interaction. We further address the energy relaxation and transport scattering rates in graphene. We find a nonmonotonic energy dependence of the inelastic relaxation rates in clean graphene, which is attributed to the resonant excitation of plasmons. Finally, we discuss the temperature dependence of the conductivity at the Dirac point in the presence of both interaction and disorder. Our results complement the kinetic equation and hydrodynamic approaches for the collision-limited conductivity of clean graphene and can be generalized to the treatment of physics of inelastic processes in strongly nonequilibrium setups.

DOI: [10.1103/PhysRevB.83.155441](https://doi.org/10.1103/PhysRevB.83.155441)

PACS number(s): 72.80.Vp, 73.23.Ad, 71.10.Ay, 71.45.-d

I. INTRODUCTION

Graphene^{1,2} is a two-dimensional (2D) material with a quasirelativistic dispersion law that has attracted outstanding attention of leading experimental as well as theoretical groups all over the world. In 2004, researchers at Manchester University first succeeded in experimental isolation of a monoatomic graphite layer—graphene—on an insulating substrate.^{3,4} This technological breakthrough was immediately followed by transport measurements^{5,6} that have shown remarkable properties related to Dirac nature of the charge carriers. In particular, a short and wide sample of clean graphene exhibits a pseudodiffusive charge transport,⁷ with the counting statistics equivalent to that of a diffusive wire.^{8,9} This equivalence has been confirmed in recent measurements of conductance and noise in ballistic graphene flakes.^{10,11} In contrast to conventional metals, ballistic graphene near the Dirac point conducts better when potential impurities are added.^{12–15}

Quantum interference in disordered graphene is also highly peculiar due to the Dirac nature of the carriers. In particular, at the Dirac point, the minimal conductivity^{5,6} $\sim e^2/h$ is “protected” from quantum localization in the absence of intervalley scattering¹⁶ or in the case of a chiral-symmetric disorder.^{17,18} Away from the Dirac point, the concentration dependence of graphene conductivity in diffusive samples depends strongly on the nature of the scatterers.¹⁷ The experimentally observed (approximately linear) dependence in most of the samples may be explained by strong impurities creating resonances near the Dirac point (“midgap states”),^{17,19} yielding $\sigma \propto n \ln^2 n$. Alternative candidates are Coulomb impurities and/or ripples, leading to^{17,20–22} $\sigma \propto n$. The dominant type (or types) of disorder and the corresponding disorder strength depend, of course, on technology of the sample preparation.

How important is the electron-electron interaction in graphene? Recent works demonstrated manifestation of the interaction in dephasing rate providing the cutoff to quantum interference phenomena,^{23–25} as well as in renormalization of conductivity.^{26,27} Recent experiments also showed that the interaction plays a particularly prominent role in suspended graphene samples.^{28,29} In such samples the splitting of integer quantum Hall transition (attributed to interaction-induced spin/valley symmetry breaking) is observed at magnetic fields as low as 2 T (i.e., an order of magnitude less than in graphene structures on a substrate). Furthermore, one observes also fractional quantum Hall plateaus, which indicates the importance of electron correlations.^{30–32} Continuous advances in the fabrication of high-quality graphene samples are expected to lead to further enhancement of the role of interaction in graphene.

On the theoretical side, interactions may have a dramatic impact on quantum electronic transport, especially in systems of reduced dimensionality.³³ Very generally, the interaction-induced phenomena may be subdivided in two big classes, related to effects of renormalization and inelastic scattering, respectively. In the case of graphene, the interaction physics becomes an even more complex problem, in view of the “relativistic” dispersion of carriers. Interaction phenomena are particularly strong near the Dirac point, where the density of states vanishes (for a clean system), the screening by intrinsic carriers becomes very inefficient, and the Drude conductivity (in the presence of disorder) is of the order of the conductance quantum.

The dimensionless bare coupling constant $\alpha_g = e^2/v_F$ describing the Coulomb repulsion in graphene samples on a SiO₂ insulating substrate is estimated to be 0.6–0.8 and can be yet larger, $\alpha_g \simeq 2.2$, in suspended graphene sheets. Transport experiments on the interaction correction to conductivity in graphene on a substrate²⁷ have found much smaller values

of the interaction constant (or, equivalently, the interaction parameter r_s) which can be attributed to the renormalization effects which we briefly overview below. Furthermore, the value of the effective fine-structure constant of free-standing graphene as inferred from the results of recent experiments on x-ray scattering in graphite³⁴ was also found to be much smaller than the bare value of α .

The interaction effects in the clean graphene have been considered in Refs. 35–41 within the weak-coupling renormalization-group (RG) scheme justified for a large number of flavors of Dirac fermions. The main result of this consideration^{35,36} is the renormalization of the Fermi velocity and hence of the interaction parameter. This perturbative renormalization group for graphene thus shows that in the clean case the Coulomb interaction is marginally irrelevant. In the disordered case, a unified ballistic RG emerges describing renormalization of disorder couplings and of the interaction.^{37–42} In Ref. 41 the corresponding one-loop RG equations are derived for time-reversal-invariant disorder and in the limit of a large number of valleys.

At sufficiently strong α_g , the Coulomb interaction in a clean graphene has been argued to give rise to various instabilities,^{43–49} in particular, to opening the gap in the Dirac spectrum of the quasiparticles (spontaneous mass generation). Physically, this instability leads to a phase transition from the semimetallic state to an insulator state, known as excitonic semimetal-insulator transition.⁴³ Recently, this type of instability has been studied by the effective mean-field-type approach,^{43,44} the renormalization-group method,³⁶ and lattice simulations.^{47,48} The mean-field consideration of the excitonic-type instability in graphene predicts a certain value of the coupling constant $\alpha_g \simeq 2$ (close to that in a suspended graphene) at which the instability occurs. The predicted semimetal-insulator transition, however, has not yet been observed in experiments in zero magnetic field. One of the possible reasons for that might be the presence of disorder.⁴⁹ In strong magnetic field—in the quantum Hall effect regime—the repulsive interaction between electrons may result in the Stoner instability^{50–52} giving rise to spontaneous breaking of spin and/or valley symmetry. Experiments do show splitting of quantum Hall plateaus in strong magnetic fields,⁵³ which is attributed to interaction effects.

It has been debated in the literature whether the Coulomb interactions in graphene can be theoretically addressed within the standard Fermi-liquid-type perturbation theory, which requires, in particular, that the particle energy is much larger than the decay rate ($\Gamma \ll \epsilon$). In order to understand to what extent graphene is a Fermi liquid, one has to explore interaction-induced inelastic collision rates. The inelastic quantum scattering rate at zero temperature (and finite quasiparticle energy) has been considered in Refs. 54 and 55. It was found in these works that the behavior at the Dirac point is rather peculiar and requires a careful incorporation of screening.

Another highly nontrivial feature of graphene is that inelastic electron-electron collisions may limit the conductivity at the Dirac point without any disorder or phonon scattering.^{56–58} This peculiarity of graphene—which should be contrasted to conventional systems where interactions do not lead (in the absence of Umklapp scattering) to finite resistivity—is a

consequence of the particle-hole symmetry and decoupling between velocity and momentum. As a result, although the total momentum of interacting particles is conserved during inelastic collisions, the total current may relax. The collision-limited conductivity of undoped graphene is found to be inversely proportional to α_g^2 and depends on temperature only through the renormalization of α_g . It was found that the energy relaxation caused by inelastic processes in graphene is fast, which allows one to treat the problem by using a relativistic hydrodynamic approach.^{59–62}

In this paper we analyze the inelastic electron-electron scattering in graphene within the Keldysh diagrammatic approach. While we focus on the equilibrium situation in this work, we have in mind to extend the treatment of physics of inelastic processes to strongly nonequilibrium setups, which explains why we prefer to work in the framework of the Keldysh formalism. More specifically, our main results are as follows:

(i) We demonstrate that finite temperature strongly affects the screening properties of graphene, which, in turn, influences the inelastic-scattering rates as compared to the zero-temperature case.

(ii) Focusing on the high-temperature regime, we calculate the quantum scattering rate, which is relevant for dephasing interference processes. We identify a hierarchy of regimes arising due to the interplay of a plasmon enhancement of the scattering and finite-temperature screening of the interaction.

(iii) We further discuss the energy relaxation and transport scattering rate in graphene. Our results complement the kinetic equation and hydrodynamic approaches^{56–58} for the collision-limited conductivity of clean graphene.

The paper is organized as follows. In Sec. II we define the model and develop the Keldysh diagrammatic formalism for treating the problem of Coulomb interaction in graphene. Section III is devoted to the analysis of polarization operator of graphene. We first review and discuss the results for the zero-temperature case. Then we turn to the case of finite temperature and discuss the properties of the polarization operator. In particular, we compare the approximate form of the dynamically screened interaction propagator with the exact numerical results. In Sec. IV we calculate the inelastic-scattering rates in the random-phase approximation. Here we use the finite-temperature polarization operator obtained in Sec. III to treat the problem analytically. We discuss the asymptotics of the inelastic rates and show that Dirac fermions show no Fermi-liquid behavior. Further, in Sec. V, we discuss the collision-limited conductivity obtained within the diagrammatic approach. In Sec. VI we conclude and summarize the main results of this paper. Technical details of the calculations are presented in four Appendices.

II. MODEL

A. Clean graphene with Coulomb interaction

In this paper we consider clean graphene near the degeneracy point. The problem is described by the following Hamiltonian, which is a sum of the Dirac Hamiltonian \hat{H}_0 (describing the physics of noninteracting electrons in

graphene at not too high energies) and the Coulomb interaction term,

$$\begin{aligned} \hat{H} &= \hat{H}_0 + \hat{V} = \sum_{\nu} \int d^2r \hat{\Psi}_{\nu}^{\dagger}(\mathbf{r}) (-i v_F \boldsymbol{\sigma} \cdot \nabla) \hat{\Psi}_{\nu}(\mathbf{r}) \\ &+ \frac{1}{2} \sum_{\nu, \nu'} \int d^2r_1 d^2r_2 \hat{\Psi}_{\nu}^{\dagger}(\mathbf{r}_1) \hat{\Psi}_{\nu'}^{\dagger}(\mathbf{r}_2) \\ &\times \frac{e^2}{\varepsilon |\mathbf{r}_1 - \mathbf{r}_2|} \hat{\Psi}_{\nu'}(\mathbf{r}_2) \hat{\Psi}_{\nu}(\mathbf{r}_1). \end{aligned} \quad (1)$$

Here ε is the dielectric constant. The spinors $\hat{\Psi}$ have two components in the sublattice space, σ_i are Pauli matrices operating in this space. The indices ν, ν' label N independent degrees of freedom (in graphene $N = 4$ accounts for spin and valleys degeneracy): The Coulomb interaction is invariant with respect to any rotations in the corresponding space. We set $\hbar = 1$. We focus on the case of undoped graphene and set the chemical potential (counted from the Dirac point) to zero, $\mu = 0$.

The retarded (advanced) Green's function of the noninteracting Hamiltonian \hat{H}_0 (the bare Green's function) in the energy-momentum space has the form

$$G_0^{R,A}(\epsilon, \mathbf{p}) = \frac{\epsilon \mathbb{1} + v_F \boldsymbol{\sigma} \cdot \mathbf{p}}{(\epsilon \pm i0)^2 - v_F^2 p^2}. \quad (2)$$

It is convenient to introduce the projection operators that distinguish between the two chiral states:

$$\mathcal{P}_{\pm}(\mathbf{p}) = \frac{\mathbb{1} \pm \boldsymbol{\sigma} \cdot \mathbf{n}}{2}, \quad (3)$$

where $\mathbf{n}_p = \mathbf{p}/p$ is the unit vector in the direction of momentum. With the help of Eq. (3) the matrix Green's function, Eq. (2), can be decomposed into the superposition of the two Green's functions corresponding to the states with + and - chiralities:

$$G_0^{R,A}(\epsilon, \mathbf{p}) = \mathcal{P}_+(\mathbf{p}) G_{0+}^{R,A}(\epsilon, \mathbf{p}) + \mathcal{P}_-(\mathbf{p}) G_{0-}^{R,A}(\epsilon, \mathbf{p}), \quad (4)$$

where

$$G_{0\pm}^R(\epsilon, \mathbf{p}) = \frac{1}{\epsilon + i0 \mp v_F p} \quad (5)$$

and $G_{0\pm}^A(\epsilon, \mathbf{p}) = [G_{0\pm}^R(\epsilon, \mathbf{p})]^*$. For later purposes we will need the quasiparticle spectral weight

$$\begin{aligned} \mathcal{A}_0(\epsilon, \mathbf{p}) &= \frac{1}{2i} [G_0^R(\epsilon, \mathbf{p}) - G_0^A(\epsilon, \mathbf{p})] \\ &= -\frac{\pi}{2\epsilon} (\epsilon \mathbb{1} + v_F \boldsymbol{\sigma} \cdot \mathbf{p}) [\delta(\epsilon - v_F p) + \delta(\epsilon + v_F p)]. \end{aligned} \quad (6)$$

Using the projection operators, we decompose the spectral weight as follows:

$$\mathcal{A}_0(\epsilon, \mathbf{p}) = \mathcal{P}_+(\mathbf{p}) \mathcal{A}_{0+}(\epsilon, \mathbf{p}) + \mathcal{P}_-(\mathbf{p}) \mathcal{A}_{0-}(\epsilon, \mathbf{p}), \quad (7)$$

$$\mathcal{A}_{0\pm}(\epsilon, \mathbf{p}) = -\pi \delta(\epsilon \mp v_F p). \quad (8)$$

It is worth noticing that for Dirac particles the spectral weight $\mathcal{A}_0(\epsilon, \mathbf{p})$ is not given by the imaginary part of the Green's function $G_0^R(\epsilon, \mathbf{p})$, because the latter contains the Pauli matrix σ_y . However, within each chirality the conventional relation holds: $\mathcal{A}_{0\pm}(\epsilon, \mathbf{p}) = \text{Im } G_{0\pm}^R(\epsilon, \mathbf{p})$.

Next, we introduce the coupling constant for Coulomb interaction in graphene,

$$\alpha_g = \frac{e^2}{\varepsilon v_F}, \quad (9)$$

which is similar to the fine-structure constant

$$\alpha = \frac{e^2}{c} \approx \frac{1}{137}$$

but is $c/\varepsilon v_F$ times larger. The bare propagator of the Coulomb interaction in Eq. (1) reads (in momentum space)

$$D_0(\mathbf{q}) = \frac{2\pi \alpha_g v_F}{|\mathbf{q}|}. \quad (10)$$

Throughout the paper we assume $\alpha_g \ll 1$. This assumption is favored by recent experiments,^{27,34} which suggested that the effective interaction constant in graphene is rather small at experimentally relevant temperatures.

B. Keldysh formalism

Although in this work we discuss only equilibrium physics, we use the Keldysh formalism^{63,64} in order to have a basis that can be generalized to the nonequilibrium situation. In particular, this Keldysh formalism will be used elsewhere for deriving kinetic equations for clean and disordered graphene.

The bare Green's function is now a matrix in the Keldysh space,

$$\check{G}_0 = \begin{pmatrix} G_0^R & G_0^K \\ 0 & G_0^A \end{pmatrix}. \quad (11)$$

Here the Keldysh component G_0^K at the equilibrium reads

$$\begin{aligned} G_0^K(\epsilon, \mathbf{p}) &= f(\epsilon) [G_0^R(\epsilon, \mathbf{p}) - G_0^A(\epsilon, \mathbf{p})] \\ &= 2if(\epsilon) \mathcal{A}_0(\epsilon, \mathbf{p}), \end{aligned} \quad (12)$$

where the fermionic thermal factor is given by $f(\epsilon) = \tanh(\epsilon/2T)$.

The full Keldysh Green's function for Eq. (1) is expressed through \check{G}_0 ,

$$\check{G} = (\check{G}_0 - \check{\Sigma})^{-1}, \quad (13)$$

where Σ is the full self-energy. In the lowest order in the fully dressed propagator of Coulomb interaction \check{D} the retarded self-energy is given by

$$\Sigma_0^R = \frac{i}{2} (D^K \circ G_0^R + D^R \circ G_0^K), \quad (14)$$

where the symbol \circ denotes integration over all internal energies and momenta.

In the equilibrium situation the Keldysh component D^K of the interaction propagator satisfies

$$D^K(\omega, \mathbf{q}) = 2ig(\omega) \text{Im } D^R(\omega, \mathbf{q}), \quad (15)$$

where $g(\omega) = \coth(\omega/2T)$ is the bosonic thermal factor. Using Eqs. (14), (15), and (12), one gets

$$\Sigma_0^R - \Sigma_0^A = -[(f+g) \text{Im } D^R] \circ (G^R - G^A). \quad (16)$$

The retarded self-energy is a matrix which contains the two terms

$$\Sigma^R = \Sigma_\epsilon^R \mathbb{1} + \Sigma_v^R \boldsymbol{\sigma} \cdot \mathbf{n}_p. \quad (17)$$

The real parts of Σ_ϵ^R and Σ_v^R give rise to the corrections to the energy and Fermi velocity, respectively:

$$\delta\epsilon = -\text{Re} \Sigma_\epsilon^R(\epsilon, \mathbf{p}), \quad (18)$$

$$\delta v_F = \text{Re} \Sigma_v^R(\epsilon, \mathbf{p})/p. \quad (19)$$

In order to fix the energy unrenormalized, we introduced the Z factor,

$$Z(\epsilon, \mathbf{p}) = 1 - \frac{\text{Re} \Sigma_\epsilon^R(\epsilon, \mathbf{p})}{\epsilon}, \quad (20)$$

so that $\epsilon - \text{Re} \Sigma_\epsilon^R(\epsilon, \mathbf{p}) = Z(\epsilon, \mathbf{p})\epsilon$. The renormalized Fermi velocity takes then the form

$$v_F^* = v_F Z(\epsilon, \mathbf{p}) \left[1 + \frac{\text{Re} \Sigma_v^R(\epsilon, \mathbf{p})}{v_F p} \right]. \quad (21)$$

Using Eqs. (20) and (21), we write the retarded Green's function as

$$G^R(\epsilon, \mathbf{p}) = Z \left[(\epsilon - iZ \text{Im} \Sigma_\epsilon^R) \mathbb{1} - \left(v_F^* + \frac{iZ \text{Im} \Sigma_v^R}{p} \right) \boldsymbol{\sigma} \cdot \mathbf{p} \right]^{-1}, \quad (22)$$

where

$$\text{Im} \Sigma_\epsilon^R = -\frac{1}{2} \text{Im} D^R(f+g) \circ \text{Tr} \mathcal{A}, \quad (23)$$

$$\text{Im} \Sigma_v^R = -\frac{1}{2} \text{Im} D^R(f+g) \circ \text{Tr} \mathcal{A} \boldsymbol{\sigma} \cdot \mathbf{n}_p. \quad (24)$$

Note that in this representation the velocity acquires a nonzero imaginary part.

The most singular terms in the renormalized velocity v_F^* , coupling constant $\alpha_g^* = e^2/\epsilon v_F^*$, and the Z factor can be summed up by means of the renormalization-group approach.³⁵ For the case of weak interaction $\alpha_g \ll 1$ the solution of one-loop RG equations has the form

$$v_F^*(\epsilon) = v_F \left(1 + \frac{\alpha_g}{4} \ln \frac{\Lambda}{\epsilon} \right), \quad (25)$$

$$\alpha_g^*(\epsilon) = \frac{\alpha_g}{1 + \frac{\alpha_g}{4} \ln \frac{\Lambda}{\epsilon}}, \quad (26)$$

$$\begin{aligned} Z(\epsilon) &= \exp \left\{ -\frac{4}{3\pi} [\alpha_g - \alpha_g^*(\epsilon)] \right\} \\ &= \exp \left[-\frac{1}{3\pi} \frac{\alpha_g^2 \ln \frac{\Lambda}{\epsilon}}{1 + \frac{\alpha_g}{4} \ln \frac{\Lambda}{\epsilon}} \right] \simeq 1. \end{aligned} \quad (27)$$

Here Λ is the ultraviolet energy cutoff (bandwidth) and the on-shell relation between ϵ and p is assumed. At finite temperature T the renormalization stops at $\max[\epsilon, T]$. Therefore for energies below T (which will be in the focus below) the renormalized velocity and the Z factor are independent of energy. Since for $\alpha_g \ll 1$ the corrections to unity in the renormalized Z factor are parametrically small, in what follows we set $Z = 1$.

Similarly to the bare Green's function, Eq. (4), the full Green's function, Eq. (22), can be represented as a sum of the two terms corresponding to \pm chiralities:

$$G^R(\epsilon, \mathbf{p}) = \mathcal{P}_+(\mathbf{p}) G_+^R(\epsilon, \mathbf{p}) + \mathcal{P}_-(\mathbf{p}) G_-^R(\epsilon, \mathbf{p}), \quad (28)$$

where

$$G_\pm^R(\epsilon, \mathbf{p}) = \frac{Z}{\epsilon \mp v_F^* p - iZ \text{Im} \Sigma_\pm^R} \quad (29)$$

with

$$\begin{aligned} \text{Im} \Sigma_\pm^R &= \text{Im} \Sigma_\epsilon^R \pm \text{Im} \Sigma_v^R \\ &= -\text{Im} D^R(f+g) \circ \text{Tr} [\mathcal{A} \mathcal{P}_\pm]. \end{aligned} \quad (30)$$

Clearly, the bare Coulomb interaction (whose propagator is purely real) does not yield the imaginary part of the self-energy, so that one has to take into account the retardation effects. In the random-phase approximation (RPA), the screened Coulomb interaction takes the form

$$D_{\text{RPA}}^R(\omega, \mathbf{q}) = \frac{D_0(q)}{1 + D_0(q) N \Pi^R(\omega, \mathbf{q})}, \quad (31)$$

where N is the number of flavors. Below we will also consider the interaction propagator at the golden rule (GR) level which corresponds to expanding Eq. (31) to the second order in D_0 ,

$$D_{\text{GR}}^R(\omega, \mathbf{q}) = -D_0^2(q) N \Pi^R(\omega, \mathbf{q}). \quad (32)$$

The dynamical screening in Eq. (31) is expressed through the bare polarization operator

$$\Pi^R = \frac{i}{2} \text{Tr} (G_0^R \circ G_0^K + G_0^K \circ G_0^A). \quad (33)$$

In Sec. III below we study Π at zero and finite temperature.

III. POLARIZATION OPERATOR

In the present section we discuss the properties of the polarization operator. This is of primary importance for understanding the screening of the electron-electron interaction and thus the physics of interaction-induced phenomena. We find it instructive to start with analyzing the zero-temperature result and describing the processes relevant for the polarization operator. Then we turn to the case of finite temperature, which is our main interest in the paper. Finally, we analyze the consequences for the RPA-screened interaction. In the end of this section we comment on the applicability of the RPA in graphene.

A. Polarization operator at zero temperature and RPA interaction propagator

The polarization operator in the energy-momentum representation reads

$$\begin{aligned} \Pi^R(\omega, \mathbf{q}) &= - \int \frac{d^2 p}{(2\pi)^2} \int \frac{d\epsilon}{2\pi} f(\epsilon) \text{Tr} \{ \mathcal{A}_0(\epsilon, \mathbf{p}) \\ &\quad \times [G_0^R(\epsilon + \omega, \mathbf{p} + \mathbf{q}) + G_0^A(\epsilon - \omega, \mathbf{p} - \mathbf{q})] \}, \end{aligned} \quad (34)$$




FIG. 1. Schematics of electron-hole (e - h) creation (left panel) vs electron-electron (e - e) scattering (right panel) near the Dirac point for $\mu = 0$. The hole-hole scattering processes are analogous to those shown in the right panel. We set $v_F = 1$ for brevity. In both panels the possible transferred momenta satisfy $|a - b| \leq |\mathbf{q}| \leq a + b$ (only the extreme cases of minimal and maximal q are shown). For e - h creation $\omega = a + b$, so that $\omega \geq qv_F$, whereas for e - e scattering the kinematic restrictions yield $\omega = a - b \leq qv_F$.

where $G_0^{R,A}$ are bare Green functions (2). The momentum integrals that appear in this expression can be conveniently evaluated using elliptic coordinates as described in Appendix A.

In the zero-temperature limit, Eq. (A1) simplifies and leads to

$$\Pi^R(\omega, q) = \frac{1}{16} \frac{q^2}{\sqrt{v_F^2 q^2 - (\omega + i0)^2}}. \quad (35)$$

Taking the imaginary part of the polarization operator, Eq. (35), yields the well-known expression³⁵

$$\text{Im } \Pi^R(\omega, \mathbf{q}) = \frac{1}{16} \text{Re} \frac{q^2 \text{sgn} \omega}{\sqrt{\omega^2 - v_F^2 q^2}}. \quad (36)$$

One can see that the imaginary part of the polarization operator is nonvanishing only if $\omega > v_F q$ and shows a divergence at the “light cone” $\omega = v_F q$.

To understand the vanishing of the imaginary part of the polarization operator at $\omega < v_F q$, it is instructive to analyze^{54,58} the kinematic restrictions for elementary processes. It is easy to see that an on-shell electron-hole pair can be created if (left panel of Fig. 1)

$$\omega \geq v_F q. \quad (37)$$

This should be contrasted to the energy and momentum conservation of on-shell electron-electron-scattering processes (right panel of Fig. 1),

$$\omega^2 + 2v_F^2 p_1 p_2 \{1 - \cos(\angle[\mathbf{p}_1, \mathbf{p}_2])\} = v_F^2 q^2, \quad (38)$$

which implies the condition

$$\omega \leq v_F q. \quad (39)$$

Thus if we restrict our consideration to on-shell particles, electron-electron-scattering processes only create electron-hole pairs under the condition $\omega = v_F q$, when $\text{Im } \Pi$ diverges. As follows from Eq. (38), scattering processes that satisfy $\omega = v_F q$ correspond to forward scattering with $\angle[p_1, p_2] = 0$.

From Eqs. (32) and (36) we obtain the GR interaction propagator at zero T :

$$D_{\text{GR}}^R(\omega, q) = -\frac{N}{16} \frac{(2\pi\alpha_g v_F)^2}{\sqrt{v_F^2 q^2 - (\omega + i0)^2}}, \quad (40)$$

$$\text{Im } D_{\text{GR}}^R(\omega, q) = -\frac{\alpha_g^2 N v_F^2 \pi^2}{4} \text{Re} \frac{\text{sgn} \omega}{\sqrt{\omega^2 - v_F^2 q^2}}.$$

After the RPA resummation [plugging Eq. (35) into Eq. (31)], the imaginary part of the zero- T interaction propagator takes the form

$$\text{Im } D_{\text{RPA}}^R(\omega, q) = -\frac{\alpha_g^2 N v_F^2 \pi^2}{4} \frac{\text{sgn} \omega \text{Re} \sqrt{\omega^2 - v_F^2 q^2}}{\omega^2 - v_F^2 q^2 (1 - \frac{\pi^2}{32} \alpha_g^2 N^2)}. \quad (41)$$

It is seen that the divergence at $\omega = v_F q$ that occurred in the GR interaction propagator, Eq. (40), is now eliminated. Moreover, the imaginary part of the propagator is zero at the light cone $\omega = v_F q$. Therefore the forward scattering divergence that arises on the GR level disappears within RPA, yielding a zero scattering rate on the RPA level. This is a manifestation of a highly singular character of the zero-temperature problem where RPA may be insufficient. We will see below that at finite temperature the thermal broadening regularizes the problem, so that scattering rates can be evaluated within the RPA.

In view of the singular character of the problem, at zero temperature the imaginary part of the self-energy is highly sensitive to changes of the electron dispersion.⁵⁸ The Fermi velocity depends logarithmically on the momentum or energy,³⁵ Eq. (25), due to the renormalization by Coulomb interaction. In Eq. (36) we have neglected the momentum dependence of v_F which leads to a separation of the two regions defined by Eqs. (37) and (39); see Fig. 2. The electron-hole creation [$\omega \geq v_F(q)q$] and electron-electron-scattering [$\omega \leq v_F(p)q$] regions are then separated due to the renormalization-induced nonlinearity of the electron dispersion, which leads to the vanishing of the zero- T scattering rate already at the GR level.

However, at finite temperature the situation is essentially different. First, conditions (37) and (39) will be smeared by temperature. Second, for energy scales smaller than T the renormalization of Fermi velocity is cut off by temperature and hence the linearity of the dispersion relations is restored: Renormalization reduces merely to a T dependence of the Fermi velocity. Therefore when discussing the finite- T physics




FIG. 2. Interaction-induced dispersion correction near the Dirac cone. The electron-hole creation [$\omega \geq v_F(q)q$] and electron-electron-scattering [$\omega \leq v_F(p)q$] regions do not overlap, so that the zero-temperature inelastic-scattering rate is zero.

of inelastic scattering on scales $\lesssim T$ we will disregard the renormalization-induced nonlinearity of the dispersion. Furthermore, below, whenever we use the notation v_F for $\epsilon \lesssim T$ we will mean the renormalized value of the velocity $v_F^*(T)$ and omit the asterisk for brevity.

B. Polarization operator at nonzero temperature

We are now ready to calculate the polarization operator at finite temperature. It turns out that the effect of finite temperature on the screening in graphene is much more pronounced than in conventional metals with finite Fermi surface and quadratic electronic dispersion. Indeed, the linearity of the spectrum of Dirac fermions gives rise to a strong (linear) energy

dependence of the density of states in two dimensions, whereas for a parabolic spectrum the density of states is constant. In the latter case, the polarization operator is essentially independent of temperature. This is not the case for Dirac particles.

Physically, a finite temperature leads to population of electronic states in an energy range $\sim T$ around the Fermi level. Let us consider the undoped graphene where the chemical potential lies at the Dirac point. The typical density of states that participate in the screening of Coulomb interaction at finite temperature is now proportional to T . In more technical terms, at finite T the integration over the fermionic energy in the polarization bubble essentially involves not only the distribution function [thermal factors $f(\epsilon)$] but also the Green's functions. This strongly changes the polarization operator at $qv_F, \omega \lesssim T$.

To simplify the notations at nonzero temperature, we introduce the dimensionless variables according to

$$Q = \frac{v_F q}{2T}, \quad \Omega = \frac{\omega}{2T}. \quad (42)$$

We use the general expression for the polarization operator, Eqs. (A1) and (A2) from Appendix A. Considering four different regions shown in Fig. 3, we simplify these equations in each of the cases, which allows us to treat the problem analytically.

The four regions are defined by the following conditions:

- region 1: $Q \ll 1$ and $Q < |\Omega|$,
- region 2: $1 \ll Q$ and $Q < |\Omega|$,
- region 3: $Q \ll 1$ and $|\Omega| < Q$,
- region 4: $1 \ll Q$ and $|\Omega| < Q$.

As shown in Appendix A, the leading-order terms with a nonvanishing imaginary part form the following simplified polarization operator:

$$\Pi^R(\Omega, Q) = \frac{T}{v_F^2} \begin{cases} \frac{\ln 2}{\pi} \left[1 - \frac{|\Omega|}{\sqrt{\Omega^2 - Q^2}} \right] + \frac{i}{8} \frac{Q^2 \tanh(\Omega/2)}{\sqrt{\Omega^2 - Q^2}}, & \text{(region 1) } |\Omega| > Q \\ \frac{\ln 2}{\pi} + \frac{i \ln 2}{\pi} \frac{\Omega}{\sqrt{Q^2 - \Omega^2}}, & \text{(region 3) } |\Omega| < Q \\ -\frac{\zeta(\frac{3}{2})(2-\sqrt{2})}{8\sqrt{\pi}} \frac{\sqrt{Q}}{\sqrt{\Omega^2 - Q^2}} + \frac{i \operatorname{sgn} \Omega}{8} \frac{Q^2}{\sqrt{\Omega^2 - Q^2}}, & \text{(region 2) } |\Omega| > Q \\ \frac{1}{8} \frac{Q^2}{\sqrt{Q^2 - \Omega^2}} + \frac{i}{\sqrt{2\pi}} \frac{\sqrt{Q} e^{-Q} \sinh \Omega}{\sqrt{Q^2 - \Omega^2}}, & \text{(region 4) } |\Omega| < Q \end{cases} \quad (43)$$

In the case of large momenta, $Q \gg 1$, we recover the zero-temperature result. For $Q \ll 1$, the polarization operator substantially differs from the zero- T expression. We will discuss this case in more detail in Sec. III C. The separation between $\Omega < Q$ and $\Omega > Q$ is dictated by the nonanalytical structure of the polarization operator; see Eq. (43).

C. RPA interaction at nonzero temperature

The real part of the polarization operator, Eq. (43), for $Q \ll 1$ is determined by temperature and leads therefore to

the screening of the Coulomb interaction:

$$\lim_{\Omega \rightarrow 0} D_{\text{RPA}}^R(\Omega, Q) = \frac{D_0(Q)}{1 + D_0(Q) N \Pi^R(0, Q)} = \frac{v_F^2 \pi}{NT \ln 2} \left(\frac{\alpha_g N \ln 2}{Q + \alpha_g N \ln 2} \right), \quad (44)$$

which yields the screening length

$$l_{\text{scr}} = \frac{v_F}{2\alpha_g NT \ln 2}. \quad (45)$$




FIG. 3. (Color online) Imaginary part of the polarization operator in the frequency-momentum plane. Four regions of dimensionless variables Ω and Q (see text) are indicated. The magnitude of $\text{Im}\Pi^R(\Omega, Q)$ is encoded by the logarithmic color scale.

Thus at finite temperature the system does screen the long-range Coulomb interaction. Note that for $Q \ll \alpha_g N$, the RPA propagator becomes independent of the interaction constant α_g ,

$$D_{\text{RPA}}^R(\Omega, Q)|_{Q \ll \alpha_g N} \simeq \frac{1}{\Pi^R(\Omega, Q)}, \quad (46)$$

as in conventional systems with Coulomb interaction. We will see, however, that the dominant contributions to relaxation rates are determined by higher transferred momenta $Q \gtrsim \alpha_g N$, where the peculiarities of the finite- T screening in graphene are crucially important.

In region 1, the real part of the polarization operator in Eq. (43) may become negative, leading to emergence of plasmon excitations. The plasmon dispersion $\Omega_p(Q)$ is determined by the zero of

$$1 + D_0 N \Pi^R = \frac{\alpha_g N Q \ln 2}{(\Omega^2 - Q^2)^{3/2}} \left[\Omega - \frac{(\alpha_g N \ln 2 + Q)\sqrt{Q}}{\sqrt{Q + \alpha_g N 2 \ln 2}} \right], \quad (47)$$

yielding

$$\Omega_p(Q) = \frac{(\alpha_g N \ln 2 + Q)\sqrt{Q}}{\sqrt{Q + 2\alpha_g N \ln 2}}. \quad (48)$$

For $Q \rightarrow 0$, this simplifies to $\Omega_p(Q) \propto \sqrt{\alpha_g N Q}$. A nonzero imaginary part of the polarization operator (43) in the corresponding region implies that these plasmons have a finite lifetime. The decay rate of plasmon excitations is given by

$$\Gamma_p(\Omega, Q) = \frac{(\Omega^2 - Q^2)^{3/2}}{\alpha_g N Q \ln 2} \text{Im} D_0(Q) \text{Im} \Pi^R(\Omega, Q), \quad (49)$$




FIG. 4. (Color online) (a) Imaginary part of RPA interaction. Left: using an analytic approximation; right: by numerical evaluation of Eqs. (A1) and (A2). Dashed lines indicate the plasmon dispersion [see Eq. (47)]. (b) cross sections through the plots of (a) from $(\Omega, Q) = (1, 0)$ to $(\Omega, Q) = (0, 1)$ (1) from $(\Omega, Q) = (0.2, 0)$ to $(\Omega, Q) = (0, 0.2)$ (2). Solid curves: analytic approximation; dashed curves: numerical integration. For all plots the interaction strength is $\alpha_g N = 1/3$.

which yields

$$\begin{aligned} \Gamma_p(\Omega, Q)|_{\Omega=\Omega_p(Q)} &= \frac{\pi \Omega}{16 \ln 2} (\Omega^2 - Q^2)|_{\Omega=\Omega_p(Q)} \\ &\leq \frac{\pi \Omega^3}{16 \ln 2} \Big|_{\Omega=\Omega_p(Q)}. \end{aligned} \quad (50)$$

Remarkably, Eq. (50) indicates a good quasiparticle behavior for plasmons, which is, as we will see in Sec. IV A, not true for electronic excitations. The situation is somewhat similar to Luttinger liquid where plasmons are almost perfect quasiparticles, whereas, from the point of view of fermionic excitations, the Luttinger liquid represents a paradigmatic example of a non-Fermi liquid.

Figure 4 demonstrates that Eq. (43) for the polarization operator yields a remarkably good approximation for evaluation of the imaginary part of the RPA interaction,

$$\text{Im } D_{\text{RPA}} = -\frac{ND_0^2 \text{Im } \Pi^R}{(1 + ND_0 \text{Re } \Pi^R)^2 + (ND_0 \text{Im } \Pi^R)^2} \quad (51)$$

[see also Eq. (B2) in Appendix B]. In Fig. 4(b) we see the plasmon peak in the RPA interaction propagator, which is strongly asymmetric and is suppressed around the light cone, so that $\text{Im } D_{\text{RPA}} = 0$ exactly at $\Omega = Q$.

Let us now discuss the status of the RPA in graphene. Above we have introduced the large number of independent flavors $N \gg 1$. At zero temperature, this is the only parameter which justifies the RPA summation in the problem of interacting Dirac fermions at $\mu = 0$. Indeed, in view of the absence of the screening, the non-RPA diagrams are parametrically the same as those included in the RPA series for $N \sim 1$. Furthermore, as discussed in the end of Sec. III A, the renormalization-induced curvature of the spectrum (which is also beyond the RPA) may dramatically affect the results obtained within the RPA.

However at finite T , the T -induced screening of the interaction, Eq. (44), restores the validity of the RPA for $q \ll T/v_F$ ($Q \ll 1$) even for $N \sim 1$. Indeed, the $1/q$ singularity of the long-range Coulomb interaction is not compensated in the denominator of Eq. (51) because the polarization operator at finite T is no longer linear in q . The situation becomes similar to that in conventional metals with a finite Fermi surface, where the RPA is justified for $q \ll k_F$. In graphene the role of k_F is played by T/v_F , which in effect establishes an analog of a finite Fermi surface. Therefore for $q \ll T/v_F$ the RPA does sum up the most singular interaction-induced terms: All other terms are nonsingular because of the screening. This means that all the observables that are dominated by the collisions with the momentum transfer smaller than T/v_F can be calculated (even with the correct numerical prefactors) within the RPA. The RPA result for those observables that are dominated by $qv_F \sim T$ is parametrically correct, but the value of the prefactor cannot be found using the RPA.

Below we employ the finite- T RPA for calculation of various scattering rates in graphene. For the sake of generality, we keep N as a parameter. In what follows we focus on the case $\alpha_g N \ll 1$, but whenever the rate under consideration is

dominated by $qv_F \lesssim T$, the condition $N \gg 1$ can be removed so that we are allowed to use the RPA for $N \sim 1$.

IV. SCATTERING RATES

In this section we calculate various inelastic-scattering rates in clean graphene at finite temperature. More specifically, we focus on the quantum scattering rate (and dephasing) and the energy relaxation rate induced by the RPA-screened Coulomb interaction. In Sec. V A below, we will also calculate the transport scattering rate due to inelastic collisions. The quantum scattering rate determines the lifetime of quantum states (plane waves) and is related to the dephasing rate. The energy relaxation rate governs the relaxation of the quasiparticle distribution function. Finally, the transport scattering rate describes the influence of the inelastic scattering on transport phenomena.

Although the origin of all these rates is the same—the inelastic electron-electron collisions, these rates may strongly differ from each other. For instance, this is exactly what happens in diffusive metals^{33,65} because of the infrared-singular collision kernel. Another prominent example of a nontrivial behavior of relaxation rates related to the infrared singularities is a Luttinger liquid (disordered or clean).^{66–69} On the other hand, within the Fermi-liquid theory of clean metals all the inelastic-scattering rates behave in the same way, since the characteristic frequency and momenta transfer in the course of electron-electron collisions is determined by temperature. The goal of this section is to understand whether or not the situation in clean graphene is similar to that of the Fermi liquid.

We have already mentioned in the Introduction (Sec. I) the previous works on the inelastic quantum scattering rate in graphene.^{35,55} These works addressed the scattering rate at the GR level at zero temperature and obtained the Fermi-liquid-type result $\tau_q^{-1} \sim \alpha_g^2 N \epsilon \ll \epsilon$. Naively, one could think that at finite temperature this consideration would lead to $\tau_q^{-1} \sim \alpha_g^2 NT$. However, since the GR result is completely determined by the “mass shell” ($\omega = v_F q$) (see discussion in Sec. III A) one concludes that the RPA resummation (which kills the on-shell interaction) would yield $\tau_q^{-1}(\epsilon) = 0$ for $T = 0$ and arbitrary ϵ . Therefore the finite- T expectation based on the GR is also doubtful.

How the quasiparticle broadening behaves in the experimentally relevant case of finite temperature is thus by far not obvious. As discussed in Sec. III B, the finite density of states at nonzero T leads to screening of the Coulomb interaction, thus justifying the use of the RPA which then sums up the most singular contributions of the interaction for small momenta, $qv_F < T$. In this section we find the behavior of inelastic rates at finite temperature within the RPA. As we have already seen in Sec. III B, the behavior of the polarization operator at finite temperature is highly nontrivial. This leads to a rather rich behavior of the scattering rates.

It turns out that, in addition to the temperature scale, two more characteristic scales appear which are relevant for relaxation rates: $\alpha_g^2 N^2 T$ and $\alpha_g NT$. In this section we will distinguish between the four regimes (I–IV) as shown in Fig. 5. The contributions to each of the scattering rates from




FIG. 5. Characteristic energy scales separating domains of distinct behavior of the rates.

different regions (1–4, see Fig. 3) of the momentum-frequency plane are calculated separately.

A. Quantum scattering rate in graphene

One of the main manifestations of the inelastic scattering in electronic systems is the interaction-induced dephasing. In order to analyze the effects of dephasing in graphene, we will follow the route suggested by earlier works on 2D (in particular, diffusive) systems.³³ A natural first step is to calculate the quantum scattering rate τ_q^{-1} which is given by the imaginary part of the quasiparticle self-energy taken at the mass shell. Indeed, for conventional metals with parabolic dispersion, in the high-temperature (ballistic) regime, the dephasing rate τ_ϕ^{-1} to leading order is given by τ_q^{-1} ; see Ref. 70. In the diffusive regime, the scattering kernel acquires an infrared singularity leading to a divergent τ_q^{-1} at finite T . This is a manifestation of the fact that the single-particle self-energy is not a gauge-invariant object; no divergencies occur in observable (gauge-invariant) quantities, such as, e.g., the dephasing rate. However, even when the quantum scattering rate diverges, the calculation of it turns out to be instructive: A parametrically correct result for the dephasing rate can be obtained from the expression for τ_q^{-1} supplemented with an appropriate infrared cutoff. It is thus useful to begin with analyzing τ_q^{-1} .

1. Quantum scattering rate: Definitions

The peculiarity of graphene is that the self-energy is a matrix in the sublattice space, which has a nonzero imaginary part, $\text{Im} \Sigma_v$ [Eq. (24)], in the off-diagonal components. Therefore the definition of the quantum scattering time in graphene is actually not unique. Indeed, one can associate with the quantum scattering rate the on-shell value of the imaginary correction to the energy in the full Green's function, $\text{Im} \Sigma_\epsilon$ [Eq. (23)], similarly to the conventional Fermi-liquid theory:

$$\frac{1}{2\tau_q(\epsilon)} = -\theta(\epsilon) \text{Im} \Sigma_\epsilon(v_F p, \mathbf{p})|_{p=\epsilon/v_F} - \theta(-\epsilon) \text{Im} \Sigma_\epsilon(-v_F p, \mathbf{p})|_{p=-\epsilon/v_F}. \quad (52)$$

Here at $\epsilon > 0$ we have taken the self-energy at the “+” mass shell corresponding to the positive energies, $\epsilon = v_F p$, and at $\epsilon < 0$ on the “−” mass shell. Clearly,

$$\frac{1}{\tau_q(\epsilon)} = \frac{1}{\tau_q(-\epsilon)} \quad (53)$$

for undoped graphene ($\mu = 0$) because of the particle-hole symmetry. Within the RPA the explicit expression for the imaginary part of the total self-energy Σ_ϵ taken at $\epsilon = v_F p$

reads

$$\begin{aligned} \text{Im} \Sigma_\epsilon(v_F p, \mathbf{p}) &= -\frac{\pi}{2} \int \frac{d\omega}{2\pi} [g(\omega) + f(v_F p - \omega)] \int \frac{d^2 q}{(2\pi)^2} \text{Im} D^R(\omega, \mathbf{q}) \\ &\quad \times [\delta(v_F p - \omega - v_F |\mathbf{p} - \mathbf{q}|) \\ &\quad + \delta(v_F p - \omega + v_F |\mathbf{p} - \mathbf{q}|)]. \end{aligned} \quad (54)$$

Alternatively, one can introduce the lifetime of the + and − chiral states through the corresponding self-energies, Eq. (30):

$$\frac{1}{2\tau_\pm(\epsilon)} = -\text{Im} \Sigma_\pm(\epsilon, \mathbf{p})|_{p=|\epsilon|/v_F}, \quad (55)$$

which yields

$$\frac{1}{\tau_+(\epsilon)} = \frac{1}{\tau_-(-\epsilon)}. \quad (56)$$

Using Eqs. (30) and (6), we get for the self-energy of electrons (+ chirality)

$$\begin{aligned} \text{Im} \Sigma_+(v_F p, \mathbf{p}) &= -\frac{\pi}{2} \int \frac{d\omega}{2\pi} [g(\omega) + f(v_F p - \omega)] \int \frac{d^2 q}{(2\pi)^2} \text{Im} D^R(\omega, \mathbf{q}) \\ &\quad \times \left[\left(1 + \frac{\mathbf{p}(\mathbf{p} - \mathbf{q})}{p|\mathbf{p} - \mathbf{q}|} \right) \delta(v_F p - \omega - v_F |\mathbf{p} - \mathbf{q}|) \right. \\ &\quad \left. + \left(1 - \frac{\mathbf{p}(\mathbf{p} - \mathbf{q})}{p|\mathbf{p} - \mathbf{q}|} \right) \delta(v_F p - \omega + v_F |\mathbf{p} - \mathbf{q}|) \right] \end{aligned} \quad (57)$$

for the “own” mass shell, and

$$\begin{aligned} \text{Im} \Sigma_+(-v_F p, \mathbf{p}) &= -\frac{\pi}{2} \int \frac{d\omega}{2\pi} [g(\omega) + f(-v_F p - \omega)] \int \frac{d^2 q}{(2\pi)^2} \text{Im} D^R(\omega, \mathbf{q}) \\ &\quad \times \left[\left(1 - \frac{\mathbf{p}(\mathbf{p} - \mathbf{q})}{p|\mathbf{p} - \mathbf{q}|} \right) \delta(-v_F p - \omega - v_F |\mathbf{p} - \mathbf{q}|) \right. \\ &\quad \left. + \left(1 + \frac{\mathbf{p}(\mathbf{p} - \mathbf{q})}{p|\mathbf{p} - \mathbf{q}|} \right) \delta(-v_F p - \omega + v_F |\mathbf{p} - \mathbf{q}|) \right] \end{aligned} \quad (58)$$

for the “wrong” (hole) mass shell.

The main formal difference between the two relaxation rates, Eqs. (52) and (55), which are related by

$$\frac{1}{\tau_q(\epsilon)} = \frac{1}{2\tau_+(\epsilon)} + \frac{1}{2\tau_-(-\epsilon)}, \quad (59)$$

is in the appearance of Dirac factors $1 \pm \cos \theta$ in Eqs. (57) and (58) where $\theta = \arccos(\mathbf{p}\mathbf{p}'/pp')$ is the scattering angle between the incoming momentum \mathbf{p} and the momentum $\mathbf{p}' = \mathbf{p} - \mathbf{q}$ after scattering. These factors are related to the additional Berry phase in the problem of Dirac particles, which arises due to the overlap of Bloch functions and, in particular, forbids the backscattering within the same chirality and valley. Note that for well-defined quasiparticles (i.e., in a Fermi-liquid situation) the self-energy at a “wrong” mass

shell would never be relevant. However, if the quasiparticle broadening is larger than the characteristic energy, this is no longer the case, so that Eq. (58) may then contribute to the observables.

In Eq. (57), the term with $1 + \cos\theta$ corresponds to the electron-electron scattering (right panel of Fig. 1), which is determined by the contribution of region 3 in Fig. 3. The term with $1 - \cos\theta$ is due to electron-hole scattering (electron-hole pairs, left panel of Fig. 1) accompanied by the excitation of plasmons and is determined by the contribution of region 1 in Fig. 3. The latter contribution is suppressed for the forward scattering $\theta = 0$ because of the Dirac factor. Furthermore, at zero temperature only the electron-electron processes are allowed by the kinematic restrictions.

At finite T , however, the situation is different for low-energy domains I and II ($\epsilon < \alpha_g NT$), where the contributions of the electron-electron and electron-hole scattering to the inelastic quantum scattering rates are of the same order. This means that the low-energy electron-type and hole-type quasiparticles are strongly correlated by the mutual inelastic scattering, whereas at higher energies ($\epsilon > \alpha_g NT$, corresponding to domains III and IV) the electronic and hole subsystems are only weakly coupled with each other, in agreement with Ref. 58.

As we will see below, depending on the energy range, Fig. 5, both τ_q^{-1} and τ_{\pm}^{-1} may be larger or smaller than energy. When the quasiparticle's broadening is small, the two rates coincide since the inelastic scattering is dominated by small scattering angles θ and the Dirac factors reduce to 0 and 1. At the lowest energies (domains I and II), both τ_q^{-1} and τ_{\pm}^{-1} exceed the energy which makes the notion of mass shell not well defined. In particular, due to the inelastic broadening, the electronic excitation (+ chirality) has tails at negative energies which overlap with the hole mass shell.

Furthermore, as shown in Sec. IV C, the characteristic rate of the energy relaxation (energy mixing due to the diffusion over energy) at low energies is of the same order as the quantum scattering rate, so that the electronic excitations constantly explore the hole mass shell and vice versa. In this situation, the broadening of quasiparticles of a given chirality is in effect described by the total scattering rate τ_q^{-1} , according to Eq. (59) rather than Eq. (55). Therefore in what follows we will mostly focus on the Fermi-liquid-type total rate τ_q^{-1} , formally defined in Eq. (52).

2. Quantum scattering rates: Results and discussion

The evaluation of integrals involved in the calculation of the total quantum scattering rate τ_q^{-1} is outlined in Appendix B. The result depends on the energy range (see Fig. 5):

$$\frac{1}{2\tau_q(\epsilon)} \approx \begin{cases} c_1 \frac{T}{N} \sqrt{\frac{\epsilon}{T}}, & \epsilon \ll \alpha_g^2 N^2 T, \text{ I} \\ c_2 \alpha_g T, & \alpha_g^2 N^2 T \ll \epsilon \ll \alpha_g NT, \text{ II} \\ c_3 \alpha_g T, & \alpha_g NT \ll \epsilon, \text{ III, IV.} \end{cases} \quad (60)$$

Here $c_1 \sim 1$, $c_2 = 3\pi/2$, and $c_3 = 2\beta(2)/\pi$ are the numerical coefficients of order unity, $\beta(x)$ is the Dirichlet beta function, and $\beta(2)$ is the Catalan constant. Note that for energies in regimes II, III, and IV, the number of independent flavors




FIG. 6. (Color online) Quantum scattering rate for $\alpha_g N = 4 \times 10^{-3}$ (double logarithmic scale). Dots: exact values obtained by numerical evaluation; solid lines: analytical asymptotics, Eq. (60).

N drops out from the expression for τ_q . The comparison of the asymptotic expressions (60) with the exact numerical evaluations is shown in Fig. 6.

The obtained rates are dominated by different values of momenta transferred during the collision. Furthermore, depending on the energy range, the main contribution may come from different regions of the Q vs Ω plane (Fig. 3) as shown in Table I.

In regime I the result is determined by momenta of order temperature and therefore all four regions contribute in the same way: $\tau_q^{-1} \sim (\epsilon T)^{1/2}/N$. All scattering angles θ contribute to the result in regime I. In order to evaluate the numerical prefactor c_1 one needs the knowledge of the screened interaction in the crossover around $qv_F \sim T$, which is beyond the analytic approximations for the polarization operator used above. For energies in regime II the dominant contributions come from regions 3 (electron-electron scattering) and 1 (electron-hole scattering); in both of them all scattering angles θ contribute. In regimes III and IV the main contribution to the quantum scattering rate stems from region 3 (electron-electron scattering) and is dominated by forward scattering ($\theta \lesssim \alpha_g NT/\epsilon \ll 1$).

An important feature of the quantum scattering rate is its nonmonotonic energy dependence; see Eq. (60) and Fig. 6: with increasing energy the quasiparticle broadening first grows in regime I, has a maximum at $\epsilon \sim \alpha_g^2 N^2 T$, then decreases in regime II, and finally becomes energy independent. The maximum of τ_q^{-1} occurs due to the resonant emission/absorption of plasmonic excitations. We will see below that the nonmonotonicity of the energy dependence is related to the peculiar properties of the dynamical screening

TABLE I. Momentum/frequency scales and regions of the Q - Ω plane that dominate the quantum scattering rate τ_q^{-1} in different domains (I, II, III, and IV) of energy ϵ .

τ_q	I	II	III	IV
$qv_F \sim \omega$	T	$(\alpha_g NT)^2/\epsilon$	$\alpha_g NT$	$\alpha_g NT$
Regions	1,2,3,4	1,3	3	3




FIG. 7. (Color online) Quantum scattering rate τ_+^{-1} for the + chirality (electrons) for $\alpha_g N = 4 \times 10^{-3}$ (double logarithmic scale). Dots: exact values obtained by numerical evaluation; solid lines: analytical asymptotics (see Appendix B).

in graphene and is a characteristic feature of all inelastic scattering rates in graphene.

In Fig. 7 we show the results for the quantum scattering rate of electrons, τ_+^{-1} , defined in Eq. (55). One sees that Fig. 7 represents an “unfolding” of Fig. 6 into the contributions of “own” ($\epsilon > 0$) and “wrong” ($\epsilon < 0$) mass shells, according to Eq. (59). The hole mass shell with $\epsilon < 0$ is probed by electrons due to the strong quasiparticle broadening. At positive energies in regimes III and IV ($\epsilon > \alpha_g N T$), the total scattering rate τ_q^{-1} is dominated by the electronic scattering rate τ_+^{-1} .

In contrast to the naive expectation ($\tau_q^{-1} \sim \alpha_g^2 N T$) based on the Fermi GR result, the quantum scattering rate calculated within the RPA is proportional to α_g (and does not depend on N) for $\epsilon \gg \alpha_g^2 N^2 T$ and is independent of α_g at the lowest energies. This enhancement of the inelastic scattering for $\alpha_g N \ll 1$ is a result of peculiar screening properties of graphene at finite T which leads to a nonanalytic behavior of the rate as a function of the natural four-fermion coupling constant α_g^2 . This behavior bears a certain similarity to the behavior of the quantum scattering rate in a spinful Luttinger liquid.^{66,67}

Since the RPA quasiparticle broadening in domain I may overlap with the higher-energy domains, where the quantum scattering rate is of the order of $\alpha_g T$, one can speculate that the characteristic strength of inelastic scattering is given by $\tau_q^{-1}(\epsilon) \sim \alpha_g T$ also at low energies. Of course, the calculation based on the lowest-order RPA diagrams for the self-energy is then insufficient. Moreover, the typical observables at finite temperatures are dominated by $\epsilon \sim T$ (the border between domains III and IV), where the quasiclassical broadening is smaller than the energy and the above calculation is justified. The low-energy inelastic relaxation may become relevant in the context of spectroscopy under strongly nonequilibrium conditions, for example, in problems related to tunneling into a nonequilibrium state (cf. Ref. 69). In this situation the inelastic effects can be treated within the quantum-kinetic approach including non-RPA contributions, similarly to one-dimensional problems.⁶⁹

B. Dephasing rate at high temperatures

Let us now discuss the dephasing rate. The result for the quantum scattering rate obtained for the clean graphene allows us to evaluate the dephasing rate relevant to weak

(anti)localization in the ballistic regime²⁴ of high temperatures $T \tau_{\text{dis}} \gg 1$, where τ_{dis} is the elastic mean free time due to scattering off impurities. It is worth noting that this condition may coexist with the condition $\tau_\phi \gg \tau_{\text{dis}}$ which allows long interfering paths in the weak-localization experiment. Indeed, the inelastic scattering is suppressed with decreasing α_g (even though in a nontrivial way) so that one expects that τ_ϕ can be made arbitrarily long.

The result for the quantum scattering rate, Eq. (60), remains intact (up to the change in prefactor) when we calculate it in a self-consistent way as appropriate for estimating the dephasing rate:

$$\tau_\phi^{-1} \propto \int_{(T \tau_\phi)^{-1}}^{\infty} dQ \dots \quad (61)$$

This happens due to the fact that the characteristic momenta dominating the integrals for the quantum scattering rate in regimes II, III, and IV is of the same order as the resulting rate [see Table I and Eq. (60)]; both are $\sim \alpha_g T$ for $N \sim 1$ (for $N \gg 1$, the characteristic momenta are much higher than $1/\tau_q$). In regime I of lowest energies, the characteristic momentum transfer is much higher than the rate. Therefore in all these regimes the infrared cutoff is redundant and the dephasing rate is given by the quantum scattering rate, Eq. (60), similarly to the case of conventional metals in the ballistic regime⁷⁰ (although, in contrast to the conventional case, the characteristic transferred frequencies are much smaller than temperature). Since the characteristic energies involved in the transport coefficients are of order of T , we conclude that in the ballistic regime the interference effects are governed by

$$\frac{1}{\tau_\phi} \sim \alpha_g T. \quad (62)$$

This prediction can be verified by transport experiments on graphene at sufficiently high temperatures (depending on the purity of the system, $T \gtrsim 10$ – 100 K for a typical setup with graphene deposited on a insulating substrate and $T \gtrsim 1$ – 10 K for suspended graphene flakes). At lower temperatures $T \tau_{\text{dis}} \ll 1$ corresponding to the “diffusive regime” with respect to interaction, one expects the conventional^{33,71} diffusive result for the dephasing rate,

$$\frac{1}{\tau_\phi} \sim \frac{T}{g} \ln g, \quad (63)$$

where g is the dimensionless Dirac-point conductance. At the Dirac point g is close to unity and hence $\tau_\phi^{-1} \sim T$.

C. Energy relaxation rate

Let us now discuss the energy relaxation time in clean graphene. In Sec. IV A we have seen that the typical momentum or energy transfer during the electron-electron collision is much smaller than temperature. In this situation, the energy relaxation occurs through multiple-scattering processes, which can be viewed as diffusion in energy space (see, e.g., Ref. 72). The characteristic energy relaxation rate is given by the diffusion coefficient of this problem, which amounts to introducing the factor $\mathcal{K}_E = \omega^2/T^2$, see Appendix B1, into the collision kernel. More rigorous calculation of




FIG. 8. (Color online) Energy relaxation rate for $\alpha_g N = 4 \times 10^{-3}$ (double logarithmic scale), obtained from the energy-diffusion consideration. Dots: exact values obtained by numerical evaluation; solid lines: analytical asymptotics, Eq. (64). Since the energy-diffusion approximation employed in the calculation is not justified in regime IV, we do not present the results in this regime for $\epsilon \gg T$.

the energy relaxation or equilibration rates can be done using the language of kinetic equation; here we only estimate the typical rate τ_E^{-1} within the energy diffusion picture.

Once the quantum scattering rate is obtained, the calculation of the energy relaxation rate can be done using the same steps as described in the previous section. Technically, the integrals in Eq. (54) are only slightly changed, which leads, however, to a substantial difference between the two rates. The detailed calculation can be found in Appendix B; here we present only the result:

$$\frac{1}{\tau_E(\epsilon)} \sim \begin{cases} \frac{T}{N} \sqrt{\frac{\epsilon}{T}}, & \text{I} \\ \frac{\alpha_g^2 N T}{\sqrt{\epsilon/T}} \ln\left(\frac{\epsilon/T}{\alpha_g^2 N^2}\right), & \text{II, III} \\ \alpha_g^2 N T \left(\frac{\epsilon}{T}\right)^{3/2} \ln\left(\frac{1}{\alpha_g N}\right), & \text{IV.} \end{cases} \quad (64)$$

In particular, we have the following estimates for the energy relaxation rate at characteristic energies separating the regimes:

$$\frac{1}{\tau_E(\epsilon)} \sim \begin{cases} \alpha_g T, & \epsilon \sim \alpha_g^2 N^2 T \\ \alpha_g^{3/2} N^{1/2} T \ln\left(\frac{1}{\alpha_g N}\right), & \epsilon \sim \alpha_g N T \\ \alpha_g^2 N T \ln\left(\frac{1}{\alpha_g N}\right), & \epsilon \sim T. \end{cases} \quad (65)$$

Again the obtained rates are dominated by different momentum scales and by contributions of different regions as shown in Table II. Due to the factor \mathcal{K}_E all contributions except in regime IV are now determined by momenta of order temperature, which does not allow us to find the numerical value of the prefactors analytically.

Furthermore, in fact, the above calculation based on the energy diffusion is not justified for “hot electrons” with high energies, $\epsilon > T$. Indeed, within our consideration, the characteristic energy transfer dominating the energy relaxation

TABLE II. Momentum/frequency scales and regions of the Q - Ω plane that dominate the energy relaxation rate τ_E^{-1} in different domains (I, II, III, and IV) of energy ϵ .

τ_E	I	II	III	IV
$qv_F \sim \omega$	T	T	T	ϵ
Regions	1,2,3,4	1,2,3,4	1,2,3,4	4

in regime IV turns out to be much higher than T , in contrast to the original assumption. Therefore the estimate Eq. (64) for regime IV cannot be trusted and another approach is needed for this regime.

The numerical results for the energy relaxation rate are shown in Fig. 8 together with the analytical asymptotics, Eq. (64). One sees that the energy relaxation has a minimum at $\epsilon \sim T$, where we recover, up to the logarithmic factor $|\ln \alpha_g|$, the GR result $\tau_E^{-1} \sim \alpha_g^2 N T$. At lower energies the inelastic scattering is enhanced due to the resonance in the RPA interaction propagator (the resonant condition corresponds to $\epsilon \sim \alpha_g^2 N^2 T$), whereas at high energies the energy relaxation is stronger because of the large phase space available for inelastic processes. We remind the reader, however, that at $\epsilon > T$ the above calculation based on the energy-diffusion approximation is not justified. In order to find the correct relaxation of the distribution functions in this regime of hot electrons, one should solve the corresponding kinetic equation which can be reduced to the Fokker-Planck equation. This will be done elsewhere.

V. TRANSPORT RATE AND CONDUCTIVITY

A. Transport scattering rate due to inelastic collisions

In this section we calculate the transport relaxation rate due to inelastic collisions. The expression for the corresponding kernel of the self-energy can be deduced from the interaction-induced correction to the conductivity, as described below. We are interested in the linear-response dc conductivity. The leading-order perturbative correction to the conductivity due to Coulomb interaction is given by the two diagrams shown in Fig. 9.

In the absence of interaction, the conductivity of a clean graphene diverges at finite temperature. Therefore we have regularized the diagrams by introducing a small broadening (δ) which mimics the finite lifetime due to weak disorder. The general analytic expression of the lowest-order interaction correction to the conductivity within the Keldysh formalism can be found in Ref. 71. This correction can be split into two parts [Figs. 9(a) and 9(b), respectively]: $\delta\sigma = \delta\sigma^s + \delta\sigma^v$,




FIG. 9. Diagrams describing the first-order interaction correction to the conductivity.

where

$$\begin{aligned} \delta\sigma_{\beta\beta}^s = & -\frac{1}{4i} \int \frac{d^2 p_1}{(2\pi)^2} \int \frac{d\epsilon_1}{2\pi} \int \frac{d^2 p_2}{(2\pi)^2} \int \frac{d\epsilon_2}{2\pi} \frac{\partial f_1}{\partial \epsilon} \text{Tr}\{j_\beta[G^R(1) - G^A(1)]j_\beta \\ & \times [(f_2 + g)[D^R - D^A][G^R(1)G^R(2)G^R(1) - G^A(1)G^A(2)G^A(1)] \\ & + f_2[G^R(1)[G^R(2)D^R - G^A(2)D^A]G^R(1) - G^A(1)[G^R(2)D^R - G^A(2)D^A]G^A(1)]\} \end{aligned} \quad (66)$$

is the self-energy contribution and

$$\begin{aligned} \delta\sigma_{\beta\beta}^v = & -\frac{1}{4i} \int \frac{d^2 p_1}{(2\pi)^2} \int \frac{d\epsilon_1}{2\pi} \int \frac{d^2 p_2}{(2\pi)^2} \int \frac{d\epsilon_2}{2\pi} \frac{\partial f_1}{\partial \epsilon} \text{Tr}\{j_\beta(f_2 + g)[D^R - D^A] \\ & \times [G^R(1)G^R(2) - G^A(1)G^A(2)]j_\beta[G^R(2)G^R(1) - G^A(2)G^A(1)] \\ & + 2j_\beta f_2[G^R(1) - G^A(1)][D^A G^R(2)j_\beta G^R(2) - D^R G^A(2)j_\beta G^A(2)]j_\beta[G^R(1) - G^A(1)]\} \end{aligned} \quad (67)$$

is due to the vertex correction. Here $j_\beta = ev_F\sigma_\beta$ is the current operator in graphene and we used the shorthand notations for the arguments of Green's functions: 1 = \mathbf{p}_1, ϵ_1 and 2 = \mathbf{p}_2, ϵ_2 . We first trace out the sublattice structure and then regularize the divergent integrals by δ . The most divergent interaction-induced correction to the dc conductivity is then proportional to δ^{-2} :

$$\begin{aligned} \delta\sigma = & -\frac{e^2 v_F^2}{4\delta^2} \int \frac{d\epsilon_1}{2\pi} \frac{\partial f}{\partial \epsilon_1} \int \frac{d\epsilon_2}{2\pi} \int \frac{d^2 p_1}{(2\pi)^2} \int \frac{d^2 p_2}{(2\pi)^2} \pi [\delta(\epsilon_1 - v_F p_1) + \delta(\epsilon_1 + v_F p_1)] \text{sgn}\epsilon_1 \\ & \times \underbrace{\left[1 - \left(\frac{\mathbf{p}_1 \cdot \mathbf{p}_2}{p_1 p_2}\right)\right]}_{\text{Transport factor}} \underbrace{\left[1 + \left(\frac{\mathbf{p}_1 \cdot \mathbf{p}_2}{p_1 p_2}\right)\right]}_{\text{Dirac factor}} K_\Sigma(\epsilon_1, \epsilon_1 - \epsilon_2, \mathbf{p}_1, \mathbf{p}_1 - \mathbf{p}_2). \end{aligned} \quad (68)$$

Here K_Σ is the integral kernel of the self-energy:

$$\text{Im} \Sigma_0^R(\epsilon, \mathbf{p}) = \int \frac{d^2 q}{(2\pi)^2} \int \frac{d\omega}{2\pi} K_\Sigma(\epsilon, \omega, \mathbf{p}, \mathbf{q}). \quad (69)$$

The correction, Eq. (68), is determined by inelastic electron-electron scattering. In the diagrams of the leading order in bare interaction shown in Fig. 9, we have $K_\Sigma \propto \text{Im} D_0 = 0$, so that the inelastic corrections are zero. What remains in the first-order conductivity correction, Fig. 9, are the contributions responsible for the renormalization of the Fermi velocity in graphene coming from the real part of the self-energy; note that these contributions are less singular in δ^{-1} .

Thus we have to consider higher-order corrections. In Fig. 10 one can see all classes of second-order skeleton




FIG. 10. Second-order skeleton diagrams for the conductivity.

diagrams. There are also the second-order diagrams of the ladder type; see Fig. 11. Such diagrams contribute only to the renormalization of v_F , similarly to the diagrams in Fig. 9, and their contribution has been already included into the calculation simply by the replacement $v_F \rightarrow v_F(T)$.

In the large N approximation diagrams in Figs. 10(d)–10(f) dominate. Diagram 10(f) is known as the Coulomb drag diagram and yields a zero contribution at the Dirac point due to electron-hole symmetry (see Refs. 73 and 74 and Appendix C). Thus we are left with diagrams 10(d) and 10(e) which correspond to the two diagrams shown in Fig. 9 with the second-order correction to the interaction instead of the bare one. For these two diagrams Eq. (68) still holds with $K_\Sigma \propto \text{Im} D_1$. This suggests to replace the bare interaction lines shown in Fig. 9 by the RPA interaction lines, which would correspond to Eq. (68) with $K_\Sigma \propto \text{Im} D_{\text{RPA}}$. Note that even with the RPA-dressed interaction, Eq. (68) still yields a divergent contribution which we have regularized with δ .




FIG. 11. Examples of second-order diagrams for the conductivity that contribute to the renormalization of velocity.

Considering the two independent scattering processes with the transport scattering rates δ and τ_{tr}^{-1} such that we find the Drude conductivity in the form

$$\sigma = e^2 \int d\epsilon \rho(\epsilon) \left(-\frac{\partial n_F}{\partial \epsilon} \right) \frac{v_F^2 \tau_{\text{tr}}(\epsilon)}{2[1 + \tau_{\text{tr}}(\epsilon)\delta]}, \quad (70)$$

where $n_F(\epsilon) = [1 - f(\epsilon)]/2$ is the thermal Fermi distribution function and $\rho(\epsilon)$ is the thermodynamic density of states. Expanding this formula in $(\delta\tau_{\text{tr}})^{-1}$, we get the ‘‘interaction-induced’’ correction:

$$\delta\sigma = \frac{e^2 v_F^2}{4\delta^2} \int d\epsilon \frac{\partial f}{\partial \epsilon} \frac{\rho(\epsilon)}{\tau_{\text{tr}}(\epsilon)}. \quad (71)$$

Comparing Eq. (71) with Eq. (68) we can identify the interaction-induced transport scattering rate in graphene. The transport scattering rate obtained from the conductivity correction corresponds to the kernel \mathcal{K}_{tr} defined in Appendix B1.

Furthermore, the connection between the kernels in the transport scattering rate and the quantum scattering rate also follows from Eqs. (68), (69), and (71). One can see from Eq. (68) that in the transport scattering rate not only is the contribution of forward scattering processes suppressed as in conventional systems but also the contribution of the backward scattering. The latter suppression is due to the Berry phase of π in graphene.

In the above derivation we have assumed $\delta \gg \tau_{\text{tr}}^{-1}$ which allowed us to extract τ_{tr}^{-1} from the expansion of the conductivity in τ_{tr}^{-1} . Using the generalized GR approach, we show in Appendix C that the expression for the transport scattering rate obtained in this way remains valid also for $\delta \rightarrow 0$.

B. Transport scattering rate: Results

Similarly to the quantum scattering rate, the transport rate is dominated by region 3 in regimes II, III, and IV. Again, this is not so for energies that are in regime II. In this case, we have an additional contribution from region 1, like in Sec. IV A. The calculation outlined in Appendix B yields

$$\frac{1}{\tau_{\text{tr}}(\epsilon)} \sim \begin{cases} \frac{T}{N} \sqrt{\frac{\epsilon}{T}}, & \text{I} \\ \alpha_g T, & \text{II} \\ \alpha_g^2 N T \left(\frac{T}{\epsilon}\right), & \text{III} \\ \alpha_g^2 N T \left(\frac{T}{\epsilon}\right)^2, & \text{IV.} \end{cases} \quad (72)$$

The characteristic momenta dominating the transport scattering rate as well as the relevant regions in the Q - Ω plane are shown in Table III. The transport scattering rate in regimes II and III is dominated by momenta much smaller than temperature, which allows us to find the numerical prefactors (given in Appendix B). Regimes I and IV are determined by momenta of order temperature, which does not allow us to find the numerical coefficient in these regimes. The analytical asymptotics are plotted alongside with the exact numerical result in Fig. 12.




FIG. 12. (Color online) Transport scattering rate for $\alpha_g N = 4 \times 10^{-3}$ (double logarithmic scale). Dots: exact values obtained by numerical evaluation; solid lines: analytical asymptotics, Eq. (72).

Similarly to Sec. IV C, we see a strong enhancement of the transport scattering rate in region II. When energy approaches temperature, the GR result

$$\tau_{\text{tr}}^{-1}(\epsilon \sim T) \sim \alpha_g^2 N T \quad (73)$$

is reproduced.

Comparing the transport scattering rate with the energy relaxation rate,

$$\frac{\tau_{\text{tr}}}{\tau_E} \propto \ln \frac{1}{\alpha_g N}, \quad (74)$$

we observe that in the limit of small α_g the relaxation of energy due to the inelastic collisions occurs much faster than the velocity relaxation, $\tau_{\text{tr}} \gg \tau_E$. The difference between the two rates comes from the fact that the forward scattering in graphene is strongly enhanced. The RPA screening suppresses the contribution of scattering angles smaller than α_g thus regularizing the logarithmically divergent contribution to the energy relaxation rate. On the other hand, the transport factor in \mathcal{K}_{tr} , see Eq. (68), kills the contribution of the forward scattering to the transport scattering rate much more efficiently than the RPA screening (this is the reason why the GR result for τ_{tr} is parametrically correct), so that no logarithmic factor arises in τ_{tr} .

The relation Eq. (74) justifies the hydrodynamic approach:^{58,60–62} the distribution functions of electrons and holes equilibrate within each type of carriers much faster than the direction of the velocity is changed. As a result, the distribution functions effectively depend only on the velocity direction. This means that the interaction-induced transport

TABLE III. Momentum/frequency scales and regions of the Q - Ω plane that dominate the transport scattering rate τ_{tr}^{-1} in different domains (I, II, III, and IV) of energy ϵ .

τ_{tr}	I	II	III	IV
$qv_F \sim \omega$	T	$\alpha_g^2 N^2 T / \epsilon$	ϵ	T
Regions	1, 2, 3, 4	1, 3	3	3

scattering rate entering the observables should be averaged over the temperature window:

$$\langle \tau_{\text{tr}}^{-1} \rangle = \frac{\int_0^\infty d\epsilon \rho(\epsilon) \frac{\partial f}{\partial \epsilon} \frac{1}{\tau_{\text{tr}}(\epsilon)}}{\int_0^\infty d\epsilon \rho(\epsilon) \frac{\partial f}{\partial \epsilon}}. \quad (75)$$

This energy-averaging procedure exactly corresponds to the solution of kinetic equation in Refs. 56 and 57; see Appendix D.

Evaluating the integral in Eq. (75) numerically with the exact expression for the RPA propagators, we find

$$\langle \tau_{\text{tr}}^{-1} \rangle \simeq 0.989 \alpha_g^2 NT. \quad (76)$$

The numerical prefactor in this result agrees with that found in Ref. 56 for the direct interaction term. As we will see below, this averaged value of the transport scattering rate can be substituted into the Drude expression for the conductivity, which yields the parametrically correct result for the collision-dominated conductivity of clean graphene.

C. Collision-limited conductivity

Above, when calculating the conductivity, we have assumed that the finite lifetime of quasiparticles is provided by some artificially introduced broadening δ , which mimics disorder. The introduction of the artificial lifetime allowed us to identify the contribution of the inelastic collisions to the transport scattering rate by considering the perturbative-in-interaction contributions to the conductivity. Let us now discuss the conductivity of clean graphene, or, more precisely, the conductivity of graphene in the regime when the inelastic collisions dominate over disorder scattering.

The conductivity of graphene at the Dirac point is a rather intricate quantity. In the absence of interaction, transport in the ballistic limit shows remarkable peculiarities in graphene.² The interplay between vanishing density of states and vanishing scattering rate leads to the nonuniversal conductivity that depends on the measurement details. In particular, the finite-size clean graphene sample of the “short-and-wide” geometry shows a behavior analogous to that of a normal diffusive metal. The zero- T conductivity of such a setup is $\sigma = 4 \times e^2/\pi h$. The same value of conductivity was predicted for an infinite sample with a large but finite electron lifetime. A different result, $\sigma = e^2/2h$, was found in the undoped graphene at a large frequency. At any nonzero value of the chemical potential, the conductivity of clean graphene is infinite. At finite temperature, energies within the temperature window contribute to the conductivity. This implies that the Dirac-point conductivity becomes infinite in the noninteracting case at any nonzero T .

As has been shown in Refs. 56–58, the conductivity of clean undoped graphene becomes finite due to the inelastic electron-electron collisions. The estimate for the collision-limited conductivity can be obtained by substituting the typical value of interaction-induced transport scattering time, Eq. (73), and the typical density of thermally populated states, $\rho(T) \sim NT/v_F^2$, into the Drude formula, which yields

$$\sigma = \frac{e^2}{h} \rho(T) v_F^2 \tau_{\text{tr}}(T) \sim \frac{e^2}{h} \frac{NT}{v_F^2} \frac{1}{\alpha_g^2 NT} \sim \frac{e^2}{h} \frac{1}{\alpha_g^2}. \quad (77)$$

Note that the explicit dependence on T , N , and v_F drops out from this formula; however, the temperature dependence appears in Eq. (77) implicitly through the renormalization of α_g . A more rigorous calculation of the collision-limited conductivity requires the analysis of the kinetic equation.^{56,57} Importantly, the fast energy relaxation discussed above not only simplifies such an analysis, but also reduces the kinetic equation to the hydrodynamic model.^{58–62}

The consideration of Sec. VA, which allowed us to find the transport scattering rate from the expression for the conductivity, relied on the perturbative treatment of the interaction. This assumes the following hierarchy of the energy scales:

$$\tau_{\text{tr}}^{-1} \ll \delta \ll T. \quad (78)$$

The first inequality implies that the broadening of the Green’s functions is due to the artificial “disorder” rate δ , whereas the second inequality establishes the ballistic regime which allows us to neglect the dressing of interaction by disorder. Since the characteristic frequency transfer in the transport scattering rate is of order of temperature, the resulting τ_{tr}^{-1} does not depend on δ as long as $\delta \ll T$. Furthermore, in Appendix C we have calculated τ_{tr}^{-1} from the generalized GR approach for $\delta = 0$ and reproduced the transport kernel \mathcal{K}_{tr} from Appendix B. Therefore we expect that the Drude formula, yielding Eq. (77), is applicable also for $\tau_{\text{tr}}^{-1} \gg \delta$, i.e., in the collision-dominated transport regime. This expectation is supported by the results of Refs. 56–58 and the comparison of our results with the solution of kinetic equation outlined in Appendix D.

Assuming the validity of the Drude formula for $\delta \ll \tau_{\text{tr}}^{-1}$, we evaluate the conductivity using the energy-averaged transport scattering rate given by Eq. (76):

$$\sigma = \frac{e^2}{h} \frac{N\pi}{\langle \tau_{\text{tr}}^{-1} \rangle} \int_0^\infty d\epsilon \rho(\epsilon) \frac{\partial f}{\partial \epsilon} \approx \frac{e^2}{h} \frac{0.7}{\alpha_g^2}. \quad (79)$$

This result agrees with that found in Refs. 56 and 57 within the kinetic equation approach [note that in Eq. (79) we included only the contribution of the direct interaction, which survives the large- N limit].

It is worth noting here that the kinetic approach of Refs. 56 and 57 was based on the GR calculation of the self-energies. As we have seen in Sec. VA, the RPA result for the transport scattering rate at relevant energies $\epsilon \sim T$ [as well as the energy-averaged characteristic rate, Eq. (75)] has the same form as given by the GR. Moreover, the averaging of the energy-dependent transport scattering rate using Eq. (75) yields the same numerical prefactor as follows from the kinetic equation with the GR collision integral,^{56,57}; see Eq. (76). Therefore we conclude that the RPA resummation does not change the numerical prefactor of conductivity in the leading order in α_g^2 . In other words, unlike the quantum scattering (or dephasing) rate considered in Secs. IVA 2 and IVB, the calculation of the conductivity can be done within the golden-rule approach. The RPA resummation yields only subleading-in- α_g corrections to the conductivity, as explained in Appendix D.




FIG. 13. (Color online) Schematic plot (only parametrical scales are given) of the temperature dependence of the Drude conductivity (in units of e^2/h ; solid line) $\alpha_g \ll 1/\ln(\Lambda\tau_{\text{dis}})$, where Λ is the bandwidth. For simplicity, the logarithmic temperature corrections to α_g , which comes from the renormalization, is not shown. Dashed lines: low- T behavior of the conductivity governed by quantum interference effects; depending on the character of disorder, the localization, antilocalization, or criticality (coincides with the solid line) may occur. Dash-dotted line: expected high- T behavior governed by electron-phonon scattering.

Finally, let us discuss on the qualitative level the effect of disorder on the Dirac-point conductivity in graphene. For simplicity we set $N = 1$ below. In the absence of interaction, effect of disorder on transport in the ballistic regime is highly unconventional and strongly depends on the type of randomness.^{13,14,18,75} On the other hand, as we have seen above, the interaction effects are crucially important for the transport already in a clean system. It is thus important to explore transport in realistic graphene structures, with both the electron-electron interaction and disorder taken into account.

The role of disorder here is twofold: (i) potential disorder introduces velocity relaxation, thus contributing to the transport scattering rate,

$$\frac{1}{\tau_{\text{tr}}} \sim \alpha_g^2 T + \frac{1}{\tau_{\text{dis}}}, \quad (80)$$

and (ii) establishes a finite density of states already in the Dirac point,

$$\rho \propto T + \frac{1}{\tau_{\text{dis}}}. \quad (81)$$

Substituting these formulas into the Drude conductivity, we obtain the following result describing the crossover between the collision-dominated and disorder-dominated regimes:

$$\sigma \sim \frac{e^2}{h} \frac{T + \frac{1}{\tau_{\text{dis}}}}{\alpha_g^2 T + \frac{1}{\tau_{\text{dis}}}}. \quad (82)$$

This expected temperature dependence of the Drude conductivity is shown in Fig. 13 for the case of weak interaction (or strong disorder), $\alpha_g \ll 1/\ln(\Lambda\tau_{\text{dis}})$ when the renormalization




FIG. 14. (Color online) Schematic plot of the temperature dependence of the Drude conductivity (in units of e^2/h) for strong interaction, $\alpha_g \gg 1/\ln(\Lambda\tau_{\text{dis}})$. The characteristic scale T_0 is given by $T_0 \sim \tau_{\text{dis}}^{-1} \ln^2(\Lambda/T_0) \sim \tau_{\text{dis}}^{-1} \ln^2(\Lambda\tau_{\text{dis}})$. For possible deviations at low and high T (interference and phonon contributions, respectively), see Fig. 13.

of α_g gives only small logarithmic corrections. For stronger interaction (or weaker disorder), $\alpha_g \ll 1/\ln(\Lambda\tau_{\text{dis}})$, the renormalization of

$$\alpha_g(T) = \frac{\alpha_g}{1 + \frac{\alpha_g}{4} \ln \frac{\Lambda}{T}}$$

becomes strong [so that the renormalized coupling “forgets” about its bare value, $\alpha_g(T) \rightarrow 4/\ln(\Lambda/T)$] already in the collision-dominated regime; see Fig. 14. One sees that, since the interaction-induced transport rate contains α_g^2 whereas the density of states of thermally excited quasiparticles does not, the two crossover T scales appear, which establishes an intermediate regime of the ballistic transport,

$$\frac{1}{\tau_{\text{dis}}} \ll T \ll \frac{1}{\alpha_g^2 \tau_{\text{dis}}}.$$

Again, a more rigorous derivation of the conductivity in the presence of both disorder and inelastic scattering is based on kinetic equation approach and will be performed elsewhere. Note that we neglected the phonon contribution to the relaxation rates (for estimate of their contribution, see, e.g., Ref. 24) which becomes relevant at sufficiently high temperatures.

At $T \sim 1/\tau_{\text{dis}}$ the Drude conductivity becomes of the order of conductance quantum and the dephasing rate becomes of the order of T . At lower temperatures, the T dependence of the conductivity is governed by interference effects: localization, antilocalization, or critical behavior may occur, depending on the symmetry of disorder.^{16,17,76} The crossover scale $1/\tau_{\text{dis}}$ in typical experiments on high-quality graphene is in the range $T \sim 1\text{--}100$ K.

VI. CONCLUSIONS

In conclusion, we have analyzed the inelastic electron-electron scattering in graphene using the Keldysh diagrammatic

approach. We have demonstrated that finite temperature strongly affects the screening properties of graphene. This, in turn, dramatically influences the inelastic-scattering rates as compared to the zero-temperature case. We have calculated the finite- T quantum scattering rate, see Eq. (60) and Fig. 6, which is relevant for dephasing of interference processes. We have identified a hierarchy of regimes, Eq. (5), arising due to the interplay of a plasmon enhancement of the scattering and finite-temperature screening of the interaction. The lifetime of quasiparticles with energies close to the Dirac point has been found to be independent of the coupling constant. We have further calculated the energy relaxation rate, Eq. (64) and Fig. 8, and transport scattering rate, Eq. (72) and Fig. 12. For all the three rates, we have found a nonmonotonic energy dependence which has been attributed to the resonant excitation of plasmons. Finally, we have discussed the collision-limited conductivity of clean graphene as well as the expected behavior of the high-temperature conductivity in the presence of disorder, see Eqs. (79) and (82), respectively. Our results complement the kinetic equation and hydrodynamic approaches for the collision-limited conductivity.

Our approach, which employs the Keldysh formalism, can be generalized for the treatment of physics of inelastic processes in strongly nonequilibrium setups. In particular, this framework is expected to allow us to investigate interaction effects on full counting statistics of the electron transport in graphene and to develop the theory

of tunneling spectroscopy in a strongly biased graphene setup.

ACKNOWLEDGMENTS

We thank S. Carr, L. Fritz, M. Müller, S. Ngo Dinh, and M. Titov for interesting and useful discussions. The work was supported by the Center for Functional Nanostructures of the Deutsche Forschungsgemeinschaft (DFG), by SPP ‘‘Graphene’’ of the DFG, by Rosnauka Grant No. 02.740.11.5072, and by the EUROHORCS/ESF EURYI Award scheme (I.V.G.).

APPENDIX A: POLARIZATION OPERATOR

In this appendix we evaluate the polarization operator, Eq. (34), at finite temperature. In a scattering process with emitting a photon, three different momenta are involved, which form in two dimensions triangles, as shown in Fig. 15. Specifically, if an electron before scattering has the momentum \mathbf{p} and the emitted photon carries momentum \mathbf{q} , the electron that is left over has to carry $\mathbf{p} - \mathbf{q}$. The angular integration over the transferred momentum \mathbf{q} becomes complicated. To proceed further it is convenient to choose elliptic coordinates defined by $\xi = p + |\mathbf{p} - \mathbf{q}|$ and $\eta = p - |\mathbf{p} - \mathbf{q}|$. The corresponding coordinate system is shown in Fig. 15. Using the elliptic coordinates, the expressions for the imaginary and real parts of the polarization bubble take the form

$$\text{Im } \Pi^R = \frac{\sinh(Q\beta)}{2\pi} \text{Re} \left[\frac{Q}{\sqrt{\beta^2 - 1}} \int_0^1 d\eta \frac{\sqrt{1 - \eta^2}}{\cosh(Q\beta) + \cosh(\eta Q)} + \frac{Q}{\sqrt{1 - \beta^2}} \int_1^\infty d\xi \frac{\sqrt{\xi^2 - 1}}{\cosh(Q\beta) + \cosh(\xi Q)} \right], \quad (\text{A1})$$

$$\text{Re } \Pi^R = -\frac{Q}{\pi^2} \int_1^\infty d\xi \int_0^1 d\eta \frac{\sinh(\xi Q) \sqrt{\frac{1 - \eta^2}{\xi^2 - 1}} \frac{\xi}{\beta^2 - \xi^2} + \sinh(\eta Q) \sqrt{\frac{\xi^2 - 1}{1 - \eta^2}} \frac{\eta}{\beta^2 - \eta^2}}{\cosh(\xi Q) + \cosh(\eta Q)}. \quad (\text{A2})$$

Here and below we introduce the notation $\beta = \Omega/Q$ to decouple expansions in small or large Q from the behavior at the singularity $Q = \Omega$. For simplicity we set $v_F = 1$ and $T = 1$. The integrals in Eqs. (A1) and (A2) are evaluated separately in the four regions shown in Fig. 3.

a. Region 1

In region 1 the condition $Q \ll 1$ and $\beta > 1$ hold, which means that in Eq. (A1) only the part with the η integral is left. Expanding the integrands in small $\eta Q \ll 1$, we get

$$\begin{aligned} \text{Im } \Pi^R &\simeq \frac{Q \sinh(Q\beta)}{2\pi \sqrt{\beta^2 - 1}} \int_0^1 d\eta \frac{\sqrt{1 - \eta^2}}{\cosh(Q\beta) + 1} \\ &= \frac{Q \tanh(Q\beta/2)}{8 \sqrt{\beta^2 - 1}} \end{aligned} \quad (\text{A3})$$

for the imaginary part and

$$\begin{aligned} \text{Re } \Pi^R &\simeq \frac{-Q}{\pi^2} \int_0^1 \frac{d\eta}{\sqrt{1 - \eta^2}} \frac{Q\eta^2}{\beta^2 - \eta^2} \int_1^\infty d\xi \frac{\sqrt{\xi^2 - 1}}{\cosh(\xi Q) + 1} \\ &\simeq -\frac{\ln 2}{\pi} \left(\frac{|\beta|}{\sqrt{\beta^2 - 1}} - 1 \right) \end{aligned} \quad (\text{A4})$$

for the real part of Π^R . The term $\propto \sinh(\xi Q)$ in Eq. (A2) yields a contribution of the order of $\sim Q^2$. The second term $\propto \sinh(\eta Q)$ results in Eq. (A4).

b. Region 2

In region 2, we have $Q \gg 1$ and $\beta > 1$. In Eq. (A1) again only the η integral is left and by neglecting $\cosh(\eta Q)$ and




FIG. 15. Sketch of elliptic coordinates.

expanding in large Q we get

$$\text{Im } \Pi^R \simeq \frac{1}{2\pi} \frac{\sinh(Q\beta)}{\sqrt{\beta^2 - 1}} \int_0^1 d\eta \frac{\sqrt{1 - \eta^2}}{\cosh(Q\beta)} = \frac{1}{8} \frac{Q \text{sgn } \beta}{\sqrt{\beta^2 - 1}}. \quad (\text{A5})$$

By expanding Eq. (A2) in large Q and resolving exponentials in the denominator by a geometric series, we see that the leading contribution comes from $\eta \simeq 1$ ($y = 1 - \eta \simeq 0$). A competition of Q and $\Omega^2 - Q^2$ arises and the limit $\Omega^2 - Q^2 \ll Q$ gives the relevant contribution. The corresponding principal value integral reads

$$\begin{aligned} \text{Re } \Pi^R &\simeq -Q^{-1/2} \sum_{n=1}^{\infty} \frac{(-1)^n}{(n\pi)^{3/2}} \underbrace{\int_0^{\infty} \frac{d\sqrt{2y} \sqrt{2y^2} e^{-Qn\sqrt{2y^2}/2}}{(\beta^2 - 1)^2 - \sqrt{2y^4}}}_{\simeq -\frac{\pi}{4} \frac{1}{\sqrt{\beta^2 - 1}}} \\ &\simeq -\frac{1}{4\sqrt{\pi}} \frac{1}{\sqrt{Q(\beta^2 - 1)}} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^{3/2}}. \end{aligned} \quad (\text{A6})$$

c. Region 3

Region 3 is characterized by the conditions $Q \ll 1$ and $\beta < 1$. In this region, only the ξ integral contributes to Eq. (A1). By expanding the integrand in small $Q\beta$ we get

$$\begin{aligned} \text{Im } \Pi^R &\simeq \frac{Q\beta}{2\pi} \frac{Q}{\sqrt{1 - \beta^2}} \int_1^{\infty} d\xi \frac{\sqrt{\xi^2 - 1}}{1 + \cosh(\xi Q)} \\ &\simeq \frac{\ln 2}{\pi} \text{Re} \frac{\beta}{\sqrt{1 - \beta^2}}. \end{aligned} \quad (\text{A7})$$

The simplification of Eq. (A2) in region 3 is similar to that leading to Eq. (A4). The contribution of the first term in

Eq. (A2) is of order $Q \ln Q$, yielding

$$\begin{aligned} \text{Re } \Pi^R &\simeq -\frac{Q}{\pi^2} \int_0^1 \frac{d\eta}{\sqrt{1 - \eta^2}} \frac{Q\eta^2}{\beta^2 - \eta^2} \int_1^{\infty} d\xi \frac{\sqrt{\xi^2 - 1}}{\cosh(\xi Q) + 1} \\ &\simeq \frac{\ln 2}{\pi}. \end{aligned} \quad (\text{A8})$$

d. Region 4

Finally, in region 4 the conditions $Q \gg 1$ and $\beta < 1$ are fulfilled. In Eq. (A1) only the ξ integral contributes and by neglecting $\cosh(Q\beta)$ and expanding the integrand in large $Q\xi$ we get

$$\begin{aligned} \text{Im } \Pi^R &\simeq \frac{\sinh(Q\beta)}{\pi} \frac{Q}{\sqrt{1 - \beta^2}} \int_1^{\infty} d\xi \underbrace{\sqrt{\xi^2 - 1} e^{-Q\xi}}_{\simeq e^{-Q} \frac{1}{Q^{3/2}} \sqrt{\frac{\pi}{2}}} \\ &\simeq \frac{\sinh(Q\beta)}{\sqrt{\beta^2 - 1}} \frac{e^{-Q}}{\sqrt{2\pi} Q}. \end{aligned} \quad (\text{A9})$$

In region 4 the simplification of Eq. (A2) differs from that in region 2 in one important point. By neglecting $\cosh(Q\eta)$ in comparison to $\cosh(Q\xi)$, the first term $[\propto \sinh(Q\xi)]$ yields no principle value integral, while β is smaller than 1. This results in

$$\begin{aligned} \text{Re } \Pi^R &\simeq -\frac{Q}{\pi^2} \int_1^{\infty} d\xi \frac{\xi}{\sqrt{\xi^2 - 1}(\beta^2 - \xi^2)} \int_0^1 d\eta \sqrt{1 - \eta^2} \\ &= \frac{1}{8} \frac{Q}{\sqrt{1 - \beta^2}}. \end{aligned} \quad (\text{A10})$$

APPENDIX B: CALCULATING THE RATES

In this appendix we calculate the integrals for inelastic scattering rates.

1. Definitions of the rates

The rates we are interested in are defined by inserting the kernel \mathcal{K}_j into the integrand for the imaginary part of the total self-energy, Eq. (54) ($j = q, +, E, \text{tr}$):

- (i) Total quantum scattering rate ($j = q$),

$$\mathcal{K}_q = 1;$$

- (ii) Energy relaxation rate ($j = E$),

$$\mathcal{K}_E = \frac{\omega^2}{T^2};$$

- (iii) Transport scattering rate ($j = \text{tr}$),

$$\mathcal{K}_{\text{tr}} = \frac{1}{2} \sin^2 \theta = \frac{1}{2} \frac{q^2 \sin^2 \gamma}{\epsilon^2 + q^2 - 2\epsilon q \cos \gamma}.$$

Here θ is the angle between incoming particle and outgoing particle and γ is the angle between \mathbf{q} and \mathbf{p} . The origin of \mathcal{K}_{tr} is explained in Sec. VA. The kernel for the chiral scattering rate τ_{\pm}^{-1} depends on the scattering channel. In particular,

for + chirality at $\epsilon > 0$ the electron-electron-scattering kernel contains $\cos^2(\theta/2)$ and the electron-hole scattering kernel contains $\sin^2(\theta/2)$ due to Dirac factors.

It is convenient to introduce the dimensionless energy $y = \epsilon/2T$. The integrals we have to handle are of the following form:

$$\tau_j^{-1}(y) = -\frac{2T^2}{v_F^2} \int_0^\infty \frac{dQ}{2\pi} Q \int_0^{2\pi} \frac{d\gamma}{2\pi} \sum_{\Omega=y \pm \sqrt{y^2+Q^2-2Qy\cos\gamma}} \{ \mathcal{K}_j(\Omega, Q, y) \text{Im} D_{\text{RPA}}^R(\Omega, Q) [\coth(\Omega) + \tanh(y - \Omega)] \}. \quad (\text{B1})$$

Since the combinations $y \pm \sqrt{y^2 + Q^2 - 2Qy \cos \gamma}$ lead to complicated integrands, we split the integrals into the parts corresponding to $Q \ll y$ and $y \ll Q$. Here γ is the angle between the transferred momentum \mathbf{q} and the initial momentum \mathbf{p} .

A. Simplifying the integrand

The imaginary part of the interaction propagator is given by

$$\begin{aligned} & \text{Im} D_{\text{RPA}}^R(\Omega, Q) \\ &= -\frac{D_0^2(Q) N \text{Im} \Pi(\Omega, Q)}{[1 + D_0(Q) N \text{Re} \Pi(\Omega, Q)]^2 + [D_0(Q) N \text{Im} \Pi(\Omega, Q)]^2}, \end{aligned} \quad (\text{B2})$$

where

$$D_0(Q) = \frac{v_F^2 \alpha_g \pi}{T Q} \quad (\text{B3})$$

is the bare Coulomb interaction.

The splitting of the Q integral leads to the following simplification:

$$\begin{aligned} \Omega &\simeq y \pm \begin{cases} y - Q \cos \gamma & Q \ll y \\ Q - y \cos \gamma & y \ll Q \end{cases}, \\ \sqrt{|Q^2 - \Omega^2|} &\simeq \begin{cases} 2y & + \\ Q |\sin \gamma| & - \\ 2\sqrt{yQ} |\sin \frac{\gamma}{2}| & + \\ 2\sqrt{yQ} |\cos \frac{\gamma}{2}| & - \end{cases} \begin{matrix} Q \ll y \\ y \ll Q \end{matrix}. \end{aligned} \quad (\text{B4})$$

Here +, - correspond to the two possible values for Ω that appear in Eq. (B1). These signs reflect the separation of integration domains into the parts above and below the mass-shell line $Q = \Omega$: + corresponds to $\Omega > Q$ and - corresponds to $\Omega < Q$. Below we simplify the integrands $\mathcal{I} = [g + f] \mathcal{K}_j \text{Im} D_{\text{RPA}}^R$ separately in each of the region in the Q vs Ω plane.

a. Region 1

In region 1, the dominant contribution to all the rates comes from the domain $y \ll Q$. Using the simplified polarization

operator, Eq. (43), we obtain

$$\begin{aligned} \mathcal{I}_{1,2}^j(y, Q, \gamma) &\simeq \frac{v_F^2}{T} \frac{1}{N} \frac{\pi^2}{8} \frac{\sqrt{yQ} |\sin \frac{\gamma}{2}|}{\left(\frac{2\sqrt{yQ}}{\alpha_g N} |\sin \frac{\gamma}{2}| - \ln 2 \right)^2 + \left(\frac{\pi}{16} Q^2 \right)^2} \\ &\times \begin{cases} 1, & j = q \\ 2 \cos^2 \frac{\gamma}{2}, & j = + \\ 4Q^2, & j = E \\ \frac{1}{2} \sin^2 \gamma, & j = \text{tr}. \end{cases} \end{aligned} \quad (\text{B5})$$

Here and below, the first digit in the subscript of the function \mathcal{I} denotes the region in the Q vs Ω plane, while the second digit is 1 for $Q \ll y$ and 2 for $Q \gg y$.

b. Region 2

The contribution to all the rates coming from region 2 is exponentially small in regime IV. In other regimes, it contains at least an extra α_g as compared to the contributions of region 3, except for the situations when results for the rate are determined by momenta q of order T/v_F . In this situation, it turns out that the asymptotics produced by region 2 is the same as the asymptotics of region 3. We remind the reader that finding the numerical value of the prefactor is beyond our analytical approach when integrals are dominated by $q \sim T/v_F$. Therefore there is no case where we need to calculate the contribution of region 2.

c. Region 3

Region 3 appears to be the most important region because most of the final results for the rates are determined by this region. In this region both small and large y compared to Q are important:

(i) $y \gg Q$:

$$\begin{aligned} \mathcal{I}_{3,1}^j(y, Q, \gamma) &\simeq \frac{v_F^2}{NT} \frac{\pi}{\ln 2} \frac{|\sin \gamma| Q^{-1}}{\left[\left(\frac{Q}{\alpha_g N \ln 2} + 1 \right)^2 - 1 \right] \sin^2 \gamma + 1} \\ &\times \begin{cases} 1, & j = q \\ 2, & j = + \\ 4Q^2 \cos^2 \gamma, & j = E \\ \frac{1}{2} \frac{Q^2}{y^2} \sin^2 \gamma, & j = \text{tr}; \end{cases} \end{aligned} \quad (\text{B6})$$

(ii) $y \ll Q$:

$$\mathcal{I}_{3,2}^j(y, Q, \gamma) \simeq \frac{v_F^2}{T} \frac{1}{N} \frac{2\pi}{\ln 2} \sqrt{y} \frac{Q^{-3/2} \cos \frac{\gamma}{2}}{\frac{4y}{Q} \left(\frac{Q}{\alpha_g N \ln 2} + 1 \right)^2 \cos^2 \frac{\gamma}{2} + 1} \times \begin{cases} 1, & j = q \\ 2 \sin^2 \frac{\gamma}{2}, & j = + \\ 4Q^2, & j = E \\ \frac{1}{2} \sin^2 \gamma, & j = \text{tr}. \end{cases} \quad (\text{B7})$$

d. Region 4

This region is only important for the energy relaxation rate ($j = E$). The corresponding contribution is governed by $Q \ll y$, so that only the energy range $y \gg 1$ is of interest, where

$$\mathcal{I}_{4,1}^E(y, Q, \gamma) \simeq \frac{v_F^2}{T} \frac{\alpha_g^2 \pi^2 4N}{\sqrt{2\pi Q}} \frac{\sin \gamma \cos^2 \gamma}{\left(\sin \gamma + \frac{\alpha_g \pi N}{8} \right)^2}. \quad (\text{B8})$$

B. Results for the rates

Finally we estimate the rates in the regimes I, II, III, and IV as defined in Fig. 5, using the simplified integrands of the previous part.

a. Region 1

The contributions of Region 1 to the quantum and transport scattering rates are only relevant for energies in regime II for $y \ll Q$, where we have

$$\tau_j^{-1}(y) = \frac{2T^2}{v_F^2} \int_{\min(y,1)}^1 \frac{dQ}{2\pi} Q \int_0^{2\pi} \frac{d\gamma}{2\pi} \mathcal{I}_{1,2}^j(y, Q, \gamma). \quad (\text{B9})$$

Performing first the angular integration over γ , we obtain the quantum scattering rate (see Table IV),

$$\tau_q^{-1}(y) \simeq 4\alpha_g T \int_0^1 da \frac{1}{\sqrt{1-a^2}} \simeq 2\pi\alpha_g T, \quad (\text{B10})$$

the chiral scattering rate (see Table V),

$$\tau_+^{-1}(y) \simeq 4\alpha_g T \int_0^1 da \sqrt{1-a^2} \simeq \pi\alpha_g T, \quad (\text{B11})$$

and the transport scattering rate (see Table VI),

$$\tau_{\text{tr}}^{-1}(y) \simeq 8\alpha_g T \int_0^1 da a^2 \sqrt{1-a^2} \simeq \frac{\pi}{2} \alpha_g T. \quad (\text{B12})$$

TABLE IV. Contribution of region 1 to the total quantum scattering rate τ_q^{-1} in regime II and the characteristic values of the transferred momentum Q , transferred frequency Ω , the angle γ between momenta \mathbf{p} and \mathbf{q} , and the scattering angle θ (the results here and in tables below are symmetric with respect to $\gamma \rightarrow -\gamma$ and $\theta \rightarrow -\theta$).

Regime	$\tau_q^{-1}(y)$	Q	Ω	γ	θ
II	$\simeq 2\pi\alpha_g T$	$\alpha_g^2 N^2 / y$	Q	$0 \leq \gamma \leq \pi$	$0 \leq \theta \leq \pi$

TABLE V. Contribution of region 1 to the chiral quantum scattering rate τ_+^{-1} in regime II and the characteristic values of Q , Ω , γ , and θ dominating this contribution.

Regime	$\tau_+^{-1}(y)$	Q	Ω	γ	θ
II	$\simeq 2\pi\alpha_g T$	$\alpha_g^2 N^2 / y$	Q	$0 < \gamma < \pi$	$0 < \theta < \pi$

Here the integrals over the variable $a = \alpha_g N \ln 2 / \sqrt{4yQ}$ correspond to the Q integration.

In all the regimes, the contribution of region 1 to the energy relaxation rate is parametrically the same as that of other regions. Since the numerical prefactor is not accessible within our calculation (the integrals are dominated by $qv_F \sim T$), we have chosen to present only the calculation of the contribution of region 3.

b. Region 2

There is no important contribution from region 2 (see Sec. VI B2 above).

c. Region 3

This is the most important region for all the rates in most of the regimes. The corresponding integrals are expressed through the kernels $\mathcal{I}_{3,i}^j$ introduced in Sec. VI as follows:

$$\tau_j^{-1}(y) = \frac{2T^2}{v_F^2} \int_0^{\min(y,1)} \frac{dQ}{2\pi} Q \int_0^{2\pi} \frac{d\gamma}{2\pi} \mathcal{I}_{3,1}^j(y, Q, \gamma) + \frac{2T^2}{v_F^2} \int_{\min(y,1)}^1 \frac{dQ}{2\pi} Q \int_0^{2\pi} \frac{d\gamma}{2\pi} \mathcal{I}_{3,2}^j(y, Q, \gamma). \quad (\text{B13})$$

Using the shorthand notations

$$x' = \frac{1}{\alpha_g N \ln 2}, \quad y' = \frac{y}{\alpha_g N \ln 2},$$

we obtain the following result for the contribution of region 3 to the quantum scattering rate (see Table VII):

$$\tau_q^{-1}(y) \simeq \frac{\alpha_g}{\pi} T \left\{ \int_0^{\min(x',y')} dx \frac{\text{arsinh}[\sqrt{x(x+2)}]}{\sqrt{x(x+2)}(x+1)} + \int_{\min(x',y')}^{x'} dx \frac{\text{arsinh}[\sqrt{4y'(\sqrt{x} + \frac{1}{\sqrt{x}})}]}{(1+x)\sqrt{1+4y'(\sqrt{x} + \frac{1}{\sqrt{x}})^2}} \right\}. \quad (\text{B14})$$

TABLE VI. Contribution of region 1 to the transport scattering rate τ_{tr}^{-1} in regime II and the characteristic values of Q , Ω , γ , and θ dominating this contribution.

Regime	$\tau_{\text{tr}}^{-1}(y)$	Q	Ω	γ	θ
II	$\simeq \pi\alpha_g T/2$	$\alpha_g^2 N^2 / y$	Q	$0 < \gamma < \pi$	$0 < \theta < \pi$

TABLE VII. Contribution of region 3 to the total quantum scattering rate τ_q^{-1} in all energy domains and the corresponding characteristic values of the transferred momentum Q , transferred frequency Ω , the angle γ between momenta \mathbf{p} and \mathbf{q} , and the scattering angle θ .

Regime	$\tau_q^{-1}(y)$	Q	Ω	γ	θ
I	$\sim \frac{\sqrt{y}}{N} T$	1	$-Q$	$0 < \gamma < \pi$	$0 < \theta < \pi$
II	$\simeq \pi \alpha_g T$	$\alpha_g^2 N^2 / y$	$-Q$	$0 < \gamma < \pi$	$0 < \theta < \pi$
III	$\simeq \frac{4\beta(2)}{\pi} \alpha_g T$	$\alpha_g N T$	$Q \cos \gamma$	$0 < \gamma < \pi$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$
IV	$\simeq \frac{4\beta(2)}{\pi} \alpha_g T$	$\alpha_g N T$	$Q \cos \gamma$	$0 < \gamma < \pi$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$

For the contribution of region 3 to the chiral scattering rate (see Table VIII) we get

$$\tau_+^{-1}(y) \simeq \frac{\alpha_g}{\pi} T \left\{ \int_0^{\min(x', y')} (x', y') dx \frac{\text{arsinh}[\sqrt{x(x+2)}]}{\sqrt{x(x+2)}(x+1)} + \int_{\min(x', y')}^{x'} dx \frac{\sqrt{1+4y'(\sqrt{x+\frac{1}{\sqrt{x}}})^2} \text{arsinh}[\sqrt{4y'(\sqrt{x+\frac{1}{\sqrt{x}})}]} - \sqrt{4y'(\sqrt{x+\frac{1}{\sqrt{x}})}]}{(1+x)^2 \frac{4y'}{x}} \right\}. \quad (\text{B15})$$

The contribution of region 3 to the energy relaxation rate (see Table IX) reads

$$\tau_E^{-1}(y) \simeq \frac{\alpha_g}{\pi(x')^2} T \left\{ \int_0^{\min(x', y')} dx \frac{x}{x+2} \left[\frac{\text{arsinh}(\sqrt{x(x+2)})}{(x+1)\sqrt{x(x+2)}} - 1 \right] + \int_{\min(x', y')}^{x'} dx \frac{x^2 \text{arsinh}[\sqrt{4y'(\sqrt{x+\frac{1}{\sqrt{x}})}]}]}{(1+x)\sqrt{1+4y'(\sqrt{x+\frac{1}{\sqrt{x}}})^2}} \right\}. \quad (\text{B16})$$

Finally, region 3 yields the following result for the transport scattering rate (see Table X) τ_{tr}^{-1} :

$$\tau_{\text{tr}}^{-1}(y) \simeq \frac{\alpha_g}{2\pi y^2} T \int_0^{\min(x', y')} dx \frac{x}{x+2} \left[1 - \frac{\text{arsinh}(\sqrt{x(x+2)})}{(x+1)\sqrt{x(x+2)}} \right] + \frac{4}{\pi} \alpha_g T \int_{\min(x', y')}^{x'} dx \sqrt{\frac{y'}{x}} \times \left[\frac{z^2+3}{3z^4} - \frac{\text{arsinh}z}{z^5(1+z^2)^{-1/2}} \right]_{z=\sqrt{4y'(\sqrt{x+\frac{1}{\sqrt{x}})}}}. \quad (\text{B17})$$

a. Region 4

This region is only relevant for the energy relaxation rate in regime IV, where the main contribution comes from $Q \ll y$,

yielding

$$\tau_E^{-1}(y) = \frac{8T^2}{v_F^2} \int_1^{\max(y, 1)} \frac{dQ}{2\pi} Q \times \int_0^{\pi/2-1/Q} \frac{d\gamma}{2\pi} T_{4,1}^{E, \gamma \ll \pi/2-1/Q}(y, Q, \gamma) \simeq \frac{12N}{\sqrt{2\pi}} \alpha_g^2 T y^{3/2} \ln\left(\frac{16}{N\pi\alpha_g\sqrt{y}}\right). \quad (\text{B18})$$

Since the result is determined by $Q \sim \Omega \sim y \gg 1$, the energy-diffusion model is, in fact, not applicable in this regime.

TABLE VIII. Contribution of region 3 to the chiral quantum scattering rate τ_+^{-1} in all energy domains and the corresponding characteristic values of Q , Ω , γ , and θ .

Regime	$\tau_+^{-1}(y)$	Q	Ω	γ	θ
I	$\sim \frac{\sqrt{y}}{N} T$	1	$-Q$	$0 < \gamma < \pi$	$0 < \theta < \pi$
II	$\simeq \pi \alpha_g T$	$\alpha_g^2 N^2 / y$	$-Q$	$0 < \gamma < \pi$	$0 < \theta < \pi$
III	$\simeq \frac{8\beta(2)}{\pi} \alpha_g T$	$\alpha_g N T$	$Q \cos \gamma$	$0 < \gamma < \pi$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$
IV	$\simeq \frac{8\beta(2)}{\pi} \alpha_g T$	$\alpha_g N T$	$Q \cos \gamma$	$0 < \gamma < \pi$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$

TABLE IX. Contribution of region 3 to the energy relaxation rate τ_E^{-1} obtained in all energy domains within the energy-diffusion approximation and the corresponding characteristic values of Q , Ω , γ , and θ . Since in all energy domains the result is dominated by $Q \sim 1$, the numerical prefactors can not be obtained from the asymptotics at $Q \ll 1$ and $Q \gg 1$.

Regime	$\tau_E^{-1}(y)$	Q	Ω	γ	θ
I	$\sim \frac{\sqrt{y}}{N} T$	1	$-Q$	$0 < \gamma < \pi$	$0 < \theta < \pi$
II	$\sim \alpha_g^2 T \frac{N}{\sqrt{y}} \ln(\frac{y}{\alpha_g^2 N^2})$	1	$-Q$	$\pi - \frac{\alpha_g N}{\sqrt{y}}$	$\frac{\alpha_g N}{\sqrt{y}} \ll 1$
III	$\sim \alpha_g^2 T \frac{N}{\sqrt{y}} \ln(\frac{y}{\alpha_g^2 N^2})$	1	$-Q$	$\pi - \frac{\alpha_g N}{\sqrt{y}}$	$\frac{\alpha_g N}{\sqrt{y}} \ll 1$
IV	$\sim \alpha_g^2 N T \ln(\frac{1}{\alpha_g^2 N^2})$	1	$-Q$	$\pi - \frac{\alpha_g N}{\sqrt{y}}$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$

APPENDIX C: SCATTERING RATES FROM THE GENERALIZED GOLDEN RULE

In this appendix we analyze the golden rule (GR) approach to calculating quantum and transport scattering rates. The quantum scattering rate for electrons was introduced in Eq. (55). Using the imaginary part of the self-energy Eq. (24), we obtain

$$\begin{aligned}
\tau_+^{-1}(\epsilon) &\propto \int d\mathbf{q} \int d\omega |D_{\text{RPA}}(q, \omega)|^2 [1 - n_F(\epsilon - \omega)] \\
&\times \int d\mathbf{p} \text{Tr} [\mathcal{P}_+(\mathbf{p}) \delta(\epsilon - v_F p) \mathcal{A}_0(\epsilon - \omega, \mathbf{p} - \mathbf{q})] \\
&\times \int dE [n_F(E - \omega/2) - n_F(E + \omega/2)] \\
&\times \int d\mathbf{k} \text{Tr} [\mathcal{A}_0(E - \omega/2, \mathbf{k} - \mathbf{q}/2) \\
&\times \mathcal{A}_0(E + \omega/2, \mathbf{k} + \mathbf{q}/2)]. \quad (\text{C1})
\end{aligned}$$

The spectral weights \mathcal{A}_0 are defined in Eq. (6). Equation (C1) is completely equivalent to Eq. (46) of the main text.

We can regard the quantum scattering rate as the probability (per unit time) of the electron decay with emitting an electron-hole pair. The amplitude of this decay process is given by the ‘‘half’’ of the self-energy diagram shown in Fig. 16. The incoming and outgoing particles in this diagram are taken at the mass shell. Equation (C1) has the form of the Fermi golden rule with the amplitude determined by the RPA-screened interaction.

We now apply the GR to calculate the transport scattering rate. This amounts to including an extra transport factor $[\mathbf{v}_i \cdot (\mathbf{v}_i - \mathbf{v}_f)]/v_F^2$ accounting for the change of the current due to scattering, in the integrand of Eq. (C1). Here \mathbf{v}_i and \mathbf{v}_f are total

velocities of incoming and outgoing particles. For the linear electronic dispersion the velocities of individual quasiparticles are determined by the relation

$$\mathbf{v} = \frac{\partial \epsilon}{\partial \mathbf{p}} = \frac{v_F^2 \mathbf{p}}{\epsilon}. \quad (\text{C2})$$

This results in the following transport factor:

$$\begin{aligned}
&\frac{\mathbf{v}_i \cdot (\mathbf{v}_i - \mathbf{v}_f)}{v_F^2} \\
&= v_F^2 \frac{\mathbf{p}}{\epsilon} \left(\frac{\mathbf{p}}{\epsilon} - \frac{\mathbf{p} - \mathbf{q}}{\epsilon - \omega} - \frac{\mathbf{k} + \mathbf{q}/2}{E + \omega/2} + \frac{\mathbf{k} - \mathbf{q}/2}{E - \omega/2} \right). \quad (\text{C3})
\end{aligned}$$

Note that in the conventional case of massive particles with the quadratic electronic dispersion, $\mathbf{v} = \mathbf{p}/m$ and hence $\mathbf{v}_i - \mathbf{v}_f \propto \mathbf{p} - (\mathbf{p} - \mathbf{q}) - (\mathbf{k} + \mathbf{q}/2) + (\mathbf{k} - \mathbf{q}/2) = 0$, which reflects the fact that because of the total momentum conservation there is no current relaxation in conventional metals due to the electron-electron interaction.

We now make use of the particle-hole symmetry of the graphene spectrum in order to simplify the expression for the transport scattering rate. Let us reverse the integration variables in the second line of Eq. (C1): $(E, \mathbf{k}) \mapsto (-E, -\mathbf{k})$. Using the symmetry $\mathcal{A}_0(\epsilon, \mathbf{p}) = \mathcal{A}_0(-\epsilon, -\mathbf{p})$, we see that the trace in the integrand is not changed. The difference of two equilibrium distribution functions in the second line of Eq. (C1) is also independent of the sign of E . Thus the contribution of particle-hole pair into the current relaxation vanishes, which corresponds to the absence of the Coulomb-drag contribution to the conductivity⁷⁷ in undoped graphene [Fig. 10(f)]. This allows us to keep only the part of the transport

TABLE X. Contribution of region 3 to the transport scattering rate τ_{tr}^{-1} in all energy domains and the corresponding characteristic values of Q , Ω , γ , and θ . Since in energy domains I and IV the result is dominated by $Q \sim 1$, the numerical prefactors in these domains are beyond the accuracy of our approximations. Furthermore, since in regime III the integral is dominated by $Q \sim y$, the splitting of the integrand according Eqs. (B6) and (B7) does not reproduce the correct prefactor.

Regime	$\tau_{\text{tr}}^{-1}(y)$	Q	Ω	γ	θ
I	$\sim \frac{\sqrt{y}}{N} T$	1	$-Q$	$0 < \gamma < \pi$	$0 < \theta < \pi$
II	$\simeq \frac{\pi}{4} \alpha_g T$	$\frac{\alpha_g^2 N^2}{y}$	$-Q$	$\pi - \frac{\alpha_g N}{\sqrt{y}}$	$\frac{\alpha_g N}{\sqrt{y}} \ll 1$
III	$\sim \frac{N}{y} \alpha_g^2 T$	y	$y(1 - 2 \sin \frac{\gamma}{2})$	$\frac{\alpha_g N}{y} \& \pi - \frac{\alpha_g N}{y}$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$
IV	$\sim \frac{N}{y^2} \alpha_g^2 T$	1	$Q \cos \gamma$	$\frac{\alpha_g N}{y}$	$\theta \lesssim \frac{\alpha_g N}{y} \ll 1$




FIG. 16. Diagram for the “elementary” inelastic-scattering amplitude.

factor Eq. (C3) that is even under reversing E and \mathbf{k} ,

$$v_F^2 \frac{\mathbf{p}}{\epsilon} \left(\frac{\mathbf{p}}{\epsilon} - \frac{\mathbf{p} - \mathbf{q}}{\epsilon - \omega} \right) = 1 - \text{sgn}(\epsilon - \omega) \cos \theta, \quad (\text{C4})$$

where θ is the scattering angle. Inserting this reduced transport factor into the first line of Eq. (C1) and calculating the trace of projection operators, we obtain

$$\begin{aligned} & [1 - \text{sgn}(\epsilon - \omega) \cos \theta] \text{Tr} [\mathcal{P}_+(\mathbf{p}) \mathcal{A}_0(\epsilon - \omega, \mathbf{p} - \mathbf{q})] \\ &= -\frac{\pi}{2} [1 - \text{sgn}(\epsilon - \omega) \cos \theta] \\ & \quad \times [(1 + \cos \theta) \delta(\epsilon - \omega - v_F |\mathbf{p} - \mathbf{q}|) \\ & \quad + (1 - \cos \theta) \delta(\epsilon - \omega + v_F |\mathbf{p} - \mathbf{q}|)] \\ &= \frac{1}{2} (1 - \cos^2 \theta) [\mathcal{A}_{0,+}(\epsilon - \omega, \mathbf{p} - \mathbf{q}) \\ & \quad + \mathcal{A}_{0,-}(\epsilon - \omega, \mathbf{p} - \mathbf{q})]. \end{aligned} \quad (\text{C5})$$

This way the transport factor $1 - \cos^2 \theta$ appears for Dirac fermions.

The GR calculation of the transport scattering rate reproduces the result obtained from the Drude conductivity Eq. (68). Note that an additional broadening δ was not used in this calculation. This shows that our result for τ_{tr}^{-1} can be actually applied in the limit $\tau_{\text{tr}}^{-1} \gg \delta$ as we argued in Sec. VC.

APPENDIX D: TRANSPORT SCATTERING RATE AND CONDUCTIVITY: DIAGRAMMATICS VS KINETIC EQUATION

In this appendix we show how the energy-averaged transport scattering rate, Eq. (75), is related to the calculation of the conductivity by using the kinetic equation approach.^{56,57} The kinetic equation in graphene governs the distribution functions $n_{\pm}(\mathbf{p})$ of + and - quasiparticles:

$$(\partial_t + \mathbf{v} \nabla_{\mathbf{R}} + e \mathbf{E} \mathbf{v} \cdot \partial_{\epsilon}) n_{\alpha} = \text{St}_{\alpha}(p)[n]. \quad (\text{D1})$$

Here is $\text{St}_{\alpha}(p)$ the collision integral (see Refs. 56 and 57 for the definition) which mixes distribution functions of electrons and holes and $\mathbf{v}(\epsilon)$ is defined as in Eq. (C2). Following Ref. 56, we use an ansatz for small electric-field-induced deviations of the distribution functions from its equilibrium form n_F (Fermi

distribution):

$$n_{\alpha} = n_F + \delta n_{\alpha} = n_F(\alpha \mathbf{p} + \alpha e \mathbf{E} \chi_{\mathbf{p}}) = n_F(\epsilon) + e \mathbf{E} \mathbf{v} \chi_{\mathbf{p}} \partial_{\epsilon} n_F. \quad (\text{D2})$$

As was shown in Refs. 56 and 57, in view of the fast energy relaxation, the conductivity is dominated by the momentum-independent part χ_0 of $\chi_{\mathbf{p}}$. Therefore we substitute Eq. (D2) with a constant χ_0 into the kinetic equation, Eq. (D1). We also insert the nonequilibrium part of the distribution function δn_{α} with momentum independent factor χ_0 into the definition of the current,

$$\begin{aligned} \mathbf{j} &= \frac{e}{\hbar} N \sum_{\alpha=\pm} \int_0^{\infty} \frac{d^2 p}{(2\pi)^2} \mathbf{v} \delta n_{\alpha}(p) \\ &= \frac{e}{\hbar v_F^2} N \int_0^{2\pi} \frac{d\phi_{\mathbf{v}}}{2\pi} \int_{-\infty}^{\infty} \frac{d\epsilon}{2\pi} \epsilon \mathbf{v}(\epsilon) \delta n(\epsilon) \\ &= \frac{e^2 \chi_0}{2\pi v_F^2 \hbar} N \int_0^{2\pi} \frac{d\phi_{\mathbf{v}}}{2\pi} \int_{-\infty}^{\infty} d\epsilon \epsilon \mathbf{v}(\epsilon) [\mathbf{E} \mathbf{v}(\epsilon)] \partial_{\epsilon} n_F(\epsilon) \\ &= \frac{e^2}{2\pi \hbar} \mathbf{E} N \chi_0 \int_{-\infty}^{\infty} d\epsilon \frac{\epsilon}{2} \partial_{\epsilon} n_F(\epsilon) \\ &= \frac{e^2}{2\pi \hbar} \mathbf{E} N \chi_0 T \ln 2. \end{aligned} \quad (\text{D3})$$

Linearizing the collision integral with respect to electric field, we simplify the kinetic equation:

$$e v_F \mathbf{E} \frac{\mathbf{p}}{\epsilon} \partial_{\epsilon} n_F(\epsilon) = \text{St}_{\text{sgn}(\epsilon)}(p) [\delta n]. \quad (\text{D4})$$

Now we observe that the left-hand side of Eq. (D4) is nothing but the integrand in the current (up to prefactors) as given by Eq. (D3). Therefore we can express the current through the collision integral as follows:

$$\begin{aligned} \mathbf{j} &= \frac{e}{2\pi v_F^2 \hbar} \chi_0 N \int_0^{2\pi} \frac{d\phi_{\mathbf{v}}}{2\pi} \int_{-\infty}^{\infty} d\epsilon \epsilon \mathbf{v}(\epsilon) \text{St}_{\text{sgn}(\epsilon)}(p) [\delta n] \\ &= \frac{e^2}{2\pi \hbar} \mathbf{E} N \chi_0^2 \int_{-\infty}^{\infty} d\epsilon \epsilon \frac{1}{\tau_{\text{tr}}(\epsilon)} \partial_{\epsilon} n_F(\epsilon). \end{aligned} \quad (\text{D5})$$

Here we have used the fact that the collision integral acting on the nonequilibrium ansatz, Eq. (D2), with $\chi_{\mathbf{p}} \rightarrow \chi_0$ yields the transport scattering rate at energy ϵ ; see Appendix C.

Comparing Eqs. (D3) and (D5), we express χ_0^{-1} through the energy-averaged transport scattering rate:

$$\chi_0^{-1} = \frac{2}{T \ln 2} \int_0^{\infty} d\epsilon \epsilon \frac{1}{\tau_{\text{tr}}(\epsilon)} \partial_{\epsilon} n_F(\epsilon), \quad (\text{D6})$$

which is equivalent to Eq. (75) and therefore allows us to identify

$$\langle \tau_{\text{tr}}^{-1} \rangle = \chi_0. \quad (\text{D7})$$

This fits perfectly to the physical picture behind the ansatz in Eq. (D2) where this rate corresponds to the time scale on which the acceleration of the electrons takes place.

We note that the above procedure is valid for the generalized GR with the RPA propagators replacing the bare interaction (see Appendix C). We have actually calculated the energy-averaged transport scattering rate using the full

RPA propagators. The resulting expression for the conductivity coincides with the second-order GR result of Refs. 56 and 57. However, the validity of the second-order calculation is not obvious *a priori*. Indeed, there are at least two possibilities to have the RPA corrections to the numerical prefactor of the conductivity:

(i) The higher-order terms in the RPA collision kernel have a resonant structure at small momentum transfer with the width governed by the interaction constant. Therefore the momentum integration over the resonance may, in principle, lead to correction to the transport scattering rate at relevant energies $\epsilon \sim T$, which is of the same order as the GR result. As an example, one may consider the following integral: $\alpha_g^3 \int dQ / [(Q - \alpha_g)^2 + \alpha_g^2]$. The expansion of the integrand in α_g starts with α_g^3 , but the result of the integration is of the order of α_g^2 . We have checked, however, that this enhancement does not occur in our calculation of $1/\tau_{tr}$ at energies $\epsilon \sim T$ relevant to the conductivity. On the contrary, the transport scattering rate at small energies ($\epsilon \sim \alpha_g T$), the RPA resummation is crucially important and does lead to

a nonanalytic enhancement of the transport scattering rate $\sim |\alpha_g|$; see Fig. 12.

(ii) The conductivity is determined by the transport scattering rate averaged over energies. We have shown in this appendix that the averaging involves the single-particle density of states $\rho(\epsilon) \propto \epsilon$; see Eq. (75). Let us assume for a moment that the energy averaging of $\tau_{tr}^{-1}(\epsilon)$ is performed with a constant weight rather than with the density of states. Since at $\epsilon < \alpha_g T$ the transport scattering rate is enhanced ($\sim |\alpha_g| T$) due to the RPA resummation, the contribution of this low-energy domain to the such averaged rate would be $\sim \alpha_g^2 T$ thus changing the numerical prefactor in the conductivity. However, since the actual energy averaging does involve the density of states, the low-energy domain ($\epsilon < \alpha_g T$), dominated by the RPA, yields only a subleading $\sim \alpha_g^3 T$ correction to the GR result for $\langle \tau_{tr}^{-1} \rangle$.

Thus our calculation based on the RPA resummation of the interaction propagators justifies the previous results for the collision-limited conductivity obtained at the GR level.

¹A. K. Geim and K. S. Novoselov, *Nat. Mater.* **6**, 183 (2007).

²A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, *Rev. Mod. Phys.* **81**, 109 (2009).

³K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, *Science* **306**, 666 (2004).

⁴K. S. Novoselov, D. Jiang, T. Booth, V. V. Khotkevich, S. M. Morozov, and A. K. Geim, *Proc. Natl. Acad. Sci. USA* **102**, 10451 (2005).

⁵K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, *Nature (London)* **438**, 197 (2005).

⁶Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, *Nature (London)* **438**, 201 (2005); Y.-W. Tan, Y. Zhang, H. L. Stormer, and P. Kim, *Eur. Phys. J. Special Topics* **148**, 15 (2007).

⁷M. I. Katsnelson, *Eur. Phys. J. B* **51**, 157 (2006).

⁸J. Tworzydło, B. Trauzettel, M. Titov, A. Rycerz, and C. W. J. Beenakker, *Phys. Rev. Lett.* **96**, 246802 (2006); C. W. J. Beenakker, *Rev. Mod. Phys.* **80**, 1337 (2008).

⁹S. Ryu, C. Mudry, A. Furusaki, and A. W. W. Ludwig, *Phys. Rev. B* **75**, 205344 (2007).

¹⁰F. Miao, S. Wijeratne, Y. Zhang, U. C. Coskun, W. Bao, and C. N. Lau, *Science* **317**, 1530 (2007).

¹¹R. Danneau, F. Wu, M. F. Craciun, S. Russo, M. Y. Tomi, J. Salmilehto, A. F. Morpurgo, and P. J. Hakonen, *Phys. Rev. Lett.* **100**, 196802 (2008).

¹²M. Titov, *Europhys. Lett.* **79**, 17004 (2007).

¹³A. Schuessler, P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin, *Phys. Rev. B* **79**, 075405 (2009).

¹⁴M. Titov, P. M. Ostrovsky, I. V. Gornyi, A. Schuessler, and A. D. Mirlin, *Phys. Rev. Lett.* **104**, 076802 (2010).

¹⁵J. H. Bardarson, J. Tworzydło, P. W. Brouwer, and C. W. J. Beenakker, *Phys. Rev. Lett.* **99**, 106801 (2007); K. Nomura, M. Koshino, and S. Ryu, *ibid.* **99**, 146806 (2007); P. San-Jose, E. Prada, and D. S. Golubev, *Phys. Rev. B* **76**, 195445 (2007);

C. H. Lewenkopf, E. R. Mucciolo, and A. H. Castro Neto, *ibid.* **77**, 081410(R) (2008); J. Tworzydło, C. W. Groth, and C. W. J. Beenakker, *ibid.* **78**, 235438 (2008).

¹⁶P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin, *Phys. Rev. Lett.* **98**, 256801 (2007); *Eur. Phys. J. Spec. Top.* **148**, 63 (2007).

¹⁷P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin, *Phys. Rev. B* **74**, 235443 (2006).

¹⁸P. M. Ostrovsky, M. Titov, S. Bera, I. V. Gornyi, and A. D. Mirlin, *Phys. Rev. Lett.* **105**, 266803 (2010).

¹⁹T. Stauber, N. M. R. Peres, and F. Guinea, *Phys. Rev. B* **76**, 205423 (2007).

²⁰T. Ando, *J. Phys. Soc. Jpn.* **75**, 074716 (2006).

²¹K. Nomura and A. H. MacDonald, *Phys. Rev. Lett.* **98**, 076602 (2007).

²²A. F. Morpurgo and F. Guinea, *Phys. Rev. Lett.* **97**, 196804 (2006).

²³R. V. Gorbachev, F. V. Tikhonenko, A. S. Mayorov, D. W. Horsell, and A. K. Savchenko, *Phys. Rev. Lett.* **98**, 176805 (2007); F. V. Tikhonenko, D. W. Horsell, R. V. Gorbachev, and A. K. Savchenko, *ibid.* **100**, 056802 (2008); D. W. Horsell, A. K. Savchenko, F. V. Tikhonenko, K. Kechedzhi, I. V. Lerner, and V. I. Fal'ko, *Solid State Commun.* **149**, 1041 (2009).

²⁴F. V. Tikhonenko, A. A. Kozikov, A. K. Savchenko, and R. V. Gorbachev, *Phys. Rev. Lett.* **103**, 226801 (2009).

²⁵E. McCann, K. Kechedzhi, V. I. Fal'ko, H. Suzuura, T. Ando, and B. L. Altshuler, *Phys. Rev. Lett.* **97**, 146805 (2006); K. Kechedzhi, V. I. Fal'ko, E. McCann, and B. L. Altshuler, *ibid.* **98**, 176806 (2007).

²⁶V. V. Cheianov and V. I. Falko, *Phys. Rev. Lett.* **97**, 226801 (2006).

²⁷A. A. Kozikov, A. K. Savchenko, B. N. Narozhny, and A. V. Shytov, *Phys. Rev. B* **82**, 075424 (2010).

²⁸X. Du, I. Skachko, A. Barker, and E. Y. Andrei, *Nat. Nanotechnol.* **3**, 491 (2008); *Int. J. Mod. Phys. B* **22**, 4579 (2008).

²⁹K. I. Bolotin, K. J. Sikes, Z. Jiang, M. Klima, G. Fudenberg, J. Hone, P. Kim, and H. L. Stormer, *Solid State Commun.* **146**, 351 (2008).

- ³⁰X. Du, I. Skachko, F. Duerr, A. Luican, and E. Y. Andrei, *Nature* **462**, 192 (2009).
- ³¹K. I. Bolotin, F. Ghahari, M. D. Shulman, H. L. Stormer, and P. Kim, *Nature (London)* **462**, 196 (2009).
- ³²I. Skachko, X. Du, F. Duerr, A. Luican, D. A. Abanin, L. S. Levitov, and E. Y. Andrei, *Phil. Trans. R. Soc. A* **368**, 5403 (2010); D. A. Abanin, I. Skachko, X. Du, E. Y. Andrei, and L. S. Levitov, *Phys. Rev. B* **81**, 115410 (2010).
- ³³B. L. Altshuler and A. G. Aronov, in *Electron-Electron Interactions in Disordered Systems*, edited by A. L. Efros and M. Pollak (Elsevier, Amsterdam, 1985), p. 1.
- ³⁴J. P. Reed, B. Uchoa, Y. I. Joe, Y. Gan, D. Casa, E. Fradkin, and P. Abbamonte, *Science* **330**, 805 (2010).
- ³⁵J. Gonzalez, F. Guinea, and M. A. H. Vozmediano, *Nucl. Phys. B* **424**, 596 (1994).
- ³⁶D. T. Son, *Phys. Rev. B* **75**, 235423 (2007).
- ³⁷J. Ye and S. Sachdev, *Phys. Rev. Lett.* **80**, 5409 (1998).
- ³⁸J. Ye, *Phys. Rev. B* **60**, 8290 (1999).
- ³⁹T. Stauber, F. Guinea, and M. A. H. Vozmediano, *Phys. Rev. B* **71**, 041406(R) (2005).
- ⁴⁰D. E. Sheehy and J. Schmalian, *Phys. Rev. Lett.* **99**, 226803 (2007).
- ⁴¹M. S. Foster and I. L. Aleiner, *Phys. Rev. B* **77**, 195413 (2008).
- ⁴²M. S. Foster and A. W. W. Ludwig, *Phys. Rev. B* **73**, 155104 (2006).
- ⁴³D. V. Khveshchenko and H. Leal, *Nucl. Phys. B* **687**, 323 (2004); D. V. Khveshchenko, *Phys. Rev. Lett.* **87**, 246802 (2001); *J. Phys.: Condens. Matter* **21**, 075303 (2009).
- ⁴⁴E. V. Gorbar, V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, *Phys. Rev. B* **66**, 045108 (2002).
- ⁴⁵I. F. Herbut, *Phys. Rev. Lett.* **97**, 146401 (2006); O. Vafek, *ibid.* **98**, 216401 (2007).
- ⁴⁶I. L. Aleiner, D. E. Kharzeev, and A. M. Tsvelik, *Phys. Rev. B* **76**, 195415 (2007).
- ⁴⁷S. J. Hands and C. G. Strouthos, *Phys. Rev. B* **78**, 165423 (2008).
- ⁴⁸J. E. Drut and T. A. Lähde, *Phys. Rev. Lett.* **102**, 026802 (2009); *Phys. Rev. B* **79**, 165425 (2009).
- ⁴⁹G.-Z. Liu, W. Li, and G. Cheng, *Phys. Rev. B* **79**, 205429 (2009).
- ⁵⁰L. Sheng, D. N. Sheng, F. D. M. Haldane, and L. Balents, *Phys. Rev. Lett.* **99**, 196802 (2007).
- ⁵¹K. Nomura and A. H. MacDonald, *Phys. Rev. Lett.* **96**, 256602 (2006).
- ⁵²D. A. Abanin, P. A. Lee, and L. S. Levitov, *Phys. Rev. Lett.* **98**, 156801 (2007); D. A. Abanin, K. S. Novoselov, U. Zeitler, P. A. Lee, A. K. Geim, and L. S. Levitov, *ibid.* **98**, 196806 (2007).
- ⁵³Z. Jiang, Y. Zhang, H. L. Stormer, and P. Kim, *Phys. Rev. Lett.* **99**, 106802 (2007).
- ⁵⁴J. Gonzalez, F. Guinea, and M. A. H. Vozmediano, *Phys. Rev. Lett.* **77**, 3589 (1996).
- ⁵⁵E. H. Hwang, Ben Yu-Kuang Hu, and S. Das Sarma, *Phys. Rev. B* **76**, 115434 (2007).
- ⁵⁶A. B. Kashuba, *Phys. Rev. B* **78**, 085415 (2008).
- ⁵⁷L. Fritz, J. Schmalian, M. Müller, and S. Sachdev, *Phys. Rev. B* **78**, 085416 (2008).
- ⁵⁸M. S. Foster and I. L. Aleiner, *Phys. Rev. B* **79**, 085415 (2009).
- ⁵⁹M. Müller and S. Sachdev, *Phys. Rev. B* **78**, 115419 (2008).
- ⁶⁰M. Müller, L. Fritz, and S. Sachdev, *Phys. Rev. B* **78**, 115406 (2008).
- ⁶¹M. Müller, J. Schmalian, and L. Fritz, *Phys. Rev. Lett.* **103**, 025301 (2009).
- ⁶²M. Müller, L. Fritz, S. Sachdev, and J. Schmalian, in *Advances in Theoretical Physics*, edited by V. Lebedev and M. Feigel'man, AIP Conf. Proc. No. 1134 (AIP, New York, 2009), p. 170.
- ⁶³A. Kamenev and A. Levchenko, *Adv. Phys.* **58**, 197 (2009).
- ⁶⁴J. Rammer and H. Smith, *Rev. Mod. Phys.* **58**, 323 (1986).
- ⁶⁵I. L. Aleiner and Ya. M. Blanter, *Phys. Rev. B* **65**, 115317 (2002).
- ⁶⁶I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. B* **75**, 085421 (2007).
- ⁶⁷A. G. Yashenkin, I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. B* **78**, 205407 (2008).
- ⁶⁸D. A. Bagrets, I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, *Semiconductors* **42**, 994 (2008); D. A. Bagrets, I. V. Gornyi, and D. G. Polyakov, *Phys. Rev. B* **80**, 113403 (2009); S. Ngo Dinh, D. A. Bagrets, A. D. Mirlin, *ibid.* **81**, 081306(R) (2010).
- ⁶⁹D. B. Gutman, Y. Gefen, and A. D. Mirlin, *Phys. Rev. Lett.* **101**, 126802 (2008); *Phys. Rev. B* **80**, 045106 (2009).
- ⁷⁰B. N. Narozhny, G. Zala, and I. L. Aleiner, *Phys. Rev. B* **65**, 180202 (2002).
- ⁷¹I. L. Aleiner, B. L. Altshuler, and M. E. Gershenson, *Waves Random Media* **9**, 201 (1999).
- ⁷²V. F. Gantmakher and Y. B. Levinson, *Carrier Scattering in Metals and Semiconductors* (North-Holland, Amsterdam, 1987).
- ⁷³B. N. Narozhny, *Phys. Rev. B* **76**, 153409 (2007).
- ⁷⁴Wang-Kong Tse, Ben Yu-Kuang Hu, and S. Das Sarma, *Phys. Rev. B* **76**, 081401(R) (2007).
- ⁷⁵A. Schuessler, P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin, *Phys. Rev. B* **82**, 085419 (2010).
- ⁷⁶A. D. Mirlin, F. Evers, I. V. Gornyi, and P. M. Ostrovsky, in *50 Years of Anderson Localization*, edited by E. Abrahams (World Scientific, Singapore, 2010); *Int. J. Mod. Phys. B* **24**, 1577 (2010).
- ⁷⁷In conventional systems with quadratic dispersion the Coulomb-drag-type contribution to the conductivity [Fig. 10(f)] cancels the contribution of the diagrams in Figs. 10(d) and 10(e); see Appendix A in I. V. Gornyi and A. D. Mirlin, *Phys. Rev. B* **69**, 045313 (2004). This is a manifestation of the total momentum conservation in inelastic processes, which in normal metals translates into the absence of current relaxation due to interactions.