

Entanglement entropy of gapped phases and topological order in three dimensions

Tarun Grover, Ari M. Turner, and Ashvin Vishwanath

Department of Physics, University of California, Berkeley, California 94720, USA

(Received 7 September 2011; revised manuscript received 22 October 2011; published 23 November 2011)

We discuss entanglement entropy of gapped ground states in different dimensions, obtained on partitioning space into two regions. For *trivial* phases without topological order, we argue that the entanglement entropy may be obtained by integrating an “entropy density” over the partition boundary that admits a gradient expansion in the curvature of the boundary. This constrains the expansion of entanglement entropy as a function of system size and points to an even-odd dependence on dimensionality. For example, in contrast to the familiar result in two dimensions, a size-independent constant contribution to the entanglement entropy can appear for trivial phases in any odd spatial dimension. We then discuss phases with topological entanglement entropy (TEE) that cannot be obtained by adding local contributions. We find that in three dimensions there is just one type of TEE, as in two dimensions, that depends linearly on the number of connected components of the boundary (the “zeroth Betti number”). In $D > 3$ dimensions, new types of TEE appear which depend on the higher Betti numbers of the boundary manifold. We construct generalized toric code models that exhibit these TEEs and discuss ways to extract TEE in $D \geq 3$.

DOI: [10.1103/PhysRevB.84.195120](https://doi.org/10.1103/PhysRevB.84.195120)

PACS number(s): 03.67.Mn, 03.65.Ud, 05.30.—d

I. INTRODUCTION

In recent years, surprising connections have emerged between error correction of quantum information and topological condensed-matter phases.^{1,2} At the same time, ideas from quantum information have proved useful in defining topological phases. Two-dimensional phases with topological order, such as those realized in the context of the fractional quantum Hall effect, are gapped phases for which the ground-state degeneracy depends on the genus of the space on which they are defined. Recent work has shown that they can be identified by the entanglement properties of their ground-state wave function.^{3–5} The entanglement entropy of a region with a smooth boundary of length L takes the form $S_A = \alpha_1 L - b_0 \gamma$, where γ is the topological entanglement entropy, b_0 is the number of connected components of the boundary of region A , and we have dropped the subleading terms. In gapped phases without topological order, such as band insulators, $\gamma = 0$ for a smooth boundary. These predictions have been verified in the context of a number of specific $D = 2$ models^{6–10} and in $D = 3$ Z_2 toric code models.¹¹ In this paper, we discuss the general structure of entanglement entropy for gapped topological and nontopological phases in $D \geq 3$.

One notices that in $D = 2$, the topological entanglement entropy depends only on a topological property of the boundary, in this case the number of connected components. There are two equivalent ways of extracting the topological entanglement entropy.^{4,5} First, via the scaling of the entropy with boundary size for smooth boundaries, so as to extract the constant term. The second is by considering a combination of entanglement entropies of three suitably chosen regions A, B, C , so that $-\gamma = S_A + S_B + S_C - S_{AB} - S_{BC} - S_{AC} + S_{ABC}$.

Here we discuss analogous questions in $D \geq 3$. In particular, consider a gapped $D = 3$ phase and a region A with a smooth boundary. (1) If the entanglement entropy S_A contains a constant term, does it necessarily reflect topological order? (2) The boundary of A is a closed two-dimensional surface that has two topological invariants associated with it: the number

of connected components and the genus (number of handles) of each component. Does this imply there are two distinct types of topological entanglement entropies (TEEs) and, correspondingly, two varieties of topological order in $D = 3$? The answer is *no* to both these questions, as we elaborate in this paper. We show that even a trivial gapped phase, with no topological order, one can have a constant term in the entanglement entropy in $D = 3$ (and any other odd dimension). Hence, this by itself does not signify topological order. Moreover, this constant is generally genus dependent, ruling out a topological origin for a genus-dependent entanglement entropy. This reduces the number of possible TEE to the same as $D = 2$. We discuss generalization of the Kitaev-Preskill scheme⁵ to extract the TEE in $D = 3$ and why some naive extrapolations fail.

A deeper understanding of TEE is obtained by considering higher dimensions. We show that at least one new TEE appears on going up every two dimensions. Thus, while $D = 2$ and 3 are similar, a new topological constant does appear in $D = 4$ (in $D = 2n$ and $2n + 1$, there are thus n constants). These are related to the Betti numbers¹² of the boundary. We construct topological phases that manifest these new TEEs and explicitly calculate their value. These are based on a discrete gauge group G . In all dimensions, the TEE for discrete gauge theories is $-\log |G|$ per connected surface component, where $|G|$ is the number of elements in G . These theories capture both Abelian (like the Z_2 toric code) and non-Abelian phases and the ground state of these theories correspond to condensate of closed loops. One can also consider more general Abelian discrete gauge theories where the fluctuating loops are readily generalized to fluctuating p -dimensional surfaces. These manifest explicitly in the TEE, through the appearance of new topological constants that depend on higher Betti numbers. Furthermore, a previously discussed duality between p and $D - p$ theories in D dimensions^{13–15} is reflected in the structure of TEE.

To isolate topological contributions it is useful to know the structure of entanglement entropy in trivial gapped phases. Since correlations are local in such phases, we propose

an expansion of entanglement entropy S_A based on adding individual contributions from patches on the surface of region A : $S_A = \sum_i S_i$. The entropy densities S_i will depend on local properties, such as the local curvature of the surface. One can then expand the entropy density in polynomials of curvature and its derivatives, similar to the Landau expansion of free-energy density.¹⁶ The contribution from higher-order terms to S_A is subdominant for large surfaces. Interestingly, not every term is allowed in this expansion. When we divide space into a region A (inside) and a region \bar{A} (outside), the entanglement entropy of both are equal; that is, $S_A = S_{\bar{A}}$. This imposes a Z_2 symmetry on the expansion that is *unique to ground-state entanglement entropy*.¹⁷ This has important consequences. Consider, for example, $D = 2$. The entropy density is not allowed to depend linearly on the boundary curvature κ , which changes sign on interchanging inside and outside. Thus, the expansion of entropy density for a trivial $D = 2$ phase is $S_i = a_0 + a_2\kappa^2(r_i) + \dots$, which when integrated around the boundary leads to $S_A = \alpha_1 L + \alpha_3/L + \dots$, where L is the length of the boundary of region A . The first term is the area law, and the next term is two orders of L down, due to the Z_2 symmetry, which eliminates the constant term in total entropy for smooth boundaries. Thus, the existence of a constant term in a gapped $D = 2$ state implies a *nontrivial* phase that is, topological order. In general, this method predicts that for an isotropic, parity-invariant state without topological order, the entanglement entropy in even (odd) spatial dimensions depends *only* on odd (even) powers of L , the linear scale of the boundary.

We assume it is possible to take the continuum limit for all the phases that we consider in this paper. This assumption excludes phases such as a layered Z_2 topological ordered phases in $D = 3$ whose topological entropy depends on the local geometry of region A .

The paper is organized as follows. In Sec. II we discuss the general structure of entanglement entropy for gapped phases and explain the basic assumptions underlying our discourse. In Sec. III we introduce the aforementioned curvature expansion for entanglement entropy of trivial gapped phases and study its consequences. In Secs. IV and V we study topological ordered phases in $D = 3$ and $D > 3$, respectively, through extracting the dependence of entanglement entropy of a region on the topology of its boundary. We also generalize the constructions for extracting topological entropy.^{4,5,11}

II. STRUCTURE OF ENTANGLEMENT ENTROPY FOR GAPPED PHASES

In this article, we assume that the entanglement entropy of region A can be decomposed into two parts:

$$S_A = S_{A,\text{local}} + S_{A,\text{topological}}. \quad (1)$$

We postpone the underpinnings of this assumption to Sec. IV when we study topologically ordered phases. Here S_{local} is defined by the property that it can be written as a sum over contributions from patches located along the boundary of region A :

$$S_{A,\text{local}} = \sum_i S_i, \quad (2)$$




FIG. 1. The local part of the entropy of region A is the sum of contributions of small patches on the boundary.

where S_i depends only on the shape of the patch i and not on the rest of the surface or how it fits with other patches (see Fig. 1), *at least* if the edge of the patch connects smoothly to all other patches.

We assume the other contribution $S_{\text{topological}}$ is topologically invariant, that is, it does not change as the boundary is deformed unless the topology of the region changes. If such a term is present and if it cannot be expressed in a local way, then the phase has long-range entanglement, which is the hallmark of topological order.^{2,4,5}

Let us consider the assumptions under which the decomposition [Eq. (2)] would be possible for a trivial (i.e., not topologically ordered) gapped phase. The reduced density matrix corresponding to region A for the ground-state wave function may be written as

$$\rho_A = e^{-H_A}/Z,$$

where H_A is the so-called *entanglement Hamiltonian* and $Z = \text{tr}(e^{-H_A})$ so that $\text{tr}(\rho_A) = 1$. Therefore, we can think of ρ_A as the thermal density matrix at temperature $T = 1$ for the Hamiltonian H_A and the von Neumann entropy $S_A = -\text{tr}(\rho_A \log \rho_A)$ as the thermal entropy for this system. Let us define $\tilde{\rho}_A(T) = e^{-H_A/T}/Z(T)$, where $Z(T) = \text{tr}(e^{-H_A/T})$. Clearly, $\rho_A = \tilde{\rho}_A(T = 1)$ and S_A obeys the following equation:

$$S_A \equiv S_{A,\text{local}} = \int_1^\infty \frac{dT}{T} \frac{\partial \langle H_A \rangle}{\partial T}, \quad (3)$$

where $\langle H_A \rangle$ denotes the thermal average of H_A at temperature T with respect to the density matrix $\tilde{\rho}(T)$. We claim that the entanglement entropy would admit an expansion such as Eq. (2) if the following conditions are satisfied.

(i) H_A can be written as a sum of local operators O 's, that is, $H_A = \sum_x O(x)$.

(ii) There is no phase transition for the Hamiltonian H_A for $T \geq 1$.

The first condition along with the fact that all correlations of local operators are short-ranged in a gapped phase imply that H_A has nonzero support only near the boundary of region A (within the distance of correlation length). In other words, the degrees of freedom inside and outside of region A are coupled only through operators that lie within a distance $\sim \xi$ from the boundary. This implies that $\langle H_A \rangle = \sum_i h_i$, where i denotes a point at the boundary of region A and the h_i 's

depend solely on the properties of the boundary in the vicinity of point i . The second condition implies that the integral in Eq. (3) does not admit any singularity so that all terms S_i in the Eq. (2) are finite. Physically, this means that the actual system of interest is smoothly connected to its $T = \infty$ zero correlation length system, where Eq. (2) holds trivially.

III. ENTANGLEMENT ENTROPY OF TRIVIAL GAPPED PHASES

In this section we focus on understanding the leading and subleading dependence of $S_{A,\text{local}}$ on L for gapped trivial phases of matter. Let us assume that the boundary of region A is smooth, and further that the phase is isotropic and parity invariant (consequences of the violation of these assumptions are discussed at the end of this section and in Appendix B). Then, as we see below, in the absence of topological order, only alternate terms in the power series expansion of $S_A(L)$ appear:

$$S_{A,\text{local}}(L) = \alpha_1 L^{D-1} + \alpha_3 L^{D-3} + \alpha_5 L^{D-5} + \dots; \quad (4)$$

that is, only those with odd codimension exponent can appear. This expansion implies a distinction between even and odd dimensions: In even dimensions, any constant contribution to the entanglement entropy must come from $S_{A,\text{topological}}$ and thus indicates topological order. In odd spatial dimensions, a constant term may appear in the local entropy, making it more difficult, though still possible, to isolate topological contributions (note that these conclusions apply only to smooth boundaries in rotationally symmetric systems; a corner can produce a constant term for even dimensions as well as odd ones).

Let us consider some instances of Eq. (4) that will motivate its derivation. First, it is well known that in $D = 2$, a constant term in $S_A(L)$ implies the presence of topological order.^{4,5} On the other hand, in $D = 3$, a constant term can appear for a non-topologically ordered phase, such as a gapped scalar field. Consider for a moment¹⁸ a massless field, where it is known that the entanglement scales as $S_A \sim L^2 + \log(L)$ for a spherical ball of radius L . Now providing a mass m to the scalar field would cut off the $\log(L)$ term and instead lead to a constant contribution proportional to $\log(1/m)$. We verified this explicitly using a numerical calculation similar to Ref. 19. Interestingly, when the surface of region A is flat, then there is no constant contribution as we show in Appendix A (a flat boundary is possible when the total system has the topology of a three torus T^3 ; then A may be taken to have the geometry $T^2 \times l$, where l is a line segment). This indicates that the presence or absence of a constant term may have something to do with the curvature of the boundary of region A . In the next section, we make this statement precise and explain the observations made above.

A. Entropy density functional

Let us consider a region in two dimensions for concreteness. We postulate that the local entropy $S_{A,\text{local}}$ is given by the following integral:

$$S_A = \sum_i S_i = \int d\sigma F(\kappa, \partial\kappa, \dots), \quad (5)$$

where $F(\kappa, \partial\kappa, \dots)$ is the ‘‘entropy density functional.’’ In a gapped phase, the entropy S_i of a patch larger than the correlation length can depend only on the properties of the patch, such as its length $\Delta\sigma_i$ and curvature κ_i , as well as derivatives of the latter, and must be proportional to $\Delta\sigma_i$. Hence, $S_i = \Delta\sigma_i F(\kappa_i, \partial^n \kappa_i)$. Taking the limit where the patches become microscopic compared to L (but greater than ξ) leads to Eq. (5).

The entropy density functional always satisfies a Z_2 symmetry, which is the key to understanding the L dependence of the entropy. The symmetry results from the fact that, if A and B are complementary regions, then $S_A = S_B$. Therefore, changing ‘‘inside’’ to ‘‘outside’’ keeps the entanglement entropy invariant. Now, under this transformation, radii of curvature clearly change sign $\kappa \rightarrow -\kappa$, and this constrains the entropy density functional (Fig. 2). As an illustration of this Z_2 symmetry, consider the form of the functional F for a gapped two-dimensional system. On a smooth boundary, one can expand the function F in a Taylor series, retaining the first few terms:

$$F(\kappa_i, \partial_\sigma^n \kappa_i) = a_0 + a_1 \kappa_i + a_2 \kappa_i^2 + b_2 \partial_\sigma \kappa_i + \dots \quad (6)$$

The first term gives the boundary law $S_A = a_0 L$. The second term, if it is present, would give a constant contribution $\oint d\sigma \kappa = 2\pi$ for the curve shown, which would be a nonuniversal constant contribution. However, such a term is, in fact, forbidden by the Z_2 symmetry, since the term is odd in κ . The term κ^2 gives the next contribution to the entropy that is proportional to $\oint d\sigma \kappa^2$. If the shape of region A is kept fixed, then this contribution scales as $1/L$. The term $\partial_\sigma \kappa$ is allowed by the Z_2 symmetry (since both the derivative and the radius of curvature change sign, assuming that the direction of the curve is set by a ‘‘right-hand rule,’’ whereby the arc length increases along a specific direction), yet it still vanishes because it is a total derivative. Generalizing these arguments, one finds that $S_A = \sum_{k=0}^\infty \alpha_{2k+1} L^{1-2k}$.

In general dimensions, we find similar results if we continue to assume rotational, parity, and translational symmetry. These assumptions imply that F can depend only on the metric tensor $g^{\alpha\beta}$ and on the extrinsic curvature of the surface. The latter tensor does not appear when considering intrinsic properties of a manifold (as in general relativity). However, entanglement entropy does depend on the embedding of the boundary ∂A since it is measuring the entanglement of the degrees of freedom in the space around the surface. The extrinsic curvature is a tensor $\kappa_{\alpha\beta}$ with two indices (see, e.g., Ref. 20 for requisite differential geometry). Thus, each term in F contains some number of factors of $\kappa_{\alpha\beta}$ and its covariant




FIG. 2. Illustration of the Z_2 symmetry for the curvature expansion discussed in the text.

derivatives, with all the indices contracted by factors of $g^{\gamma\delta}$ (if parity is broken, the antisymmetric volume tensor $\gamma^{\alpha_1, \dots, \alpha_{D-1}}$ is allowed as well; Appendix B).

The inside/outside symmetry further limits the form of the terms in F : It implies that each term in F includes an even number of factors n_κ of κ . The total order of all the derivatives n_D must also be even. This follows from rotational symmetry. For rotational symmetry to be respected, one has to contract all the lower indices with the tensor $g^{\gamma\delta}$. This leads to an even number of lower indices, which includes the derivatives as well as the curvature indices. Since the curvature tensor is of even rank, the number of derivatives n_D has to be even. Putting everything together, one finds that the contribution to the entropy density F scales as $L^{-(n_\kappa + n_D)}$ that clearly has an even exponent, explaining why only alternate terms appear in the entropy [Eq. (4)].

When rotational or inversion symmetries are broken spontaneously or by applying a field, additional terms appear in the entropy, as summarized in Table I. We provide the details leading to these results in Appendix B.

The local entropy can also contain topology-dependent terms, for example, the term $\int G dA = 4\pi\chi$ in three dimensions where G is the Gaussian curvature which is the determinant of the matrix $\kappa_{\alpha\beta}$. Note that this term is compatible with the symmetry $\kappa \rightarrow -\kappa$, since G is quadratic in κ . Hence, as mentioned earlier, the presence of a term in the entropy that is proportional to the Euler characteristic does not necessarily correspond to topological order. In general dimensions, $\int \det \kappa dA$ is topological, but it is only symmetric in odd dimensions, where it is proportional to the Euler characteristic of the boundary (in general, it is proportional to the Euler characteristic of the region itself²¹).

IV. TOPOLOGICAL ENTANGLEMENT ENTROPY IN $D = 3$

We now turn to the topological part of the entanglement entropy. Our starting point is Eq. (1), which we rewrite here for convenience:

$$S_A = S_{A,\text{local}} + S_{A,\text{topological}}. \quad (7)$$

This decomposition is what enables the extraction of topological entropy using Kitaev-Preskill⁵ or Levin-Wen⁴ constructions for two-dimensional topological ordered phases. The assumptions underlying this equation are somewhat tricky. Though Eq. (7) holds for toric code models in all dimensions and there is strong numerical evidence that it also holds for many interesting two-dimensional topological ordered states such as Z_2 spin liquids, quantum dimer models, and various quantum Hall states,^{6-10,22} Eq. (7) surreptitiously rules out a

layered Z_2 topologically ordered state. Such a state would lead to a correction in entanglement entropy $\Delta S_A = -\gamma_{2D} L_z$, for layering perpendicular to the z direction. Here γ_{2D} is the TEE associated with the theory living in each layer. Clearly, ΔS_A is not topologically invariant. The assumptions underlying Eq. (7) most likely also do not apply to the self-correcting code state of Ref. 23; this state (whose ground state degeneracy depends on the divisibility of system size by powers of 2, for example) illustrates why we need to make an assumption of this type. Nevertheless, we briefly discuss the TEE of layered Z_2 state in Appendix D.

Independent contributions to $S_{\text{topological}}$. The boundary ∂A of a three-dimensional region A is a compact manifold that is characterized by Betti numbers b_0 and b_1 (note that for compact manifolds $b_2 = b_0$). As we show in Appendix E in three dimensions $S_{A,\text{topological}}$ is a linear function of b_0, b_1 , say, $S_{\text{topological}} = -\gamma_0 b_0 - \gamma_1 b_1$ (we assume that the space in which region A is embedded has the topology of \mathbb{R}^3 ; otherwise, more complicated dependence is possible in principle). This might lead one to suspect that there are two different kinds of topological orders in three dimensions, namely, those corresponding to a nonzero γ_0 and γ_1 , respectively. However, b_0, b_1 are related to the Euler characteristic χ through $2b_0 - b_1 = \chi$. Thus, one may redefine $S'_{A,\text{local}} = S_{A,\text{local}} + \alpha\chi$ and $S'_{A,\text{topological}} = S_{A,\text{topological}} - \alpha\chi$ without changing the entropy and α may be adjusted so that the b_1 dependence of $S_{\text{topological}}$ is canceled out. Here the term $\alpha\chi$ may be thought of as both local and topological. It is local, because the Euler's formula, $\chi = V - E + F$, gives a local expression for this term, where V , E , and F are the number of vertices, edges, and faces into which ∂A is divided (alternatively, in a continuum theory, $\alpha\chi$ can be incorporated into the entropy density F since χ is the integral of the Gaussian curvature). It is also topological, because $\alpha\chi$ is independent of how the surface is divided up into regions. The upshot of this discussion is that there is only one kind of topological entropy in three dimensions.

Z_2 string and Z_2 membrane models. As an alternative way to understand the above result, let us study specific models whose topological entanglement potentially depends on different Betti numbers. Consider the following model of Z_2 gauge theories consisting of spin-1/2 degrees of freedom that live on the links of a three-dimensional cubic lattice:

$$H_{\text{string}} = - \sum_{\square} \prod_{l \in \square} \tau_{z,l} - h \sum_l \tau_{x,l}, \quad (8)$$

where \square denotes a plaquette of the cubic lattice and the operators $\tau_{x,l}, \tau_{z,l}$ live on the links l of the lattice. The above Hamiltonian is supplemented with the constraint (“Gauss law”) $\prod_{l \in \text{vertex}} \tau_{x,l} = 1$ to impose the absence of Z_2 charges

TABLE I. Terms in the entropy forbidden by symmetries. The three columns describe systems with rotational and parity symmetry, no spatial symmetry, and rotational but not parity symmetry. The entries list the scaling of terms L^k that are forbidden from appearing in the entropy.

	Full symmetry	All symmetries broken	Broken parity alone
$D = 2$	All even terms L^0, L^{-2}	Constant term	All even terms
$D \geq 3$ odd	All odd terms L^{D-2}, L^{D-4}, \dots	Nothing forbidden	Odd terms with positive exponents and also L^{-1} if $D \equiv 3 \pmod{4}$
$D \geq 4$ even	Even terms L^{D-2}, L^{D-4}, \dots	Nothing forbidden	Even terms

in the theory. Because of this constraint the gauge-invariant degrees of freedom in this model consist of closed loops \mathcal{C} on the edges of the lattice. In the deconfined phase of the gauge theory, $|h| \ll 1$, the loops condense because they do not cost much energy. The entanglement entropy of this model for region A depends only on the Betti number b_0 of ∂A , since each component of the boundary places a separate constraint on the loops that intersect the boundary ∂A . Let us take Kitaev's "toric code limit" of the above model² by setting $h = 0$. In this limit, the constraint commutes with the Hamiltonian and can be included as a part of it. Hence, the model may be written as

$$H_{\text{string}, h=0} = - \sum_{\square} \prod_{l \in \square} \tau_{z,l} - \sum_{\text{vertex}} \prod_{\text{vertex} \in l} \tau_{x,l}. \quad (9)$$

Interestingly, the ground state of Eq. (9) may be reinterpreted as a superposition of closed *membranes*. This is seen as follows. The first term in the Hamiltonian may be regarded as the constraint $\prod_{l \in \square} \tau_{z,l} = 1$. Now consider the dual lattice each of whose plaquettes is pierced by a link "l" of the original cubic lattice. A surface can be defined by the plaquettes of the dual lattice pierced by $\tau_{z,l} = -1$ bonds. Due to the constraint, this surface is closed. Thus, there is no distinction between condensed loops and condensed membranes in this case,^{13,15} consistent with the fact that in three dimensions there is only one kind of TEE.

Discrete gauge theories in $D = 3$. Before moving on to the discussion of topological entropy in general dimensions, let us derive the entanglement entropy corresponding to a discrete gauge theory with general gauge group G for Kitaev model² on a cubic lattice²⁴

$$H = -t \sum_p \delta(g_1 g_2 g_3 g_4 = e) - V \sum_{s,g} L_g^1 L_g^2 L_g^3 L_g^4. \quad (10)$$

Here "p" stands for a plaquette, "s" for a star (i.e., six links emanating from a vertex), while g 's are the elements of group G with size of group being $|G|$. For non-Abelian groups one needs to choose an orientation of the links so that for opposite orientations, the group element on a link is g and g^{-1} . The operators L_g^s live on the links and their action is described by $L_{g_1} |g_2\rangle = |g_1 g_2\rangle$ or $L_{g_1} |g_2\rangle = |g_2 g_1^{-1}\rangle$, depending on whether g_1 points away from or toward the vertex at which the action of L_g is being considered. The ground state of $|\Phi\rangle$ of H is given by

$$|\Phi\rangle = \sum_{\{g\}, g_1 g_2 g_3 g_4 = e \forall \text{ plaquettes}} |\{g\}\rangle. \quad (11)$$

Let us divide the entire system into regions A and B and assume that the boundary is made up of plaquettes of the lattice. The links along the boundary are labeled by the group elements h_1, h_2, \dots, h_n . The Schmidt decomposition of $|\Phi\rangle$ reads

$$|\Phi\rangle = \sum_{\{h\}} |\phi\}_{\text{in}}^{\{h\}} \otimes |\phi\}_{\text{out}}^{\{h\}}, \quad (12)$$

where

$$|\phi\}_{\text{in}}^{\{h\}} = \sum_{\substack{\{g\}, g_1 g_2 g_3 g_4 = e \forall \text{ plaquettes} \in A, \\ g_i = h_i \text{ for } i \in \partial A}} |\{g\}\rangle \quad (13)$$

and $|\phi\}_{\text{out}}^{\{h\}}$ is defined similarly. All the states in the Schmidt decomposition enter with the same weight and are orthogonal;

therefore, the entanglement entropy is the logarithm of the number of states. These may be counted by finding all the configurations for $\{h\}$ that satisfy the following constraint: The product of the $\{h\}$'s around any closed loop on the boundary must equal the identity.^{3,4} This includes contractible as well as noncontractible loops on the surface, and each independent loop reduces the total number of configurations by a factor of $|G|$, leading to

$$S = \log(|G|^{V-1}) = V \log(|G|) - \gamma, \quad (14)$$

where V is the number of vertices on the boundary and $\gamma = \log(|G|)$ is the TEE. This result for TEE is identical to that for discrete gauge theories in $D = 2$.

A. Extracting topological entanglement entropy in $D = 3$

In the spirit of Refs. 4 and 5 we would like to combine the total entanglement entropies of certain regions in such a way that the local part of the entropy cancels out while topological part survives.

The Kitaev-Preskill construction, which succeeds in this task in two dimensions can be modified so that it works for three dimensions as well (Ref. 11 describes an extension of the Levin-Wen scheme to $D = 3$). The construction involves three regions A, B, C embedded inside region D :

$$-\gamma_{\text{topo}} = S_A + S_B + S_C - S_{AB} - S_{BC} - S_{CA} + S_{ABC}. \quad (15)$$

In two dimensions, the regions are taken to be three 120° segments of a circle. In three dimensions, a direct generalization of the two-dimensional construction [dividing a *cylinder* into three sectors as in Fig. 3(c)] fails to be topologically invariant because the changes in the entropy near the points at the top and bottom of the cylinder where A, B, C , and D all meet do not cancel. However, the regions such as the two shown in Figs. 3(a) and 3(b) can be used where such points do not exist. For example, if one deforms the circle at which regions A, B, D all meet (in either geometry), then

$$\begin{aligned} \Delta \gamma_{\text{topo}} &= -[\Delta(S_A - S_{CA}) + \Delta(S_B - S_{BC}) + \Delta(S_{ABC} - S_{AB})] \\ &= 0. \end{aligned} \quad (16)$$

The last equation follows because each of the three terms in the brackets could be thought of as the difference between entropies of two regions that differ by addition of region C (that is located far from the point where A, B , and D meet). Since each region has a single boundary component, $\gamma_{\text{topo}} = S_{\text{topological}}$. In Appendix C, we detail the general requirements for a construction that would always yield a topological invariant.

Based on our earlier discussion of curvature expansion for entanglement entropy, we note that in the special case of a completely flat boundary between a region and the rest of the system, the constant term in the entanglement entropy corresponding to that region can indeed be identified with TEE.²⁵ This can be realized by taking the total system to be T^3 and region A as $T^2 \times l$, where l is a line segment (similar to the calculation of entanglement entropy for a free scalar in Appendix A).




FIG. 3. Panels (a) and (b) show two valid ABC constructions [Eq. (15)] in three dimensions that can be used to extract the TEE. In (a) the cross section of a torus has been divided into three tori A , B , and C , while in (b) a torus that has been divided into three cylinders A , B , and C . Panel (c) shows an *invalid* construction as explained in the text. In all three figures, we define region D to be the rest of the system.

V. TOPOLOGICAL ENTANGLEMENT ENTROPY IN $D > 3$

Independent terms in $S_{\text{topological}}$ in arbitrary dimensions. Following our discussion of TEE in $D = 3$, in this section we study the independent contributions to $S_{\text{topological}}$ in a general dimension $D > 3$. The boundary ∂A of a D -dimensional region A is a compact manifold that is characterized by Betti numbers, b_0, \dots, b_{D-1} that describe various orders of connectivity of the surface (see, e.g., Ref. 26).

We assume a linear relationship, $S_A = -\sum_{k=0}^{D-1} \gamma_k b_k$. In principle, in higher dimensions the entanglement entropy could depend on more subtle topological properties of the boundary, but we focus only on this form. Further, as we see below, this form turns out to be sufficient for Kitaev models that describe discrete p -form gauge theories ($p \geq 1$) in arbitrary dimensions.

To see how many types of topological entropy can exist in higher dimensions, first note that for compact manifolds, the Betti numbers have a symmetry, $b_k = b_{D-1-k}$ and hence the sum may be cut short, at $k = \lfloor \frac{D-1}{2} \rfloor$. Furthermore, owing to the relation $\chi = \sum_{k=0}^{D-1} (-1)^k b_k$, in all odd dimensions a part of the topological entropy may be absorbed into the local entropy, reducing the number of coefficients by one more. Hence, there are n topologically nontrivial contributions to the entanglement entropy in $2n$ and $2n + 1$ dimensions:

$$S_{A, \text{topological}} = \begin{cases} -\gamma_0 b_0 - \gamma_1 b_1 - \dots - \gamma_{\frac{D}{2}-1} b_{\frac{D}{2}-1}, & \text{if } D \text{ is even,} \\ -\gamma_0 b_0 - \gamma_1 b_1 + \dots - \gamma_{\frac{D-3}{2}} b_{\frac{D-3}{2}}, & \text{if } D \text{ is odd.} \end{cases} \quad (17)$$

Precisely such a hierarchy of states associated with different Betti numbers has been arrived at by Ref. 13 by constructing

a sequence of Kitaev “toric-code” type models where the ground state is a superposition of all p -dimensional manifolds on a lattice (for $1 \leq p \leq D - 1$). This state is dual to the superposition of all $q = D - p$ dimensional manifolds, so the number of distinct models is $\lfloor \frac{D}{2} \rfloor$, the same as the number of types of topological entropies.

$S_{\text{topological}}$ for gauge theories in arbitrary dimensions. Similar to three dimensions, one may study models of discrete gauge theories to understand these results. For example, on a hypercubic lattice in $D = 4$, the string and membrane theories describe very different ground states¹⁴ and unlike $D = 3$, the membrane theory is now dual to itself, not to the string phase. Explicitly, in the “toric code limit”^{2,14} these two theories are given by

$$H_{\text{string}} = -\sum_{\square} \prod_{l \in \square} \tau_{z,l} - \sum_{\text{vertices}} \prod_{\text{vertex} \in l} \tau_{x,l}, \quad (18)$$

$$H_{\text{membrane}} = -\sum_l \prod_{l \in \square} \sigma_{z,\square} - \sum_{\text{cubes}} \prod_{\square \in \text{cube}} \sigma_{x,\square}. \quad (19)$$

As we show now, the entanglement entropy of the model in Eq. (18) in four dimensions depends on the Betti number b_0 of ∂A , while that corresponding to the model in Eq. (19) depends on the difference $b_1 - b_0$. For the sake of generality, let us derive the entanglement entropy of a generalized toric model in arbitrary spatial dimensions D whose ground state is given by sum over all closed d_g dimensional membranes. This ground state describes deconfined phase of a d_g -form Abelian gauge theory. These membranes intersect the boundary ∂A of region A in closed membranes of dimension $d_g - 1$, with the restriction that these intersections are always boundaries of a membrane of dimension d_g contained in ∂A . For example, consider the entanglement of membrane model in Eq. (19) in $D = 3$ when the boundary of region A is a torus T^2 (note that the form of Hamiltonian for membrane theory is identical in $D = 3$ and $D = 4$). When a closed membrane intersects $\partial A = T^2$, one sees that one can only obtain an even number of closed loops along any noncontractible cycle of T^2 , which would therefore form the boundary of two-dimensional membrane. Returning to the general case, let us denote the number of independent n -dimensional membranes that belong to ∂A by C_n and those that are boundary of a $n + 1$ -dimensional membrane by B_n .

Using the definition of Betti numbers²⁶ and simple linear algebra, one finds that the entanglement entropy S_A

$$S_A \propto \sum_{n=0}^{d_g-1} (-1)^{d_g-1+n} C_n - \sum_{n=0}^{d_g-1} (-1)^{d_g-1+n} b_n. \quad (20)$$

Since the C_n are expressed in terms of local quantities such as the number of edges, vertices, etc., that lie on the boundary without any additional constraint, we identify the first sum as S_{local} and the second as $S_{\text{topological}}$. The proportionality constant depends on the gauge group and akin to three dimensions equals $\log(|G|)$, where $|G|$ is the number of elements in the Abelian gauge group (note that the calculation of TEE in $D = 3$

[Eq. (14)] applies to Abelian as well as non-Abelian discrete gauge theories). Therefore,

$$S_{\text{topological}} = -\log(|G|) \sum_{n=0}^{d_g-1} (-)^{d_g-1+n} b_n. \quad (21)$$

Extracting topological entropy in four dimensions. We restrict our discussion of extracting $S_{\text{topological}}$ to four dimensions. For a given region A , from Eq. (17) one has $S_A = S_{A,\text{local}} - b_0\gamma_0 - b_1\gamma_1$ and one would like to have a construction similar to Levin-Wen⁴ and/or Kitaev-Preskill⁵ that enables one to extract the topological numbers γ_0 and γ_1 .

We extract γ_1 by a generalization of the construction in Ref. 4. Let region A have the topology of $B^2 \times S^2$. Region B is A with a channel cut in it and has topology of B^4 . Finally, region C has a second identical channel cut out opposite to the first one and has topology $S^1 \times B^3$.

Now $S_A - 2S_B + S_C$ is topologically invariant just as in two dimensions and the Betti numbers of the bounding surfaces are

$$\begin{aligned} b_0(\partial A) &= 1, & b_1(\partial A) &= 1, \\ b_0(\partial B) &= 1, & b_1(\partial B) &= 0, \\ b_0(\partial C) &= 1, & b_1(\partial C) &= 1. \end{aligned}$$

Hence, $(S_A - S_B) - (S_B - S_C) = -2\gamma_1$. Since $\gamma_1 \neq 0$ for the membrane Kitaev model H_{membrane} [Eq. (19)] while it is zero for the string model H_{string} [Eq. (18)] in $d = 4$, this construction measures membrane correlations.

To isolate γ_0 , the analogous procedure, but with A being $B^3 \times S^1$, suffices. The combination $(S_A - S_B) - (S_B - S_C)$ gives $\gamma_0 + \gamma_1$, and this may be combined with the previous construction to extract both γ_0 and γ_1 . This construction selectively measures string correlations since $\gamma_0 + \gamma_1$ is zero for H_{membrane} .

The Kitaev-Preskill construction of dividing a disk into three triangles that meet at the center is readily extended to any even dimension. In $D = 4$ consider dividing the ball B^4 into five ‘‘pentahedra’’ that meet at the center. The combination

$$\Delta(\{S\}) = \sum_i S_i - \sum_{i < j} S_{ij} + \dots + S_{12345} \quad (22)$$

is topologically invariant and gives $-\gamma_0$. Here S_{i_1, \dots, i_n} denotes the entanglement entropy corresponding to the region $A_{i_1} \cup A_{i_2} \dots \cup A_{i_n}$.

VI. DISCUSSION AND CONCLUSION

In this paper, we discussed the qualitative structure of the entanglement entropy for gapped phases. We introduced the concept of ‘‘entanglement entropy density’’ whose integral over the boundary of region A yields the entanglement entropy of region A . For gapped trivial phases the symmetry constraints on the entropy density, including the inside-outside exchange symmetry $S_A = S_{\bar{A}}$, naturally lead to the leading and subleading dependence of the entanglement entropy on the linear size of a given region.

In the second half of the paper, we studied the TEE $S_{\text{topological}}$ of topologically ordered systems in various dimensions. A key result was that in $D = 3$ there is a single category of TEE, as in $D = 2$, that depends linearly on the number of

connected components of the boundary. This constrains the possible forms of topological order in $D = 3$.

We briefly discussed TEE in higher dimensions, using generalized Kitaev toric code like models (i.e., deconfined phases of p -form discrete gauge theories) to realize various topologically ordered phases. In $D = 4$ we find two categories of TEE. In general, one new category of TEE appears each time the dimension is raised by two. This even-odd effect is understood as follows. $S_{\text{topological}}$ depends on the Betti numbers of the boundary of region A . In odd spatial dimensions, the Gauss-Bonnet theorem relates Betti numbers to the curvature of the boundary of region A . This implies that there is one linear combination of Betti numbers that can be expressed as an integral of a local property of the boundary (such as curvature) and is thus not an independent topological contribution to the entanglement entropy. We also mentioned how to extract $S_{\text{topological}}$ by a generalization of the $D = 2$ Kitaev-Preskill and Levin-Wen constructions.

Potentially, in $D \geq 4$, $S_{\text{topological}}$ may depend not only on Betti numbers of the boundary manifold, but on more subtle topological properties such as its homotopy group. If such phases do exist, then entanglement entropy could shed light on the classification of manifolds. Lattice three-dimensional models realize a richer variety of topological phases than the isotropic phases considered here. For example, there exist layered Z_2 topologically ordered phases, which retain a two-dimensional character despite coupling between layers. Another example is the self-correcting quantum memory of Ref. 23. For these, the separation between the topological and the local part of the entanglement entropy is not obvious. General statements about entanglement in such topological phases remain for future work.

One might also consider a curvature expansion for the fluctuations of a conserved quantity such as particle number or total spin inside region A . Intuitively, these would be a property of the boundary of region A .²⁷⁻³⁰ Indeed, akin to entanglement entropy, one has $F_A = F_{\bar{A}}$, where $F_A = \sqrt{\langle (\sum_{r \in A} O_r)^2 \rangle - \langle \sum_{r \in A} O_r \rangle^2}$ is the variance of O inside region A . Therefore, a curvature expansion for F_A would inherit many of the arguments we used to derive the leading and subleading behavior of the quantity F_A and can provide a framework to understand known results.²⁷⁻³²

Finally, it may be possible to learn more about the systematics of the size dependence of entanglement entropy in *gapless* phases by a generalization of the curvature expansion under certain conditions. Many gapless systems such as massless scalar/Dirac fermion also follow an area law and have an expression for entropy with interesting parallels to Eq. (4).

ACKNOWLEDGMENTS

We thank Michael Levin for illuminating discussions and Matt Hastings for helpful comments on the manuscript. Support from NSF DMR-0645691 is acknowledged.

APPENDIX A: ABSENCE OF A CONSTANT TERM FOR MASSIVE SCALAR IN THE ABSENCE OF CURVATURE

We are interested in the entanglement entropy of a massive scalar field in three dimensions when region A has a geometry

$T^2 \times l$, where the torus T^2 extends along the directions 1,2 while l is a line segment of length l along the direction 3. In particular, we want to show that the constant part of the entanglement entropy is, in fact, exactly zero; that is, $S = AL^2 + O(1/L)$. This is consistent with the curvature expansion of entanglement entropy [Eq. (5)] since now region A does not have any extrinsic curvature.

The Euclidean action S is given by

$$S = \int |\phi(k_1, k_2, k_3, \omega)|^2 \{m^2 + \omega^2 + \gamma^2[3 - \cos(k_1) - \cos(k_2) - \cos(k_3)]\}, \quad (\text{A1})$$

where we impose periodic boundary conditions in all directions and we have set the lattice spacing to unity. Periodic boundary conditions imply momenta k_1, k_2 remain good quantum numbers even after making the partition. The total entanglement entropy may therefore be written as

$$S = \sum_{k_1, k_2} S_{1D}(M(k_1, k_2, m)), \quad (\text{A2})$$

where $S_{1D}(M)$ is the entanglement entropy of a one-dimensional massive scalar theory with mass $M = \sqrt{m^2 + \gamma^2[2 - \cos(k_1) - \cos(k_2)]}$. Using³³ $S_{1D}(M) \propto -\log(M^2)$,

$$S \propto - \sum_{k_1, k_2} \log\{m^2 + \gamma^2[2 - \cos(k_1) - \cos(k_2)]\}. \quad (\text{A3})$$

Using Euler-Maclaurin formula, one finds the following expression for S , correct to $O(L^0)$:

$$S \propto I_1 + I_2, \quad (\text{A4})$$

where

$$-I_1 \simeq L^2 \log(m^2) + L^2 \int_0^1 dt \log\{1 + \gamma^2/m^2[1 - \cos(2\pi t)]\}$$

and

$$\begin{aligned} -I_2 &\simeq 2\pi\gamma^2 L^2 \int_{u=1/L}^1 \int_{t=0}^1 \frac{t \sin(2\pi t)}{m^2 + \gamma^2[2 - \cos(2\pi t) - \cos(2\pi u)]} \\ &\quad + 2\pi\gamma^2 L \int_0^1 \frac{t \sin(2\pi t)}{m^2 + \gamma^2[1 - \cos(2\pi t)]} \\ &\simeq 2\pi\gamma^2 L^2 \int_{u=0}^1 \int_{t=0}^1 \frac{t \sin(2\pi t)}{m^2 + \gamma^2[2 - \cos(2\pi t) - \cos(2\pi u)]}. \end{aligned}$$

Clearly, neither I_1 nor I_2 contribute to a constant term in the entanglement entropy. Therefore, as anticipated from the curvature expansion, the entanglement entropy $S = I_1 + I_2$ does not contain a constant term for the $T^2 \times l$ geometry and is proportional to L^2 up to $O(L^0)$.

APPENDIX B: ADDITIONAL TERMS IN ENTANGLEMENT ENTROPY IN THE ABSENCE OF ROTATIONAL/PARITY SYMMETRY

A. Broken rotational symmetry

When rotational symmetry is broken, all powers of L (after the area-law term) are present, except in two dimensions. In

two dimensions, there is no constant term when the boundary of the region is smooth. So if rotational symmetry is broken, it is not as easy to recognize a topological phase in even dimensions higher than two without resorting to the ABC construction [Eq. (15)] or its analogs.

In one dimension, when symmetry is broken, it is convenient to express the entropy in terms of $x(s)$ and $y(s)$, the parametric equation for the boundary. Because rotational symmetry is broken, there is no requirement that these terms appear symmetrically. However, because of translational symmetry, the entropy is a function only of the derivatives of these functions. The only requirement is that the entropy density must be symmetric under $s \rightarrow -s$. Otherwise, the entropy depends on whether s is measured clockwise or counterclockwise around the region, and this violates the symmetry between the inside and outside. (Clockwise is defined relative to a choice of the region's inside, as is familiar from using residues to evaluate integrals in the complex plane.) The expression $\int (\frac{d^2x}{ds^2})^3 ds$ is symmetric and it scales as $\frac{1}{L^2}$. One can check that the integral is nonzero for an ellipse (assume the ellipse has a small eccentricity so that the integral can be evaluated; one can then expand it to first order in the eccentricity).

To see that there is no scale-independent term, notice that such a term would have to result from integrating an entropy density with units of $1/L$. Such terms are of the form

$$F(s) = f\left(\frac{dx}{ds}, \frac{dy}{ds}\right) \left(\frac{d^2x}{ds^2}\right) + g\left(\frac{dx}{ds}, \frac{dy}{ds}\right) \frac{d^2y}{ds^2}, \quad (\text{B1})$$

where s is the arclength.

These terms are total derivatives: Let α define the angle of the tangent vector. Then $\frac{dx}{ds} = \cos \alpha$, $\frac{dy}{ds} = \sin \alpha$. The term equals $[-f(\cos \alpha, \sin \alpha) \sin \alpha + g(\cos \alpha, \sin \alpha) \cos \alpha] \frac{d\alpha}{ds}$.

Now the total entropy, $\int F(s) ds$ can be rewritten as an integral with respect to α :

$$\int [-f(\cos \alpha, \sin \alpha) \sin \alpha + g(\cos \alpha, \sin \alpha) \cos \alpha] d\alpha. \quad (\text{B2})$$

By the symmetry $s \rightarrow -s$, f and g must be even functions of cosine and sine. Therefore, the integral is equal to zero since the contributions from α and $\alpha + \pi$ cancel one another.

While there are no constant terms for smooth regions in two dimensions, shapes with corners do have scale-independent terms that can be attributed to the corners (this can happen even if the rotational symmetry is not broken, as shown for the quantum Hall state in Ref. 22).

The first anomalous term in two dimensions scales as $\frac{1}{L^2}$; one example of such a term is

$$\int ds \left(\frac{\partial^2 x}{\partial s^2}\right)^3. \quad (\text{B3})$$

This expression does not vanish identically, as can be seen by an example of a region which is nearly circular, that is, described by the polar coordinates

$$r = 1 + \epsilon(\theta), \quad (\text{B4})$$

where $\epsilon \ll 1$. Evaluating the integral to linear order in ϵ , one obtains

$$\int ds \left(\frac{\partial^2 x}{\partial s^2} \right)^3 \approx - \int ds (\cos^3 \theta) [1 - 6\epsilon(\theta) - 3\epsilon'(\theta) \tan \theta], \quad (\text{B5})$$

which is nonzero for $\epsilon(\theta) \propto \cos 3\theta$.

In higher dimensions, the entropy can have terms that depend on $\frac{\partial x^i}{\partial u^\alpha}$ and higher derivatives. When rotational symmetry is broken, the Cartesian indices do not have to be contracted but the indices for the coordinates on the surface still do, because the entropy has to be independent of the coordinate system. The first allowed correction to the entropy scales as $\frac{L^{D-1}}{L}$ and a simple term in the entropy density that leads to such a correction

$$\left(\frac{D^2 x}{Du^\alpha Du^\beta} g^{\alpha\beta} \right) \left(\frac{\partial x}{\partial u^\gamma} \frac{\partial x}{\partial u^\delta} g^{\gamma\delta} \right). \quad (\text{B6})$$

The first factor gives the proper scaling, $\frac{1}{L}$. The other factors ensure that the expression is not a total derivative in dimensions above two. (The simplest term that scales as $\frac{1}{L}$, $\frac{D^2 x}{Du^\alpha Du^\beta} g^{\alpha\beta} = \nabla^2 x$, is a total derivative.) The indices are all contracted in pairs so the expression is coordinate invariant.

Once a term of order L^{D-2} has appeared, one would expect that all terms of lower orders appear too, and that is what one finds. Since the L^0 term is especially important, we checked explicitly that in even dimensions greater than two, a scale-invariant term L^0 is allowed as long as there is no symmetry. As an example, take Eq. (B6) and multiply it by a power of the mean curvature, $(g^{\alpha\beta} \kappa_{\alpha\beta})^{D-2}$. This has the units L^{1-D} ; therefore, it gives a scale-invariant contribution when integrated (note that since D is even, there are an even number of powers of κ as required). Both Eq. (B6) and this term with the extra factors of κ give a nonzero entropy for a generic region. As an example, consider the D -dimensional surface of revolution obtained by rotating Eq. (B4) (which can be regarded as a curve in the x_1, x_2 plane in D -dimensions) around the x_1 axis [explicitly, $x_1 = r(\theta) \cos \theta, \sqrt{(x_2)^2 + \dots + (x_D)^2} = r(\theta) \sin \theta$]. Both integrals are nonzero for $\epsilon(\theta) \propto \cos 3\theta$.

B. Broken parity symmetry

Breaking just parity can also lead to terms whose exponents deviate from Eq. (4) even if rotational symmetry still exists, when the dimension is odd. However, these additional terms all vary as a negative power of L . In even dimensions, parity does not lead to any additional terms.

The extra terms arise from an additional tensor, γ , that is allowed when parity is broken. This tensor is related to the antisymmetric tensor ϵ :

$$\gamma^{\alpha_1, \dots, \alpha_{D-1}} = \frac{1}{\sqrt{g}} \epsilon^{\alpha_1 \alpha_2, \dots, \alpha_{D-1}}, \quad (\text{B7})$$

where \sqrt{g} is the square root of the determinant of the metric. (This coefficient is necessary for ensuring that γ transforms as a tensor.)

To understand these results, first note that the γ tensor, like the κ tensor, depends on how one chooses the normal to




FIG. 4. Defining the orientation of a hypersurface from an orientation of space, illustrated in three dimensions. A pair of axes on the surface \hat{a}, \hat{b} is defined to be right-handed if the triad $\hat{a}, \hat{b}, \hat{n}$ is right-handed. Formally speaking, γ is defined by contracting the D -dimensional epsilon tensor with the normal \hat{n} and then transforming to curvilinear coordinates, $\gamma^{\alpha_1 \alpha_2, \dots, \alpha_{D-1}} = n^{i_d} \epsilon_{i_1 i_2, \dots, i_d} \frac{\partial x^{i_1}}{\partial u^{\beta_1}} \dots \frac{\partial x^{i_{D-1}}}{\partial u^{\beta_{D-1}}} g^{\alpha_1 \beta_1} g^{\alpha_2 \beta_2} \dots g^{\alpha_{D-1} \beta_{D-1}}$.

the surface. The sign of γ depends on how the orientation of the surface is chosen, and this, in turn, depends not only on the orientation of space (which is determined by the parity-violating ground state) but also on the normal to the surface (see Fig. 4). Since γ is odd under changing the sign of \hat{n} , if a factor of γ appears in the entropy density, an *odd* number n_κ of factors of κ must appear as well,

$$n_\kappa \equiv 1 \pmod{2}. \quad (\text{B8})$$

Now the requirement that all the indices of the κ 's and its derivatives can be contracted with the upper indices of the g 's and the factor of the γ still implies that n_D , the number of covariant derivatives is even if D is odd, because then γ has an even number of upper indices. So a term that includes a factor of γ has units of $L^{-n_\kappa - n_D}$, which is an *odd* power of $\frac{1}{L}$.

In an even number of dimensions, the entropy still goes down by two powers of L at a time because in this case, there must be an odd number of derivatives as well as an odd number of factors of κ to respect both the rotational symmetry and the Z_2 symmetry between the inside and the outside of the region.

The first anomalous term in the entropy is, however, very small, and scales as $\frac{1}{L}$ or $\frac{1}{L^3}$ depending on whether the dimension of space is 1 or 3 modulo 4, respectively. All the terms between the area law term, L^{D-1} , and the constant term go in steps of L^2 . This is essentially because the anomalous terms include factors of γ , which has $D-1$ upper indices which all need to be contracted with something, forcing the term to have at least D factors of κ or covariant derivatives. (One might think that $\frac{D}{2}$ factors of κ should be enough since each κ has two indices; however, since κ is symmetric and γ is antisymmetric, contracting both indices of κ with γ gives 0.)

For illustration, here are some examples of nonzero terms:

$$I_3 = \int dA g^{\sigma\tau} \gamma^{\alpha_1 \alpha_2} D_{\alpha_1} \kappa_{\alpha_2 \sigma} \partial_\tau (\text{tr } \kappa)^2,$$

$$I_5 = \int dA g^{\sigma_1 \sigma_2} \gamma^{\alpha_1 \alpha_2 \alpha_3 \alpha_4} D_{\alpha_1} \kappa_{\alpha_2 \sigma_1} D_{\alpha_3} \kappa_{\alpha_4 \sigma_2} \text{tr } \kappa,$$

for three and five dimensions, respectively, where $\text{tr } \kappa$ is the mean curvature times $D - 1$, $g^{\alpha\beta} \kappa_{\alpha\beta}$. These both scale as $\frac{1}{L^3}$. Both expressions can be generalized to higher dimensions by introducing extra factors of $D_{\alpha_i} \kappa_{\alpha_{k+1} \sigma_{\frac{k+1}{2}}}$ and adding factors of $g^{\sigma_i \sigma_j}$ to contract all the σ 's.

APPENDIX C: VALID CONSTRUCTIONS TO EXTRACT TOPOLOGICAL ENTROPY IN $D = 3$

Let us define $\mathfrak{D}[X(A, B, C)]$, where X denotes some property of a manifolds A, B, C to be

$$\begin{aligned} \mathfrak{D}[X(A, B, C)] &= X_A + X_B + X_C \\ &\quad - X_{AB} - X_{BC} - X_{CA} + X_{ABC}. \end{aligned}$$

For example, in this notation $\gamma_{\text{topo}} = \mathfrak{D}[S(A, B, C)]$. We note that there is one obvious constraint for the Eq. (15) to be useful, which is that the number $\mathfrak{D}[b_0(\partial A, \partial B, \partial C)] \neq 0$. This is because $\gamma_{\text{topo}} \propto \mathfrak{D}[b_0(\partial A, \partial B, \partial C)]$ where the proportionality constant is the universal topological constant associated with the phase of matter.

We claim that in $D = 3$, Eq. (15) still holds as long as the following condition is satisfied:

$$\mathfrak{D}[\chi(\partial A, \partial B, \partial C)] = 0, \quad (\text{C1})$$

where χ denotes the Euler characteristic. This is because this Eq. (C1) guarantees that any dependence on local curvature cancels out on the right-hand side of Eq. (15). Since $\chi = 2 - 2g$, the above equation may be reexpressed as $\mathfrak{D}[g(\partial A, \partial B, \partial C)] = 1$. Just to illustrate this point, consider the case when $A \cap B \cap C \neq 0$. Generically, three regions in $D = 3$ would intersect along a line. Therefore, there are two possibilities: The region C wraps around the surface defined by $A \cap B$ or it does not. First consider the former possibility. In this case, one may easily check that

$$\begin{aligned} g[\partial(AC)] &= g(\partial A) + g(\partial C) - 1, \\ g[\partial(BC)] &= g(\partial B) + g(\partial C) - 1, \\ g[\partial(ABC)] &= g[\partial(AB)] + g(\partial C) - 1. \end{aligned}$$

This implies that $\mathfrak{D}[g(\partial A, \partial B, \partial C)] = 1$ and therefore this is a valid construction. An example is provided by the construction in Fig. 3(a). On the other hand, when C does not wrap around $A \cap B$, then

$$\begin{aligned} g[\partial(AC)] &= g(\partial A) + g(\partial C), \\ g[\partial(BC)] &= g(\partial B) + g(\partial C), \\ g[\partial(ABC)] &= g[\partial(AB)] + g(\partial C), \end{aligned}$$

and therefore $\mathfrak{D}[g(\partial A, \partial B, \partial C)] = 0$, which implies that this is an invalid construction.

APPENDIX D: ENTANGLEMENT ENTROPY OF LAYERED Z_2 TOPOLOGICAL PHASES

Here we briefly discuss the layered topologically ordered states mentioned at the beginning of Sec. II. These phases would lead to a correction $\Delta S_A = -\gamma_{2D} L_z$, for layering perpendicular to the z direction, where γ_{2D} is the TEE




FIG. 5. Extracting entanglement entropy when a three-dimensional topological ordered state coexists with a layered two-dimensional topological order. (a) The ABC construction for this geometry yields $\gamma = L_z \gamma_{2D}$. (b) The ABC construction for this geometry yields $\gamma = \gamma_{3D}$.

associated with the theory living in each layer. For a generic geometry, the total TEE γ may be written as

$$\gamma(L_z) = 2\gamma_{2D} L_z + \gamma_{3D}. \quad (\text{D1})$$

Here L_z is the dimension of regions A, B, C in the z direction. Figure 5 shows two different geometries for which the application of ABC formula [Eq. (15)] yields γ_{2D} and γ_{3D} separately.

APPENDIX E: LINEAR DEPENDENCE OF S_{topo} ON BETTI NUMBERS IN THREE DIMENSIONS

Recapitulating the results from the previous section, in $D = 3$, the topological part of the entanglement entropy is proportional to b_0 and is independent of b_1 since the genus dependence can be obtained by patching local Gaussian curvature. Implicit in these statements is an important assumption which is that entanglement entropy depends *linearly* on b_0, b_1 . This form for the entropy also assumes that the entropy does not depend on knotting or linking of the toroidal surfaces. In this section we provide a proof of these two statements. Our only assumption is that the space in which region A is embedded is flat (i.e., has the topology of \mathbb{R}^3).

We begin by proving the linearity of the entropy for the simplest geometries shown in Figs. 6(a) and 7(a). Figure 6(a) can be used to show that the entropy is linear in b_1 . Consider first the decomposition shown in Fig. 6(b), where the boundary of region A is a k torus (i.e., a torus with genus k) and B and C are three-balls B^3 that join together so that the boundary of region $B \cup C$ is a torus with genus one. We denote by $S_{\text{topo}}(k)$ the topological part of the entanglement entropy corresponding to a region whose boundary has genus k . Now apply the strong subadditivity inequality, $S_{A \cup B} + S_{B \cup C} \geq S_{A \cup B \cup C} + S_B$. The *local* parts of these entropies satisfy an *equality*, $S_{\text{local}}(A \cup B) + S_{\text{local}}(B \cup C) = S_{\text{local}}(A \cup B \cup C) + S_{\text{local}}(B)$ for the configuration shown, and any configuration where A and C do not meet. The reason is that each patch on the boundaries of the regions occurs an equal number of times on both sides of the equation. Hence, the *topological* parts of the entropy also satisfy

$$S_{\text{topo}}(k) + S_{\text{topo}}(1) \geq S_{\text{topo}}(k+1) + S_{\text{topo}}(0). \quad (\text{E1})$$




FIG. 6. Illustration for proving that the entanglement entropy is linear in the first Betti number. Panel (a) shows a genus $k + 1$ torus ($k = 3$), and (b) and (c) give two ways of dividing it up into A, B, C regions. In (b) A is a k torus, B is a ball, and C is also a ball that has been stretched out. In (c) A is a k -torus, B a 1 torus, and C a ball filling up the hole of the 1 torus. Applying strong subadditivity to both configurations gives a recursion formula for $S(k + 1)$ in terms of $S(k)$. Strong subadditivity gives inequalities, but the left- and right-hand sides of the inequality in (b) and (c) are reverses of one another, leading to an exact relation.

Similarly, we construct a different geometry as shown in Fig. 6(c) where region A is the same as in Fig. 6(b), region B has a boundary with genus one, and region C is topologically a ball B^3 . For this topology, adding C to $A \cup B$ decreases the number of handles from $k + 1$ to k , so the strong subadditivity inequality yields the opposite conclusion:

$$S_{\text{topo}}(k + 1) + S_{\text{topo}}(0) \geq S_{\text{topo}}(k) + S_{\text{topo}}(1). \quad (\text{E2})$$

The above two equations imply that $S_{\text{topo}}(k + 1) - S_{\text{topo}}(k)$ is independent of k and hence S_{topo} is linear in k . This genus dependence can be traded for a nontopological contribution plus a b_0 dependence.

To prove that the topological term is linear in the Betti number b_0 (= number of connected components of the boundary) we repeat the above argument by replacing k tori with a region with k spherical cavities cut in it (see Fig. 7). This yields the result that $S_{\text{topo}} = \alpha b_0 + \beta$. Since S_{topo} should vanish for $b_0 = 0$, this suggests that S_{topo} is strictly proportional to b_0 .

Now we show that the entropy of any three-dimensional region, no matter how knotted, is given by the same formula in terms of the Betti numbers. First, we assume that $b_0 = 1$. The theory of surfaces shows that every connected surface embedded in three space is topologically equivalent to one of the k -holed tori. This torus may be built up by attaching handles repeatedly to increase the genus. However, there are many ways to attach a handle. For example, one could add a knotted handle, like in Fig. 8(a). The argument in Fig. 6(c) does not apply to this handle because there is no way to define region C ; for the argument above to work it should fill in the hole in the handle so that the genus decreases by 1, but this is not possible since the handle is knotted through itself.

To generalize the argument we therefore construct a proof of the upper bound that does not require filling the hole in. We choose regions A, B, C that are all subregions of the initial




FIG. 7. Linear dependence of the Betti number on b_0 . (a) A spherical region with $k + 1$ hollow cavities in it ($b_0 = k + 2$ including the outer surface). (b),(c) Two sets of regions for applying strong subadditivity in order to prove that the entropy for $k + 1$ holes minus the entropy for k holes is a constant. (b) The region is divided up into C , containing $k - 1$ of the cavities, a slab B , and a region A containing the remaining cavity. (c) C again contains $k - 1$ of the holes, B is the rest of the region, and A is a solid sphere that fills one of the cavities.

region and the handle itself, so that the argument works if there is some linking.

Ways of building up a region. First we describe all the ways a region whose boundary is a connected genus k surface can be built up starting from a sphere R_0 , so we can be sure that




FIG. 8. Illustration of gluing and drilling. (a) Gluing on a general handle: The torus region is R_k , and the region H is the handle. The handle meets the boundary of region R in two disks. Attaching the handle gives $R_{k+1} = R_k \cup H$. (b) Drilling a hole: The initial region here, R_k , is a solid sphere, and H is a knot that is hollowed out from the inside of the ball to obtain $R_{k+1} = R_k - H$. The boundary of this region is topologically a genus one torus, although it is not possible to continuously deform it into one. Hence drilling increases the genus by one just as adding a handle does.

our argument applies to all of them. Given a region R_k whose boundary is a k torus, one may either glue on a handle H as illustrated in Fig. 8(a) or drill out a hole, as in Fig. 8(b). A handle is a solid cylinder outside region R_k whose two ends are on the boundary of R_k . Attaching H means defining $R_{k+1} = R_k \cup H$.

At some stages it may be necessary to drill a hole out of the region instead. Drilling a hole is an inverted version of the previous process [see Fig. 8(b)]: Let H also have the topology of a cylinder, but $H \subset R_k$, with its circular faces on the surface of R_k . Then drilling the hole H is passing from R_k to $R_{k+1} = R_k - H$. For example, applying this process to a ball, as in Fig. 8(b), gives a “knot complement,” a manifold that is very interesting to knot theorists, since it encodes the structure of the knot.

We now prove that $S_{\text{topo}}(R_{k+1}) - S_{\text{topo}}(R_k) = S_{\text{topo}}(\text{torus}) - S_{\text{topo}}(\text{sphere})$ for the general case. We focus on the case where R_{k+1} is constructed by adding a handle. (The other case is similar: If R_{k+1} is obtained by drilling, then the complements, R_{k+1}^c and R_k^c , are related by handle *adding*, so we can apply our arguments to these complements instead.)

Change of entropy on adding a handle not affected by links passing through H. We first show that the entropy changes by $S_{\text{torus}} - S_{\text{ball}}$ even if the handle H is linked with other portions of R_k . In this argument, we assume that the handle is *not* knotted with itself; the next argument shows that knotting does not affect the entropy.

Consider the regions in Fig. 8(a). We first prove this inequality:

$$S_{\text{topo}}(R_{k+1}) - S_{\text{topo}}(R_k) \geq S_{\text{torus}} - S_{\text{ball}}, \quad (\text{E3})$$

where S_{torus} and S_{ball} are the topological entropies of an unknotted torus and a ball, respectively. Begin by sliding the two ends of the handle along the surface of R_k so that they are right next to each other, as shown in the Fig. 8(a). The proof of this inequality has two steps, using the decompositions illustrated in Figs. 9(b) and 9(c).

First, as in Fig. 9(b), let $B = R_k$, and let A and C be narrow strips along the left and right side of H . $A \cup B$ and $C \cup B$ are two regions that can both be deformed into the same topology




FIG. 9. Figure for the proof that links passing through the handle do not affect the entropy. Panel (a) shows region R_k and the handle (in gray) that is added to it to get R_{k+1} . As in Fig. 6, the goal is to relate the entropy of a $k + 1$ -holed torus to a k -holed one, where $k = 3$. The dotted circle indicates the portion of the figure that is enlarged in the later panels of this figure. The portion of R_k that passes through the hole is not shown in the subsequent frames for clarity. Parts (b) and (c) show the two decompositions that are used to prove the lower bound on $S_{\text{topo}}(R_{k+1}) - S_{\text{topo}}(R_k)$. In (b), there are three regions: Region B (checkered) is R_k . The handle is flattened so that it is ribbonlike, and then striplike regions A and C are demarcated along its edges. These together form $A \cup B \cup C := R_{\text{double}}$ which has the topology of R_k with *two* handles attached. In (c) we decompose R_{k+1} into A , the center of the ribbon, B (checkered), the border of the ribbon closed up with parts of R_k to form a loop, and C , the rest of R_k . In this figure, $B \cup C = R_{\text{double}}$. Part (D) shows the decomposition used to prove the upper bound, which corresponds to the division in Fig. 6(b).




FIG. 10. Showing that the entropy of a knotted handle is the same as if it were an unknotted. (a) Shows the knotted handle H (the gray region) that is attached to region R_k (the white region). Passing one strand through the other so the topology changes to panel (e) causes the handle to become unknotted: The loops in (e) can be untwisted, giving a simple handle. Any knot can become unknotted if the right strands are passed through each other. The intermediate panels show that this process does not change the entropy. Panel (b) is a magnification of panel (a). In panel (c) the topology is changed so that one strand passes through an eyelet in the other. This configuration is obtained by adding a handle along the dotted lines in (b); hence, the entropy changes by $S_{\text{torus}} - S_{\text{sphere}}$ (the handle is unknotted). Panel (d) is obtained by *removing* a handle from (c); the entropy therefore decreases back to its original value. Thus, the change from (a) to (e) does not affect the topological entropy.

(R_{k+1}) ; altogether, $A \cup B \cup C$ forms a region R_{double} with two handles attached to R_k . The strong subadditivity implies

$$2S_{\text{topo}}(R_{k+1}) \geq S_{\text{topo}}(R_k) + S_{\text{topo}}(R_{\text{double}}). \quad (\text{E4})$$

Next, as in Fig. 9(c), let A be the interior of the strip H . Let B be the border of this strip, marked with a checkered pattern in the figure. This border is closed by adding some small parts of R_k so that it surrounds region A completely. Let C be the rest of R_k . In this decomposition, $C \cup B = R_{\text{double}}$, while $A \cup B \cup C = R_{k+1}$, and $A \cup B$ and B are a ball and a torus, respectively. Then

$$S_{\text{topo}}(\text{ball}) + S_{\text{topo}}(R_{\text{double}}) \geq S_{\text{torus}} + S_{\text{topo}}(R_k). \quad (\text{E5})$$

Adding the previous two equations and canceling R_{double} gives Eq. (E3).

In this argument it has not been necessary to add on regions *external* to the handle, so the argument still works if parts of R_k are linked through it.

Now the reverse inequality $S_{\text{topo}}(R_{k+1}) \leq S_{\text{topo}}(R_k) + S_{\text{torus}} - S_{\text{ball}}$ is proved just as in the argument for the simple k tori using the division shown in Fig. 9(d), which is no different than the original construction in Fig. 6(b). Hence,

$S_{\text{topo}}(R_{k+1}) - S_{\text{topo}}(R_k) = S_{\text{torus}} - S_{\text{ball}}$, and we conclude that $S_{\text{topo}}(R_k) = kS_{\text{torus}} - (k-1)S_{\text{ball}}$; that is, the entropy is linear in the genus of the surface.

For the case where the surface has more than one boundary component we just start with a region whose boundary has many components, which are all spheres. The entropy of this region is proportional to b_0 by an argument similar to the one illustrated in Fig. 7 (this applies even when the spheres are nested in each other). It is known that any region may be built up from such a region by either attaching handles or drilling holes. It is even possible to choose the handles and holes so that each one starts and ends on the same component of the boundary of R_k . Then the calculation of the entropy as the handles are attached proceeds just as above.

Change of entropy not affected by knotting of the handle. If the handle is knotted, this argument does not immediately work, because the entropy change $S_{\text{torus}} - S_{\text{ball}}$ could be different when the torus is knotted. However, a knotted handle may be transformed to an unknotted one by repeatedly adding and then removing of *unknotted* handles as illustrated in Fig. 10. By the previous argument, the entropy returns to its original value after this is done. Hence, the knotted and unknotted handles have the same entropy.

¹C. Nayak, S. Simon, A. Stern, M. Freedman, and S. Sarma, *Rev. Mod. Phys.* **80**, 1083 (2008).

²A. Kitaev, *Ann. Phys.* **303**, 2 (2003).

³A. Hamma, R. Ionicioiu, and P. Zanardi, *Phys. Lett. A* **337**, 22 (2005); *Phys. Rev. A* **71**, 022315 (2005).

⁴M. Levin and X.-G. Wen, *Phys. Rev. Lett.* **96**, 110405 (2006).

⁵A. Kitaev and J. Preskill, *Phys. Rev. Lett.* **96**, 110404 (2006).

⁶S. Furukawa and G. Misguich, *Phys. Rev. B* **75**, 214407 (2007).

⁷M. Haque, O. Zozulya, and K. Schoutens, *Phys. Rev. Lett.* **98**, 060401 (2007); O. S. Zozulya, M. Haque, K. Schoutens, and E. H. Rezayi, *Phys. Rev. B* **76**, 125310 (2007); A. M. Lauchli, E. J. Bergholtz, and M. Haque, *New J. Phys.* **12**, 075004 (2010).

⁸Shiyong Dong, Eduardo Fradkin, Robert G. Leigh, and Sean Nowling, *J. High Energy Phys.* **05** (2008) 016.

⁹S. Isakov, M. Hastings, and R. Melko, *Nat. Phys.* **7**, 772 (2011).

¹⁰Y. Zhang, T. Grover, and A. Vishwanath, *Phys. Rev. B* **84**, 075128 (2011).

¹¹C. Castelnovo and C. Chamon, *Phys. Rev. B* **78**, 155120 (2008).

¹²Heuristically, the Betti number b_k measures the dimension of k -dimensional closed surfaces within a manifold that are not boundaries of a $k+1$ -dimensional surface. See, for example, Chap. 3 of Ref. 26 for an introduction.

¹³H. Bombin and M. A. Martin-Delgado, *Phys. Rev. B* **75**, 075103 (2007).

¹⁴Eric Dennis, Alexei Kitaev, Andrew Landahl, and John Preskill, *J. Math. Phys.* **43**, 4452 (2002).

¹⁵A. Hamma, P. Zanardi, and X.-G. Wen, *Phys. Rev. B* **72**, 035307 (2005).

¹⁶See, e.g., P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (Cambridge University Press, Cambridge, 2000).

¹⁷The same symmetry is responsible for relating corner contributions to the entanglement entropy with angles θ and $2\pi - \theta$ (Refs. 5 and 22).

¹⁸H. Casini and M. Huerta, *J. Phys. A* **42**, 504007 (2009).

¹⁹M. Srednicki, *Phys. Rev. Lett.* **71**, 666 (1993).

²⁰Manfredo Do Carmo, *Differential Geometry of Curves and Surfaces* (Prentice Hall, Englewood Cliffs, NJ, 1976).

²¹C. B. Allendoerfer, *Bull. Am. Math. Soc.* **54**, 249 (1948).

²²I. D. Rodriguez and G. Sierra, *J. Stat. Mech.* **1012**, P12033 (2010).

²³J. Haah, *Phys. Rev. A* **83**, 042330 (2011).

²⁴We are grateful to Michael Levin for an illuminating discussion on entanglement entropy of discrete gauge theories.

²⁵Note that in this case there is a ground-state degeneracy and the TEE in general will depend on the particular linear combination of the ground states for which the entanglement is being calculated (see, e.g., Ref. 8).

²⁶M. Nakahara, *Geometry, Topology, and Physics* (A. Hilger, London, 1990).

²⁷H. Francis Song, Stephan Rachel, and Karyn Le Hur, *Phys. Rev. B* **82**, 012405 (2010); H. Francis Song, Christian Flindt, Stephan Rachel, Israel Klich, and Karyn Le Hur, *ibid.* **83**, 161408(R) (2011).

²⁸Israel Klich, Gil Refael, and Alessandro Silva, *Phys. Rev. A* **74**, 032306 (2006).

²⁹Israel Klich, Leonid Levitov, in *Advances in Theoretical Physics: Landau Memorial Conference*, edited by V. Lebedev and M. V. Feigelman, AIP Conf. Proc. No. 1134 (AIP, New York, 2009), pp. 36–45.

³⁰D. Gioev and I. Klich, *Phys. Rev. Lett.* **96**, 100503 (2006).

³¹R. Helling, H. Leschke, and W. Spitzer, *Int. Math. Res. Not.* **2011**, 1451 (2011).

³²B. Swingle, e-print [arXiv:1007.4825](https://arxiv.org/abs/1007.4825).

³³P. Calabrese and J. Cardy, *J. Stat. Mech.* **06** (2004) P06002.