# **Optical conductivity for a dimer in the dynamic Hubbard model**

G. H. Bach<sup>1,2</sup> and F. Marsiglio<sup>1</sup>

<sup>1</sup>Department of Physics, University of Alberta, Edmonton, Alberta, Canada T6G 2G7 <sup>2</sup>Faculty of Physics, Hanoi University of Science, Vietnam National University, 334 Nguyen Trai Street, Hanoi, Vietnam (Received 22 September 2011; revised manuscript received 15 February 2012; published 20 April 2012)

The dynamic Hubbard model represents the physics of a multiband Hubbard model by using a pseudospin degree of freedom to dynamically modify the on-site Coulomb interaction. Here we use a dimer system to obtain analytical results for this model. The spectral function and the optical conductivity are calculated analytically for any number of electrons, and the distribution of optical spectral weight is analyzed in great detail. The impact of polaronlike effects due to overlaps between pseudospin states on the optical spectral weight distribution is derived analytically. Our conclusions support results obtained previously with different models and techniques: holes are less mobile than electrons.

DOI: 10.1103/PhysRevB.85.155134

PACS number(s): 74.20.-z, 74.25.Gz, 74.72.-h

# I. INTRODUCTION

The occurrence of electron-hole asymmetry in tunneling spectra<sup>1</sup> and the anomalous behavior in the optical conductivity sum rule at the superconducting transition temperature<sup>2–6</sup> both contribute to the possibility that the superconductivity in the cuprate materials is unusual in several respects. In particular, the notion of "kinetic energy-driven superconductivity" is implied by the optical experiments, as predicted almost ten years in advance of these experiments.<sup>7</sup>

These experiments indicated that a significant transfer of spectral weight occurs in the cuprates,<sup>8</sup> both in the normal state as a function of temperature and as a result of the superconducting transition. More importantly, perhaps, is the range of frequencies affected by the transition, as a significant amount of spectral weight is transferred from very high frequencies to very low frequencies. Thus there is an apparent violation of the Ferrell-Glover-Tinkham optical sum rule, as an examination of the low-frequency region alone shows a spectral weight discrepancy. This indicates that physics beyond the usual paradigm of "energy lowering due to potential energy considerations" is at work; in particular, the anomalous sign of the change in low-frequency optical spectral weight indicates that some mechanism involving the kinetic energy of the carriers is at work.

Earlier modeling of cuprate superconductivity<sup>9</sup> includes some of this physics-this is what motivated the relatively early theoretical discussion of optical spectral weight transfer-but a more recent theoretical model, advanced more than ten years ago,<sup>10</sup> goes further to explain some of the anomalous features in the spectroscopic measurements of the cuprates; this is the so-called dynamic Hubbard model. This model utilizes a phenomenological pseudospin degree of freedom at each lattice site designed to mimic orbital relaxation effects which necessarily occur in real atoms. As far as optical spectral weight transfer is concerned, this model includes higher frequency excitations (here modeled by the pseudospins), and therefore, while we do not address superconductivity or temperature effects in this paper, by using the dynamical Hubbard model, we can study how spectral weight transfer occurs as a function of doping.

The Hamiltonian for the dynamical Hubbard model is<sup>10</sup>

$$H_{\text{DHB}} = -t \sum_{\langle i,j \rangle \sigma} (c_{i\sigma}^{\dagger} c_{j\sigma} + c_{j\sigma}^{\dagger} c_{i\sigma}) - \mu \sum_{i,\sigma} n_{i\sigma} + \sum_{i} \left( \omega_0 \sigma_i^x + g \omega_0 \sigma_i^z \right) + \sum_{i} \left( U - 2g \omega_0 \sigma_i^z \right) n_{i\uparrow} n_{i\downarrow},$$
(1)

where the pseudospin degree of freedom is here represented by a Pauli operator  $\sigma_i$  at each site; it interacts with the electron charge through the double occupancy term, and contributes a dynamical interaction in addition to the usual Hubbard interaction. The rest of the Hamiltonian is as follows: the first term represents the electron kinetic energy within a tight-binding model with one orbital per site. Note that we are really trying to understand physics that originates in processes involving multiple orbitals. It is desirable to minimize the complexity by retaining a single orbital, and it is then the pseudospin that acts to mimic the physics of carriers undergoing transitions between multiple orbitals when the local occupation changes. The second term determines the electron density through the chemical potential, the third term defines the two-level system for the pseudospin degree of freedom at each site, and the last term is the on-site interaction which, in addition to the short-range Coulomb repulsion represented by U, is also modulated through a coupling constant  $g\omega_0$  by the state of the pseudospin.

When the double occupancy is high, the pseudospin will reside in its excited state for the sake of minimizing the Coulomb repulsion, much like the phenomenon in real atoms, where two electrons will sacrifice having a minimal electronion energy and spread out amongst the excited orbitals in order to minimize their Coulomb repulsion. In the opposite limit, when the double occupancy is very low, electrons will tend to stay in the lowest energy state available in the given atom (loosely, the Wannier state which is being modeled in the tight-binding Hamiltonian), and the Coulomb energy will be high, though irrelevant, since only rarely will two electrons occupy the same site.

The dynamic Hubbard model contains at least some of the phenomenology of hole superconductivity,<sup>9,11–13</sup> proposed

more than 20 years ago. In particular, the model contains electron-hole asymmetry, where holes at the top of a band are heavier than electrons at the bottom of a band. Electronhole asymmetry can arise in a number of ways: just having further than nearest-neighbor hopping can result in a band mass asymmetry, upon which polaronic mass asymmetry can build.<sup>14</sup> Even with just nearest-neighbor hopping, lattice geometry can also result in electron-hole asymmetry, and again be amplified through polaron effects.<sup>15</sup> Here, the electron-hole asymmetry arises through realistic interactions, and is strongly connected to the fact that holes pair more readily than electrons by lowering their kinetic energy in the superconducting phase, a phenomenon supported by the anomalous observations in the optical conductivity sum rule mentioned above. More recently, the dynamic Hubbard model has been explored with dynamical mean-field theory (DMFT).<sup>16</sup> In particular, the spectral function and the optical conductivity were calculated, in the normal state, to illustrate the electron-hole asymmetry present in the model.

In this paper, we will focus on the optical conductivity, and provide a complementary calculation involving a simple dimer. Such a small system does not constitute a very realistic system; however, dimer calculations have an illustrious history for providing insight into models.<sup>17–19</sup> Furthermore, following Ref. 7, a dimer calculation provides good insight into the processes that contribute to the conductivity. We will briefly review the optical conductivity and the sum rule in the next section, and describe the details of the dimer calculation in the third section, in perturbation theory; we have also performed exact diagonalizations to delineate the regime of validity of the perturbative results. This is followed with a discussion of the results, particularly in light of the DMFT results reported in Ref. 16. We then conclude with a summary.

# II. OPTICAL CONDUCTIVITY AND RELATED SUM RULES

The real part of the optical conductivity  $\sigma_1(\omega)$ , as a function of frequency,  $\omega$ , at finite temperature *T* can be written as

$$\sigma_{1}(\omega) = \frac{\pi}{Z} \sum_{n,m} \frac{e^{-\beta E_{n}} - e^{-\beta E_{m}}}{E_{m} - E_{n}} |\langle m|J|n \rangle|^{2} \\ \times \delta \left( \omega - \frac{E_{m} - E_{n}}{\hbar} \right),$$
(2)

where  $|m\rangle$  and  $|n\rangle$  are eigenstates of the Hamiltonian of the system, Z is the partition function,  $\beta = 1/k_BT$ , and J is the current operator, obtained through the polarization operator.<sup>20</sup> As formulated by Kubo in 1957,<sup>21</sup> the optical conductivity satisfies the general sum rule,

$$\int_0^{+\infty} \operatorname{Re}[\sigma_{\mu\nu}(\omega)] d\omega = \frac{\pi}{2} \sum_r \frac{e_r^2 n_r}{m_r} \delta_{\mu\nu}, \qquad (3)$$

where *r* denotes the type of charge carrier,  $n_r$ ,  $e_r$ , and  $m_r$  are the number density, charge, and mass, respectively, of the *r*-type carrier, and  $\mu$ ,  $\nu$  are the indices of the conductivity tensor. For an isotropic electron system, the sum rule (3) is rewritten as

$$\int_0^{+\infty} \sigma_1(\omega) d\omega = \frac{\pi e^2 n}{2m},\tag{4}$$

with *m* the bare mass of electrons and *n* the total electron density.<sup>22</sup>

In condensed matter systems we often work with effective Hamiltonians, for example, formulated for a single band within tight binding. One can then formulate a sum rule restricted to that single band, and  $obtain^{23-25}$ 

$$\int_{0}^{+\infty} \operatorname{Re}[\sigma_{xx}(\omega)] d\omega = \frac{\pi e^2}{4\hbar^2} \left\{ \frac{4}{N} \sum_{k} \frac{\partial^2 \epsilon_k}{\partial k_x^2} n_k \right\}, \quad (5)$$

where  $n_k$  is the single electron occupation number, and  $\epsilon_k$  is the dispersion relation for the noninteracting electrons. In reality, the integration in Eq. (5) is taken up to a cutoff frequency  $\omega_c$  determined experimentally in order to allow only intraband transitions, and to avoid the inevitable interband transitions which are *not* part of the sum rule Eq. (5). Theoretically, the right-hand side (RHS) of Eq. (5) is often used,<sup>26</sup> as this is much simpler to calculate. Furthermore, when only nearest-neighbor hopping is allowed on a hypercubic lattice, the sum rule Eq. (5) reduces to<sup>23</sup>

$$\int_{0}^{\omega_{c}} \sigma_{1}(\omega) d\omega = -\frac{\pi e^{2} a^{2}}{2\hbar^{2}} \langle K \rangle, \qquad (6)$$

with  $K = -t \sum_{\langle ij \rangle, \sigma = \uparrow \downarrow} (c_{i\sigma}^{\dagger} c_{j\sigma} + \text{H.c.})$  and *a* the lattice constant. From the RHS of Eq. (6), it is clear that the optical sum depends not only on the external parameters (such as the temperature) but also on the electronic structure of the system. The validity of using the kinetic energy on the RHS instead of the expression in Eq. (5), even when the dispersion is not just nearest-neighbor hopping has been explored in Refs. 28 and 27, to which the reader is referred. In the following, we assume Eq. (6) holds.

Measurements of the optical conductivity sum rule in a number of the cuprate superconductors generally show an increase of spectral weight in the low-frequency regime in the superconducting state, at least in the underdoped and optimally doped materials.<sup>3–6</sup> This enhancement of the optical sum at the superconducting transition temperature conflicts with the result from BCS-like superconductivity where an increase in the kinetic energy (and therefore a decrease in the single band optical sum) is expected instead. This can be explained through a number of different scenarios, examples of which are preformed pairs<sup>29</sup> and phase fluctuations.<sup>30–32</sup> In contrast to the model considered here, many of these calculations have, as a key ingredient, proximity to a nearby Mott insulating state.<sup>31,33</sup> Note that some authors<sup>26,34</sup> have attributed the anomalous temperature dependence of the lowfrequency optical spectral weight to a cutoff effect (required in the experimental analysis). Karakozov et al.<sup>34</sup> have also attributed the anomalous change in spectral weight at the superconducting transition temperature to a cutoff effect, though this has been refuted in Ref. 25. See Ref. 35 for a brief review.

On the other hand, models like the hole mechanism of "kinetic energy driven" superconductivity support the idea of minimizing the total energy by reducing the kinetic energy, and therefore the optical sum has an anomalous temperature dependence below  $T_c$ .<sup>36</sup> When the system goes superconducting, the missing optical spectral weight is predicted to be distributed over the whole range of frequencies, i.e., weight is transferred

from high frequency to low frequency and the low energy sum rule appears to increase as a consequence.<sup>7,37</sup>

## **III. OPTICAL CONDUCTIVITY IN A DIMER**

We proceed now with a brief discussion of the site Hamiltonian followed by a detailed description of the dimer.

#### A. Properties of the Hamiltonian

Following Refs. 16 and 38, we begin with the on-site Hamiltonian *for electrons*:

$$H_{\rm DHM}^{(i)} = \omega_0 \sigma_x^i + g \omega_0 \sigma_z^i + \left[ U - 2g \omega_0 \sigma_z^i \right] n_{i\uparrow} n_{i\downarrow}.$$
 (7)

The solutions are provided in detail in Refs. 16 and 38; for *n* electrons the ground state  $(|n\rangle)$  and the first excited state  $(|\bar{n}\rangle)$  are

$$|n\rangle = u(n)|+\rangle - v(n)|-\rangle, \quad |\bar{n}\rangle = v(n)|+\rangle + u(n)|-\rangle, \quad (8)$$
  
with

$$u(0) = u(1) = v(2), \quad v(0) = v(1) = u(2),$$
 (9)

and

$$u(0) = \sqrt{\frac{1}{2} \left( 1 - \frac{g}{\sqrt{1 + g^2}} \right)}, \quad v(0) = \sqrt{\frac{1}{2} \left( 1 + \frac{g}{\sqrt{1 + g^2}} \right)}.$$
(10)

The eigenvalues [ground state  $\epsilon(n)$  and excited state  $\bar{\epsilon}(n)$ ] are

$$\epsilon(n) = \delta_{n,2}U - \omega_0\sqrt{1+g^2}, \quad \bar{\epsilon}(n) = \delta_{n,2}U + \omega_0\sqrt{1+g^2}.$$
(11)

Especially important for the hopping processes is the overlap of background spin states with different numbers of electron; these are

$$T = \langle 0|1 \rangle = u(0)u(1) + v(0)v(1) = u(0)^{2} + v(0)^{2} = 1,$$
  

$$S = \langle 1|2 \rangle = u(1)u(2) + v(1)v(2) = 2u(1)v(1) = \frac{1}{\sqrt{1+g^{2}}}.$$
(12)

These parameters play an important role for the spectral function; these are defined in Ref. 39 as  $A_{n+1,n}$  for electron destruction in a system of n + 1 electrons (and  $A_{n,n+1}$  for electron creation in a system of *n* electrons. For a single site these single-particle spectral functions are<sup>39</sup>

$$A_{10}(\omega) = A_{01}(\omega) = \delta(\omega),$$
  

$$A_{12}(\omega) = A_{21}(\omega) = S^2 \delta(\omega) + (1 - S^2) \delta(\omega - \Omega_0).$$
(13)

Even though the spectral weight is calculated for a single site, it is clear that there is a reduction of the weight at zero frequency if the second electron is added to the one-electron ground state. The reason is because there are two possibilities: the pseudospin can remain in the same state as the first electron with a probability  $S^2 < 1$ , or it can become excited with an energy cost  $\Omega_0 = 2\omega_0\sqrt{1+g^2}$ . In the thermodynamic limit, this effect is known as the reduction of quasiparticle weight by transferring part of the coherent contribution (at  $\omega = 0$ ) to the incoherent part (at large  $\omega$ ), resulting in a one-particle spectral weight, z < 1. Since the quasiparticle weight is inversely proportional to the effective mass,  $z \sim m/m^*$ , this statement means that holes are heavier (or more "dressed"<sup>40</sup>) than electrons.

For calculating the optical conductivity in perturbation theory, the Hamiltonian is divided into two parts:

$$H = H_0 + H',$$
 (14)

$$H_0 = \sum_{i} \left( \omega_0 \sigma_i^x + g \omega_0 \sigma_i^z \right) + \left( U - 2g \omega_0 \sigma_i^z \right) n_i \uparrow n_i \downarrow, \quad (15)$$

$$H' = K = -t \sum_{\langle i,j \rangle \sigma} (c^{\dagger}_{i\sigma} c_{j\sigma} + c^{\dagger}_{j\sigma} c_{i\sigma}), \qquad (16)$$

where  $H_0$  is the site Hamiltonian and H' is the hopping part which is considered as a perturbation under the following conditions. Based on the definition of the overlaps between the pseudospin ground states in Eq. (12), we can define a pseudospin state for a given number of electrons in terms of the eigenstates involving a different number of electrons. These overlaps contain the background deformations (modeled by the pseudospin) that must be "dragged"along as the electron hops. Thus, following Ref. 7,

$$|1\rangle = S|2\rangle - \bar{S}|\bar{2}\rangle, \qquad (17)$$

$$|\bar{1}\rangle = \bar{S}|2\rangle + S|\bar{2}\rangle, \qquad (18)$$

$$|2\rangle = S|1\rangle + \bar{S}|\bar{1}\rangle, \qquad (19)$$

$$|\bar{2}\rangle = -\bar{S}|1\rangle + S|\bar{1}\rangle, \qquad (20)$$

where

$$\bar{S} = \sqrt{1 - S^2} = \frac{g}{\sqrt{1 + g^2}}.$$
 (21)

We wish to solve this problem in all number sectors (one, two, and three electrons). We cover these in the following subsections.

#### B. Three electron sector

Beginning with three electrons on the dimer (a holelike configuration), the (non-normalized) ground-state wave function is given in first-order perturbation theory as

$$\left|\psi_{0}^{(3)}\right\rangle = |1\rangle_{o} + \frac{\sqrt{2}tS\bar{S}}{\Omega_{0}}|3\rangle_{o} - \frac{t\bar{S}^{2}}{2\Omega_{0}}|4\rangle_{o}, \qquad (22)$$

where

$$\begin{split} |1\rangle_{a}^{e} &= \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |2\rangle_{a} \otimes c_{b\uparrow}^{\dagger} |1\rangle_{b} \pm c_{a\uparrow}^{\dagger} |1\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |2\rangle_{b}], \\ |2\rangle_{a}^{e} &= \frac{1}{2} [\pm c_{a\uparrow}^{\dagger} |1\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |\bar{2}\rangle_{b} \pm c_{a\uparrow}^{\dagger} |\bar{1}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |2\rangle_{b} \\ &+ c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |2\rangle_{a} \otimes c_{b\uparrow}^{\dagger} |\bar{1}\rangle_{b} + c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |\bar{2}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} |1\rangle_{b}], \\ |3\rangle_{a}^{e} &= \frac{1}{2} [\mp c_{a\uparrow}^{\dagger} |1\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |\bar{2}\rangle_{b} \mp c_{a\uparrow}^{\dagger} |\bar{1}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |2\rangle_{b} \\ &+ c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |2\rangle_{a} \otimes c_{b\uparrow}^{\dagger} |\bar{1}\rangle_{b} + c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |\bar{2}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} |1\rangle_{b}], \\ |4\rangle_{a}^{e} &= \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |\bar{2}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} |\bar{1}\rangle_{b} \pm c_{a\uparrow}^{\dagger} |\bar{1}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |\bar{2}\rangle_{b}], \end{aligned}$$

are the eight basis states required to span the Hilbert space for the three electron sector. Here the subscript e refers to the even states, and a and b are the indices of the first and second site, respectively, in the dimer. Note that kets followed by the subscript a or b have numbers 0, 1, or 2 (with or without bars on top) that refer to the pseudospin eigenstates defined in Eq. (8), whereas kets followed by e (for even) or o (for odd) refer to linear combinations of product states of electrons and pseudospin eigenstates, as, for example, in Eq. (23).

The three-particle ground-state energy, to first order in the hopping perturbation, is given by

$$E_0^{(3)} \equiv {}_o\langle 1|H|1\rangle_o = U - tS^2 - \Omega_0,$$
(24)

and the excited-state energies for the three electron sector are

$$E_{1}^{(3)} \equiv {}_{e}\langle 1|H|1\rangle_{e} = U + tS^{2} - \Omega_{0},$$
  

$$E_{2}^{(3)} \equiv {}_{o}^{e}\langle 2|H_{0}|2\rangle_{o}^{e} = U, \quad E_{3}^{(3)} \equiv {}_{o}^{e}\langle 3|H_{0}|3\rangle_{o}^{e} = U, \quad (25)$$
  

$$E_{4}^{(3)} \equiv {}_{o}^{e}\langle 4|H_{0}|4\rangle_{o}^{e} = U + \Omega_{0}.$$

Note that there are degeneracies at zeroth order between even and odd states; these are broken in first-order perturbation theory, as is explicitly written in Eq. (24) and the first of Eqs. (25). Only the zeroth-order energies (and wave functions) are needed for the other excited states, and that is what is written here. Also, the Hamiltonian will only couple states of a given parity, whereas the conductivity will couple only states of opposite parity.

The optical conductivity for the dimer at zero temperature can be calculated for the three electron sector as

$$\sigma_1(\omega) = \pi \sum_{m \neq 0} \frac{\left| \left( \psi_0^{(3)} |J|m \right)_e \right|^2}{E_m^{(3)} - E_0^{(3)}} \delta\left( \omega - \frac{E_m^{(3)} - E_0^{(3)}}{\hbar} \right), \quad (26)$$

where  $|m\rangle_e$  are the excited states of the system [only even parity is required since the ground state has odd parity—these are given in Eq. (23)], and

$$J = \frac{iet}{\hbar} \sum_{\sigma} (c^{\dagger}_{a\sigma} c_{b\sigma} - c^{\dagger}_{b\sigma} c_{a\sigma})$$
(27)

is the current operator. By acting with the *J* operator on the ground state  $|\psi_0^{(3)}\rangle$ , we connect to three of the excited states (all even parity) given in Eq. (23). Note that, since the current operator is already of order *t*, only zeroth-order wave functions are required, but for the first even parity excited state, first-order corrections to the energy are required in the denominator of Eq. (26) to break the degeneracy.

Operating with the current operator on the unperturbed ground state gives

$$J|1\rangle_0 = \frac{iet}{\hbar} (S^2|1\rangle_e - \sqrt{2}S\bar{S}|3\rangle_e - \bar{S}^2|4\rangle_e), \qquad (28)$$

so that the optical conductivity for three electrons includes three peaks. These three transitions are shown schematically in Fig. 1; the analytical expression for the optical conductivity (for three electrons) is

$$\sigma_{1}^{(3)}(\omega) = \frac{\pi e^{2}t}{2\hbar^{2}} \left[ S^{2}\delta\left(\omega - \frac{2tS^{2}}{\hbar}\right) + 4S^{2}\bar{S}^{2}\frac{t}{\Omega_{0}}\delta\left(\omega - \frac{\Omega_{0}}{\hbar}\right) + \bar{S}^{4}\frac{t}{\Omega_{0}}\delta\left(\omega - \frac{2\Omega_{0}}{\hbar}\right) \right].$$
(29)



FIG. 1. Schematic depiction of optical transitions in a dimer with three electrons. The lines (both solid and dashed) show the two available levels of the pseudospin energy at each site; the solid lines correspond to the occupied pseudospin state and the dashed lines correspond to the unoccupied state. Transitions between states with the same pseudospin energy levels are diagonal; these contribute to the intraband conductivity, while nondiagonal transitions between states with different pseudospin energy levels modify the interband conductivity. State labels are those found in Eq. (23) in the text, where they are given in full even or odd form.

The optical sum rule can be checked by calculating the expectation value of the K operator in the ground state,

$$\left\langle \psi_{0}^{(3)} \right| - K \left| \psi_{0}^{(3)} \right\rangle = t \left( S^{2} + 4S^{2} \bar{S}^{2} \frac{t}{\Omega_{0}} + \bar{S}^{4} \frac{t}{\Omega_{0}} \right),$$
 (30)

which is precisely the combination of weights given in Eq. (29). The first contribution comes from intraband transitions—this would correspond to the Drude weight for an extended system. This Drude response is, however, weighted by the overlap  $S = \langle 1|2 \rangle$  between the respective ground states of the pseudospin with one and two electrons. This is referred to as a "diagonal"transition in Fig. 1, since the background (here, the pseudospin) doesn't become excited in the transition. The second and the third peaks involve transitions corresponding to one and two pseudospin excitations,  $\Omega_0$ ; these are recognized as interband transitions in the language of multiple band models. This first-order perturbation approximation result remains valid as long as  $t/\Omega_0 \ll 1$ . Comparisons with exact results will be shown below.

### C. Two electron sector

The same procedure can be performed with the more difficult case of two electrons. In this case there are 16 basis states, and again they can be divided into eight even and eight odd states. We use a slightly different notation—there are now states involving double occupation of a single site, and those involving only single occupation. The states with double occupation are

$$\begin{aligned} |d_{1}\rangle_{c}^{\epsilon} &= \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |2\rangle_{a} \otimes |0\rangle_{b} \pm |0\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |2\rangle_{b}], \\ |d_{2}\rangle_{c}^{\epsilon} &= \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |2\rangle_{a} \otimes |\bar{0}\rangle_{b} \pm |\bar{0}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |2\rangle_{b}], \\ |d_{3}\rangle_{c}^{\epsilon} &= \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |\bar{2}\rangle_{a} \otimes |0\rangle_{b} \pm |0\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |\bar{2}\rangle_{b}], \\ |d_{4}\rangle_{c}^{\epsilon} &= \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger} c_{a\downarrow}^{\dagger} |\bar{2}\rangle_{a} \otimes |\bar{0}\rangle_{b} \pm |\bar{0}\rangle_{a} \otimes c_{b\uparrow}^{\dagger} c_{b\downarrow}^{\dagger} |\bar{2}\rangle_{b}], \end{aligned}$$
(31)

where the *d* in the ket stands for double, and the subscripts 1,2,3,4 simply enumerate the states, starting with the lowest,  $|d_1\rangle$ , where both pseudospins are in their ground states. The three other basis states correspond to the first site having an excited pseudospin state, the second site having an excited pseudospin state, and both sites having excited pseudospin states, respectively.

Similarly, for the basis states involving only singly occupied sites, the *s* in the ket stands for single; these are

$$|s_{1}\rangle_{o}^{\epsilon} = \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger}|1\rangle_{a} \otimes c_{b\downarrow}^{\dagger}|1\rangle_{b} \pm c_{a\downarrow}^{\dagger}|1\rangle_{a} \otimes c_{b\uparrow}^{\dagger}|1\rangle_{b}],$$

$$|s_{2}\rangle_{o}^{\epsilon} = \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger}|1\rangle_{a} \otimes c_{b\downarrow}^{\dagger}|\overline{1}\rangle_{b} \pm c_{a\downarrow}^{\dagger}|\overline{1}\rangle_{a} \otimes c_{b\uparrow}^{\dagger}|1\rangle_{b}],$$

$$|s_{3}\rangle_{o}^{\epsilon} = \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger}|\overline{1}\rangle_{a} \otimes c_{b\downarrow}^{\dagger}|1\rangle_{b} \pm c_{a\downarrow}^{\dagger}|1\rangle_{a} \otimes c_{b\uparrow}^{\dagger}|\overline{1}\rangle_{b}],$$

$$|s_{4}\rangle_{o}^{\epsilon} = \frac{1}{\sqrt{2}} [c_{a\uparrow}^{\dagger}|\overline{1}\rangle_{a} \otimes c_{b\downarrow}^{\dagger}|\overline{1}\rangle_{b} \pm c_{a\downarrow}^{\dagger}|\overline{1}\rangle_{a} \otimes c_{b\uparrow}^{\dagger}|\overline{1}\rangle_{b}].$$
(32)

The first has both pseudospin states in the ground state, with the three others having excited pseudospin states as in the case of the doubly occupied states. Note that an equally valid set of states combines  $|s_2\rangle$  and  $|s_3\rangle$  symmetrically and antisymmetrically [as we did in the middle two basis states of Eq. (23) for the three electron sector].

We want the unperturbed ground state to reside in the space of states involving only ground-state pseudospin states. Confining ourselves to this sector only, the ground-state wave function is

$$|\psi_0^{(2)}\rangle = a_0|d_1\rangle_e + b_0|s_1\rangle_o,$$
 (33)

with

$$a_0^2 = \frac{1}{2} \left( 1 - \frac{U/2}{\sqrt{(U/2)^2 + 4t^2 S^2}} \right), \tag{34}$$

$$b_0^2 = \frac{1}{2} \left( 1 + \frac{U/2}{\sqrt{(U/2)^2 + 4t^2 S^2}} \right).$$
(35)

This will be an accurate ground-state wave function as long as the pseudospin excitation energy remains the largest energy scale in the problem, i.e.,  $U \ll \Omega_0$ , along with the restriction already used,  $t \ll \Omega_0$ . When  $U \ll 2tS$  the ground state consists of nearly equal amplitudes of the two basis states; on the other hand, when  $U \gg 2tS$ ,  $a_0 \sim 0$  and  $b_0 \sim 1$ , and the singly occupied basis state dominates the ground state, as expected. With these same assumptions the two electron ground-state energy is given as

$$E_0^{(2)} \approx -\Omega_0 + U/2 - \sqrt{(U/2)^2 + (2tS)^2}.$$
 (36)



FIG. 2. Schematic depiction of the optical conductivity with two electrons in a dimer. Parts (a) and (b) refer to the two basis states that make up the ground-state wave function given by Eq. (33) with ground-state energy given by Eq. (36). The first has only (two) nondiagonal transitions (the second transition, to states given by  $|s_2\rangle_e$ , is qualitatively the same as the one shown), while the second basis state has both diagonal and nondiagonal transitions, as shown. The state representations are schematic only; formulas in the text represent the full (even or odd) state.

To determine the optical conductivity we need the result of the current operator on each component of the ground state; the result is

$$J|d_1\rangle_e = \frac{ietS}{\hbar}(|s_2\rangle_e - |s_3\rangle_e),$$
  

$$J|s_1\rangle_o = \frac{2iet}{\hbar}(S|d_1\rangle_o - \bar{S}|d_3\rangle_o).$$
(37)

Figure 2 summarizes these transitions schematically. The unperturbed energies associated with these states are readily determined by inspection from Eqs. (31) and (32). Using the analog of Eq. (26) appropriate to two electrons, with the ground state now a linear superposition of basis states given by Eq. (33), the optical conductivity for two electrons is

obtained as

$$\sigma_{1}^{(2)}(\omega) = \frac{\pi e^{2}t}{2\hbar^{2}} \bigg\{ \frac{4tS^{2}}{\sqrt{(U/2)^{2} + (2tS)^{2}}} \delta[\omega - \epsilon/\hbar] \\ + \frac{4a_{0}^{2}t\bar{S}^{2}}{\Omega_{0} - U + \epsilon} \delta[\omega - (\Omega_{0} - U + \epsilon)/\hbar] \\ + \frac{8b_{0}^{2}t\bar{S}^{2}}{\Omega_{0} + \epsilon} \delta[\omega - (\Omega_{0} + \epsilon)/\hbar] \bigg\},$$
(38)

where

$$\epsilon \equiv U/2 + \sqrt{(U/2)^2 + (2tS)^2}$$
 (39)

plays the role of a low energy scale, i.e., t (as  $U \rightarrow 0$ ) or U (as  $t \rightarrow 0$ ).

The two electron optical conductivity also has three peaks, as seen in Eq. (38) or schematically in Fig. 2. In the perturbative approach we have used, however, the two higher frequency peaks should be viewed as one occurring at a frequency scale of order  $\Omega_0$  that has been split by a low energy scale of order U or t. The analog of the third peak in the three electron case is absent here; there is no peak at  $2\Omega_0$ .

The first (low energy) peak corresponds to the diagonal transition from state  $|s_1\rangle_o$  to  $|d_1\rangle_o$ , where, as in the three electron case, the hopping of an electron between sites occurs without modification of the pseudospin background. This diagonal transition corresponds to the Drude-like (or coherent) part of the optical conductivity, though it may extend to a range of (low) frequencies on the scale of *U* as well as *t* (as represented by  $\epsilon$ ).

The second peak at frequency  $\omega = \Omega_0 - U + \epsilon$  is given by the two transitions:  $|d_1\rangle_e$  to both  $|s_2\rangle_e$  and  $|s_3\rangle_e$ . The third peak at frequency  $\Omega_0 + \epsilon$  is obtained by the  $|s_1\rangle_o$  to  $|d_3\rangle_o$  state transition, as detailed in Fig. 2. The second and third peaks correspond to nondiagonal transitions, which means that the pseudospin background is excited in the transition; this in turn corresponds to transitions involving higher energy bands, not included in our starting Hamiltonian. There is also an overall factor of 2 enhancement because there are now two carriers instead of the one carrier present in the three electron case presented above (or the one electron case shown below).

One can again verify the conductivity sum rule; for this we need the ground-state wave function given to first order in  $t/\Omega_0$ . Straightforward calculation<sup>41</sup> gives

$$\begin{aligned} \left|\psi_{0}^{(2)}\right\rangle \ \approx a_{0}|d_{1}\rangle_{e} + b_{0}|s_{1}\rangle_{0} + \frac{a_{0}t\bar{S}}{\Omega_{0} - U + \epsilon}(|s_{2}\rangle_{0} + |s_{3}\rangle_{0}) \\ - \frac{2t\bar{S}b_{0}}{\Omega_{0} + \epsilon}|d_{3}\rangle_{e}, \end{aligned} \tag{40}$$

and an evaluation of the kinetic energy expectation value gives

$$-\langle \psi_0^{(2)} | K | \psi_0^{(2)} \rangle = \frac{4t^2 S^2}{\sqrt{(U/2)^2 + (2tS)^2}} + \frac{4t^2 \bar{S}^2 a_0^2}{\Omega_0 - U + \epsilon} + \frac{8t^2 \bar{S}^2 b_0^2}{\Omega_0 + \epsilon},$$
(41)

again in agreement with the weights in Eq. (38).

#### D. One electron sector

Finally, similar calculations in the one electron sector are particularly simple, because the single site pseudospin ground states for zero and one electron are identical. This means that there are no pseudospin excitations arising from application of the current operator to the single electron ground state. The optical conductivity for one electron is given by the simple expression

$$\sigma_1^{(1)}(\omega) = \frac{\pi e^2 t}{2\hbar^2} \delta(\omega - 2t/\hbar), \qquad (42)$$

which contains only a Drude contribution, with no normalization (as required by the sum rule).

# **IV. DISCUSSION**

We wish to show the relative contributions to the conductivity for the three different electron densities that we can access. Because this is a tight-binding model it will not conserve total oscillator strength. This makes comparisons for different numbers of electrons and/or different parameter values difficult.<sup>7</sup> Here, for a given number of electrons, we will normalize the conductivity to the overall spectral weight in the conductivity for that number of electrons. Figure 3 shows the two-site optical conductivity with one, two, and three electrons, using U = 0 and g = 3, with  $\omega_0 = 1$ , as a "standard"set of parameters. While not necessarily realistic, they are chosen specifically to remove Mott complications at half filling; here the presence of U will result in a significant decrease in low-frequency spectral weight, in spite of the increase in the number of available carriers. These parameters will serve to illustrate the spectral weight transfer physics inherent in this model. At the same time, it is clear that the



FIG. 3. (Color online) Normalized optical conductivity as a function of frequency in a dimer obtained by perturbation theory with U = 0, g = 3, and  $\omega_0 = 1$ . The  $\delta$  functions in the formulas in the text are represented here as broadened Lorentzians (with a width 0.05). Note the decrease in relative low-frequency spectral weight (labeled as  $p_1$ ) as one goes from one to two to three electrons, indicating a reduced mobility as the number of electrons increases. Also note as the number of electrons decreases the higher frequency relative weight decreases: for the two electron case spectral weight is entirely absent at  $2\Omega_0$  (labeled as  $p_3$ ), while for the one electron case weight is absent even for  $\omega = \Omega_0$  (labeled as  $p_2$ ).

perturbation calculation is valid if  $\omega_0$  is large enough so that  $\Omega_0$  is much larger than U,t as assumed above.

The simplest case is clearly that of one electron. Equation (42) and Fig. 3 show that a single peak is present; it is located at  $\omega = 2t/\hbar$ , a nonzero value only because we are dealing with a dimer, and not the thermodynamic limit. In the thermodynamic limit this would be a  $\delta$  function at zero frequency, representing the Drude contribution. Normally this peak would be broadened through, for example, impurity scattering, but here (and even in the DMFT study of Ref. 16) it remains a  $\delta$  function, broadened in Fig. 3 artificially by hand, so as to be visible. For small but nonzero electron densities U would contribute as well, but for the most part this picture would remain unchanged. In particular, excited pseudospin states are essentially absent (see Fig. 3 in Ref. 16, which shows that the the expectation value of the pseudospin operator is essentially its ground-state value for low densities). Because the electron density is relatively dilute, there are very few optical transitions involving doubly occupied sites, and therefore it is not possible to excite the pseudospin excited state. For the dimer the only representative sector for this physics is the one electron sector (two electrons already constitutes a somewhat crowded lattice).

In contrast, the three electron sector represents the most "crowded" situation for a dimer, while the two electron sector is somewhat in between, and, as mentioned previously, the absence of a Coulomb repulsion (U = 0) aids to highlight the pseudospin physics, and suppress the Mott-related physics (which, from our point of view, is not essential, and will complicate the analysis). Referring to Fig. 3, note that the three electron conductivity has a significant relative contribution at high frequency  $(2\Omega_0)$ ; this is entirely absent in the two electron conductivity-it has been pushed down to lower frequency  $(\Omega_0)$ . The reason for this is as follows: with three electrons, the ground state consists of a doubly occupied and a singly occupied site, with the respective pseudospins at each site in its ground state—see the first of Eqs. (23). An optical transition can result in one of the three states shown in Fig. 1; one of these, state  $|4\rangle_e$ , has two excited pseudospin states, corresponding to an energy  $2\Omega_0$ . One of these excitations comes from the site with a single electron-before the transition this site was doubly occupied, and the ground state for this configuration required a pseudospin ground state corresponding to two electrons. Since one has left, there is now a component of the pseudospin which corresponds to an excited state for the one electron configuration. Similar remarks apply for the site that was previously singly occupied and is now doubly occupied. For two electrons this cannot happen—see Fig. 2 and note the absence of an alternative involving the two excited pseudospin state. This is because the pseudospin ground state is the same for an empty and singly occupied site—see Eq. (9) or the first of Eqs. (12), where T = 1.

This accounts for the peak structure for the various electron sectors in Fig. 3. The dimer calculations have an "all or nothing" character to them—no high frequency  $(2\Omega_0)$  peak for the two electron sector, and not even an intermediate  $(\Omega_0)$  peak for the one electron sector. Of course, in the DMFT calculations the changes from one electron density to another are smoothed out, as one can see in Fig. 13 of Ref. 16. The other feature that is apparent in Fig. 3 is the decrease of spectral

weight in the relative Drude (low-frequency) portion as one goes from the one electron to the two and then three electron sector. This is due to the polaronlike hopping renormalization already discussed. The relative weight of the Drude portion is indicative of the coherence of the carriers, so again, in the dilute limit, electrons can hop while the background pseudospin degree of freedom remains in the same ground state at both the site from which the electron hops, and at the site to which the electron hops, because only empty or singly occupied sites are involved. In the more crowded lattice limit (here represented by the three electron sector), doubly occupied sites are necessarily involved, and then the background pseudospin has to adjust according to whether a singly or doubly occupied site is involved.

The progression of spectral weight with electron number is perhaps best exemplified by examining the conductivity formulas, Eqs. (29), (38), and (42) for three, two, and one electron(s), respectively, for U = 0. Then the low-frequency spectral weights are [omitting the common factor  $\pi e^2 t/(2\hbar^2)$ ]  $S^2$  for three electrons (but one hole carrier), 2S/2 = S for two electrons, and unity for one electron; these weights steadily increase by reducing the number of electrons, since S < 1always, and this illustrates the principle that holes are less mobile than electrons.

We analyze in more detail our results for the frequency dependence of the optical conductivity. The three electron optical conductivity has three distinct peaks from low to high frequency, one at  $\omega \approx t$ , one at  $\omega \approx \Omega_0$ , and one at  $\omega \approx 2\Omega_0$ , whose weights we denote p<sub>1</sub>, p<sub>2</sub>, and p<sub>3</sub>, respectively. In the two electron conductivity, there are again three peaks, but as explained above, the two high frequency ones are at the same characteristic frequency (identical if U = 0), so we will combine the weight from these two and denote it as  $p_2$ ; we will continue to use  $p_1$  for the lowest-frequency peak, and of course for the one electron conductivity, there is only a low-frequency Drude-like peak, which we will also denote as  $p_1$ . In Fig. 4 we plot these weights to show how the spectral distribution of the optical conductivity varies with the strength of coupling gfor 3 (a) and for 2 (b) electrons; in (c) and (d) we show the corresponding results as a function of  $\omega_0$ , and in (e) and (f) results are shown for a variation of U. In all cases, the optical conductivity has been normalized to the total spectral weight for the parameters used in Fig. 3, separately for each electron number.

As expected, increasing the coupling strength *g* between the electron and the background (pseudospins) reduces the mobility of the electron as obtained in the spectral weight of the first peak p<sub>1</sub> (which would correspond to the Drude weight for an infinite lattice). This is simply due to the polaron effect mentioned above; with increased coupling, the amount of "background adjustment"required as the electron hops increases. Physically, the actual coupling in a given lattice is given by the amount of multiorbital involvement required to minimize the energy locally when two electrons try to accommodate one another on the same site. Since we model this process with the pseudospin degree of freedom, we span a considerable parameter range in the figures. The absolute weight p<sub>2</sub> of the second peak for the three electron case is given analytically as  $4t \frac{g^2}{(1+g^2)^2} \frac{t}{\Omega_0}$  [see Eq. (29)], and achieves its maximum value at  $g = \sqrt{2/3} \approx 0.8$ , which is independent



FIG. 4. (Color online) Dependence of the various normalized spectral weight contributions to the optical conductivity on electron-pseudospin coupling g for three electrons (a) and two electrons (b). In (c) and (d) we show the same quantities as a function of  $\omega_0$ , while in (e) and (f) they are shown as a function of U. Perturbation results are shown as curves, while exact results (for the dimer) are shown by symbols, as indicated. Note that usually the Drude weight dominates; however, for sufficiently large g (and U in the case of two electrons) the Drude weight is significantly reduced, indicative of reduced mobility, especially for the highly (electron) doped regime.

of U. For the two electron case, the off-diagonal transition contributions, represented by p2, are quite negligible compared with the low-frequency weight p<sub>1</sub> at weak coupling, but they play a more important role at strong coupling. Note that results arising from a complete diagonalization of all the dimer states are also shown, and, for these parameter regimes, the agreement is excellent, as expected. The transitions denoted by p<sub>2</sub> and p<sub>3</sub> represent incoherent processes; they may well correspond to the midinfrared band that seems to feature so prominently in a wide variety of cuprate superconductors.<sup>42</sup> The other experimental feature to which we can make contact with these dimer calculations is the dependency on doping. Experimentally, the anomalies at the superconducting transition are most pronounced in the low hole regime,<sup>5</sup> consistent with the fact that the pseudospin physics in these calculations plays a large role precisely in this regime as well.

Comparison with the results obtained from DMFT calculations<sup>16</sup> is also possible. For example, in Fig. 13 of Ref. 16, we show the conductivity as a function of frequency for various electron densities. Note that the parameters used in the DMFT calculation are in the more weak to intermediate coupling regime. Nonetheless, the calculations here are semiquantitatively consistent with those. The first panel there refers to the very dilute limit (n = 0.1), and, as suggested here, there is a single low-frequency Drude peak. Of course, there it is centered around zero frequency, while here it is at 2t, for reasons already explained. In the last panel in the same Fig. 13, n = 1.9, corresponding more to our present three electron calculation. We expect weight at  $\Omega_0 = 2\omega_0\sqrt{1+g^2} \approx$ 5.7, which is very close to the one shown there. Furthermore, the Drude-like peak has reduced spectral weight (clearer in Fig. 15 of Ref. 16) compared to the result at n = 0.1. The expected peak at  $2\Omega_0$  is, however, barely present, and at a higher frequency than expected. It is not clear what the cause of this latter discrepancy is, especially in light of the quantitative accuracy of the other peaks.

For completeness we have included plots to show the variation with  $\omega_0$  and U, where the expected behavior occurs. Note that at half filling (two electrons) the exact results differ considerably from the perturbation theory results, as Mott physics becomes more prevalent (this is not surprising since this was not considered in the perturbative approach we took). As  $\omega_0$  increases, the results for three and two electrons become dominated by the Drude-like peak near the origin. Again, this is entirely expected, since pseudo-spin excitations become more and more energetically costly, and so, as seen explicitly in our perturbative expressions, energy denominators increasingly suppress these transitions requiring excited pseudospin states, so that these play much less of a role as  $\omega_0$  increases. As is clear from panels (c) and (d), exact diagonalization results support these perturbative calculations.

### **V. CONCLUSIONS**

We have investigated spectral properties of the dynamic Hubbard model on a dimer, primarily to gain a qualitative understanding of the physics of electron-hole asymmetry, and polaronlike mobility inherent in real atoms. Primarily we have investigated the spectral features of the optical conductivity with different numbers of electrons. The physics we are trying to capture is that when electron movement results in a change from a doubly occupied site to a singly occupied site, or vice versa, a considerable amount of "background" adjustment needs to take place. In real atoms this is apparent in that the orbitals occupied by a single electron are considerably modified when two electrons occupy that same orbital. In the dynamic Hubbard model, these modifications are simulated by a pseudospin degree of freedom, at each site; an excited pseudospin state corresponds to an electron (partially) occupying an orbital that does not minimize the electron-ion energy, but does minimize the (local) electron-electron repulsion.

Such processes will impact the optical sum rule; in particular, weight will be transferred over a considerable range of energies, as a function of temperature and as a result of a phase transition. A considerable variation is expected as a function of electron concentration, and it is this aspect on which we have focused in the dimer calculations presented here. If the electron concentration is low, the pseudospin degree of freedom will be rarely excited, and the electrons will be highly coherent. However, if the electron concentration is high, then electron movement will be accompanied by pseudospin excitations. There is considerable experimental evidence for such incoherent processes in the cuprates, namely the midinfrared band. Our calculations clearly indicate that the Drude-like portion for holes has reduced mobility compared to that of electrons. The connection of the optical sum rule to the kinetic energy and how this probe can demonstrate this physics has been worked out in great detail for the dimer system considered here. More detailed comparison to experiment will have to rely on DMFT calculations<sup>16</sup> that provide answers in the thermodynamic limit.

The results of the dimer calculations presented here agree with the physics originally obtained in a model in which the pseudospin degree of freedom impacted the on-site energy of an electron.<sup>7</sup> Here, the pseudospin degree of freedom alters the effective Coulomb interaction between two electrons through a dynamical change in the on-site electron-electron interaction epitomized by U.<sup>10</sup> The qualitative picture obtained here also provides a better understanding of the conclusions obtained for an infinite lattice in Ref. 16: holes are less mobile than electrons, and the optical spectral weight distribution is significantly different for holes than for electrons.

# ACKNOWLEDGMENTS

We thank Jorge Hirsch for helpful discussions. This work was supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC), by ICORE (Alberta), and by the Canadian Institute for Advanced Research (CIfAR).

- <sup>1</sup>See, for example, A. N. Pasupathy, A. Pushp, K. K. Gomes, C. V. Parker, J. Wen, Z. Xu, G. Gu, S. Ono, Y. Ando, and A. Yazdani, Science **320**, 196 (2008).
- <sup>2</sup>D. N. Basov, S. I. Woods, A. S. Katz, E. J. Singley, R. C. Dynes, M. Xu, D. G. Hinks, C. C. Homes, and M. Strongin, Science 283, 49 (1999).
- <sup>3</sup>A. F. Santander-Syro, R. P. S. M. Lobo, N. Bontemps, Z. Konstantinovic, Z. Z. Li, and H. Raffy, Europhys. Lett. **62**, 568 (2003).
- <sup>4</sup>H. J. A. Molegraaf, C. Presura, D. van der Marel, P. H. Kes, and M. Li, Science **295**, 2239 (2002).
- <sup>5</sup>F. Carbone, A. B. Kuzmenko, H. J. A. Molegraaf, E. van Heumen, V. Lukovac, F. Marsiglio, D. van der Marel, K. Haule, G. Kotliar, H. Berger, S. Courjault, P. H. Kes, and M. Li, Phys. Rev. B 74, 064510 (2006).
- <sup>6</sup>E. van Heumen, R. Lortz, A. B. Kuzmenko, F. Carbone, D. van der Marel, X. Zhao, G. Yu, Y. Cho, N. Barisic, M. Greven, C. C. Homes, and S. V. Dordevic, *Phys. Rev. B* **75**, 054522 (2007).

<sup>7</sup>J. E. Hirsch, Physica C **201**, 347 (1992).

- <sup>8</sup>Note that the weight transfer observed in the cuprates in Refs. 2–6 has also been observed in the pnictides. See A. Charnukha, P. Popovich, Y. Matiks, D. L. Sun, C. T. Lin, A. N. Yaresko, B. Keimer, and A. V. Boris, Nature Commun. **2**, 1 (2011).
- **D.** Keinier, and *K*. V. Boris, Mature Commun. 2, 1 (2011).
- <sup>9</sup>J. E. Hirsch and F. Marsiglio, Phys. Rev. B **39**, 11515 (1989).
- <sup>10</sup>J. E. Hirsch, Phys. Rev. Lett. **87**, 206402 (2001).
- <sup>11</sup>J. E. Hirsch, Phys. Lett. A **134**, 451 (1989).
- <sup>12</sup>J. E. Hirsch, Physica C 158, 326 (1989).
- <sup>13</sup>F. Marsiglio and J. E. Hirsch, Physica C 171, 554 (1990); Phys. Rev. B 41, 6435 (1990).
- <sup>14</sup>For a recent example, see M. Chakraborty, A. N. Das, and A. Chakrabarti, J. Phys.: Condens. Matter **23**, 025601 (2011).
- <sup>15</sup>J. P. Hague, P. E. Kornilovitch, A. S. Alexandrov, and J. H. Samson, Phys. Rev. B **73**, 054303 (2006).
- <sup>16</sup>G. H. Bach, J. E. Hirsch, and F. Marsiglio, Phys. Rev. B 82, 155122 (2010).
- <sup>17</sup>A. B. Harris and R. V. Lange, Phys. Rev. **157**, 295 (1967).

# G. H. BACH AND F. MARSIGLIO

- <sup>18</sup>J. Ranninger and U. Thibblin, Phys. Rev. B **45**, 7730 (1992).
- <sup>19</sup>A. Avella, F. Mancini, and T. Saikawa, Eur. Phys. J. B **36**, 445 (2003).
- <sup>20</sup>G. D. Mahan, *Many-Particle Physics* (Kluwer Academic/Plenum, New York, 2000).
- <sup>21</sup>R. Kubo, J. Phys. Soc. Jpn. **12**, 570 (1957).
- <sup>22</sup>See, for example, D. Y. Smith and E. Shiles, Phys. Rev. B 17, 4689 (1978) and E. Shiles, T. Sasaki, M. Inokuti, and D. Y. Smith, *ibid*. 22, 1612 (1980), for an informative description of the fulfillment of this sum rule through a hierarchy of energy scales in the case of aluminum.
- <sup>23</sup>P. F. Maldague, Phys. Rev. B 16, 2437 (1977).
- <sup>24</sup>A. J. Millis, in *Strong Interactions in Low Dimensions*, edited by D. Baeriswyl and L. Degiorgi (Kluwer Academic Publishers, Amsterdam, 2004), p. 195.
- <sup>25</sup>F. Marsiglio, E. van Heumen, and A. B. Kuzmenko, Phys. Rev. B 77, 144510 (2008).
- <sup>26</sup>Some exceptions are found in M. R. Norman, A. V. Chubukov, E. van Heumen, A. B. Kuzmenko, and D. van der Marel, Phys. Rev. B 76, 220509(R) (2007), and Refs. 25 and 27, where the conductivity [LHS of Eq. (5)] is integrated to explore the sum rule in more detail.
- <sup>27</sup>A. Toschi and M. Capone, Phys. Rev. B 77, 014518 (2008).
- <sup>28</sup>F. Marsiglio, F. Carbone, A. B. Kuzmenko, and D. van der Marel, Phys. Rev. B **74**, 174516 (2006).

- <sup>29</sup>A. S. Alexandrov and N. F. Mott, *High Temperature Superconduc*tors and Other Superfluids (Taylor and Francis, London, 1994).
- <sup>30</sup>T. Eckl, W. Hanke, and E. Arrigoni, Phys. Rev. B 68, 014505 (2003).
- <sup>31</sup>A. Toschi, M. Capone, M. Ortolani, P. Calvani, S. Lupi, and C. Castellani, Phys. Rev. Lett. **95**, 097002 (2005); A. Toschi, M. Capone, and C. Castellani, Phys. Rev. B **72**, 235118 (2005).
- <sup>32</sup>B. Kyung, A. Georges, and A.-M. S. Tremblay, Phys. Rev. B 74, 024501 (2006).
- <sup>33</sup>K. Haule and G. Kotliar, Europhys. Lett. 77, 27007 (2007).
- <sup>34</sup>A. E. Karakozov, E. G. Maksimov, and O. V. Dolgov, Solid State Commun. **124**, 119 (2002); A. E. Karakozov and E. G. Maksimov, *ibid.* **139**, 80 (2006).
- <sup>35</sup>F. Marsiglio, J. Supercond. Nov. Magn. 22, 269 (2009).
- <sup>36</sup>J. E. Hirsch, Science **295**, 2226 (2002).
- <sup>37</sup>J. E. Hirsch and F. Marsiglio, Phys. Rev. B 62, 15131 (2000).
- <sup>38</sup>J. E. Hirsch, Phys. Rev. B **65**, 214510 (2002).
- <sup>39</sup>J. E. Hirsch, Phys. Rev. B **66**, 064507 (2002).
- <sup>40</sup>J. E. Hirsch, Phys. Rev. B **62**, 14487 (2000).
- <sup>41</sup>Note that we have already diagonalized the Hamiltonian inside the two state subspace in which the pseudospin is always in the ground state. The excited state within this subspace is therefore omitted in the perturbation expansion.
- <sup>42</sup>D. B. Tanner and T. Timusk, in *Physical Properties of High Temperature Superconductors III*, edited by D. M. Ginsberg (World Scientific, Singapore, 1992), p. 363.