

$U(1)$ slave-spin theory and its application to Mott transition in a multiorbital model for iron pnictides

Rong Yu and Qimiao Si

Department of Physics & Astronomy, Rice University, Houston, TX 77005

(Received 25 April 2012; revised manuscript received 9 July 2012; published 2 August 2012)

A $U(1)$ slave-spin representation is introduced for multiorbital Hubbard models. As with the Z_2 form of de'Medici *et al.* [*Phys. Rev. B* **72**, 205124 (2005)], this approach represents a physical electron operator as the product of a slave spin and an auxiliary fermion operator. For nondegenerate multiorbital models, our $U(1)$ approach is advantageous in that it captures the noninteracting limit at the mean-field level. For systems with either a single orbital or degenerate multiple orbitals, the $U(1)$ and Z_2 slave-spin approaches yield the same results in the slave-spin-condensed phase. In general, the $U(1)$ slave-spin approach contains a $U(1)$ gauge redundancy, and properly describes a Mott insulating phase. We apply the $U(1)$ slave-spin approach to study the metal-to-insulator transition in a five-orbital model for parent iron pnictides. We demonstrate a Mott transition as a function of the interactions in this model. The nature of the Mott insulating state is influenced by the interplay between the Hund's rule coupling and crystal-field splittings. In the metallic phase, when the Hund's rule coupling is beyond a threshold, there is a crossover from a weakly correlated metal to a strongly correlated one, through which the quasiparticle spectral weight rapidly drops. The existence of such a strongly correlated metallic phase supports the incipient Mott picture of the parent iron pnictides. In the parameter regime for this phase and in the vicinity of the Mott transition, we find that an orbital selective Mott state has nearly as competitive a ground-state energy.

DOI: [10.1103/PhysRevB.86.085104](https://doi.org/10.1103/PhysRevB.86.085104)

PACS number(s): 71.10.Hf, 71.27.+a, 71.30.+h, 74.70.Xa

I. INTRODUCTION

Many important questions remain on the physics of the iron pnictides and related iron-based high- T_c superconductors.¹⁻⁶ One central issue is the strength of electron correlations these systems contain. The metallic nature and the collinear antiferromagnetic (AFM) ground state⁷ in the parent pnictides can arise within a weak-coupling approach, in which the Fermi surface nesting plays an important role.⁸ On the other hand, the large room-temperature electrical resistivity, showing the bad metal behavior with the electron mean-free path on the order of the interelectron spacing, as well as the suppression of Drude weight in optical conductivity,⁹⁻¹¹ the well-defined zone-boundary spin waves,¹² and the renormalization of local density approximation (LDA) band structure in the angle-resolved photoemission spectroscopy (ARPES) measurements¹³ provide evidence for sufficiently strong electron correlations such that the system is in close proximity to a Mott transition with dominant contributions to the spin spectral weight from quasilocal moments.¹⁴⁻²⁸

This incipient Mott picture is further supported by the properties of iron chalcogenides.⁴⁻⁶ In either the 11- or the 122-chalcogenides, the magnetic ordering wave vector and the large magnetic moment²⁹⁻³² (ranging from 2 to 3.4 μ_B) can hardly be explained by a Fermi surface nesting mechanism, but is readily understood within a quasilocal moment model.³³⁻³⁶ It has also been shown that the band-narrowing effect, either due to the expansion of the iron lattice unit cell in an iron oxychalcogenide³⁷ or from the ordered iron vacancies in 122 iron selenides,^{35,38,39} may drive the system through the Mott transition to a Mott insulator.

In general, the degree of electron correlations can be measured by the ratio U/D , where U refers to a characteristic interaction strength, such as the Coulomb repulsion in a Hubbard model, and D is the full bandwidth, a scale of the

kinetic energy of the system. A Mott transition separating the metallic and the Mott insulating phases takes place at $U_c \sim D$. In the metallic phase, electron correlations can alternatively be measured by the quasiparticle spectral weight Z , which is unity at the noninteracting limit $U = 0$, and vanishes at the Mott transition $U = U_c$. The incipient Mott picture relies on the existence of a (putative) Mott transition, and assume that the system is not too far from this transition so that Z is relatively small.

The above considerations suggest that it is very important to theoretically investigate how a metal-to-Mott-insulator transition takes place in a model which is applicable to the parent iron pnictides and/or chalcogenides. In a previous paper, by studying a two-orbital and a four-orbital model, the authors have shown that a Mott transition generally exists in these models.⁴⁰ But, both models are at half-filling. By contrast, in the parent iron pnictides, six electrons occupy five $3d$ orbitals of each Fe. This means that the system is away from half-filling. It would be important to investigate the extent to which the Mott insulating states persist in such a situation.

Historically, the Mott transition in a single-orbital Hubbard model has been studied by using various techniques, such as dynamical mean-field theory⁴¹ (DMFT) and Gutzwiller approximation.⁴² Among these methods, the Kotliar-Ruckenstein slave-boson method⁴³ has been broadly used. In this theory, a Mott transition at finite U/D is obtained already at the mean-field level. However, it is difficult to apply this approach to multiorbital systems as it would introduce 4^M slave-boson fields for a model with M orbitals; the number of variational variables is already huge even for $M = 5$. Recently, several other slave-particle theories following the idea of charge-spin separation have been proposed.⁴⁴⁻⁴⁶ Among these theories, two of them, the slave-rotor theory⁴⁵ and the slave-spin theory,⁴⁴ have been applied to multiorbital

systems. In both theories, a slave-bosonic variable [a quantum $O(2)$ rotor in the slave-rotor theory, and a quantum $S = 1/2$ spin in the slave-spin theory] is introduced to carry the electric charge, and an auxiliary fermion (the spinon) carries the spin of the electron. The metallic phase corresponds to the state that the slave particles are Bose condensed, so that charge excitations are gapless along with the spin excitations. By contrast, the Mott insulator corresponds to the state that the slave particles are disordered and gapped; charge excitations are gapped, while the spin excitations remain gapless. The slave-rotor method is very economical because it introduces only one rotor per site taking account of the total charge. It is also very efficient if the interactions have an $SU(2M)$ symmetry for an M -orbital system. However, it can not be easily applied to systems with a nonzero Hund's rule coupling which breaks the $SU(2M)$ symmetry. It is also not convenient to handle systems that exhibit strong orbital dependence, such as the orbital selective Mott transition (OSMT). The slave-spin theory overcomes these drawbacks by introducing a slave spin for each orbital and spin flavor. Compared with the Kotliar-Ruckenstein slave-boson theory, it is still very economical because it introduces only $2M$ slave spins per site. It has been successfully used to study the Mott transition and OSMT in multiorbital systems with a nonzero Hund's rule coupling.⁴⁷ There are, however, several issues with the slave-spin representation. First, it has difficulties when applied to a multiorbital system in which one or more orbitals are away from half-filling due to crystal-field splitting. For instance, it is hard to properly describe the noninteracting limit in such nondegenerate multiorbital models. Second, in its original construction, the slave-spin representation has a Z_2 gauge redundancy. This makes the spinons carry both spin and charge currents, and causes difficulties in describing a Mott insulating phase.⁴⁸

In this paper, we propose an XY slave-spin theory that is free of these issues. This slave-spin theory has a $U(1)$ gauge redundancy, and properly describes Mott insulating phases. We develop a mean-field theory, which can be applied to multiorbital systems with a nonzero crystal-field splitting. For a model with a single orbital or degenerate multiorbitals, on the other hand, our $U(1)$ slave-spin mean-field theory and the Z_2 slave-spin theory⁴⁹ give the same results in their spin-condensed phases. We then apply our formulation to study the Mott transition in a five-orbital model for the parent iron pnictides. We establish the existence of a Mott transition in this model. Both the nature of the metallic and Mott insulating phases are strongly affected by the interplay of Hund's rule coupling and the crystal-field splitting. A crossover to a strongly correlated metallic state exists when the Hund's rule coupling is beyond a threshold. The existence of this state is in agreement with the incipient Mott picture.

The rest of the paper is organized as follows. In Sec. II, we first introduce our construction of the $U(1)$ slave-spin theory, and develop a mean-field theory based on this new construction. We also compare our construction with the slave-rotor and Z_2 slave-spin theories. In Sec. III, we apply the $U(1)$ slave-spin mean-field theory to study the Mott transition in a five-orbital model for the parent iron pnictides, and show how the transition is affected by the interplay between Hund's

rule coupling and the crystal-field splitting. Finally, Sec. IV contains some concluding remarks.

II. METHOD

In this paper, we are interested in the metal-to-insulator transition in a multiorbital system, for which the Hamiltonian reads as

$$H = H_0 + H_{\text{int}}. \quad (1)$$

H_0 contains the tight-binding parameters among the multiple orbitals,

$$H_0 = \frac{1}{2} \sum_{ij\alpha\beta\sigma} t_{ij}^{\alpha\beta} d_{i\alpha\sigma}^\dagger d_{j\beta\sigma} + \sum_{i\alpha\sigma} (\Delta_\alpha - \mu) d_{i\alpha\sigma}^\dagger d_{i\alpha\sigma}, \quad (2)$$

where $d_{i\alpha\sigma}^\dagger$ creates an electron in orbital α with spin σ at site i , Δ_α is the onsite energy reflecting the crystal-field splitting, and μ is the chemical potential. H_{int} contains onsite Hubbard interactions

$$\begin{aligned} H_{\text{int}} = & \frac{U}{2} \sum_{i,\alpha,\sigma} n_{i\alpha\sigma} n_{i\alpha\bar{\sigma}} \\ & + \sum_{i,\alpha<\beta,\sigma} \{U' n_{i\alpha\sigma} n_{i\beta\bar{\sigma}} + (U' - J) n_{i\alpha\sigma} n_{i\beta\sigma} \\ & - J(d_{i\alpha\sigma}^\dagger d_{i\alpha\bar{\sigma}} d_{i\beta\bar{\sigma}}^\dagger d_{i\beta\sigma} - d_{i\alpha\bar{\sigma}}^\dagger d_{i\alpha\sigma} d_{i\beta\sigma}^\dagger d_{i\beta\bar{\sigma}})\}. \end{aligned} \quad (3)$$

where $n_{i\alpha\sigma} = d_{i\alpha\sigma}^\dagger d_{i\alpha\sigma}$. In this model, U , U' , and J , respectively, denote the intraorbital repulsion, the interorbital repulsion, and the Hund's rule exchange coupling. In the following, we will take $U' = U - 2J$.⁵⁰

A. $U(1)$ slave-spin theory

To study this multiorbital model, we implement the idea of charge and spin separation of the d electrons by using the slave-spin approach. For each orbital and spin flavor, we rewrite the electron creation operator to be

$$d_{i\alpha\sigma}^\dagger = S_{i\alpha\sigma}^+ f_{i\alpha\sigma}^\dagger, \quad (4)$$

where $S_{i\alpha\sigma}^+$ is a ladder operator of the slave quantum $S = 1/2$ spin carrying the charge degree of freedom of the electron, and $f_{i\alpha\sigma}^\dagger$ is a spinon creation operator. We further enforce a constraint for each site,

$$S_{i\alpha\sigma}^z = f_{i\alpha\sigma}^\dagger f_{i\alpha\sigma} - \frac{1}{2}, \quad (5)$$

which restricts the Hilbert space to the physical one.

Note that our slave-spin formulation in Eq. (4) is different from that introduced in Refs. 44 and 51, in which

$$d_{i\alpha\sigma}^\dagger = O_{i\alpha\sigma}^\dagger f_{i\alpha\sigma}^\dagger, \quad (6)$$

where

$$O_{i\alpha\sigma}^\dagger = S_{i\alpha\sigma}^+ + c_{i\alpha\sigma} S_{i\alpha\sigma}^-, \quad (7)$$

with $c_{i\alpha\sigma}$ being a complex number. In that formulation, the gauge redundancy is reduced from $U(1)$ to Z_2 in Eqs. (6) and (7) due to the mixing between $S_{i\alpha\sigma}^+$ and $S_{i\alpha\sigma}^-$ (hence they are referred to as a Z_2 slave-spin representation); as a consequence, the slave spins can not carry the $U(1)$ charge.

In our Eqs. (4) and (5), there is a $U(1)$ gauge redundancy corresponding to $f_{i\alpha\sigma}^\dagger \rightarrow f_{i\alpha\sigma}^\dagger e^{-i\theta_{i\alpha\sigma}}$ and $S_{i\alpha\sigma}^+ \rightarrow S_{i\alpha\sigma}^+ e^{i\theta_{i\alpha\sigma}}$. The slave spins carry the $U(1)$ charge, similarly as the slave rotors.⁴⁵ We will refer to this as a $U(1)$ slave-spin theory. In our construction, the phase that corresponds to disordered slave spins [preserving the $U(1)$ symmetry] but with gapless spinons corresponds to a Mott insulator.

Next, we develop a mean-field theory based on the construction of Eqs. (4) and (5). A naive mean-field theory based on Eq. (4) would not produce the correct quasiparticle spectral weight in the noninteracting limit.⁵¹ To make progress, we rewrite the slave-spin operators in their Schwinger boson representation: $S_{i\alpha\sigma}^+ = a_{i\alpha\sigma}^\dagger b_{i\alpha\sigma}$, $S_{i\alpha\sigma}^- = b_{i\alpha\sigma}^\dagger a_{i\alpha\sigma}$, and $S_{i\alpha\sigma}^z = (a_{i\alpha\sigma}^\dagger a_{i\alpha\sigma} - b_{i\alpha\sigma}^\dagger b_{i\alpha\sigma})/2$. The constraint in Eq. (5) then becomes $a_{i\alpha\sigma}^\dagger a_{i\alpha\sigma} - b_{i\alpha\sigma}^\dagger b_{i\alpha\sigma} = 2f_{i\alpha\sigma}^\dagger f_{i\alpha\sigma} - 1$. Here, we need to introduce an extra constraint $a_{i\alpha\sigma}^\dagger a_{i\alpha\sigma} + b_{i\alpha\sigma}^\dagger b_{i\alpha\sigma} = 1$ so that the Schwinger bosons represent $S = 1/2$ spins. We then see that $a_{i\alpha\sigma}$ and $b_{i\alpha\sigma}$ are hard-core bosons. In light of the Kotliar-Ruckenstein slave-boson mean-field theory,⁴³ we now define a dressed operator in the Schwinger boson representation which automatically takes into account this constraint:

$$z_{i\alpha\sigma}^\dagger = P_{i\alpha\sigma}^+ a_{i\alpha\sigma}^\dagger b_{i\alpha\sigma} P_{i\alpha\sigma}^-, \quad (8)$$

where $P_{i\alpha\sigma}^\pm = 1/\sqrt{1/2 + \delta \pm (a_{i\alpha\sigma}^\dagger a_{i\alpha\sigma} - b_{i\alpha\sigma}^\dagger b_{i\alpha\sigma})/2}$, and δ is an infinitesimal positive number to regulate $P_{i\alpha\sigma}^\pm$. $z_{i\alpha\sigma}^\dagger$ and $a_{i\alpha\sigma}^\dagger b_{i\alpha\sigma}$ are equivalent in the physical Hilbert space. In the Schwinger boson representation, Eq. (4) becomes

$$d_{i\alpha\sigma}^\dagger = z_{i\alpha\sigma}^\dagger f_{i\alpha\sigma}^\dagger. \quad (9)$$

At the mean-field level, we treat the constraint (5) on average by introducing a Lagrange multiplier, and decompose the boson and spinon operators. We obtain two mean-field Hamiltonians, respectively, for the spinons and the Schwinger bosons:

$$H_f^{\text{mf}} = \frac{1}{2} \sum_{ij\alpha\beta\sigma} t_{ij}^{\alpha\beta} \langle z_{i\alpha\sigma}^\dagger z_{j\beta\sigma} \rangle f_{i\alpha\sigma}^\dagger f_{j\beta\sigma} + \sum_{i\alpha\sigma} (\Delta_\alpha - \lambda_{i\alpha\sigma} - \mu) f_{i\alpha\sigma}^\dagger f_{i\alpha\sigma}, \quad (10)$$

$$H_S^{\text{mf}} = \frac{1}{2} \sum_{ij\alpha\beta\sigma} t_{ij}^{\alpha\beta} \langle f_{i\alpha\sigma}^\dagger f_{j\beta\sigma} \rangle z_{i\alpha\sigma}^\dagger z_{j\beta\sigma} + \sum_{i\alpha\sigma} \frac{\lambda_{i\alpha\sigma}}{2} (\hat{n}_{i\alpha\sigma}^a - \hat{n}_{i\alpha\sigma}^b) + H_{\text{int}}^S, \quad (11)$$

where $\langle \dots \rangle$ denotes the mean-field value, $\hat{n}_{i\alpha\sigma}^a = a_{i\alpha\sigma}^\dagger a_{i\alpha\sigma}$, and $\lambda_{i\alpha\sigma}$ is the Lagrange multiplier to handle the constraint in Eq. (5). The quasiparticle spectral weight is defined as $Z_{i\alpha\sigma} = |\langle z_{i\alpha\sigma} \rangle|^2$. In Eq. (11), H_{int}^S refers to the interaction Hamiltonian in the Schwinger boson representation. It can be obtained by rewriting Eq. (3) in the slave-spin representation $H_{\text{int}} \rightarrow H_{\text{int}}(S)$,⁴⁰ then substitute the Schwinger bosons for the spin operators. The mean-field Hamiltonian H_S^{mf} has an internal $U(1)$ symmetry of the bosons. For a single orbital, it is a Bose-Hubbard model for two species of bosons, and is equivalent to a model of interacting XY spins in a magnetic field. At commensurate fillings, by breaking the internal $U(1)$ symmetry, this model has a phase transition from a bosonic

Mott insulator to a superfluid with decreasing the interactions. These two phases correspond to the Mott insulating and metallic states in the original d -electron problem. We then approach the Mott transition from the ordered phase. This can be done by assuming the Bose condensation takes place in the composite boson field $z_{i\alpha\sigma}$, and further adopting a single-site approximation to Eqs. (10) and (11) with the decoupling $z_{i\alpha\sigma}^\dagger z_{j\beta\sigma} \approx \langle z_{i\alpha\sigma}^\dagger \rangle z_{j\beta\sigma} + z_{i\alpha\sigma}^\dagger \langle z_{j\beta\sigma} \rangle - \langle z_{i\alpha\sigma}^\dagger \rangle \langle z_{j\beta\sigma} \rangle$. For simplicity, we focus on the paramagnetic phase, and assume translational symmetry. These allow us to drop the spin and site indices in the formulas. The mean-field boson Hamiltonian then reads as

$$H_S^{\text{mf}} \approx \sum_{\alpha\beta} \epsilon^{\alpha\beta} (\langle z_\alpha^\dagger \rangle z_\beta + \langle z_\beta \rangle z_\alpha^\dagger) + \sum_{\alpha} \frac{\lambda_\alpha}{2} (\hat{n}_\alpha^a - \hat{n}_\alpha^b) + H_{\text{int}}^S, \quad (12)$$

where $\epsilon^{\alpha\beta} = \sum_{ij\sigma} t_{ij}^{\alpha\beta} \langle f_{i\alpha\sigma}^\dagger f_{j\beta\sigma} \rangle / 2$. In Eq. (12), we Taylor expand z_α and z_α^\dagger in terms of $\hat{A} - \langle \hat{A} \rangle$ (where $\hat{A} = \hat{n}_\alpha^a, \hat{n}_\alpha^b, a_\alpha^\dagger b_\alpha$), and keep up to the linear terms in $\hat{A} - \langle \hat{A} \rangle$, obtaining

$$z_\alpha^\dagger \approx \bar{z}_\alpha^\dagger + \langle \bar{z}_\alpha^\dagger \rangle \eta_\alpha [\hat{n}_\alpha^a - \hat{n}_\alpha^b - (2n_\alpha^f - 1)], \quad (13)$$

where $\bar{z}_\alpha^\dagger = \langle P_\alpha^+ \rangle a_\alpha^\dagger b_\alpha \langle P_\alpha^- \rangle$, $\eta_\alpha = (2n_\alpha^f - 1)/[4n_\alpha^f(1 - n_\alpha^f)]$, $n_\alpha^f = \frac{1}{N} \sum_k \langle f_{k\alpha}^\dagger f_{k\alpha} \rangle$, and $n_\alpha^f = \langle \hat{n}_\alpha^a \rangle = 1 - \langle \hat{n}_\alpha^b \rangle$ from the constraints. We find that Eq. (13) already gives good mean-field results, and will hence drop the higher-order terms in the expansion. Note that the approximate form of z_α^\dagger in Eq. (13) is only used to simplify H_S^{mf} , but can not be fed into Eq. (9) to calculate the electron Green's functions since the approximate operator behaves differently from the original one in the physical Hilbert space. Nevertheless, $\langle z_\alpha \rangle = \langle \bar{z}_\alpha \rangle$. With Eq. (13), Eq. (12) is then approximated to be

$$H_S^{\text{mf}} \approx \sum_{\alpha\beta} \epsilon^{\alpha\beta} (\langle \bar{z}_\alpha^\dagger \rangle \bar{z}_\beta + \langle \bar{z}_\beta \rangle \bar{z}_\alpha^\dagger) + \sum_{\alpha} \left(\frac{\lambda_\alpha}{2} + \bar{\epsilon}_\alpha \eta_\alpha \right) (\hat{n}_\alpha^a - \hat{n}_\alpha^b) + H_{\text{int}}^S, \quad (14)$$

where $\bar{\epsilon}_\alpha = \sum_{\beta} (\epsilon^{\alpha\beta} \langle \bar{z}_\alpha^\dagger \rangle \langle \bar{z}_\beta \rangle + \text{c.c.})$. Further using the constraint (5), we can move the term proportional to η_α to H_f^{mf} by introducing an effective onsite potential $\tilde{\mu}_\alpha = 2\bar{\epsilon}_\alpha \eta_\alpha$. The resulting mean-field Hamiltonians are then

$$H_f^{\text{mf}} = \sum_{k\alpha\beta} [\epsilon_k^{\alpha\beta} \langle \bar{z}_\alpha^\dagger \rangle \langle \bar{z}_\beta \rangle + \delta_{\alpha\beta} (\Delta_\alpha - \lambda_\alpha + \tilde{\mu}_\alpha - \mu)] f_{k\alpha}^\dagger f_{k\beta}, \quad (15)$$

$$H_S^{\text{mf}} = \sum_{\alpha\beta} \left[\epsilon^{\alpha\beta} (\langle \bar{z}_\alpha^\dagger \rangle \bar{z}_\beta + \langle \bar{z}_\beta \rangle \bar{z}_\alpha^\dagger) + \delta_{\alpha\beta} \frac{\lambda_\alpha}{2} (\hat{n}_\alpha^a - \hat{n}_\alpha^b) \right] + H_{\text{int}}^S, \quad (16)$$

where $\epsilon_k^{\alpha\beta} = \frac{1}{N} \sum_{ij} t_{ij}^{\alpha\beta} e^{ik(r_i - r_j)}$, and $\delta_{\alpha\beta}$ is Kronecker's delta function. Equations (15) and (16) are the main formulation of our slave-spin mean-field theory. Note that although the mean-field Hamiltonian in Eq. (16) contains quartic terms of boson operators from the H_{int}^S term, it is defined on a single

site. Given that the bosons are hard core (or, equivalently, recognizing that they can be transformed back to the slave-spin representation), we can diagonalize the Hamiltonian exactly. The mean-field parameters $\langle \tilde{z}_\alpha \rangle$ and λ_α can then be solved self-consistently. In the noninteracting limit, it is easy to check that $Z_\alpha = |\langle \tilde{z}_\alpha \rangle|^2 = 1$ can be achieved by taking $\lambda_\alpha = \tilde{\mu}_\alpha$; the quasiparticle weights are equal to 1 as it should be.

B. Comparison with the Z_2 slave-spin theory

Here, we compare our $U(1)$ slave-spin theory with the Z_2 slave-spin theory. One advantage of the $U(1)$ slave-spin theory over the Z_2 slave-spin approach is that it can be directly generalized to the multiorbital systems with nonzero crystal-field splitting and/or away from half-filling. As an example, we consider a two-orbital model at half-filling and with equal bandwidth but with a finite crystal-field splitting. In the noninteracting limit, we expect the quasiparticle spectral weight $Z_{\alpha=1,2} = 1$ and the spinon band structure is identical to the tight-binding dispersion of the d electrons. The Z_2 slave-spin mean-field theory failed to obtain these results. This is easy to see: Both of the two orbitals are away from half-filling in the presence of the crystal-field splitting. According to Eq. (17), to obtain $Z_{1(2)} = |\langle O_{1(2)} \rangle|^2 = 1$, it is necessary that $\lambda_1 = -\lambda_2 \neq 0$. In absence of the potential $\tilde{\mu}$ in Eq. (15), this already distorts the band structure of the spinons from the original tight-binding form. Hence, one can not obtain the desired spinon filling as required by the constraint. However, in our theory, $Z_\alpha = 1$ is guaranteed by the condition $\lambda_\alpha = \tilde{\mu}_\alpha$. These two potentials cancel out exactly as seen in Eq. (15). Therefore, the spinon band structure is identical to the tight-binding one, and the noninteracting limit is properly recovered.

However, we find that at the mean-field level and in the symmetry-broken phases of the bosons/spins, the two theories have very similar forms. To see this explicitly, we compare the mean-field Hamiltonians of the two theories in the slave-spin representation. In this representation, the mean-field Hamiltonian of the $U(1)$ slave-spin theory can be obtained by performing a Schwinger-boson-to-spin mapping to the Hamiltonian H_S^{mf} in Eq. (16). H_S^{mf} then reads as

$$H_S^{\text{mf}} = \sum_{\alpha\beta} [\epsilon^{\alpha\beta} (\langle O_\alpha^\dagger \rangle O_\beta + \langle O_\beta \rangle O_\alpha^\dagger) + \delta_{\alpha\beta} \lambda_\alpha S_\alpha^z] + H_{\text{int}}(\mathbf{S}), \quad (17)$$

where

$$O_\alpha^\dagger = \langle P_\alpha^+ \rangle S_\alpha^+ \langle P_\alpha^- \rangle, \quad (18)$$

$P_\alpha^\pm = 1/\sqrt{1/2 + \delta \pm S_\alpha^z}$, and $Z_\alpha = |\langle O_\alpha \rangle|^2$. Surprisingly, we see that the mean-field Hamiltonian of the Z_2 theory takes exactly the same form as in Eq. (17) if we define

$$O_\alpha^\dagger = (\langle P_\alpha^- \rangle \langle P_\alpha^+ \rangle - 1) S_\alpha^- + S_\alpha^+. \quad (19)$$

Interestingly, the two definitions in Eqs. (18) and (19) give the same quasiparticle spectral weight $Z_\alpha = |\langle O_\alpha \rangle|^2$. In models with a single orbital or degenerate multiple orbitals, $\tilde{\mu}_\alpha$ in Eq. (15) becomes orbital independent and can thus be absorbed into the chemical potential. Therefore, in these cases, in the

metallic phase, the $U(1)$ slave-spin mean-field theory and the Z_2 theory give the same results (up to a constant in the free energies). It should be stressed that, even for single-orbital or degenerate multiorbital models, the agreement between the two theories is obtained only in the ordered phase of the slave spins. Generally, the constructions in the two formulations are different in the sense already mentioned. In the $U(1)$ slave-spin theory, the operator equation (4) has a $U(1)$ gauge redundancy, and allows a proper Mott insulating phase.

III. MOTT TRANSITION IN A FIVE-ORBITAL MODEL FOR IRON PNICTIDES

In a previous paper,^{40,52} we have discussed the metal-to-insulator transition in two- and four-orbital models. For both models, we find a metal-to-Mott-insulator transition at finite U . We also find that the nature of the phase transition and the critical value U_c can be strongly influenced by the Hund's rule coupling. These results are based on the assumption that the system under study is at half-filling. But, the parent iron pnictides are away from half-filling, with six d electrons occupying five orbitals in each iron atom. To consider this more involved case, here we use the $U(1)$ slave-spin mean-field theory to study the five-orbital Hubbard model for iron pnictides. We take the tight-binding parameters as those proposed in Ref. 53 for the parent LaOFeAs. The interaction part of the Hamiltonian is as given in Eq. (3). For simplicity, here we consider only the density-density interactions and drop the spin-flip and pair-hopping terms in Eq. (3). The results with the full interactions are qualitatively similar.⁴⁰

Figure 1 shows the ground-state phase diagram at electron filling $n = 6$ of the five-orbital model in the $J-U$ plane. We find a single transition from a metal to an insulator. At $J = 0$, $U_c \approx 13.1$ eV. In this case, we have also independently




FIG. 1. (Color online) Ground-state phase diagram of the five-orbital model at $n = 6.0$. The solid curve with symbols shows the boundary between the metallic and Mott insulating phases. The dotted line shows a crossover in the metallic phase where the system changes from a weakly correlated metal to a strongly correlated metal. The dashed line indicates a low- to high-spin transition in the Mott insulating phase. The atomic configurations corresponding to the low- and high-spin Mott states are illustrated on the right side.

determined $U_c^{SR} \approx 11.0$ eV from the slave-rotor mean-field theory.⁵⁴ For a model with M -fold-degenerate orbitals, the ratio $U_c/U_c^{SR} = (M+1)/M$.^{44,45} This relation approximately holds in the five-orbital model, in which we find $U_c/U_c^{SR} \approx 1.19$. It can be understood by the fact that in this model, the largest crystal-field splitting ($\Delta \approx 0.5$ eV) is relatively small compared to the full bandwidth ($D \approx 4.0$ eV), so the orbitals are nearly degenerate.

A nonzero Hund's rule coupling strongly affects the metal insulator transition (MIT) and the nature of both the insulating and metallic states. In the insulating phase, we find that the degenerate xz and yz orbitals are always at half-filling, and hence in a Mott insulating state, while the $x^2 - y^2$ orbital is fully occupied, in a band insulating state. The transition is therefore an orbital selective MIT.⁴⁰ Due to the interplay of J and crystal-field splitting Δ , the other two orbitals can be either in a Mott insulating ($J > \Delta$) or in a band insulating state ($J < \Delta$). Accordingly, the five-orbital model can accommodate either a high-spin ($S = 2$) or a low-spin ($S = 1$) Mott state, as illustrated in Fig. 1. These two states are separated by a low-spin-to-high-spin transition inside the insulating phase.

A general effect of the Hund's coupling is to reduce the orbital fluctuations, as it partially lifts the degeneracy of the ground-state configurations.^{40,47} This effect is more significant when J is small and the system is at or close to half-filling, where the orbital fluctuations are strong. It is known that orbital degeneracy effectively increases the kinetic energy.^{55,56} By reducing the orbital fluctuations, the Hund's coupling among the degenerate orbitals will effectively reduce the kinetic energy, which is proportional to D . As a result, it is expected that U_c decreases with increasing J . In the models at half-filling, U_c indeed decreases monotonically with increasing J/U , consistent with the above argument. But, the phase diagram of the five-orbital model shows a significant difference: as J/U is increased from zero, U_c first decreases for $J/U \lesssim 0.1$, but then increases with J/U for $J/U \gtrsim 0.1$. To understand the different behaviors of U_c in the four- and five-orbital models, we estimate and compare the Mott gaps in these two models at two cases: $0 < J \lesssim \Delta$ and $J \gtrsim \Delta$. The Mott gap G^M measures the distance between the upper and lower Hubbard bands in a Mott insulator. It can be approximated by $G^M \approx G^A - D$; here, $G^A = E^{n+1} + E^{n-1} - 2E^n$ is the excitation gap in the atomic limit, and E^n is the energy of the atomic configuration with n electrons. A reasonable estimate of U_c can be obtained from $G^A \sim D$. For $0 < J \lesssim \Delta$, the dominant configuration for the undoped compound is the $S = 1$ low-spin state. For this configuration, $G^A = U + J$ in both the four- and five-orbital models. Hence, when J/U is small, in both models we expect $U_c \sim D/(1 + J/U)$, decreasing with J/U . But, for $J \gtrsim \Delta$, the dominant configuration is the $S = 2$ high-spin state, and G^A depends on the electron filling of the high-spin state. In the four-orbital model, the system is at half-filling, $G^A \sim U + 3J$, and $U_c \sim D/(1 + 3J/U)$, still decreasing with J/U . But, in the five-orbital model with $n = 6$, $G^A \sim U - 3J$. This gives $U_c \sim D/(1 - 3J/U)$. When $J \ll U$, the behavior of U_c is still dominant by D , which decreases with J . But, further increasing J , the J dependence of D becomes weak, and the factor $1/(1 - 3J/U)$ will finally make U_c increasing with J .




FIG. 2. (Color online) (a) Evolution of the quasiparticle spectral weight with U at $n = 6.0$ and $J/U = 0.1$. (b) Same as (a) but at $J/U = 0.25$. The inset shows the difference of free energies (Δf) between three competing states for the same model parameters, with f_m , f_{MI} , and f_{OS} , respectively, denoting the free energies of the metallic, Mott insulating, and orbital selective Mott states.

Therefore, we expect a nonmonotonic behavior of U_c with increasing J/U in the five-orbital model, which is seen in the numerical results shown in Fig. 1. Interestingly, similar behavior has also been reported in a recent study on the different parameter regime of a different model (with three degenerate orbitals).⁴⁷

The Hund's rule coupling also affects the properties in the metallic state. In Fig. 2, we compare the evolution of quasiparticle spectral weight Z_α for the same model, but at two different J/U values. In both cases, the insulating phase is the $S = 2$ high-spin Mott state. On the other hand, Z_α behaves very differently in the metallic phases. At $J/U = 0.1$, $U_c \approx D$, and Z_α drops rapidly down to zero only at $U \approx U_c$. The orbital dependence of Z_α is weak. At $J/U = 0.25$, the Mott transition takes place at $U \approx 3D$. But, Z_α drops rapidly to small but nonzero values at $U \approx 2.7$ eV. This rapid drop allows us to define a crossover scale U^* in the metallic state. In Fig. 1, we plot this crossover line in the phase diagram. For large J/U , U^* can be smaller than D . U^* increases with decreasing J/U and the crossover line ends when it crosses the MIT phase boundary at $J/U \approx 0.11$. At $U < U^*$, the spectral properties of the system are similar to its noninteracting limit, with weakly renormalized quasiparticle spectral weights. But, for $U > U^*$, the quasiparticle spectral weights are strongly suppressed. In this regime, electron correlations are sufficiently strong in the metallic phase even at $U \lesssim D$, and the system is close to a Mott insulator. So, U^* roughly separates the regimes of a weakly correlated metal and a strongly correlated metal. The strongly correlated metallic phase has the features prescribed in the

incipient Mott picture. As another remarkable observation, we find that for $U > U^*$, Z_α become significantly orbital dependent. For example, at $J/U = 0.25$ and for $U \gtrsim 4$ eV, $Z_{xy} \lesssim 0.1$ and is much smaller than that of the other orbitals. This implies that the system is close to an orbital selective Mott phase (OSMP). We have calculated the ground-state energy of an OSMP with xy orbital insulating but all other orbitals metallic, and compare it with the energies of the metallic and insulating solutions in the inset of Fig. 2. Although the OSMP never becomes the true ground state, it is indeed energetically close. The stabilization of an OSMP requires a high-spin configuration.⁵⁷ This is consistent with the observation that a threshold is needed to trigger the strongly correlated metallic state.

IV. CONCLUSION

We have developed a $U(1)$ slave-spin theory, which allows the study of metal-to-Mott-insulator transition in both single-orbital and multiorbital systems. For models with a single orbital or multiple degenerate orbitals, we show that the mean-field theory in the slave-spin-condensed phase is mathematically equivalent to that of the previous Z_2 slave-spin mean-field theory. For models with multiple nondegenerate

orbitals, our $U(1)$ slave-spin formulation provides proper descriptions for both the metal and Mott insulating phases.

We have applied the $U(1)$ slave-spin approach to study a five-orbital Hubbard model for the parent iron pnictides. We find that the model exhibits a metal-to-Mott-insulator transition. The interplay between the Hund's rule coupling and crystal-field splittings strongly affect the Mott transition and the associated phases. The insulating phase can be either an $S = 1$ low-spin Mott state or an $S = 2$ high-spin Mott state, depending on the strength of the Hund's rule coupling. In the metallic phase, a crossover between a weakly correlated to a strongly correlated metallic phase exists when the Hund's rule coupling is beyond a threshold. Inside the strongly correlated metallic phase, the quasiparticle spectral weights are strongly suppressed, in agreement with the incipient Mott picture. In this phase and in the vicinity of the Mott transition, we find that an orbital selective Mott phase has ground-state energies which are nearly as competitive as those of the metallic and Mott insulating states.

ACKNOWLEDGMENTS

This work has been supported in part by NSF Grant No. DMR-1006985 and the Robert A. Welch Foundation Grant No. C-1411.

-
- ¹Y. Kamihara, T. Watanabe, H. Hirano, and H. Hosono, *J. Am. Chem. Soc.* **130**, 3296 (2008).
²Z. A. Ren *et al.*, *Chin. Phys. Lett.* **25**, 2215 (2008).
³M. Rotter, M. Tegel, and D. Johrendt, *Phys. Rev. Lett.* **101**, 107006 (2008).
⁴F.-C. Hsu *et al.*, *Proc. Natl. Acad. Sci. USA* **105**, 14262 (2008).
⁵J. Guo, S. Jin, G. Wang, S. Wang, K. Zhu, T. Zhou, M. He, and X. Chen, *Phys. Rev. B* **82**, 180520(R) (2010).
⁶M. Fang *et al.*, *Europhys. Lett.* **94**, 27009 (2011).
⁷C. de la Cruz *et al.*, *Nature (London)* **453**, 899 (2008).
⁸J. Dong *et al.*, *Europhys. Lett.* **83**, 27006 (2008).
⁹M. M. Qazilbash *et al.*, *Nat. Phys.* **5**, 647 (2009).
¹⁰W. Z. Hu, J. Dong, G. Li, Z. Li, P. Zheng, G. F. Chen, J. L. Luo, and N. L. Wang, *Phys. Rev. Lett.* **101**, 257005 (2008).
¹¹Q. Si, *Nat. Phys.* **5**, 629 (2009).
¹²J. Zhao *et al.*, *Nat. Phys.* **5**, 555 (2009).
¹³M. Yi *et al.*, *Phys. Rev. B* **80**, 024515 (2009).
¹⁴Q. Si and E. Abrahams, *Phys. Rev. Lett.* **101**, 076401 (2008).
¹⁵Q. Si, E. Abrahams, J. Dai, and J.-X. Zhu, *New J. Phys.* **11**, 045001 (2009).
¹⁶K. Haule, J. H. Shim, and G. Kotliar, *Phys. Rev. Lett.* **100**, 226402 (2008).
¹⁷A. Kutepov, K. Haule, S. Y. Savrasov, and G. Kotliar, *Phys. Rev. B* **82**, 045105 (2010).
¹⁸T. Yildirim, *Phys. Rev. Lett.* **101**, 057010 (2008).
¹⁹F. Ma, Z.-Y. Lu, and T. Xiang, *Phys. Rev. B* **78**, 224517 (2008).
²⁰C. Fang, H. Yao, W. F. Tsai, J. P. Hu, and S. A. Kivelson, *Phys. Rev. B* **77**, 224509 (2008).
²¹C. Xu, M. Muller, and S. Sachdev, *Phys. Rev. B* **78**, 020501(R) (2008).
²²J. Dai, Q. Si, J.-X. Zhu, and E. Abrahams, *Proc. Natl. Acad. Sci. USA* **106**, 4118 (2009).
²³G. S. Uhrig, M. Holt, J. Oitmaa, O. P. Sushkov, and R. R. P. Singh, *Phys. Rev. B* **79**, 092416 (2009).
²⁴K. Seo, B. A. Bernevig, and J. Hu, *Phys. Rev. Lett.* **101**, 206404 (2008).
²⁵W.-Q. Chen, K.-Y. Yang, Y. Zhou, and F.-C. Zhang, *Phys. Rev. Lett.* **102**, 047006 (2009).
²⁶P. Goswami, P. Nikolic, and Q. Si, *Europhys. Lett.* **91**, 37006 (2010).
²⁷A. Moreo, M. Daghofer, J. A. Riera, and E. Dagotto, *Phys. Rev. B* **79**, 134502 (2009).
²⁸E. Berg, S. A. Kivelson, and D. J. Scalapino, *New J. Phys.* **11**, 085007 (2009).
²⁹O. J. Lipscombe, G. F. Chen, C. Fang, T. G. Perring, D. L. Abernathy, A. D. Christianson, T. Egami, N. Wang, J. Hu, and P. Dai, *Phys. Rev. Lett.* **106**, 057004 (2011).
³⁰W. Bao *et al.*, *Chin. Phys. Lett.* **28**, 086104 (2011).
³¹W. Bao *et al.*, arXiv:1102.3674.
³²F. Ye, S. Chi, W. Bao, X. F. Wang, J. J. Ying, X. H. Chen, H. D. Wang, C. H. Dong, and M. Fang, *Phys. Rev. Lett.* **107**, 137003 (2011).
³³F. Ma, W. Ji, J. Hu, Z.-Y. Lu, and T. Xiang, *Phys. Rev. Lett.* **102**, 177003 (2009).
³⁴C. Fang, B. A. Bernevig, and J. Hu, *Europhys. Lett.* **86**, 67005 (2009).
³⁵C. Cao and J. Dai, *Phys. Rev. Lett.* **107**, 056401 (2011).
³⁶R. Yu, P. Goswami, and Q. Si, *Phys. Rev. B* **84**, 094451 (2011).
³⁷J.-X. Zhu, R. Yu, H. Wang, L. L. Zhao, M. D. Jones, J. Dai, E. Abrahams, E. Morosan, M. Fang, and Q. Si, *Phys. Rev. Lett.* **104**, 216405 (2010).
³⁸R. Yu, J.-X. Zhu, and Q. Si, *Phys. Rev. Lett.* **106**, 186401 (2011).

- ³⁹Y. Zhou, D.-H. Xu, F.-C. Zhang, and W.-Q. Chen, *Europhys. Lett.* **95**, 17003 (2011).
- ⁴⁰R. Yu and Q. Si, *Phys. Rev. B* **84**, 235115 (2011).
- ⁴¹A. Georges, G. Kotliar, W. Krauth, and M. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (1996).
- ⁴²W. F. Brinkman and T. M. Rice, *Phys. Rev. B* **2**, 4302 (1970).
- ⁴³G. Kotliar and A. E. Ruckenstein, *Phys. Rev. Lett.* **57**, 1362 (1986).
- ⁴⁴L. de'Medici, A. Georges, and S. Biermann, *Phys. Rev. B* **72**, 205124 (2005).
- ⁴⁵S. Florens and A. Georges, *Phys. Rev. B* **70**, 035114 (2004).
- ⁴⁶A. Vaezi and X.-G. Wen, [arXiv:1010.5744](https://arxiv.org/abs/1010.5744); M. Mardani, M.-S. Vaezi, and A. Vaezi, [arXiv:1111.5980](https://arxiv.org/abs/1111.5980).
- ⁴⁷L. de'Medici, *Phys. Rev. B* **83**, 205112 (2011).
- ⁴⁸R. Nandkishore, M. A. Metlitski, and T. Senthil, *Phys. Rev. B* **86**, 045128 (2012).
- ⁴⁹In this paper, we refer as Z_2 slave-spin theory the formulation initially proposed in Ref. 44 and further developed in Ref. 51. There is some difference between this approach and the one proposed in A. Rüegg, S. D. Huber, and M. Sgrist, *Phys. Rev. B* **81**, 155118 (2010), although both formulations involve a Z_2 invariance.
- ⁵⁰C. Castellani, C. R. Natoli, and J. Ranninger, *Phys. Rev. B* **18**, 4945 (1978).
- ⁵¹S. R. Hassan and L. de'Medici, *Phys. Rev. B* **81**, 035106 (2010).
- ⁵²In our earlier paper (Ref. 40), the two- and four-orbital models are already studied using the $U(1)$ slave-spin mean-field method for the five-orbital model, although the method itself was not explicitly introduced there.
- ⁵³S. Graser, T. A. Maier, P. J. Hirschfeld, and D. J. Scalapino, *New J. Phys.* **11**, 025016 (2009).
- ⁵⁴T. Einav, R. Yu, and Q. Si (unpublished).
- ⁵⁵O. Gunnarsson, E. Koch, and R. M. Martin, *Phys. Rev. B* **56**, 1146 (1997).
- ⁵⁶S. Florens, A. Georges, G. Kotliar, and O. Parcollet, *Phys. Rev. B* **66**, 205102 (2002).
- ⁵⁷L. de'Medici, S. R. Hassan, M. Capone, and X. Dai, *Phys. Rev. Lett.* **102**, 126401 (2009).