

Symmetry-protected topological phases with charge and spin symmetries: Response theory and dynamical gauge theory in two and three dimensions

Peng Ye¹ and Juven Wang^{2,1}¹*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada N2L 2Y5*²*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

(Received 9 July 2013; revised manuscript received 11 November 2013; published 10 December 2013)

A large class of symmetry-protected topological phases (SPT) in boson/spin systems have been recently predicted by the group cohomology theory. In this work, we consider bosonic SPT states *at least* with charge symmetry $[U(1) \text{ or } Z_N]$ or spin- S^z rotation symmetry $[U(1) \text{ or } Z_N]$ in two (2D) and three dimensions (3D) and the surface of 3D. If both are $U(1)$, we apply external electromagnetic field/“spin gauge field” to study the charge/spin response. For the SPT examples we consider {i.e., $U_c(1) \times Z_2^T$, $U_s(1) \times Z_2^T$, $U_c(1) \times [U_s(1) \times Z_2]$; subscripts c and s are short for charge and spin; Z_2^T and Z_2 are time-reversal symmetry and π rotation about S^y , respectively}, many variants of Witten effect in the 3D SPT bulk and various versions of *anomalous* surface quantum Hall effect are defined and systematically investigated. If charge or spin symmetry reduces to Z_N by considering charge- N or spin- N condensate, instead of the linear response approach, we gauge the charge/spin symmetry, leading to a *dynamical gauge theory* with some remaining global symmetry. The 3D dynamical gauge theory describes a symmetry-enriched topological phase (SET), i.e., a topologically ordered state with global symmetry which admits nontrivial ground-state degeneracy depending on spatial manifold topology. For the SPT examples we consider, the corresponding SET states are described by dynamical topological gauge theory with topological BF term and axionic Θ term in 3D bulk. In addition, the surface of SET is described by the chiral boson theory with quantum anomaly.

DOI: [10.1103/PhysRevB.88.235109](https://doi.org/10.1103/PhysRevB.88.235109)

PACS number(s): 73.43.Cd, 75.10.Jm

I. INTRODUCTION

Searching for exotic quantum many-body states is one of the main goals in condensed matter physics. Physically, all states (or “phases” interchangeably) are either gapped states or gapless states, depending on the bulk energy gap between the ground state and first excitation. Recently, considerable attentions have been drawn by both kinds of states. Some gapless states with strong correlations emerge at quantum critical points without long-lived quasiparticles, in which some new approaches based on the holographic principle in string theory are introduced into condensed matter physics. On the other hand, the notion of “quantum entanglement” becomes an important viewpoint for understanding gapped states.^{1,2} By utilizing the well-defined “local unitary transformation (LU)” suggested by Chen *et al.*, all gapped states are classified into two categories: “short-range entangled states (SRE)” and “long-range entangled states (LRE).”^{3–5}

For LRE states, there is no canonical LU to connect LRE to a trivial direct product state (a state with zero entanglement range). The well-known fractional quantum Hall states⁶ (FQH) are a class of LRE states with highly long-range entanglement. A LRE state generally admits an intrinsic topological order (TO) or “topological order” for short.^{7–9} A TO state is defined by the following features: ground-state degeneracy in a topologically nontrivial closed manifold,^{7–10} or emergent fermionic/anyonic excitations,^{11,12} or chiral gapless edge excitations.^{13,14} If, in addition to a topological order, the ground state also has a global symmetry, such a state will be referred to as a “symmetry-enriched topological (SET) phase.”

In contrast, a SRE state can always be adiabatically deformed to a trivial direct product state and thus generically

has no TO, which looks quite boring. However, recent rapid progress indicates that some SRE states are quite special for the reason that these SRE states can not be adiabatically deformed to a direct product state *unless* a certain onsite global symmetry group (i.e., a global symmetry operation that is a direct product of the operation on each lattice site) is broken, unveiling the existence of nontrivial symmetry-protected properties in such SRE states.^{3–5} This fact leads to the notion of “symmetry-protected topological phases” (SPT), which are currently under considerable investigation. The well-known three-dimensional noninteracting fermionic topological insulator (TI) state is a fermionic SPT in which the surface single massless Dirac fermion is protected by the symmetry group $U(1) \times Z_2^T$ where $U(1)$ is charge symmetry related to the fermionic particle-number conservation and Z_2^T is time-reversal symmetry.^{15–21} The Haldane phase in an antiferromagnetic Heisenberg spin chain with integer spin and $SO(3)$ spin-rotation symmetry is a bosonic SPT.²² At each edge of the spin chain, a free spin- $\frac{1}{2}$ degree of freedom arises, which is protected by $SO(3)$ symmetry group. In the following, we will only discuss SPT states in boson/spin systems. In boson/spin systems, interactions are crucial to realize nontrivial SPT states or TO states. Some useful theoretical approaches are recently proposed, such as exactly soluble models,^{23–30} the fusion category approach,^{31,32} projective construction,^{33–35} K -matrix Chern-Simons approach^{10,36–40} and other field theory approaches,^{41–45} the group cohomology approach,^{3–5} and modular invariance of edge conformal field theory (CFT).⁴⁶ In the group cohomology classification theory, a SPT state with symmetry group “ G ” in d -dimensional spatial lattice is classified by the group cohomology $\mathcal{H}^{d+1}[G, U(1)]$.^{3–5} The famous Haldane phase is classified by $\mathcal{H}^2[SO(3), U(1)] = Z_2$ indicating that there is only one nontrivial Haldane phase in

addition to one trivial phase. In 3D, the “bosonic topological insulator” (BTI) with $U(1) \times Z_2^T$ are studied by Refs. 34, 35, 41, 47, and 48.

The group cohomology theory⁴⁹ provides the elements of the cohomology group to label SPT states, while it is also interesting to build up the connection between the abstract labels and physical properties (such as electromagnetic response). The following two general approaches are mainly applied in the community: *response theory* and *dynamical gauge theory*.

For $U(1)$ SPT states in 2D which are labeled by $\mathcal{H}^3[U(1), U(1)] = \mathbb{Z}$, an applicable way to understand the ground states is to add an external $U(1)$ gauge field to study the *response theory*. Here, the $U(1)$ gauge field is nondynamical. The resultant response action is a Chern-Simons term with a quantized Hall conductance $\sigma = 2k \times \frac{e^*{}^2}{2\pi}$ where $k \in \mathbb{Z}$ and e^* is the fundamental gauge charge carried by bosons.^{33,50-53} This integer k is also the integer label defined in the group cohomology classification theory as a one-to-one correspondence.⁵⁴

On the other hand, a pioneer work by Levin and Gu²³ leads to a breakthrough. They study the 2D Ising paramagnets (SPT states with Z_2 symmetry) by fully gauging the Z_2 symmetry, which results in a *dynamical gauge theory* where the gauge field variables become new degrees of freedom. Their gauging procedure indicates that a trivial paramagnet maps to a dynamical Z_2 gauge theory (i.e., Z_2 toric code), while a nontrivial paramagnet maps to a nontrivial dynamical Z_2 gauge theory (i.e., Z_2 doubled semions). This correspondence gives Z_2 classification which is consistent with the group cohomology classification $\mathcal{H}^3[Z_2, U(1)] = Z_2$, i.e., one nontrivial SPT state and one trivial state. In a more complicated symmetry group, one can partially gauge a normal subgroup of the global symmetry group of SPT. The resultant dynamical gauge theory may describe a SET state since some global symmetries remain in the gauged theory.^{37,38,55,56}

In this work, we will elaborate the two approaches “response theory” and “dynamical gauge theory” in many SPT examples which at least have charge symmetry [$U(1)$ or Z_N] or spin- S^z rotation symmetry [$U(1)$ or Z_N] in two and three dimensions. The symmetry groups in this work are more relevant to realistic physical systems in condensed matter. More specifically, if both charge and spin symmetry are the simplest continuous $U(1)$, we apply the response theory to study the charge and spin dynamics of the three-dimensional SPT bulk (denoted by Σ^3), the surface (denoted by $\partial\Sigma^3$) of the Σ^3 bulk, and the two-dimensional SPT bulk (denoted by Σ^2). We stress that, instead of looking into microscopic lattice models and utilizing sophisticated mathematical techniques or physical arguments, in the whole discussions of this work we will attempt to start with the minimal necessary physical input (such as gauge invariance principle, absence of topological order, etc.) to extract the response theory. We will also apply the so-called “top-down approach” based on the “ K_G -matrix” Chern-Simons term (a gauged version of K -matrix Chern-Simons field theory by adding the external electromagnetic field A_μ^c minimally coupled to charge current and the external “spin gauge field” A_μ^s minimally coupled to spin current). From this response approach, we will construct many variants of the celebrated Witten effect and also many different versions of quantum Hall effects, depending on the choice of probe

fields (external electromagnetic field or external spin gauge field) and the choice of response current (charge current or spin current). By studying the response phenomena case by case, we emphasize that, although both $\partial\Sigma^3$ and Σ^2 are two dimensional, the response theory on $\partial\Sigma^3$ is realized in an *anomalous* fashion in a sense that an extra spatial dimension (deep into the bulk) is required, in sharp contrast to the response theory of SPT defined on Σ^2 with the same symmetry implementation. The above results will be discussed in Sec. II. A summary of Witten effects and quantum Hall effects is shown in Table II.

If charge or spin symmetry reduces to Z_N by considering charge- N or spin- N condensate, we change our strategy due to the “Meissner effect” for charge and spin. Instead of the response approach, we diagnose the SPT states by gauging a normal subgroup (i.e., the charge symmetry and spin- S^z rotation symmetry) of the whole symmetry of SPT, resulting in a dynamical gauge theory with both gauge symmetry and global symmetry. In the examples we will consider, each dynamical gauge theory in Σ^3 describes a SET state in which ground-state degeneracy is nontrivially dependent on the spatial manifold topology. More specifically, the generic form of the dynamical gauge theory in Σ^3 is a topological gauge theory and consists of two topological terms: topological BF terms and variants of axionic Θ terms. In addition, its surface ($\partial\Sigma^3$) theory is a chiral boson matter field theory that admits quantum anomaly and is meant to cancel the anomaly from Σ^3 due to the existence of topological BF term. By studying the dynamical gauge theory description of SPT states, we emphasize that, although both $\partial\Sigma^3$ and Σ^2 are two dimensional, the surface dynamical theory on $\partial\Sigma^3$ after promoting the bulk SPT to a SET state can not be realized in a dynamical gauge theory defined on Σ^2 after promoting the Σ^2 SPT to a dynamical gauge theory. The strategy of dynamical gauge theory provides the connection between SPT and SET. Most importantly, if we gauge a normal subgroup and place gauge connection on the lattice links,²³ we can in principle study the corresponding dynamical gauge theory with boson matter put on lattice in a numerical simulation approach, in order to extract the nature of the underlying SPT state. The above results will be discussed in Sec. III. Some key results are collected in Table III for the reader’s convenience. Section IV is devoted to the conclusions of the paper.

II. CONTINUOUS $U_c(1)$ CHARGE SYMMETRY AND $U_s(1)$ SPIN SYMMETRY

A. General response theory based on gauge invariance

1. Θ -matrix response theory in Σ^3

Let us begin with a three-dimensional bulk of spin-1 and charge-1 boson systems where *at least* $U_c(1)$ and $U_s(1)$ are unbroken. Here, the conserved charge corresponding to $U_s(1)$ symmetry is the spin density S^z . Let us probe the charge and spin dynamics by gauging these two $U(1)$ symmetries rendering two *nondynamical* gauge fields: the “spin gauge field” A_μ^s and the usual electromagnetic gauge field A_μ^c . Note that the spin gauge field A_μ^s directly couples to spin density and spin current.

TABLE I. Four different quantum Hall effects.

Quantum Hall effects	Hall conductance	Probe fields	Response current
Quantum charge Hall effect	$\sigma^c, \tilde{\sigma}^c$	\mathbf{E}^c	\mathbf{J}^c
Quantum spin Hall effect	$\sigma^s, \tilde{\sigma}^s$	\mathbf{E}^s	\mathbf{J}^s
Quantum charge-spin Hall effect	$\sigma^{cs}, \tilde{\sigma}^{cs}$	\mathbf{E}^s	\mathbf{J}^c
Quantum spin-charge Hall effect	$\sigma^{sc}, \tilde{\sigma}^{sc}$	\mathbf{E}^c	\mathbf{J}^s

According to the general principle of *gauge invariance*, the linear response theory obtained by integrating out bosons should be the following general form in the long-wavelength limit:

$$\mathcal{Z}[A_\mu^s, A_\mu^c] = e^{iS_{\text{top}}}, \quad S_{\text{top}} = \int d^4x \mathcal{L}, \quad (1)$$

$$\mathcal{L} = \frac{\Theta_{IJ}}{8\pi^2} \partial_\mu A_\nu^I \partial_\lambda A_\rho^J \epsilon^{\mu\nu\lambda\rho},$$

where the partition function \mathcal{Z} is a functional of the two nondynamical gauge fields. The indices $I, J = 1, 2$. A_μ^1, A_μ^2 denote A_μ^c, A_μ^s , respectively. The usual Maxwell terms are not written explicitly for the reason that their physical effects are to renormalize electromagnetic parameters (dielectric constant and permeability) of the bulk systems. The coefficient Θ_{IJ} forms a symmetric matrix

$$\Theta = \begin{pmatrix} \theta_c & \theta_0 \\ \theta_0 & \theta_s \end{pmatrix}. \quad (2)$$

The Lagrangian \mathcal{L} can be written in terms of three parts:

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_c + \mathcal{L}_s + \mathcal{L}_0, \\ \mathcal{L}_c &= \frac{\theta_c}{8\pi^2} \partial_\mu A_\nu^c \partial_\lambda A_\rho^c \epsilon^{\mu\nu\lambda\rho}, \\ \mathcal{L}_s &= \frac{\theta_s}{8\pi^2} \partial_\mu A_\nu^s \partial_\lambda A_\rho^s \epsilon^{\mu\nu\lambda\rho}, \\ \mathcal{L}_0 &= \frac{\theta_0}{4\pi^2} \partial_\mu A_\nu^c \partial_\lambda A_\rho^s \epsilon^{\mu\nu\lambda\rho}. \end{aligned} \quad (3)$$

In the following, the “electric field” \mathbf{E}^c and “magnetic field” \mathbf{B}^c are constructed from the gauge field A_μ^c in the usual convention. And, the new notions of “spin-electric field” \mathbf{E}^s and “spin-magnetic field” \mathbf{B}^s are specified to the gauge field A_μ^s .

If the spin current and charge current are well defined on the surface $\partial\Sigma^3$ of the three-dimensional bulk Σ^3 or on a strictly two-dimensional plane Σ^2 , we define four kinds of quantum Hall effects as shown in Table I. For example, in the quantum charge-spin Hall effect, the external spin gauge field’s electric field \mathbf{E}^s drives a perpendicular charge current \mathbf{J}^c . The Hall conductance is denoted by σ^{cs} ($\tilde{\sigma}^{cs}$) if the Hall effect is on Σ^2 ($\partial\Sigma^3$).

B. $U_c(1) \times Z_2^T$ in Σ^3

Let us consider the Σ^3 bulk with $U_c(1) \times Z_2^T$ symmetry, where $U_c(1)$ and Z_2^T are charge-conservation symmetry and time-reversal symmetry, respectively. A bosonic system with this symmetry in Σ^3 is a *bosonic topological insulator* (BTI) which has been recently studied.^{34,35,41,47,48} By applying A_μ^c to probe the topological electromagnetic properties, the resultant

response theory gives rise to the topological magnetoelectric effect with θ_c quantized at $2\pi \bmod(4\pi)$. In the following, we will study the response theory along two approaches. First, based on the response current and the definition of SPT states, we will derive the quantization of θ_c , charge Witten effect^{20,21,57,58} in bosonic topological insulator in Σ^3 , and the quantum charge Hall effect on the Z_2^T -broken surface $\partial\Sigma^3$ and Z_2^T -broken 2D plane Σ^2 . Partially along the lines of the physical arguments of Ref. 35, we shall elaborate the derivation in details in order for the generalization to other symmetry groups (Table II) in the remaining parts of Sec. II. Second, we shall rederive these results through the top-down approach by comparing the K_G matrix on Σ^2 and its anomalous realization on $\partial\Sigma^3$ (“ K_G ” will be defined later).

1. Charge Witten effect in Σ^3

BTI admits surface charge Hall effect by breaking Z_2^T on the surface. The formation of surface can be viewed as an interface between vacuum and BTI bulk where the derivative of θ_c forms a two-dimensional domain wall. Let us study the response equation of A^c in the bulk θ_c term

$$J_\mu^c \equiv \frac{\delta \mathcal{L}_c}{\delta A_\mu^c} = 2 \frac{\theta_c}{8\pi^2} \partial_\nu \partial_\lambda A_\rho^c \epsilon^{\mu\nu\lambda\rho} = \frac{\theta_c}{4\pi^2} \partial_\nu \partial_\lambda A_\rho^c \epsilon^{\mu\nu\lambda\rho}, \quad (4)$$

where J_μ^c is (3+1)D response charge current. The prefactor 2 comes from twice variations with respect to A_μ^c . The zero component J_0^c denotes the response charge density probed by external field A_μ^c :

$$J_0^c = \frac{\theta_c}{4\pi^2} \nabla \cdot \mathbf{B}^c, \quad (5)$$

where \mathbf{B}^c is the magnetic field variable. If the gauge field A^c is smooth everywhere, $\nabla \cdot \mathbf{B}^c = 0$ due to absence of magnetic charge. However, if singular configuration is allowed, the divergence may admit singularities in the bulk and its total contribution in the bulk is quantized due to the Dirac quantization condition (or more general Schwinger-Zwanziger quantization condition^{59,60}). In a simplest configuration, let us consider one magnetic monopole which is located at the origin of the three-dimensional space:

$$\int d^3x \nabla \cdot \mathbf{B}^c = 2\pi N_m^c, \quad (6)$$

where $N_m^c \in \mathbb{Z}$ is an integer-valued “magnetic charge” in the A_μ^c gauge group. Therefore, the corresponding response charge is

$$N^c = \int d^3x J_0^c = \frac{\theta_c}{2\pi} N_m^c, \quad (7)$$

which indicates that a nonzero theta term supports a “polarization charge cloud” in the presence of magnetic monopole.

TABLE II. Charge and spin response of spin-1 and charge-1 boson systems (the case of Θ_{22} is generalized to spin systems of any integer spin s). $q_i = s, s-1, s-2, \dots$ and $q_i > 0$ where s is total spin defined by $\mathbf{S}^2 = s(s+1)$. The full units of σ^c , σ^s , σ^{cs} , σ^{sc} are $\frac{e^2}{h}$, \hbar , e , and e , respectively, where e is elementary electric charge and \hbar is reduced Planck constant. $U_c(1)$ and $U_s(1)$ denote the $U(1)$ symmetry of charge and spin, respectively. Z_2 symmetry in “ $U_s(1) \times Z_2$ ” is the π rotation about S^y . Z_2^T is time-reversal symmetry. $k \in \mathbb{Z}$, and, “ \times ” stands for “semidirect product.”

Axionic theta angle	Full symmetry group	3D bulk (Σ^3) response	Surface ($\partial\Sigma^3$) anomalous response	2D plane (Σ^2) response
$\Theta_{11} \equiv \theta_c = 2\pi + 4\pi k$, charge-1 boson system	$U_c(1) \times Z_2^T$	Charge Witten effect: $N^c = n^c + N_m^c$	Quantum charge Hall effect on Z_2^T broken $\partial\Sigma^3: \tilde{\sigma}^c = (1+2k)\frac{1}{2\pi}$	Quantum charge Hall effect on Z_2^T broken $\Sigma^2: \sigma^c = 2k\frac{1}{2\pi}$
$\Theta_{22} \equiv \theta_s = 2\pi + 4\pi k$, spin or boson systems with integer spins	$U_s(1) \times Z_2^T$	Spin Witten effect: $N^s = \sum_i q_i n_i^s + N_m^s \sum_i q_i^2$	Quantum spin Hall effect on Z_2^T broken $\partial\Sigma^3: \tilde{\sigma}^s = (1+2k)\frac{1}{2\pi} \sum_i q_i^2$	Quantum spin Hall effect on Z_2^T broken $\Sigma^2: \sigma^s = 2k\frac{1}{2\pi} \sum_i q_i^2$
$\Theta_{12} = \Theta_{21} \equiv \theta_0$, $= \pi + 2\pi k$, boson system of charge-1 and spin-1	$U_c(1) \times [U_s(1) \times Z_2]$	Mutual Witten effect: $N^c = n^c + \frac{1}{2}N_m^c$; $N^s = n_+^s - n_-^s + \frac{1}{2}N_m^c$	Quantum charge-spin/ spin-charge Hall effects on Z_2 broken $\partial\Sigma^3: \tilde{\sigma}^{cs} = \tilde{\sigma}^{sc} = (\frac{1}{2} + k)\frac{1}{2\pi}$	Quantum charge-spin/ spin-charge Hall effects on Z_2 broken $\Sigma^2: \sigma^{cs} = \sigma^{sc} = k\frac{1}{2\pi}$

A monopole can also trivially attach integer number (n^c) of charge-1 bosons in the bulk. Therefore, the whole formula of the so-called *charge Witten effect*^{57,58} in Table II can be expressed as

$$N^c = n^c + \frac{\theta_c}{2\pi} N_m^c. \quad (8)$$

2. Quantum charge Hall effect on Z_2^T -broken $\partial\Sigma^3$ and Z_2^T -broken Σ^2

The bulk θ_c term \mathcal{L}_c can be written as a surface term

$$\mathcal{L}_{c,\partial\Sigma^3} = \frac{\theta_c}{8\pi^2} A_\mu^c \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}, \quad (9)$$

which leads to the surface response current

$$J_\mu^{c,\partial\Sigma^3} \equiv \frac{\delta \mathcal{L}_{c,\partial\Sigma^3}}{\delta A_\mu^c} = \frac{\theta_c}{4\pi^2} \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}. \quad (10)$$

The surface charge Hall conductance σ^c is defined by Ohm's equation $J_x^{c,\partial\Sigma^3} = \tilde{\sigma}^c E_y^c$ where E_y^c is the electric field along the y direction (assuming that $\partial\Sigma^3$ is parametrized by x - y coordinates):

$$\tilde{\sigma}^c = \frac{\theta_c}{4\pi^2}. \quad (11)$$

To understand the quantization of the surface charge Hall conductance $\tilde{\sigma}^c$, we need to first understand the quantization on the θ_c angle and σ^c in a strictly 2D system (i.e., Σ^2) which is defined as a $U_c(1)$ SPT. Let us write the Chern-Simons term in Z_2^T broken Σ^2 which describes the response theory of $U_c(1)$ SPT on Σ^2 :

$$\mathcal{L}_{c,\Sigma^2} = \frac{\sigma^c}{2} A_\mu^c \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}. \quad (12)$$

Upon adiabatically piercing Σ^2 by 2π magnetic flux ($\Phi^c = \int d^2x \nabla \times \mathbf{A}^c = 2\pi$), the total response charge $\int d^2x J_0^{c,\Sigma^2} = \sigma^c \int d^2x \nabla \times \mathbf{A}^c = 2\pi\sigma^c$. In SPT states where topological order is trivial by definition (*at least* no exotic fractional charge), this pumped charge in the center of the vortex core must be quantized at integer carried by charge-1 bosons of

underlying microscopic model, such that $2\pi\sigma^c \in \mathbb{Z}$. This condition is enough for the free-fermion system. However, for the bosonic system [i.e., $U_c(1)$ SPT we are considering], we need further forbid quasiparticles which carry nonbosonic statistics (i.e., fermionic statistics and anyonic statistics) in order to obtain states without topological order. To achieve this goal, one can spatially exchange two vortex cores of 2π fluxes each of which traps $2\pi\sigma^c$ quasiparticles. The quasiparticles in the first vortex core will perceive a π phase as half a magnetic flux of the second vortex core, and vice versa. The total Aharonov-Bohm phase “ Φ_{AB} ” in the Chern-Simons theory, however, is only *half* of the totally accumulated quantum phases: $\Phi_{AB} = \frac{1}{2} \times (2\pi\sigma^c \times \pi + 2\pi\sigma^c \times \pi) = 2\pi^2\sigma^c$.⁶¹ In order to forbid nonbosonic statistics, a new condition should be satisfied: $\Phi_{AB}/2\pi \in \mathbb{Z}$. Overall, $2\pi\sigma^c/2 \in \mathbb{Z}$, i.e.,

$$\sigma^c = 2k\frac{1}{2\pi}, \quad (13)$$

where $k \in \mathbb{Z}$. After this preparation, let us move on to the θ_c angle quantization and its periodicity. Generally, a theta term is odd under Z_2^T and thus breaks Z_2^T symmetry and thus results in CP violation in the context of high-energy physics⁵⁷ because under Z_2^T , $\mathbf{E}^c \rightarrow \mathbf{E}^c$, $\mathbf{B}^c \rightarrow -\mathbf{B}^c$, $\frac{\theta_c}{8\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\mu A_\nu^c \partial_\lambda A_\rho^c = \frac{\theta_c}{4\pi^2} \mathbf{E}^c \cdot \mathbf{B}^c \rightarrow -\frac{\theta_c}{4\pi^2} \mathbf{E}^c \cdot \mathbf{B}^c$. However, if θ_c admits a periodic shift such that $-\theta_c$ can be shifted back to θ_c , the action eventually is time-reversal invariant. Therefore, the minimal theta value should be one half of its periodicity (say, P), and the symmetry group for the bulk is indeed $U_c(1) \times Z_2^T$ as we defined at the beginning of this section. Physically, the periodicity can be understood as trivially depositing arbitrary copies of Σ^2 Hall systems onto the surface.^{35,41} As such, a P shift in θ_c leads to an additional term in the surface charge Hall conductance formula (11):

$$\tilde{\sigma}^{c'} - \tilde{\sigma}^c = \frac{P}{4\pi^2}, \quad (14)$$

which is contributed by deposited Σ^2 layers which are described by Eq. (13). A minimal choice is $\frac{P}{4\pi^2} = 2 \times \frac{1}{2\pi}$,

so that $P = 4\pi$, and the minimal choice of θ_c is $\frac{P}{2} = 2\pi$, i.e.,

$$\theta_c = 2\pi + 4\pi k, \quad (15)$$

where the integer k is the same k defined in Eq. (13).

Substituting Eq. (15) into (11) leads to

$$\tilde{\sigma}^c = (1 + 2k) \frac{1}{2\pi}. \quad (16)$$

The most anomalous phenomenon on $\partial\Sigma^3$ is that the surface quantum charge Hall conductance $\tilde{\sigma}^c$ admits a $\frac{1}{2\pi}$ value which can not be realized in Σ^2 where σ^c is always even integer copies of $1/2\pi$. In addition, substituting Eq. (15) into (8) leads to

$$N^c = n^c + N_m^c, \quad (17)$$

where $k = 0$ is selected for simplicity. $\theta_c = 2\pi$ is topologically distinct from $\theta_c = 0$ trivial vacuum once the symmetry group $U_c(1) \rtimes Z_2^T$ is unbroken. Different choices of k actually correspond to the *same phase*.⁴¹

3. Anomalous K_G matrix on $\partial\Sigma^3$

In the above discussion, we obtained the charge Hall conductance in Σ^2 and its anomalous realization in Z_2^T broken $\partial\Sigma^3$ based on the general principle of gauge invariance, response definition, and the definition of SPT states. In those derivations, the microscopic degree of freedoms is not explicitly written in terms of Lagrangian or Hamiltonian. In the following, we shall start with microscopic degrees of freedom, which is called the ‘‘top-down approach.’’

By definition, the fundamental elements of a given 2D SPT microscopic model are spins or bosonic particles. In the hydrodynamical approach, however, the low-energy modes dominating the partition function can be effectively replaced by a SET of *statistical* one-form $U(1)$ gauge fields $\{a_\mu^I\}$ or two-form $U(1)$ gauge fields $\{b_{\mu\nu}^I\}$ ($I = 1, 2, \dots$) or higher-form gauge fields. In the following, these gauge field variables are dubbed ‘‘intrinsic/statistical gauge fields’’ interchangeably. Especially in a 2D system, the current operator J_μ of a point particle can be expressed as $J^\mu = \frac{1}{2\pi} \epsilon^{\mu\nu\lambda} \partial_\nu a_\lambda$ which automatically resolves the current conservation equation $\partial_\mu J^\mu = 0$.

What is the generic low-energy theory of the SPT state in terms of these dynamical gauge fields? If we only focus on the topological properties of the SPT ground state (or a general ground state with Abelian topological order), the renormalization group flows to an infrared fixed-point field theory in the Chern-Simons form in $(2+1)D$.⁶² In other words, the low-energy field theory of the microscopic SPT model is effectively described by a generic Chern-Simons theory of $\{a_\mu^I\}$ with a K -matrix coefficient: $\mathcal{L}_{\text{SPT}} = \frac{1}{4\pi} K_{IJ} a_\mu^I \partial_\nu a_\lambda^J \epsilon^{\mu\nu\lambda}$ where, $I, J = 1, 2, \dots$

Thus, it is effective to describe the internal microscopic degrees of freedom of SPT states by a generic K -matrix Chern-Simons field theory of $\{a_\mu^I\}$. Based on this low-energy field theory, the response theory is straightforward by adding a minimal coupling term $J^\mu A_\mu$ where A_μ is an external gauge field and J^μ is the current operator carrying gauge charge (i.e., the Noether current related to a global symmetry before gauging it) in terms of $\{a_\mu^I\}$.

This top-down approach starts with intrinsic gauge field a_μ to construct SPT with Lagrangian $\mathcal{L}_{\text{SPT}}(a)$ and then probe it by external field A_μ . The Lagrangian becomes $\mathcal{L}_{\text{SPT+Gauge}}(a, A)$. By integrating out $\{a_\mu^I\}$ to obtain the low-energy physics of external field A_μ , we obtain an effective theory $\mathcal{L}_{\text{SPT+Gauge}}(A)$. This is exactly what we would like to do in the following: to confirm our previous result (response theory of electromagnetic field A_μ^c which couples to charge current and ‘‘spin gauge field’’ A_μ^s which couples to spin current) by comparing with the top-down approach (starting from the intrinsic $\{a_\mu\}$ statistical gauge fields). It should be noted that the field variables A_μ^s and A_μ^c are always treated as nondynamical background fields in the whole Sec. II due to the standard definition of linear response theory.

The above procedure is known and applied in the literature,^{23,36–40,63} however, to be self-contained and make this method more accessible to the research community, in Appendix A we carry out an explicit derivation for the following steps:

$$\mathcal{L}_{\text{SPT}}(a) \rightarrow \mathcal{L}_{\text{SPT+Gauge}}(a, A) \rightarrow \mathcal{L}_{\text{SPT+Gauge}}(A). \quad (18)$$

We use the K -matrix Chern-Simons effective field theory approach to understand this procedure:

$$\mathcal{L}_{\text{SPT}}(a) = \frac{1}{4\pi} K_{S,IJ} \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J \quad (19)$$

and we end up with

$$\mathcal{L}_{\text{SPT+Gauge}}(A) = \frac{1}{4\pi} K_{G,IJ} \epsilon^{\mu\nu\rho} A_\mu^I \partial_\nu A_\rho^J. \quad (20)$$

We denote K_S and K_G as K matrices for $\mathcal{L}_{\text{SPT}}(a)$ and $\mathcal{L}_{\text{SPT+Gauge}}(A)$, respectively. a_μ^I ($I = 1, 2, \dots$) represent a set of intrinsic fields a_μ in a general case. Each of A_μ^I ($I = 1, 2, \dots$) represents an external field which couples to the matter current carrying $U_I(1)$ gauge charge. Our inspiration is from earlier pioneer works. In Ref. 36, Lu and Vishwanath focus on $\mathcal{L}_{\text{SPT}}(a)$. In Ref. 39, Cheng and Gu attempt to apply the braiding statistics in $\mathcal{L}_{\text{SPT+Gauge}}(A)$ to determine the classification of SPT. Our approach is analog to the work by Hung and Wan⁴⁰ who had carried out the simplest gauging procedure for Z_N SPT in 2D. On the other hand, our key focus is to bridge Ref. 36 to Refs. 39 and 40 by directly gauging the global symmetry current and then apply to more complicated symmetry groups (in Appendix A).

Let us explicitly work out the response theory on Z_2^T broken $\partial\Sigma^3$ of 3D bulk with $U_c(1) \rtimes Z_2^T$, and on Z_2^T broken Σ^2 . Here, we save detailed derivations to Appendix A and list key results directly. We first study the response theory on Σ^2 with $U_c(1)$ global symmetry. What we start with is the intrinsic SPT’s $K_S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. By gauging the $U(1)$ global symmetry current coupling to A^c , we obtain $K_G = 2p$ and response action

$$\mathcal{L}_{\text{SPT+Gauge}}(A^c) = \frac{2p}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^c \partial_\nu A_\rho^c \quad (21)$$

with $p \in \mathbb{Z}$ labeling the \mathbb{Z} class of the cohomology group $\mathcal{H}^3[U(1), U(1)] = \mathbb{Z}$. The charge Hall conductance as the quantum feature of the response of this $\mathcal{L}_{\text{SPT+Gauge}}(A^c)$ is

$$\sigma^c = 2p \frac{1}{2\pi}. \quad (22)$$

This result matches exactly as Eq. (13). On the other hand, the anomalous K_G matrix in $\partial\Sigma^3$ is harder to obtain from the top-down approach because the full classification of (3+1)D SPT from intrinsic topological field theory is not yet known to be complete (not even fully matching with the group cohomology). In principle, we should have topological terms (such as $g_1\epsilon^{\mu\nu\rho\tau}\partial_\mu a_\nu\partial_\rho a_\tau + g_2\epsilon^{\mu\nu\rho\tau}b_{\mu\nu}\partial_\rho a_\tau + \text{etc.}$) to generate all classes of $\mathcal{H}^3[\text{U}(1)\times\text{Z}_2^T, \text{U}(1)] = \text{Z}_2^2$ (or Z_2^3 according to other field theory approaches⁴¹). Here, we simply adopt the result from the previous section to state the effective $K_{G,\partial\Sigma^3} = 2p + \theta_c/2\pi$, which means

$$\mathcal{L}_{\text{SPT+Gauge}}(A^c) = \frac{2p + \theta_c/2\pi}{4\pi}\epsilon^{\mu\nu\rho}A_\mu^c\partial_\nu A_\rho^c \quad (23)$$

with surface charge Hall conductance

$$\bar{\sigma}^c = (2p + \theta_c/2\pi)\frac{1}{2\pi}. \quad (24)$$

Therefore, we have Eqs. (22) and (24) written in a consistent manner as Eqs. (13) and (16). Now, we have two K_G matrices: one is for the surface and one is for the 2D SPT. We dubbed $K_{G,\partial\Sigma^3}$ the ‘‘anomalous K_G matrix’’ since it has the same symmetry as the 2D SPT but different response theory. The matrices in the present example are nothing but a number; in the example $\text{U}_c(1)\times[\text{U}_s(1)\times\text{Z}_2]$, the matrices are two dimensional.

C. $\text{U}_s(1)\times\text{Z}_2^T$ in Σ^3

1. Quantum spin Hall effect on Z_2^T broken $\partial\Sigma^3$ and Z_2^T broken Σ^2

Now, let us move on to the bulk θ_s term which is the response action of three-dimensional SPT protected by $\text{U}_s(1)\times\text{Z}_2^T$. Under Z_2^T , $\mathbf{E}^s \rightarrow -\mathbf{E}^s$, $\mathbf{B}^s \rightarrow \mathbf{B}^s$ in contrast to $\text{U}_c(1)$ gauge fields. And, the gauge charge in $\text{U}_s(1)$ gauge group will also change sign under Z_2^T due to its nature of pseudoscalar. As bosons in a realistic material system carry integer spins, we start with spin-1 boson system in three dimensions with $\text{U}_s(1)\times\text{Z}_2^T$. For this simplest case, we can derive all quantities through the same way as in $\text{U}_c(1)\times\text{Z}_2^T$. In the following, we shall derive the quantities ($\theta^s, \sigma^s, \bar{\sigma}^s$, spin Witten effect, etc.) through a different way which is helpful for higher integer spin.

Let us embed $\text{U}_s(1)$ into $\text{SU}(2)$ full spin-rotation symmetry. Then, we shall only focus on the Lie group representations labeled by integer spin s , which forms a complete set of irreducible representations of $\text{SO}(3)$ spin-rotation symmetry group. It turns out that the physical consideration, i.e., ‘‘large gauge invariance,’’ is sufficient to obtain the results.

$\text{SU}(2)$ SPT on Σ^2 admit nonchiral edge states, but the symmetry group acts chirally. This edge profile is stable under symmetry-allowed perturbation. For the edge of $\text{SU}(2)$ SPT, the projective Kac-Moody algebra description is $\int dx \frac{2\pi}{3}(v_L : \mathbf{J}_L \cdot \mathbf{J}_L : + v_R : \mathbf{J}_R \cdot \mathbf{J}_R :)$ where v_L/v_R is velocity, and, the left mover \mathbf{J}_L is $\text{SU}(2)$ doublet and the right mover \mathbf{J}_R is $\text{SU}(2)$ singlet.^{33,45} After gauging $\text{SU}(2)$, the resultant gauge theory after integrating out all matter field fluctuations must be a generic Chern-Simons form in (2+1)D space-time, i.e. (coupling constant $g = 1$),

$$\mathcal{L}_{s,\Sigma^2} = \frac{k}{4\pi}\epsilon^{\mu\nu\lambda}\text{Tr}\left[A_\mu\partial_\nu A_\lambda + \frac{2}{3}A_\mu A_\nu A_\lambda\right], \quad (25)$$

where k is quantized at integer since it is the winding number labeling the nontrivial homotopic mapping $\pi_3[\text{SU}(2)] = \mathbb{Z}$. A_μ is the matrix-valued gauge vector defined as $A_\mu = T^a A_\mu^a$ ($a = 1, 2, 3$), where T^a are the generators of $\text{SU}(2)$ in a given representation (labeled by total spin s), and A_μ^a are the real-number-valued gauge potentials. The field strength $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]$. The Lie algebra structure constant $f^{abc} = \epsilon^{abc}$, i.e., $[T^a, T^b] = i\epsilon^{abc}T^c$. And, most importantly, the trace $\text{Tr}[T^a T^b]$ depends on the choice of representation, i.e., the total spin s :

$$\text{Tr}[T^a T^b] = \frac{1}{3}s(s+1)(2s+1)\delta_{ab}. \quad (26)$$

Such a normalization condition as well as the structure constant leads to the fact that the generators T^a are precisely identical to spin operators S^a of spin s [$(T^1)^2 = (T^2)^2 = (T^3)^2 = \frac{1}{3}s(s+1)\mathbb{I}$ with \mathbb{I} the $(2s+1)$ -dimensional identity matrix]. As a result, the gauge field A_μ^a precisely couples to spin current of spin s along the a -spin direction in a correct unit.

In order to probe the quantum spin Hall effect of the bosonic system, it is sufficient to merely consider a $\text{U}(1)$ subgroup of $\text{SU}(2)$ in a given representation. For example, let us study the Hall current generated by A_μ^3 (which is identical to A_μ^s):

$$J_\mu^{s,\Sigma^2} \equiv \frac{\delta\mathcal{L}_{s,\Sigma^2}}{\delta A_\mu^s}. \quad (27)$$

For convenience, let us express Chern-Simons action explicitly in terms of A_μ^a :

$$\begin{aligned} \mathcal{L}_{s,\Sigma^2} &= \frac{k}{4\pi}\epsilon^{\mu\nu\lambda}\text{Tr}\left[A_\mu\partial_\nu A_\lambda + \frac{2}{3}A_\mu A_\nu A_\lambda\right] \\ &= \frac{k}{4\pi}\epsilon^{\mu\nu\lambda}\text{Tr}\left[A_\mu^a\partial_\nu A_\lambda^b T^a T^b + \frac{2}{3}A_\mu A_\nu A_\lambda\right] \end{aligned} \quad (28)$$

since we only consider the response action of A_μ^3 ($\equiv A_\mu^s$). Let us drop all terms containing A_μ^1, A_μ^2 :

$$\mathcal{L}_{s,\Sigma^2} = \frac{k}{4\pi}\epsilon^{\mu\nu\lambda}A_\mu^s\partial_\nu A_\lambda^s\text{Tr}[T^3 T^3]. \quad (29)$$

Therefore, the spin Hall current is given by

$$J_\mu^{s,\Sigma^2} = \frac{k}{2\pi}\text{Tr}[T^3 T^3]\epsilon^{\mu\nu\lambda}\partial_\nu A_\lambda^s. \quad (30)$$

The spin Hall conductance σ^s is readily given by

$$\begin{aligned} \sigma^s &= \frac{k}{2\pi}\frac{1}{3}s(s+1)(2s+1) \\ &= 2k\left(\frac{1}{6}s(s+1)(2s+1)\right)\frac{1}{2\pi} = 2k\frac{1}{2\pi}\sum_i q_i^2, \end{aligned} \quad (31)$$

where $q_i = s, s-1, s-2, \dots$ and $q_i > 0$. In the presence of $\sum_i q_i^2$, we find that the $\sum_i q_i^2 \frac{1}{2\pi}$ is nothing but the unit of spin Hall conductance when ‘‘spin charge q_i ’’ is generically a number other than 1. For a spin- s boson system, there are $\mathcal{N}_f = 2s+1$ flavors of bosons. Each of them contributes the same even number $2k$ in total spin Hall response. We stress that in the present approach, the even integer ‘‘ $2k$ ’’ arises naturally as a result of large gauge invariance. As a simple check, when $s = \frac{1}{2}$, $\sigma^s = \frac{k}{4\pi} = 2k \times (\frac{1}{2})^2 \frac{1}{2\pi}$. When $s = 1$, $\sigma^s = \frac{k}{\pi} = 2k \times (1)^2 \frac{1}{2\pi}$. When $s = \frac{3}{2}$, $\sigma^s = \frac{5k}{2\pi} = 2k \times [(\frac{3}{2})^2 + (\frac{1}{2})^2] \frac{1}{2\pi}$. The first two results ($s = \frac{1}{2}, 1$) were derived

by Liu and Wen⁴⁵ through the principal chiral nonlinear sigma models in which $s = \frac{1}{2}$ and 1 are discussed in SU(2) and SO(3) SPT states, respectively. Actually, for the response theory itself, it turns out the different results between SU(2) and SO(3) actually depend on the trace normalization of generators ($\text{Tr}[T^a T^b]$). Indeed, the result that σ^s of SO(3) is four times of σ^s of SU(2) originates from trace normalization ($\text{Tr}[T^a T^b] = \frac{1}{2}\delta_{ab}$, if $s = \frac{1}{2}$; $\text{Tr}[T^a T^b] = 2\delta_{ab}$, if $s = 1$). We comment that a more mathematical exposition on this $\frac{1}{4}$ factor quantization difference between SU(2) and SO(3) is found in Sec. 4 of Ref. 49 and Sec. 2 of Ref. 64. Here, we use a rather physical language instead, comparing to the more mathematically oriented formalism in Refs. 49 and 64.

Let us focus on integer spin s . In the $s = 1$ case, all results can also be derived if the method in the derivation of σ^c is adopted. In a generic s case, the even integer $2k$ directly leads to the absence of topological order in Σ^2 in a sense that

(i) all spin excitations carry integer-valued spin- S^z quantum number, no fractional quantum number;

(ii) all spin excitations are bosonic with the total Aharonov-Bohm phase $\Phi_{\text{AB}} = 2\pi k$ accumulated by spatially exchanging two vortex cores. The spin angular momentum does not contribute fermionic sign since s is integer.

Next, let us consider a SPT with $\text{SU}(2) \times Z_2^T$ in Σ^3 in the integer spin- s representation and then only consider the $U_s(1)$ subgroup through which we will obtain the surface Chern-Simons term $\mathcal{L}_{s, \partial\Sigma^3}$. With Z_2^T symmetry, the bulk should be a non-Abelian theta term. Generically, the bulk non-Abelian theta action is $\Theta \times \mathcal{P}$ where \mathcal{P} is the integer-valued Pontryagin index [the bar in the symbol $\bar{\Theta}$ is to distinguish the $\bar{\Theta}$ and Θ matrices defined in Eq. (1)]

$$S = \bar{\Theta} \times \mathcal{P} = \int d^4x \bar{\Theta} \frac{1}{16\pi^2} \text{Tr}[F_{\mu\nu} \tilde{F}_{\mu\nu}], \quad (32)$$

where the dual tensor $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\lambda\rho} F_{\lambda\rho}$. $F_{\mu\nu}$ can be further written as $F_{\mu\nu} = T^a F_{\mu\nu}^a$, $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + \epsilon^{abc} A_\mu^b A_\nu^c$. Therefore, the Lagrangian form of Eq. (32) is reformulated to

$$\mathcal{L}_{s, \Sigma^3} = \frac{2\bar{\Theta}}{16\pi^2} \sum_i q_i^2 F_{\mu\nu}^a \tilde{F}_{\mu\nu}^a. \quad (33)$$

The dual tensor for each component is defined as $\tilde{F}_{\mu\nu}^a = \frac{1}{2}\epsilon^{\mu\nu\lambda\rho} F_{\lambda\rho}^a$. In Eq. (33), despite that the summation over the three spin directions is independent, the three spin directions actually couple to each other due to the last *nonlinear* term in $F_{\mu\nu}$. As a matter of fact, the $\bar{\Theta}$ term is a total-derivative term which does not generate topological bulk response if A_μ is smooth without singularities. Let us consider a material with a surface. Suppose that the surface breaks Z_2^T explicitly or spontaneously, the surface has spin-rotational symmetry, and the response theory is again a non-Abelian Chern-Simons theory. Mathematically, starting from Eq. (32), we find its surface action

$$S = \frac{2\bar{\Theta}}{16\pi^2} \epsilon^{\mu\nu\lambda} \text{Tr} \left[A_\mu \partial_\nu A_\lambda + \frac{2}{3} A_\mu A_\nu A_\lambda \right]. \quad (34)$$

Completion of trace operation leads to

$$S = \frac{2\bar{\Theta}}{8\pi^2} \sum_i q_i^2 \epsilon^{\mu\nu\lambda} A_\mu^a \partial_\nu A_\lambda^a + \dots, \quad (35)$$

where “...” are nonlinear response terms. Let us only consider the $A_\mu^3 \equiv A_\mu^s$ response theory, so that $2\bar{\Theta} \equiv \theta_s$ leading to the surface spin Hall conductance

$$\tilde{\sigma}^s = \frac{\theta_s}{2\pi} \sum_i q_i^2 \frac{1}{2\pi}. \quad (36)$$

To derive the periodicity and minimal value of θ_s for the nontrivial phase with $U_s(1) \times Z_2^T$ in Σ^3 , we apply the same strategy in the derivation of θ_c . The minimal theta value should be one half of its periodicity (say, P), and the symmetry group for the bulk is $U_s(1) \times Z_2^T$. Physically, the periodicity can be understood as trivially depositing arbitrary copies of Σ^2 Hall systems onto the surface. As such, a P shift in θ_s leads to an additional term in the surface spin Hall conductance formula (36):

$$\tilde{\sigma}^{s'} - \tilde{\sigma}^s = \frac{P}{4\pi^2} \sum_i q_i^2, \quad (37)$$

which is contributed by deposited Σ^2 layers which are described by Eq. (31). A minimal choice is $\frac{P}{4\pi^2} = 2 \times \frac{1}{2\pi}$, so that $P = 4\pi$. And, the minimal choice of θ_s is $\frac{P}{2} = 2\pi$, i.e.,

$$\theta_s = 2\pi + 4\pi k, \quad (38)$$

where the integer k is the same k defined in Eq. (31). Substituting Eq. (38) into (36) leads to

$$\tilde{\sigma}^s = (1 + 2k) \frac{1}{2\pi} \sum_i q_i^2. \quad (39)$$

2. Spin Witten effect in Σ^3

In order to derive the so-called *spin Witten effect* in Table II for generic spin s , by noting that $2\bar{\Theta} = \theta_s$, let us drop all terms irrelevant to A_μ^z (i.e., A_μ^s) in Eq. (33), resulting in

$$\mathcal{L}_s = \frac{\theta_s}{8\pi^2} \sum_i q_i^2 \partial_\mu A_\nu^s \partial_\lambda A_\rho^s \epsilon^{\mu\nu\lambda\rho}. \quad (40)$$

Once $s = 1$, we obtain $\sum_i q_i^2 = 1$ and above Lagrangian \mathcal{L}_s is back to the original version of the bulk θ_s term defined in Eq. (3). Here, we would like to consider a generic s which leads to the following response equation for A_μ^s in Σ^3 :

$$J_\mu^s \equiv \frac{\delta \mathcal{L}_s}{\delta A_\mu^s} = \frac{\theta_s}{4\pi^2} \sum_i q_i^2 \epsilon^{\mu\nu\lambda\rho} \partial_\nu \partial_\lambda A_\rho^s, \quad (41)$$

where J_μ^s is the (3+1)D response spin current. The zero component J_0^s denotes the response charge density probed by external spin gauge field A_μ^s :

$$J_0^s = \frac{\theta_s}{4\pi^2} \sum_i q_i^2 \nabla \cdot \mathbf{B}^s, \quad (42)$$

where \mathbf{B}^s is the spin-magnetic-field variable. If the gauge field \mathbf{A}^s is smooth everywhere, $\nabla \cdot \mathbf{B}^s = 0$ due to the absence of magnetic charge. However, if singular configuration is

allowed, the divergence may admit singularities in the bulk and its total contribution in the bulk is quantized due to the Dirac quantization condition (or the more general Schwinger-Zwanziger quantization condition). Let us consider one magnetic monopole (of A_μ^s gauge group) located at the origin of the three-dimensional space:

$$\int d^3x \nabla \cdot \mathbf{B}^s = 2\pi N_m^s, \quad (43)$$

where $N_m^s \in \mathbb{Z}$ is an integer-valued ‘‘magnetic charge’’ of the A_μ^s gauge group. Therefore, the corresponding response total spin is

$$N^s = \int d^3x J_0^s = \frac{\theta_s}{2\pi} \sum_i q_i^2 N_m^s, \quad (44)$$

which indicates that a nonzero theta term supports a ‘‘polarization spin cloud’’ in the presence of magnetic monopole of A_μ^s gauge group. A monopole of A_μ^s gauge group can also trivially attach integer number ($n_i^s \in \mathbb{Z}$) of bosons with $S^z = q_i$ in the bulk (as $q_i > 0$, a negative n_i^s implies that $|n_i^s|$ bosons in the spin state ‘‘ $S^z = -q_i$ ’’). Therefore, the whole formula of the so-called *spin Witten effect* can be expressed as

$$N^s = \sum_i n_i^s q_i + \frac{\theta_s}{2\pi} N_m^s \sum_i q_i^2. \quad (45)$$

Substituting (38) into (45) leads to

$$N^s = \sum_i n_i^s q_i + N_m^s \sum_i q_i^2, \quad (46)$$

where $k = 0$ is selected for simplicity. Different choices of k actually correspond to the *same phase*.

3. Anomalous K_G matrix on $\partial\Sigma^3$

Similar to Sec II B3, let us use the top-down approach explicitly working out the external field theory on $\partial\Sigma^3$ of $U_s(1) \times Z_2^T$ global symmetry with Z_2^T symmetry broken, and the external field theory on Σ^2 of $U_s(1)$ global symmetry. Here, we save detailed derivations to Appendix A and list key results directly. We first study on Σ^2 with $U_s(1)$ global symmetry: what we start with is the intrinsic SPT's $K_S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and gauging the $U(1)$ global symmetry current coupling to A^s , we obtain $K_G = 2p$. The only difference from Sec. II B3 is that the spin gauge field contributes a factor of $\sum_i q_i^2$, which simply sums over all the spin contribution $q_i = s, s-1, s-2, \dots$ and $q_i > 0$,

$$\mathcal{L}_{\text{SPT+Gauge}}(A^s) = \frac{2p}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^s \partial_\nu A_\rho^s \sum_i q_i^2 \quad (47)$$

with $p \in \mathbb{Z}$ labeling the \mathbb{Z} class of the cohomology group $\mathcal{H}^3[U(1), U(1)] = \mathbb{Z}$. The Hall conductance as the response of this $\mathcal{L}_{\text{SPT+Gauge}}(A^c)$ is

$$\sigma^s = 2p \frac{1}{2\pi} \sum_i q_i^2. \quad (48)$$

This result matches exactly as Eq. (31). On the other hand, for the anomalous K_G matrix in $\partial\Sigma^3$, so far the intrinsic topological field theory of Σ^3 in (3+1)D is not yet known to be completed, we simply adopt the result from the previous

section to state the effective $K_{G, \partial\Sigma^3} = 2p + \theta_s/2\pi$, which means

$$\mathcal{L}_{\text{SPT+Gauge}}(A^s) = \frac{2p + \theta_s/2\pi}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^s \partial_\nu A_\rho^s \sum_i q_i^2 \quad (49)$$

with Hall conductance

$$\tilde{\sigma}^s = (2p + \theta_s/2\pi) \frac{1}{2\pi} \sum_i q_i^2. \quad (50)$$

Therefore, we have Eqs. (48) and (50) written in a consistent manner as Eqs. (31) and (39). $K_{G, \partial\Sigma^3}$ is the anomalous K_G matrix of the present surface state.

D. $U_c(1) \times [U_s(1) \times Z_2]$ in Σ^3

1. Quantum charge-spin/spin-charge Hall effect on Z_2 broken $\partial\Sigma^3$ and Z_2 broken Σ^2

In the above discussions, we have studied the bulk θ_c and θ_s terms, each of which is constructed by one kind of gauge field. In the following, we shall consider the bulk θ_0 term \mathcal{L}_0 in which A_μ^s and A_μ^c are both involved. The *minimal* symmetry requirement of this term is $U_c(1) \times [U_s(1) \times Z_2]$, where Z_2 can be viewed as a π rotation about spin S^y . The additional Z_2 is required by the following observation. Under Z_2 operation, $A_\mu^s \rightarrow -A_\mu^s$, and thus $\mathbf{E}^s \rightarrow -\mathbf{E}^s, \mathbf{B}^s \rightarrow -\mathbf{B}^s, \mathbf{E}^c \rightarrow \mathbf{E}^c, \mathbf{B}^c \rightarrow \mathbf{B}^c$, such that $\mathcal{L}_0 = \frac{\theta_0}{4\pi^2} \partial_\mu A_\nu^c \partial_\lambda A_\rho^s \epsilon^{\mu\nu\lambda\rho} = \frac{\theta_0}{4\pi^2} (\mathbf{E}^c \cdot \mathbf{B}^s + \mathbf{B}^c \cdot \mathbf{E}^s) \rightarrow -\mathcal{L}_0$. However, if a periodicity in θ_0 is allowed, $-\theta_0$ will be shifted back to θ_0 leading to the invariance of \mathcal{L}_0 under Z_2 spin rotation. Due to the existence of the periodicity, we expect that Z_2 symmetry plays a similar role in determining quantization conditions of θ_0 and related Hall effects, in comparison with the role of Z_2^T in the bulk θ_c and θ_s terms. Note that a theta term with only one kind of gauge field (such as θ_c and θ_s terms) in a three-dimensional insulator can be formally viewed as an expectation value of divergence of chiral current in the context of the Adler-Bardeen-Bell-Jackiw anomaly^{65,66} in a chiral gapless system.

For simplicity, we restrict our attention on a spin-1 and charge-1 boson system in Σ^3 with symmetry $U_c(1) \times [U_s(1) \times Z_2]$. Since \mathcal{L}_0 is a total derivative term, the bulk response is trivial unless at least one of the gauge field configurations admits singularities. Let us first consider a Z_2 -broken surface $\partial\Sigma^3$. The Lagrangian \mathcal{L}_0 can be written as a *surface mutual Chern-Simons* term

$$\mathcal{L}_{0, \partial\Sigma^3} = \frac{\theta_0}{4\pi^2} A_\mu^s \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}. \quad (51)$$

Recently, a mutual Chern-Simons term with dynamical or non-dynamical gauge fields has been studied in other contexts.⁶⁷⁻⁷² The surface mutual Chern-Simons term (51) leads to the following two different surface response currents:

$$J_\mu^{s, \partial\Sigma^3} \equiv \frac{\delta \mathcal{L}_{0, \partial\Sigma^3}}{\delta A_\mu^s} = \frac{\theta_0}{4\pi^2} \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}, \quad (52)$$

$$J_\mu^{c, \partial\Sigma^3} \equiv \frac{\delta \mathcal{L}_{0, \partial\Sigma^3}}{\delta A_\mu^c} = \frac{\theta_0}{4\pi^2} \partial_\nu A_\lambda^s \epsilon^{\mu\nu\lambda}. \quad (53)$$

Here, the surface spin current $J_\mu^{s, \partial\Sigma^3}$ is induced by applying external electromagnetic gauge field A_μ^c rendering the quantum spin-charge Hall effect, while the surface charge current $J_\mu^{c, \partial\Sigma^3}$

is induced by applying external spin gauge field A_μ^s rendering the quantum charge-spin Hall effect. The corresponding surface spin-charge/charge-spin Hall conductance formulas are

$$\tilde{\sigma}^{sc} = \tilde{\sigma}^{cs} = \frac{\theta_0}{4\pi^2}. \quad (54)$$

We note that the two quantum Hall effects share the same Hall conductance and the same unit “ e .” To understand the quantization of the surface spin-charge and charge-spin Hall conductance $\tilde{\sigma}^{sc}, \tilde{\sigma}^{cs}$, we need to first understand the quantization on the θ_0 angle and σ^{sc}, σ^{cs} in strictly 2D systems (i.e., Σ^2). The Σ^2 system will be used to determine the periodicity and the minimal value of θ_0 by depositing it onto the Z_2 -broken surface $\partial\Sigma^3$, so that the minimal symmetry requirement on Σ^2 is $U_c(1) \times U_s(1)$. Let us write a generic topological response theory in Σ^2 :

$$\mathcal{L}_{0,\Sigma^2} = \frac{1}{2} (A_\mu^c, A_\mu^s) \begin{pmatrix} \sigma^c & \sigma^{cs} \\ \sigma^{sc} & \sigma^s \end{pmatrix} \partial_\nu \begin{pmatrix} A_\lambda^c \\ A_\lambda^s \end{pmatrix} \epsilon^{\mu\nu\lambda}, \quad (55)$$

which leads to three independent Chern-Simons terms. Therefore, in general, we need three quantities $\sigma^c, \sigma^s, \sigma^{cs}$ ($\sigma^{cs} = \sigma^{sc}$) to label the quantum Hall states in Σ^2 . But, the state we are considering will be applied to be deposited onto the surface $\partial\Sigma^3$ where only the mutual Chern-Simons term exists as shown in Eq. (51). In other words, we shall consider the Σ^2 system with $\sigma^s = 0$ and $\sigma^c = 0$, rendering a mutual Chern-Simons term for Σ^2 :

$$\mathcal{L}_{0,\Sigma^2} = \sigma^{sc} A_\mu^s \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}. \quad (56)$$

Thus, the response currents are

$$J_\mu^{s,\Sigma^2} \equiv \frac{\delta \mathcal{L}_{0,\Sigma^2}}{\delta A_\mu^s} = \sigma^{sc} \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda}, \quad (57)$$

$$J_\mu^{c,\Sigma^2} \equiv \frac{\delta \mathcal{L}_{0,\Sigma^2}}{\delta A_\mu^c} = \sigma^{cs} \partial_\nu A_\lambda^s \epsilon^{\mu\nu\lambda}. \quad (58)$$

Upon adiabatically piercing Σ^2 by 2π magnetic flux ($\Phi^c = \int d^2x \nabla \times \mathbf{A}^c = 2\pi$), the total spin accumulation $\int d^2x J_0^{s,\Sigma^2} = \sigma^{sc} \int d^2x \nabla \times \mathbf{A}^c = 2\pi \sigma^{sc}$. In SPT states where topological order is trivial by definition, this pumped spin in the center of the vortex core must be quantized at integer since the fundamental spin is carried by spin-1 bosons (for example, a spin- $\frac{1}{2}$ quasiparticle is not allowed), such that $2\pi \sigma^{sc} \in \mathbb{Z}$. On the other hand, let us consider the condition under which all the quasiparticles are bosonic in order to forbid topological order. To achieve this goal, one can spatially exchange two vortex cores of 2π fluxes each of which traps $2\pi \sigma^{sc}$ quasiparticles with pure spins and neutral charge. The quasiparticles in the first vortex core will perceive a π phase as half a magnetic flux of the second vortex core, and vice versa. Unlike the Chern-Simons theory, the total Aharonov-Bohm phase “ Φ_{AB} ” in the mutual Chern-Simons theory is the totally accumulated quantum phases $\Phi_{AB} = (2\pi \sigma^{sc} \times \pi + 2\pi \sigma^{sc} \times \pi) = 4\pi^2 \sigma^{sc}$. In order to forbid nonbosonic statistics, $\Phi_{AB}/2\pi \in \mathbb{Z}$. Overall, combining the “absence of fractional spin” and “absence of nonbosonic statistics” leads to, still, $2\pi \sigma^{sc} \in \mathbb{Z}$, i.e.,

$$\sigma^{sc} = \sigma^{cs} = k \frac{1}{2\pi}, \quad (59)$$

where $k \in \mathbb{Z}$ and the unit is the fundamental electric charge e .

After this preparation, let us move on to the θ_0 angle quantization and its periodicity P . As mentioned, the periodicity can be understood as trivially depositing arbitrary copies of Σ^2 Hall systems onto the surface $\partial\Sigma^3$. As such, a P shift in θ_0 leads to an additional term in the surface spin-charge/charge-spin Hall conductance formula (54):

$$\tilde{\sigma}^{sc'} - \tilde{\sigma}^{sc} = \frac{P}{4\pi^2}, \quad (60)$$

which is contributed by deposited Σ^2 layers which are described by Eq. (59). A minimal choice is $\frac{P}{4\pi^2} = 1 \times \frac{1}{2\pi}$, so that $P = 2\pi$. And, the minimal choice of θ_0 is $\frac{P}{2} = \pi$, i.e.,

$$\theta_0 = \pi + 2\pi k, \quad (61)$$

where the integer k is the same k defined in Eq. (59).

Substituting Eq. (61) into (54) leads to

$$\tilde{\sigma}^{sc} = \tilde{\sigma}^{cs} = \left(\frac{1}{2} + k \right) \frac{1}{2\pi}. \quad (62)$$

The most anomalous phenomenon in the surface charge Hall effect is that the $\tilde{\sigma}^{sc}$ and $\tilde{\sigma}^{cs}$ admit a $\frac{1}{4\pi}$ value which can not be realized in Σ^2 where σ^{sc} and σ^{cs} are always integer copies of $1/2\pi$.

2. Model construction on Σ^2

The quantum charge-spin/spin-charge Hall effects can be modeled as a similar two-component boson model proposed by Senthil and Levin,⁵⁰ but with slight modification as follows. The first-component bosons are charge neutral but carry spin-1, while the second-component bosons are spinless but carry electric charge-1. Then, an external “spin-magnetic field” \mathbf{B}^s and an external real “magnetic field” \mathbf{B}^c are applied to the first- and second-component bosons, respectively, each of which forms a bosonic Landau level with filling $\nu = 1$. It implies that the gauge fields \mathbf{A}^s and \mathbf{A}^c play the roles of \mathbf{A}_1 and \mathbf{A}_2 defined in the Senthil-Levin paper, respectively.

Therefore, by using the flux-attachment Chern-Simons theory, one can realize a many-body state which has quantum spin-charge/charge-spin Hall effects with $\sigma^{sc} = \sigma^{cs} = \frac{1}{2\pi}$.

3. Mutual Witten effect in Σ^3

In order to derive the so-called *mutual Witten effect* in Table II, let us write the response equation in the bulk Σ^3 :

$$J_\mu^s \equiv \frac{\delta \mathcal{L}_0}{\delta A_\mu^s} = \frac{\theta_0}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\nu \partial_\lambda A_\rho^c, \quad (63)$$

$$J_\mu^c \equiv \frac{\delta \mathcal{L}_0}{\delta A_\mu^c} = \frac{\theta_0}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\nu \partial_\lambda A_\rho^s, \quad (64)$$

where J_μ^s and J_μ^c are (3+1)D response spin and charge currents, respectively. The zero components J_0^s and J_0^c denote the response spin and charge density probed by external spin

gauge field A_μ^s and external electromagnetic field A_μ^c :

$$J_0^s = \frac{\theta_0}{4\pi^2} \nabla \cdot \mathbf{B}^c, \quad (65)$$

$$J_0^c = \frac{\theta_0}{4\pi^2} \nabla \cdot \mathbf{B}^s. \quad (66)$$

If the gauge fields A^c and A^s are smooth everywhere, $\nabla \cdot \mathbf{B}^c = 0$, $\nabla \cdot \mathbf{B}^s = 0$, due to absence of magnetic charge and spin-magnetic charge. However, if singular configuration is allowed, the divergences may admit singularities in the bulk and the total contribution in the bulk is quantized due to the Dirac quantization condition (or the more general Schwinger-Zwanziger quantization condition). For example, let us consider one magnetic monopole (of A_μ^c gauge group) located at the origin of the three-dimensional space. $\int d^3x \nabla \cdot \mathbf{B}^c = 2\pi N_m^c$ where $N_m^c \in \mathbb{Z}$ is an integer-valued ‘‘magnetic charge.’’ Therefore, the corresponding response total spin $N^s = \int d^3x J_0^s = \frac{\theta_0}{2\pi} N_m^c$, which indicates that a nonzero theta term supports a ‘‘polarization spin cloud’’ in the presence of magnetic monopole of the A_μ^c gauge group. A monopole of the A_μ^c gauge group can also trivially attach integer numbers ($n_+^s, n_-^s \in \mathbb{Z}$) of bosons with $S^z = 1, -1$ in the bulk, respectively. Therefore, the whole formula of the so-called *mutual Witten effect* can be expressed as

$$N^s = n_+^s - n_-^s + \frac{\theta_0}{2\pi} N_m^c. \quad (67)$$

Substituting (61) into (67) leads to

$$N^s = n_+^s - n_-^s + \frac{1}{2} N_m^c, \quad (68)$$

where $k = 0$ is selected for simplicity. Likewise, we can place a magnetic monopole of the A_μ^s gauge group. Then, a *polarization charge cloud* arises. The corresponding mutual Witten effect is

$$N^c = n^c + \frac{\theta_0}{2\pi} N_m^s, \quad (69)$$

where n^c is the number of bosons trivially attached to the magnetic charge of the A_μ^s gauge group, i.e., spin-magnetic charge. Substituting (61) into (69) leads to

$$N^c = n^c + \frac{1}{2} N_m^s. \quad (70)$$

In Fig. 1, we illustrate the mutual Witten effect, motivated by Ref. 73. The ferromagnetic (FM) thin film *uniformly* breaks Z_2 spin rotation symmetry (π rotation about spin S^y) of the surface of $\theta_0 = \pi$ nontrivial bulk. We assume that the width of the FM film is sufficiently small, i.e., $z_3 \gtrsim z_2$. In Fig. 1(a), a spinful but charge-neutral impurity (the red ball) is located at $(x, y, z) = (0, 0, z_3)$ near the surface ($z = z_2$). To solve the static electromagnetic problem of both U(1) gauge groups in the region $(z_2, z_4) \cup (z_4, \infty)$, one can use the trick of ‘‘image charge.’’⁷⁴ One can introduce an image ‘‘spin’’ (with spin N^s in unit of \hbar) located at $(0, 0, z_1)$ with $z_2 = (z_1 + z_3)/2$ which effectively takes the boundary condition on the $z = z_2$ plane into consideration. At the same position, the *mutual Witten effect* admitted by the nontrivial bulk ($z < z_2$) induces an image magnetic monopole (with magnetic charge $N_m^c = 2N^s$ in unit of h/e , denoted by the blue ball) of the A_μ^c gauge group. This image monopole determines the magnetic field \mathbf{B}^c inside the region (z_3, z_4) (the solid blue lines with arrows). Since these magnetic lines are formed by the electromagnetic field A_μ^c , the




FIG. 1. (Color online) Illustration of the experimental setup to realize the mutual Witten effect. The x , y , and z axes form the three spatial directions. In both (a) and (b), a ferromagnetic thin film (FM) is located between a trivial $U_c(1) \times [U_s(1) \times Z_2]$ state with $\theta_0 = 0$ and the nontrivial state with $\theta_0 = \pi$. The width of the film is sufficiently small. In (a), the red and blue balls stand for a spin impurity and an image magnetic monopole of electromagnetic A_μ^c gauge field, respectively. The solid blue lines in the region $(z_2, z_4) \cup (z_4, \infty)$ represent the magnetic field \mathbf{B}^c induced by the spin impurity. In (b), the blue and red balls stand for an electric charge impurity and an image magnetic monopole of A_μ^s gauge field (i.e., the ‘‘spin gauge field’’), respectively. The solid red lines in the region (z_2, z_4) represent the magnetic field \mathbf{B}^s induced by the electric charge impurity (see text).

magnetic lines can also penetrate the $z = z_4$ plane, i.e., the top surface of the $\theta_0 = 0$ bulk, and flows into the vacuum (z_4, ∞) where both dielectric constant ϵ_0 and permeability μ_0 are nonzero forming nonzero static charge-charge correlation. The radial magnetic distribution provides a Lorentz force acting on electrically charged currents. The latter can be excited if temperature is nonzero inside the trivial gapped bulk. It should be noted that the roles of other boundaries (such as the $z = z_4$ plane) are ignored since they are irrelevant to the formation of the mutual Witten effect discussed here.

There is one remaining issue to be stressed in the following. Different from Ref. 73 where only electromagnetic field A_μ^c is considered, we must add a trivial bulk above the FM film. The reason is following. The vacuum medium is really ‘‘empty’’ for the A_μ^s gauge field in a sense that the vacuum has no background dynamics admitting communication between two spinful particles (namely, particle-1 with spin q_1 and particle-2 with spin q_2), which is in contrast to A_μ^c that has background $\epsilon_0 < \infty$ and $\mu_0 > 0$. In other words, in the vacuum the two spinful particles can not interact with each other via the A_μ^s gauge field. Classically, it indicates that the ‘‘electromagnetic force’’ felt by particle-1 is vanishing: $\mathbf{F}_{12} \equiv q_1 \mathbf{E}^s + q_1 \mathbf{v}_1 \times \mathbf{B}^s = 0$, where \mathbf{v}_1 is the velocity vector of particle-1, and \mathbf{E}^s and \mathbf{B}^s are spin-electric field and spin-magnetic field formed by particle-2 current. On the other hand, the trivial bulk on the top of FM film provides a nontrivial dynamical background where spin-spin correlation is well formed and thus two spinful particles can talk to each other via the A_μ^s gauge field.

Likewise, in Fig. 1(b), we may place an electric charge impurity (the blue ball) which finally induces a magnetic monopole (the red ball) of the A_μ^s gauge group. The monopole determines the magnetic field \mathbf{B}^s distribution in the trivial state. As explained above, \mathbf{B}^s magnetic lines (the solid red line with arrows) are confined inside the trivial bulk and can not penetrate into the vacuum, in contrast to \mathbf{B}^c . The radial

magnetic distribution provides a Lorentz force acting on spin currents. The latter can be excited if temperature is nonzero inside the trivial gapped bulk.

4. Anomalous K_G matrix on $\partial\Sigma^3$

Similar to Secs. II B3 and II C3, let us use the top-down approach explicitly working out the external field theory on $\partial\Sigma^3$ of $U_c(1) \times [U_s(1) \times Z_2]$ global symmetry with Z_2^T symmetry broken, and the external field theory on Σ^2 with $U_c(1) \times U_s(1)$ global symmetry. Here, we save detailed derivations to Appendix A and list key results directly. We first study on Σ^2 with $U_c(1) \times U_s(1)$ global symmetry; what we start with is the intrinsic SPT's $K_S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and gauging the $U(1) \times U(1)$ global symmetry current coupling to A^s , we obtain $K_G = \begin{pmatrix} 2p_1 & p_{12} \\ p_{12} & 2p_2 \end{pmatrix}$ in the gauge charge sectors of $U(1) \times U(1)$. In the case of a charge-1 and spin-1 bosonic system, we have spin sum $\sum q_i^2 = 1$:

$$\mathcal{L}_{\text{SPT+Gauge}}(A^c, A^s) = \frac{1}{4\pi} \epsilon^{\mu\nu\lambda} (A_\mu^c, A_\mu^s) \begin{pmatrix} 2p_1 & p_{12} \\ p_{12} & 2p_2 \end{pmatrix} \partial_\nu \begin{pmatrix} A_\lambda^c \\ A_\lambda^s \end{pmatrix} \quad (71)$$

with $p_1, p_2, p_{12} \in \mathbb{Z}$ labeling the class of the cohomology group $\mathcal{H}^3[U(1) \times U(1), U(1)] = \mathbb{Z}^3$. We comment that this result is more general than Eq. (56) because there $U_c(1) \times [U_s(1) \times Z_2]$ symmetry restricts $\sigma^c = \sigma^s = 0$ so $p_1 = 0$ and $p_2 = 0$. However, the Hall conductance as the response of this $\mathcal{L}_{\text{SPT+Gauge}}(A^c)$ is the same:

$$\sigma^{sc} = \sigma^{cs} = p_{12} \frac{1}{2\pi}. \quad (72)$$

On the other hand, as the intrinsic topological field theory of Σ^3 in $(3+1)\text{D}$ is not yet known to be completed, for the anomalous K_G matrix in $\partial\Sigma^3$ we simply adopt the result from the previous section to modify the effective K_G matrix on $\partial\Sigma^3$ as

$$K_{G, \partial\Sigma^3} = \begin{pmatrix} 2p_1 & p_{12} + \frac{\theta_0}{2\pi} \\ p_{12} + \frac{\theta_0}{2\pi} & 2p_2 \end{pmatrix} \quad (73)$$

in Eq. (71). With Hall conductance,

$$\tilde{\sigma}^{sc} = \tilde{\sigma}^{cs} = (p_{12} + \theta_0/2\pi) \frac{1}{2\pi}. \quad (74)$$

Therefore, we have Eqs. (72) and (74) written in a consistent manner as Eqs. (59) and (62). The top-down approach here shows the consistency to Sec. II D1.

III. DISCRETE Z_N CHARGE SYMMETRY AND Z_N SPIN SYMMETRY

A. Dynamical gauge theory: A general discussion

In Sec. II, we discussed the response theory of SPT states with charge/spin continuous symmetry. In the examples we considered, we found exotic Σ^3 bulk response phenomena which are descendants of the original Witten effect although the bulk is fully gapped insulators. We also found exotic $\partial\Sigma^3$ surface Hall effects which contain many variants, and all of them can not be realized in a strictly two-dimensional SPT

state with the same symmetry as the $\partial\Sigma^3$. The statement is that, given a SPT with symmetry G in Σ^3 , its surface ($\partial\Sigma^3$) with symmetry G' (as a subgroup of G) can not be realized in a SPT with the same symmetry G' but defined in Σ^2 .

In this section, we will consider discrete charge/spin symmetry. Recall that, in a BCS superconductor, the charge symmetry reduces to Z_2 from $U(1)$ due to Cooper pairs. As a result, the magnetic flux inside a type-II BCS superconductor is quantized to π . The charge response current in terms of Ohm's equation is screened in a sense that in the linear response regime the external electromagnetic field can not be adiabatically turned on from zero due to photon mass. Likewise, in a SPT state with discrete charge/spin symmetry group which can be achieved by charge- N condensate/spin- N condensate, the response phenomenon in the linear regime is always dominated by the Meissner effect.

Instead of the response approach utilized in Sec. II, in the following, we will study these states by gauging the charge/spin symmetry which results in the *dynamical gauge theory* description of SPT states. We stress that all field variables in the dynamical gauge theory description are now dynamical and appear in the path-integral measure. Most importantly, this dynamical gauge theory is *not* the low-energy theory of the SPT state but a new window/tool to diagnose the SPT states. By studying the dynamical gauge theory, we will find the anomalous surface ($\partial\Sigma^3$), in a sense that the gauged theory on the surface with symmetry G' is different from the gauged theory of a two-dimensional SPT with the same G' . And, the dynamical gauge theory bridges SPT to a topological ordered state, namely, "symmetry-enriched topological phase" (SET) in which the fingerprint of SPT is hidden.

1. Gauging SPT to SET

The more precise statement of gauging the subgroup G' of the full global symmetry group G is that we convert partially the global symmetry group to the gauge symmetry group, with a leftover global symmetry (Z_2 or Z_2^T). In other words, what we really do is convert SPT states to the symmetry-enriched topological (SET) states, i.e., topologically ordered states enriched with a global symmetry. Let us illuminate this relation as follows.

Basics of SET. We first set up the SET picture in Refs. 37, 38, 55, and 56. Following pioneer works,^{38,55,56} let us define the global symmetry group as G_s , the gauge symmetry group as G_g . This SET picture in Ref. 38 considers the exact sequence in SET,

$$1 \rightarrow G_g \rightarrow \text{PSG} \rightarrow G_s \rightarrow 1, \quad (75)$$

which says that projective symmetry group (PSG) is an extension of the global symmetry group G_s by the gauge symmetry group G_g . From the exact sequence, G_g is a normal subgroup. The global symmetry group G_s is regarded as the quotient group $G_s = \text{PSG}/G_g$, from the full symmetry group PSG mod out a gauge symmetry group G_g .

Promote SPT to SET. We now clarify that in all of our examples, the full symmetry group has the form

$$G = G' \rtimes G'' \text{ or } G = G' \times G'', \quad (76)$$

TABLE III. The dynamical gauge theory description of spin-1 and charge-1 boson SPT systems with discrete spin symmetry and/or discrete charge symmetry. By following Sec. III and specifically Sec. III A1, we connect the concept of gauging symmetry-protected topological (SPT) states to symmetry-enriched topological (SET) states. The first column: projective symmetry group (PSG) in SET corresponds to the full symmetry group in SPT. The second column: G_g in SET corresponds to the gauged symmetry group in SPT. The third column: G_s in SET corresponds to the remaining ungauged symmetry group in SPT. Z_2 in this column is π rotation about S^y . The fourth column shows the effective dynamical gauge theory description of SET (more details in Sec. III A2), which includes the topological BF term and the Θ term $F \wedge F$. The fifth column shows the surface gapless anomalous boson theory on G_s broken $\partial\Sigma^3$ of Σ^3 bulk SET (more details in Sec. III A3) The last column is filled with the dynamical gauge theory of G_g -symmetry SPT (but with no G_s symmetry) on intrinsic Σ^2 surface; by gauging the symmetry G_g , it compares with the fifth column $\partial\Sigma^3$. The central messages are as follows: (1) After gauging a normal subgroup of the symmetry of 3D SPT, we obtain a SET state described by a dynamical gauge theory with a remaining global symmetry. (2) The surface (with Z_2^T or Z_2 broken) of 3D SET is described by a gapless boson matter field with quantum anomaly. (3) After fully gauging a SPT on Σ^2 , we obtained a dynamical gauged field theory described by K_G -matrix Chern-Simons theory. (4) The resultant states on $\partial\Sigma^3$ and Σ^2 are different although both are two-dimensional space manifolds. Be aware that B^c and B^s are external antisymmetric 2-form $B_{\mu\nu}^c$ and $B_{\mu\nu}^s$; we should *not* misunderstand its meaning mixed with magnetic field $\mathbf{B}^c, \mathbf{B}^s$.

Full symmetry group PSG	Gauge symmetry group G_g	Global symmetry group G_s	3D bulk (Σ^3) dynamical gauge theory	Surface ($\partial\Sigma^3$) boson theory with anomaly	2D plane (Σ^2) dynamical gauge theory with K_G matrix
$Z_N \times Z_2^T$	Z_N	Z_2^T	$\frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^c \partial_\lambda A_\rho^c + \frac{\theta_c}{8\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\mu A_\nu^c \partial_\lambda A_\rho^c$	Z_2^T broken $\partial\Sigma^3$: $\frac{N}{2\pi} \partial_0 \phi^c \epsilon^{ij} \partial_i \lambda_j^c$	Z_2^T broken Σ^2 : $\begin{pmatrix} 2p & N \\ N & 0 \end{pmatrix}$
$Z_N \times Z_2^T$	Z_N	Z_2^T	$\frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^s \partial_\lambda A_\rho^s + \frac{\theta_s}{8\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\mu A_\nu^s \partial_\lambda A_\rho^s$	Z_2^T broken $\partial\Sigma^3$: $\frac{N}{2\pi} \partial_0 \phi^s \epsilon^{ij} \partial_i \lambda_j^s$	Z_2^T broken Σ^2 : $\begin{pmatrix} 2p & N \\ N & 0 \end{pmatrix}$
$Z_N \times [U_s(1) \times Z_2]$	$Z_N \times U_s(1)$	Z_2	$\frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^c \partial_\lambda A_\rho^c + \frac{\theta_0}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\mu A_\nu^s \partial_\lambda A_\rho^c$	Z_2 broken $\partial\Sigma^3$: $\frac{N}{2\pi} \partial_0 \phi^c \epsilon^{ij} \partial_i \lambda_j^c$	Z_2 broken Σ^2 : $\begin{pmatrix} 2p_1 & N & p_{12} & 0 \\ N & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$
$U_c(1) \times [Z_N \times Z_2]$	$U_c(1) \times Z_N$	Z_2	$\frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^s \partial_\lambda A_\rho^s + \frac{\theta_0}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\mu A_\nu^s \partial_\lambda A_\rho^c$	Z_2 broken $\partial\Sigma^3$: $\frac{N}{2\pi} \partial_0 \phi^s \epsilon^{ij} \partial_i \lambda_j^s$	Z_2 broken Σ^2 : $\begin{pmatrix} 2p_1 & 0 & p_{12} & 0 \\ 0 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & N \\ 0 & 0 & N & 0 \end{pmatrix}$
$Z_{N_1} \times [Z_{N_2} \times Z_2]$	$Z_{N_1} \times Z_{N_2}$	Z_2	$\frac{N_1}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^c \partial_\lambda A_\rho^c + \frac{N_2}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^s \partial_\lambda A_\rho^s + \frac{\theta_0}{4\pi^2} \epsilon^{\mu\nu\lambda\rho} \partial_\mu A_\nu^s \partial_\lambda A_\rho^c$	Z_2 broken $\partial\Sigma^3$: $\frac{N_1}{2\pi} \partial_0 \phi^c \epsilon^{ij} \partial_i \lambda_j^c + \frac{N_2}{2\pi} \partial_0 \phi^s \epsilon^{ij} \partial_i \lambda_j^s$	Z_2 broken Σ^2 : $\begin{pmatrix} 2p_1 & N_1 & p_{12} & 0 \\ N_1 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & N_2 \\ 0 & 0 & N_2 & 0 \end{pmatrix}$

with G' as the symmetry group being gauged and $G'' = Z_2$ (i.e., the π rotation about S^y) or $G'' = Z_2^T$ is the leftover global symmetry group. This specific form of G implies that G' is always a normal subgroup of G . Thus, the gauging process for all of our five examples in this section (these are $Z_N \times Z_2^T$, $Z_N \times [U_s(1) \times Z_2]$, $U_c(1) \times [Z_N \times Z_2]$, $Z_{N_1} \times [Z_{N_2} \times Z_2]$; see, also, Table III) guarantee the forms as Eq. (76) corresponding to the Eq. (75) in the SET picture.³⁸

Let us now tie everything together. In Refs. 38 and 56, the full symmetry group G is a projective symmetry group (PSG). Before gauging G' , what we have is SPT state with $\text{PSG} = G_s = G$ and $G_g = Z_1 = 1$. After partially gauging the subgroup $G_g = G'$, we can view this as choosing a normal subgroup in the PSG. The leftover global symmetry group is indeed the quotient group as $G_s = \text{PSG}/G_g$.

To be more precise, based on the relation

$$\text{Gauging SPT: } \frac{G}{G'} = G'' \Leftrightarrow \text{SET: } \frac{\text{PSG}}{G_g} = G_s, \quad (77)$$

the gauging process for all of our five examples in this section can be regarded as converting SPT to SET:

$$\text{SPT} \begin{cases} \text{PSG} = G \\ G_g = 1 \\ G_s = \frac{\text{PSG}}{G_g} = G \end{cases} \xrightarrow{\text{gauging}} \text{SET} \begin{cases} \text{PSG} = G \\ G_g = G' \\ G_s = \frac{\text{PSG}}{G_g} = \frac{G}{G'} \end{cases} \quad (78)$$

In the following and the remaining parts of Sec. III, we will base on this principle: gauging SPT to SET, try to distinguish the features of SPT from the dynamical gauge theory viewpoint of SET.

2. Dynamical BF term in Σ^3

In this section, we demonstrate that gauging a discrete Z_N symmetry group will introduce a new topological term, namely, BF term in which all field variables are dynamical. To achieve a Z_N gauge theory, we start with a $U(1)$ gauge theory with a gauge field A_μ (A_μ will be replaced by A_μ^s and A_μ^c later) and couple A_μ to charge- N bosonic condensate χ . Adding a Higgs potential $U(\chi)$, the resultant condensate

$\langle\chi\rangle$ (or vacuum expectation value) will spontaneously break $U(1)$ symmetry down to Z_N symmetry. The Lagrangian for Z_N gauge theory⁷⁵ is

$$\begin{aligned} & |(\partial - iNA)\chi|^2 + U(\chi) + \dots \\ & = \langle\chi\rangle^2 |(\partial\varphi - NA)|^2 + U(\langle\chi\rangle) + \dots, \end{aligned} \quad (79)$$

where ... stands for other terms that already exist in the $U(1)$ gauge theory $\chi = \langle\chi\rangle e^{i\varphi}$. Following, we show the dual description of this Z_N gauge theory is BF theory. This argument works in arbitrary space-time dimension D , so let us demonstrate more conveniently in differential form. Here, $F = dA$ is the 2-form field strength of A , with A is the 1-form gauge field, while B is another $(D-2)$ form with independent gauge degree of freedom different from A . The trick is dualizing φ by introducing a Lagrangian multiplier B . Since $d^2\varphi = 0$, let us name $\rho = d\varphi$, we can impose $d\rho = 0$ by a Lagrangian multiplier B :

$$\begin{aligned} & \langle\chi\rangle^2 (d\varphi - NA) \wedge *(d\varphi - NA) \\ & = \langle\chi\rangle^2 (\rho - NA) \wedge *(\rho - NA) + \frac{1}{2\pi} B \wedge d\rho. \end{aligned} \quad (80)$$

In the first line of the above equation, the integral measure of the path integral is $\mathcal{D}A\mathcal{D}\varphi$, while in the second line it is changed to $\mathcal{D}A\mathcal{D}\rho\mathcal{D}B$. By writing $B \wedge d\rho = (-1)^{D-2} dB \wedge \rho$ and redefining field $(\rho - NeA) \rightarrow \rho$, we may directly integrate out the configuration $\mathcal{D}\rho$ rendering

$$\frac{N}{2\pi} B \wedge dA + \frac{(-1)^{(D-1)}}{(4\pi\langle\chi\rangle)^2} dB \wedge *dB, \quad (81)$$

where the first term is the topological BF term while the second term is the Maxwell term for $(D-2)$ -form $U(1)$ gauge field B . The path integral of this action has the measure $\mathcal{D}A\mathcal{D}B$ since we integrated out $\mathcal{D}\rho$. We assume that the superfluid density $\langle\chi\rangle$ is sufficiently large forming an ultraviolet energy scale and thus the Maxwell term which is quadratic becomes irrelevant in the low-energy field theory.

In closing, the derivation of the BF term as the dual description of the Z_N gauge theory, we need to confirm that the prefactor of the BF term “ $\frac{N}{2\pi} B \wedge dA$ ”, i.e., $N/2\pi$ is the correct normalization by examining whether the *statistical angles* of this BF theory exactly recover the result in the Z_N gauge theory or not.

In the famous Kitaev’s Z_2 toric code⁷⁶ (as Z_2 topological order or Z_2 gauge theory⁶¹), there are two kinds of excitations e and m anyons. When doing a full winding (or twice exchange) between e and m , the e - and m -wave functions gain a π phase, so-called statistical angle π . In general, it is known that a $2\pi/N$ statistical angle can be obtained from doing a full winding (or twice exchange) between certain two excitations of Z_N gauge theory. On the other hand, in BF theory, these two excitations are a point-particle with space-time trajectory described by a one-dimensional worldline J which minimally couples to A via $A \wedge *J$ and a higher-dimensional object (such as string or membrane) with space-time trajectory described by a worldsheet or worldvolume Σ which minimally couples to B via $B \wedge *\Sigma$. To test the statistical angle, we determine the statistical interaction between the two matter field space-time

trajectories J and Σ by studying the following Lagrangian:

$$\begin{aligned} & \frac{N}{2\pi} B \wedge dA + A \wedge *J + B \wedge *\Sigma \\ & = \frac{N}{2\pi(D-2)!} \epsilon^{\dots} B_{\dots} \partial A d^D x + A_{\mu} J^{\mu} d^D x \\ & + \frac{1}{(D-2)!} B_{\dots} \Sigma^{\dots} d^D x. \end{aligned} \quad (82)$$

Here, we skip the apparent indices as ... abbreviation. By noticing that the path integral of the above Lagrangian has integral measure $\mathcal{D}A\mathcal{D}B\mathcal{D}J\mathcal{D}\Sigma$, we integrate out $\mathcal{D}A\mathcal{D}B$ to deduce the statistical interaction, resulting in a Hopf term⁶²

$$\frac{2\pi}{N} J \cdot \frac{\epsilon^{\dots} \partial}{\partial^2} \Sigma_{\dots}. \quad (83)$$

This is the phase appearing in the exponent of the partition function $\exp[i\frac{2\pi}{N} J \cdot \frac{\epsilon^{\dots} \partial}{\partial^2} \Sigma_{\dots}]$, as this $\frac{2\pi}{N}$ factor implies a $\frac{2\pi}{N}$ statistical angle when braiding J around Σ by 2π , which reassures our $\frac{2\pi}{N}$ normalization is correct.

Let us express the differential form explicitly:

$$B \wedge dA = \frac{1}{(D-2)!} \epsilon^{\mu_1 \mu_2 \dots \mu_D} B_{\mu_1 \dots \mu_{D-1}} \partial_{\mu_D} A_{\mu_D} d^D x. \quad (84)$$

In Σ^3 bulk (i.e., space-time dimension $D=4$), the BF term is explicitly expressed by

$$\mathcal{L} = \frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu} \partial_{\lambda} A_{\rho}, \quad (85)$$

where $A_{\mu} \equiv A_{\mu}^c$ (A_{μ}^s) and $B_{\mu\nu} \equiv B_{\mu\nu}^c$ ($B_{\mu\nu}^s$) if Z_N symmetry originates from $U_c(1)$ charge symmetry [$U_s(1)$ spin symmetry]. The path-integral measure is $\mathcal{D}A\mathcal{D}B$.

If the Z_N gauge theory is defined on Σ^2 (i.e., space-time dimension $D=3$), the BF term reduces to a mutual Chern-Simons term

$$\mathcal{L} = \frac{N}{2\pi} \epsilon^{\mu\nu\lambda} A_{\mu} \partial_{\nu} \bar{A}_{\lambda}, \quad (86)$$

where a general $(D-2)$ -form gauge field B reduces to a simplest 1-form gauge field denoted by \bar{A}_{μ} , and the path-integral measure is $\mathcal{D}A\mathcal{D}\bar{A}$. $A_{\mu} \equiv A_{\mu}^c$ (A_{μ}^s) and $\bar{A}_{\mu} \equiv \bar{A}_{\mu}^c$ (\bar{A}_{μ}^s) if Z_N symmetry originates from $U_c(1)$ charge symmetry [$U_s(1)$ spin symmetry].

We comment that if we view both B and A fields in BF theory as dynamical gauge fields, the overall theory is a *dynamical topological field theory* in the sense that the ground-state degeneracy of BF theory will depend on the topology of the spatial manifold. For example, $(3+1)D$ BF theory has N^3 ground-state degeneracy, where N is the coefficient “level N ” of the BF term. Another example, $(2+1)D$ mutual Chern-Simons theory, has N^2 ground-state degeneracy, where N is the coefficient level N of the mutual Chern-Simons term.⁷⁷ As a side note, recently the BF term has been applied to other contexts with different interpretation in condensed matter physics.^{41,77,78}

3. Surface theory with quantum anomaly

In this section, we briefly preview the procedure carried out in Secs. III B, C 1, and C 2: the comparison of two kinds of dynamical gauge theory on the anomalous surface $\partial\Sigma^3$ and on the intrinsic Σ^2 bulk, and both have 2D spatial dimensions. The

philosophy is that we will treat the external fields appearing in Sec. II as dynamical gauge fields to gauge the 3D bulk Σ^3 and study its gauged surface theory $\partial\Sigma^2$ (with Z_2 or Z_2^T symmetry broken), and compare it to the gauged intrinsic 2D bulk Σ^2 (without Z_2 or Z_2^T symmetry). We find the gauged theory on the anomalous surface $\partial\Sigma^3$ with symmetry G' is different from the gauged theory of a two-dimensional SPT with the same G' in the intrinsic Σ^2 bulk.

We call this $\partial\Sigma^3$ surface anomalous because the boundary field theory on $\partial\Sigma^3$ is meant to cancel the anomaly contributed from the dynamical gauge theory in Σ^3 . Here, the situation is similar to the case that (2+1)D bulk topological Chern-Simons theory requires (1+1)D Wess-Zumino-Witten model on the boundary to cancel the anomaly.⁷⁹⁻⁸¹ A more familiar case is the (2+1)D bulk Abelian Chern-Simons theory, which requires (1+1)D edge theory of chiral bosons to preserve the gauge invariance on the manifold with boundary.^{61,81} Similarly, the (3+1)D bulk topological BF theory requires the (2+1)D anomalous edge theory of chiral bosons to preserve the gauge invariance on the manifold with boundary. We comment that the interpretation of (2+1)D electromagnetism in Ref. 41 is improper. Instead, we interpret the surface theory of BF theory as an anomalous chiral boson theory and leave the details to Appendix B. More on the understanding on the anomaly of topological phase or topological field theory, and their relation to bulk-edge correspondence, can be found in Refs. 54, 82, 83, and in particular Sec. VI of Ref. 84.

B. Derivation of surface chiral boson theory:

An example with $Z_N \times Z_2^T$

Physically, a SPT state with $Z_N \times Z_2^T$ in Σ^3 can be viewed as a time-reversal-symmetric bosonic superconductor with charge- N bosonic condensate. By collecting Eq. (85) and the θ_c term in Eq. (3), we obtain the following dynamical gauge theory with path-integral measure $\mathcal{D}A^c \mathcal{D}B^c$:

$$\mathcal{L} = \frac{\theta_c}{8\pi^2} \partial_\mu A_\nu^c \partial_\lambda A_\rho^c \epsilon^{\mu\nu\lambda\rho} + \frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^c \partial_\lambda A_\rho^c, \quad (87)$$

where $\theta_c = 2\pi + 4\pi k$ ($k \in \mathbb{Z}$). Its Z_2^T -broken surface $\partial\Sigma^3$, however, is meant to cancel the anomaly contribution from the bulk topological BF theory. This derivation is mentioned in Ref. 41 in a different context, but let us still walk through the logic to have coherent discussion. A convenient way to derive the chiral (vector and scalar) bosons is to choose a temporal gauge choice $A_0^c = 0$, $B_{0i}^c = 0$. The gauge choice itself should not affect the overall physics and thus should only be based on the convenience. The equations of motion (EOM) of A_0^c and B_{0i}^c impose the following constraints: $\epsilon^{0ijk} \partial_i B_{jk}^c = 0$ and $\epsilon^{0ijk} \partial_j A_k^c = 0$ which imply $B_{jk}^c = \partial_j \lambda_k - \partial_k \lambda_j$ and $A_k^c = \partial_k \phi$ as pure gauge forms. One interprets λ_k as vector bosons and ϕ as a scalar boson. Let us consider a $\partial\Sigma^3$ formed by x_1 - x_2 (i.e., x - y) plane at $x_3 = 0$ (i.e., $z = 0$) and then collect the term on $\partial\Sigma^3$ to be $\frac{1}{2} \frac{N}{4\pi} 4 \int (B_{12} F_{03}^c + B_{23} F_{01}^c + B_{31} F_{02}^c) = \frac{N}{2\pi} \int dx_3 \partial_3 (-\lambda_2 F_{01}^c + \lambda_1 F_{02}^c) + \dots$, so the surface theory is described by the action

$$\frac{N}{2\pi} \int d^3x (\partial_1 \lambda_2 - \partial_2 \lambda_1) \partial_0 \phi. \quad (88)$$

By choosing a light-cone gauge⁷⁹ $A_0^c + v_1 A_1^c + v_2 A_2^c = 0$, we can add velocity⁸¹ (so the Hamiltonian is not zero) to the boson theory, so the action becomes

$$\frac{1}{2\pi} \int d^3x \epsilon^{ij} \partial_i \lambda_j (N \partial_0 \phi - v_1 \partial_1 \phi - v_2 \partial_2 \phi) \quad (89)$$

with i, j running in 1,2. One can massage this surface action into a more symmetric form

$$\frac{1}{4\pi} \int d^3x \epsilon^{ij} \partial_i \lambda_j (N \partial_0 \phi - v_1 \partial_1 \phi - v_2 \partial_2 \phi) + \epsilon^{ij} \partial_i \phi (k \partial_0 \lambda_j - v_1 \partial_1 \lambda_j - v_2 \partial_2 \lambda_j). \quad (90)$$

The pure gauge forms also affect the θ_c term on $\partial\Sigma^3$, $\frac{\theta_c}{4\pi^2} \epsilon^{\nu\lambda\rho} A_\nu^c \partial_\lambda A_\rho^c = 0$ because of $A_k^c = \partial_k \phi$. So, the θ_c term becomes strictly zero on the surface. In this sense, Eq. (89) is the only leftover term, which is required to cancel the anomaly from the bulk BF theory in Σ^3 .

For intrinsic Z_N symmetry SPT on Z_2 broken Σ^2 , collecting Eqs. (12), (13) and (86) leads to the dynamical gauge theory

$$\begin{aligned} \mathcal{L} &= \frac{2p}{4\pi} A_\mu^c \partial_\nu A_\lambda^c \epsilon^{\mu\nu\lambda} + \frac{N}{2\pi} \epsilon^{\mu\nu\lambda} A_\mu^c \partial_\nu \bar{A}_\lambda^c \\ &= \frac{1}{4\pi} (A_\mu^c, \bar{A}_\mu^c) \begin{pmatrix} 2p & N \\ N & 0 \end{pmatrix} \partial_\nu \begin{pmatrix} A_\lambda^c \\ \bar{A}_\lambda^c \end{pmatrix} \epsilon^{\mu\nu\lambda} \end{aligned} \quad (91)$$

with path-integral measure $\mathcal{D}A^c \mathcal{D}\bar{A}^c$, integer $p = k$.

The $Z_N \times Z_2^T$ symmetry group is similar and the results are shown in Table III. Derivations of other symmetry groups are straightforward. (Details of derivations can be found in Appendix C.)

IV. CONCLUSIONS

In summary, in this work we study the response theory and dynamical gauge theory approach of bosonic symmetry-protected topological states (SPT) at least with charge symmetry [$U(1)$ or Z_N] or spin- S^z symmetry [$U(1)$ or Z_N] in 2D bulk, 3D bulk, and the surface of 3D bulk. The response theory applied in the case of the continuous $U(1)$ spin or charge symmetry group is based on the minimal physical input (such as the principle of gauge invariance, absence of topological order) without relying on lattice microscopic models. The 3D examples contain $U_c(1) \times Z_2^T$, $U_s(1) \times Z_2^T$, and $U_c(1) \times [U_s(1) \times Z_2]$, where $U_c(1)$ is charge-conservation symmetry, $U_s(1)$ is spin-rotation symmetry about S^z , Z_2^T is time-reversal symmetry, and Z_2 is specified to the π -rotation symmetry about spin S^y . Z_2^T -broken and Z_2 -broken surfaces are focused. The symmetry implementation in 2D examples is the same as the surfaces of 3D examples. By studying the 3D bulk response, we define many variants of the celebrated Witten effects, i.e., charge Witten effect, spin Witten effect, and mutual Witten effect. The last one is especially discussed in details which exhibit a very exotic experimental phenomenon. Through a case-by-case comparison of the quantum Hall effects between the surface and 2D bulk with the symmetry implementation, we emphasize that the surface of 3D SPT is anomalous and its existence requires the existence of an extra spatial dimension. The systematical study on the response theory of these SPT states with simple spin and charge symmetry implementation sheds light on the realistic charge

and spin response properties of underlying SPT states which will be possibly synthesized in condensed matter materials or cold-atom experiments in the near future.

On the other hand, the dynamical gauge theory description is also studied through the concrete examples at least with discrete Z_N spin symmetry or discrete Z_N charge symmetry. The latter can be viewed as bosonic topological superconductors. The 3D examples contain $Z_N \times Z_2^T$ (Z_N is charge symmetry), $Z_N \times Z_2^T$ (Z_N is spin- S^z symmetry), $U_c(1) \times [Z_N \times Z_2]$ (Z_2 is π rotation about spin S^y), $Z_N \times [U_s(1) \times Z_2]$, and, $Z_{N_1} \times [Z_{N_2} \times Z_2]$ (Z_{N_1} and Z_{N_2} are charge and spin symmetries, respectively). Z_2^T -broken and Z_2 -broken surfaces are focused. The symmetry implementation in 2D examples is the same as the surfaces of 3D examples. The dynamical gauge theory in 3D bulk is a topological gauge theory with topological BF term + variant of axionic Θ term. Its surface theory is gapless boson matter field theory with quantum anomaly. The dynamical gauge theory in 2D bulk is described by multicomponent dynamical Chern-Simons gauge theory with K_G -matrix coefficient. By studying the dynamical gauge theory, we explicitly show the connection between a SPT in 3D and a symmetry-enriched topological phase (SET) in 3D through the concrete examples.

There are several open questions to be stressed in future work.

1. *Symmetry implementation on the surface.* It will be quite interesting to study different symmetry-breaking patterns on the surface other than Z_2^T breaking and Z_2 breaking. For $U_c(1) \times Z_2^T$ and $U_s(1) \times Z_2^T$ SPT states in 3D, Refs. 41 and 48 have discussed many possible symmetry implementations on the surface based on field theory approach. $U_c(1) \times [U_s(1) \times Z_2]$ in 3D will be an interesting SPT state by studying different symmetry-breaking patterns on the surface. For all discrete groups we considered, their surface anomalous theory will be also interesting to be investigated with other symmetry implementation.

2. *Classification.* The Z_2 nature of the theta angles ($\theta_c, \theta_s, \theta_0$) gives one nontrivial state and one trivial state. There are more classes within the group cohomology level³ and possibly some classes beyond group cohomology,^{25,26,41} especially those with Z_2^T symmetry. In our case where we only consider the Θ and BF terms, it will be interesting to search for a complete set of topological terms to obtain more nontrivial states.

3. *Lattice realization and exactly solvable model.* The microscopic model, lattice model, and exact solvable model can help to determine more physical properties. Several works along this direction can be found in Refs. 23–26 and 28–30. It is noteworthy that the Z_N , $U(1)$ symmetry of the charges and spins can be implemented as the rotor angles in a quantum rotor model.^{28,29} In particular, the SPT state with Z_2 symmetry has been constructed where the Z_2 symmetry is realized as the Z_2 spin degree of freedom of σ_x, σ_z .²³ SPT states with Z_N symmetry have been constructed in Ref. 30. For a detailed lattice construction of the SPT edge states with a Z_N symmetry can be found in Ref. 29. Apparently, the experimental relevant materials for realizing these SPT states will be mostly significant. Various (charge, spin, mutual) Witten effects we proposed may shed light on the identification of these materials. Overall, further connections from our approach (on the response theory and the dynamical gauge

theory) to an explicit lattice/experimental realization will be desirable.

4. *Numerical simulation.* Our dynamical gauge theory formulation has numerical simulation implications. For example, we can apply the procedure in Ref. 23 by adding dynamical gauge field variables on the link, while global symmetry acts on the boson/spin on the sites, and then investigate the subsequent gauged SPT. This can also be done by cocycles formulation^{38,54,63,85} from a group cohomology viewpoint. A recent tensor network approach in Ref. 86 using quantum-state renormalization^{87–91} is applied in identifying Affleck-Kennedy-Lieb-Tasaki (AKLT) states in one and two dimensions. Based on the construction of SPT lattice models^{23–26} and further gauging the SPT by adding gauge field variable on the links,^{23,92} it will be applicable to apply similar numerical simulations to identify the gauged SPT (or SET), and further pin down the original SPT.

ACKNOWLEDGMENTS

We would like to thank X.-G. Wen and G. Vidal's organization of Emergence & Entanglement II conference during which part of the work was initiated. We are grateful to the hospitality of the Institute for Advanced Study in Tsinghua University. J.W. thanks D. Mazac, L. Santos, and D. Gaiotto for helpful discussion. This research is supported by NSF Grants No. DMR-1005541, No. NSFC 11074140, and No. NSFC 11274192 (J.W.). Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development & Innovation (P.Y. and J.W.).

APPENDIX A: K -MATRIX CHERN-SIMONS THEORY FOR SPT AND DERIVATION OF RESPONSE THEORY

Here, we derive the detailed K -matrix construction for SPT order (symmetry-protected topological order) and its response theory. Motivated by pioneer works,^{23,36–40,63} however, we still keep our following discussion self-contained and accessible. In $(2+1)D$, it is believed that a large class of SPT orders, especially Abelian SPT orders, can be classified and categorized by Abelian K -matrix Chern-Simons theory.^{61,93} The intrinsic field theory description of SPT has the following action:

$$S_{\text{SPT}, \Sigma^2} = \int dt d^2x \frac{1}{4\pi} K_{S, IJ} \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J, \quad (\text{A1})$$

where a is the intrinsic gauge field (or so-called statistical gauge field), and K_S is the K matrix which classifies and categorizes the SPT orders.

The SPT order is symmetry protected, so by definition its order is protected by global symmetry, say, some global symmetry group G_s . The distinct features of SPT from trivial insulator are its boundary edge states. The effective degree of freedom of SPT edges is the chiral boson field Φ , where Φ is introduced to preserve action invariance on the boundary under gauge transformation of the field a .⁶¹ The boundary action is

$$S_{\text{SPT}, \partial\Sigma^2} = \frac{1}{4\pi} \int dt dx K_{S, IJ} \partial_t \Phi_I \partial_x \Phi_J - V_{IJ} \partial_x \Phi_I \partial_x \Phi_J. \quad (\text{A2})$$

When G_s symmetry is preserved, the SPT edge states are gapless (otherwise it has degenerated ground states when adding symmetry-allowed gapping term). The SPT has ground-state degeneracy (GSD) on the torus as $\text{GSD} = |\det K| = 1$,^{10,36,61} this leads to the constrained canonical form of K_S . In this paper, we focus on the bosonic Abelian SPT. Due to its bosonic statistics, the quadratic form has all even integer coefficient, the canonical form^{10,94} is known to be the K matrix $K_{N \times N}^{b\pm}$, composed by blocks of $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and a set of all positive (or negative) coefficients E_8 lattices K_{E_8} . We can explicitly write $K_{N \times N}^{b\pm}$ as

$$K_{N \times N}^{b+} = K^{b0} \oplus K_{E_8} \oplus K_{E_8} \oplus \dots \quad (\text{A3})$$

and

$$K_{N \times N}^{b-} = K^{b0} \oplus (-K_{E_8}) \oplus (-K_{E_8}) \oplus \dots, \quad (\text{A4})$$

where

$$K_{N \times N}^{b0} = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right) \oplus \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right) \oplus \dots \quad (\text{A5})$$

and

$$K_{E_8} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}. \quad (\text{A6})$$

In our paper, however, we will not need the K_{E_8} state for our SPT examples. While it has been discussed in Ref. 36 that many classes of SPT can be realized by rank-2 K matrix, here we will show some SPT examples in our study need to have K matrix of large ranks, such as rank-4.

The implementation of this global symmetry can be explicitly shown by the symmetry transformation on the chiral bosons of the edge states

$$g : \{W^g, \delta\Phi^g, \eta_g\}. \quad (\text{A7})$$

The group element g of symmetry group G_s acts on chiral boson fields as

$$\Phi \rightarrow \eta_g (W^g)^{-1} \Phi + \delta\Phi^g, \quad (\text{A8})$$

$$K \rightarrow \eta_g (W^g)^T K W^g, \quad (\text{A9})$$

where $\eta_g = \pm 1$, with $+$ for the unitary symmetry and $-$ for the antiunitary symmetry transformation.

We will use the structure of the G_s to constrain the allowed g as $\{W^g, \delta\Phi^g, \eta_g\}$. The constraint is as follows: under any $\prod_i g_i = \mathbf{e}$, we have chiral boson field as a quantum phase unchanged up to module 2π ,

$$\Phi \xrightarrow{\prod_i g_i = \mathbf{e}} \Phi \text{ mod } 2\pi. \quad (\text{A10})$$

We should also allow the gauge equivalence to identify the same phases disguised by seemingly different transformations. Those gauge transformations are represented by some $\{X, \Delta\Phi\}$ where X obeys $X^T K X = K$ and $X \in GL(N, \mathbb{Z})$ as a general linear group of degree N over integer \mathbb{Z} , and $\Phi \rightarrow \Phi + \Delta\Phi$, such that we identify

$$W_g \rightarrow X^{-1} W_g X, \quad (\text{A11})$$

$$\delta\Phi^g \rightarrow X^{-1} (\Delta\Phi + \delta\Phi^g - \eta_g W_g^{-1} \Delta\Phi). \quad (\text{A12})$$

To gauge the theory, we need to couple the global symmetry current to the (dynamical or external) gauge field A . In the specific examples we study [such as $U(1)$, Z_N , $U(1) \times U(1)$, $Z_{N_1} \times U(1)$, $Z_{N_1} \times Z_{N_2}$], all the global symmetry can be restricted to g as $g = \{W^g = \mathbb{I}, \delta\Phi^g, \eta_g = +1\}$, so the global symmetry current is fully determined by $\delta\Phi^g$. Therefore, the global symmetry current on $(1+1)\text{D}$ (here $\partial\Sigma_2$) is $\epsilon^{\mu\nu} \partial_\nu \Phi / 2\pi$, coupled to the external gauge field A as

$$\mathbf{q}_J^I \frac{1}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^I \partial_\nu \Phi^J, \quad (\text{A13})$$

the global symmetry current in $(2+1)\text{D}$ (here Σ_2) is $\epsilon^{\mu\nu\rho} \partial_\nu a_\rho^J / 2\pi$, coupled to the external gauge field A as

$$\mathbf{q}_J^I \frac{1}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^I \partial_\nu a_\rho^J. \quad (\text{A14})$$

The t_J^I vector is proportional to $\delta\Phi_J^g$, with the I of t_J^I specifying the I th-independent generator of the group G_s . The Lagrangian of the SPT order with intrinsic a coupled to the external gauge field A will be

$$\mathcal{L}_{\text{SPT+Gauge}} = \frac{1}{4\pi} K_{S,IJ} \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J + \mathbf{q}_J^I \frac{1}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^I \partial_\nu a_\rho^J. \quad (\text{A15})$$

To integrate out a , we adopt EOM as a constraint:

$$\begin{aligned} \frac{1}{2\pi} K_{S,IJ} \epsilon^{\mu\nu\rho} \partial_\nu a_\rho^J + \mathbf{q}_J^I \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \partial_\nu A_\rho^J &= 0 \\ \Rightarrow a_\rho^{J'} &= -K_{S,I'J'}^{-1} \mathbf{q}_J^I A_\rho^J. \end{aligned} \quad (\text{A16})$$

We get the gauged version description, left with only external gauge field A ,

$$\begin{aligned} \mathcal{L}_{\text{SPT+Gauge}}(A) &= \frac{1}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^{I'} (-\mathbf{q}_J^I K_{S,IJ}^{-1} \mathbf{q}_J^{J'}) \partial_\nu A_\rho^{J'} \\ &\equiv \frac{1}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^{I'} K_{G,I'J'} \partial_\nu A_\rho^{J'}, \end{aligned} \quad (\text{A17})$$

$$\text{where } K_{G,I'J'} \equiv -\mathbf{q}_J^I K_{S,IJ}^{-1} \mathbf{q}_J^{J'}.$$

We will work through examples shown in our main text, relevant to the response study of Σ_2 , $\partial\Sigma_3$. To reiterate the examples below only requires $W^g = \mathbb{I}$ and $\eta_g = +1$, so below we only list $\delta\Phi$ to specify the symmetry transformation. As a side remark, our group elements g representation also form a faithful representation.^{36,85}

1. U(1)

A rank-2 K matrix suffices to exhaust all classes of group cohomology $\mathcal{H}^3[\text{U}(1), \text{U}(1)] = \mathbb{Z}$ with U(1) symmetry,

$$K_{\text{SPT}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A18})$$

where the symmetry transformation of U(1) with an angle θ specifies the group element g ,

$$g_\theta : \delta\Phi^{U(1)\theta} = \theta\mathbf{q} = \theta \begin{pmatrix} 1 \\ -p \end{pmatrix}. \quad (\text{A19})$$

Since a U(1) group only requires one generator, there is only one kind of charge vector $\mathbf{q} = (1, p)$. Here, p labels the \mathbb{Z} class of the cohomology group $\mathcal{H}^3[\text{U}(1), \text{U}(1)] = \mathbb{Z}$, while $K_G = -\mathbf{q}_I K_{S, IJ}^{-1} \mathbf{q}_J = 2p$. So the topological term in the gauged theory is

$$\mathcal{L}_{\text{SPT+Gauge}}(A) = \frac{2p}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^I \partial_\nu A_\rho^I. \quad (\text{A20})$$

2. \mathbb{Z}_N

Similarly as the U(1) symmetry case, the \mathbb{Z}_N symmetry only requires a rank-2 K matrix, which exhausts all $\mathcal{H}^3[\mathbb{Z}_N, \text{U}(1)] = \mathbb{Z}_N$:

$$K_{\text{SPT}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A21})$$

$$g_n : \delta\Phi = \frac{2\pi}{N} n \begin{pmatrix} 1 \\ -p \end{pmatrix}. \quad (\text{A22})$$

Here, p labels the \mathbb{Z}_N class of the cohomology group $\mathcal{H}^3[\mathbb{Z}_N, \text{U}(1)] = \mathbb{Z}_N$. Both p and n have module N structure as elements in \mathbb{Z}_N .

However, the main difference from U(1) gauged theory is that for the \mathbb{Z}_N case, the gauge charge and gauge flux are quantized by module N , which can be captured by a mutual Chern-Simons term $\frac{N}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^I \partial_\nu A_\rho^{II}$ (or more generally a BF theory, see Sec. III A2), where the statistics angle of a full wave function gains a $2\pi/N$ phase after a full winding between a unit gauge charge and a unit gauge flux:

$$\mathcal{L}_{\text{SPT+Gauge}}(A) = \frac{1}{4\pi} \epsilon^{\mu\nu\rho} A_\mu^{I'} \begin{pmatrix} 2p & N \\ N & 0 \end{pmatrix}_{I'J'} \partial_\nu A_\rho^{J'}. \quad (\text{A23})$$

3. U(1) \times U(1)

We require a rank-4 K matrix to obtain all classes of group cohomology $\mathcal{H}^3[\text{U}(1) \times \text{U}(1), \text{U}(1)] = \mathbb{Z}^3$ with U(1) symmetry

$$K_{\text{SPT}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A24})$$

$$g_\theta : \delta\Phi = \delta\Phi^{U(1)\theta_1} + \delta\Phi^{U(1)\theta_2} = \theta_1 \mathbf{q}^1 + \theta_2 \mathbf{q}^2 \quad (\text{A25})$$

$$\text{with } \mathbf{q}^1 = \begin{pmatrix} 1 \\ -p_1 \\ 0 \\ -p_{12} \end{pmatrix}, \quad \mathbf{q}^2 = \begin{pmatrix} 0 \\ -p_{21} \\ 1 \\ -p_2 \end{pmatrix} \quad (\text{A26})$$

with $\theta_1, \theta_2 \in \text{U}(1)$.

The terms with gauge fields coupling to the symmetry current are

$$\mathbf{q}_J^1 \frac{1}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^1 \partial_\nu a_\rho^J + \mathbf{q}_J^2 \frac{1}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^3 \partial_\nu a_\rho^J. \quad (\text{A27})$$

Here, we couple the two generators of symmetry group to different gauge fields, and purposefully choose them to be A^1 and A^3 to represent the charge sector^{38,40} of gauge fields, while the meaning of this choice will be revealed in the next section in Sec. A4. It is easy to see $p_{12} + p_{21}$ identify the same index from the gauged coupling term $(p_{12} + p_{21}) \epsilon^{\mu\nu\rho} A_\mu^1 \partial_\nu A_\rho^3 / 2\pi$. So, we may identify $p_{12} + p_{21} \rightarrow p_{12}$, with p_1, p_2, p_{12} each labeling a \mathbb{Z} in \mathbb{Z}^3 . The gauged theory has this K matrix

$$K_G = \begin{pmatrix} 2p_1 & 0 & p_{12} & 0 \\ 0 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (\text{A28})$$

or simply in the gauge charge sectors of $\text{U}(1) \times \text{U}(1)$: $K_G = \begin{pmatrix} 2p_1 & p_{12} \\ p_{12} & 2p_2 \end{pmatrix}$.

4. $\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}$

We require a rank-4 K matrix to obtain all classes of group cohomology $\mathcal{H}^3[\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}, \text{U}(1)] = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{\text{gcd}(N_1, N_2)}$ with U(1) symmetry,

$$K_{\text{SPT}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A29})$$

$$g : \delta\Phi = \delta\Phi^1 + \delta\Phi^2 = \frac{2\pi}{N_1} n_1 \mathbf{q}^1 + \frac{2\pi}{N_2} n_2 \mathbf{q}^2 \quad (\text{A30})$$

$$\text{with } \mathbf{q}^1 = \begin{pmatrix} 1 \\ -p_1 \\ 0 \\ -p_{12} \end{pmatrix}, \quad \mathbf{q}^2 = \begin{pmatrix} 0 \\ -p_{21} \\ 1 \\ -p_2 \end{pmatrix} \quad (\text{A31})$$

with $n_1 \in \mathbb{Z}_{N_1}, n_2 \in \mathbb{Z}_{N_2}$.

Again, $p_{12} + p_{21}$ identify the same index from the gauged coupling term $(p_{12} + p_{21}) \epsilon^{\mu\nu\rho} A_\mu^1 \partial_\nu A_\rho^3 / 2\pi$. So, we may identify $p_{12} + p_{21} \rightarrow p_{12}$. However, the two gauged sectors of \mathbb{Z}_{N_1} symmetry and \mathbb{Z}_{N_2} symmetry share the same index p_{12} . Therefore, we should emphasize the number p_{12} of different topological phases is identified by $p_{12} \sim p_{12} + c_1 N_1 + c_2 N_2 = p_{12} + c_{12} \text{gcd}(N_1, N_2)$ for any integer c_1, c_2 . There is a corresponding integer c_{12} from the Chinese remainder theorem. This means

$$p_{12} = p_{12} \text{ mod } [\text{gcd}(N_1, N_2)]. \quad (\text{A32})$$

All together, we have that p_1, p_2, p_{12} each labels $\mathbb{Z}_{N_1}, \mathbb{Z}_{N_2}, \mathbb{Z}_{\text{gcd}(N_1, N_2)}$. While our argument is based on the symmetry transformation from the SPT side, this relation can also be confirmed by a different argument from the statistics angle³⁹ of the gauged theory side. The main difference from U(1) \times U(1) gauged theory is that for the $\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}$ case, the gauge charge and gauge flux are quantized by module N , which can be captured by two mutual Chern-Simons terms $\frac{N_1}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^1 \partial_\nu A_\rho^2 + \frac{N_2}{2\pi} \epsilon^{\mu\nu\rho} A_\mu^3 \partial_\nu A_\rho^4$, where the statistics angle of a full wave function gains a $2\pi/N_1$ (or $2\pi/N_2$) phase after a full winding between a unit gauge charge and a unit gauge flux of \mathbb{Z}_{N_1} symmetry (or \mathbb{Z}_{N_2} symmetry). The gauged theory

has this K matrix

$$K_G = \begin{pmatrix} 2p_1 & N_1 & p_{12} & 0 \\ N_1 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & N_2 \\ 0 & 0 & N_2 & 0 \end{pmatrix}. \quad (\text{A33})$$

5. $\mathbb{Z}_{N_1} \times \text{U}(1)$

Similar to Secs. A 3 and A 4, we require a rank-4 K matrix to obtain all classes of group cohomology $\mathcal{H}^3[\mathbb{Z}_{N_1} \times \text{U}(1)] = \mathbb{Z}_{N_1} \times \mathbb{Z} \times \mathbb{Z}_{N_1}$ with $\text{U}(1)$ symmetry

$$K_{\text{SPT}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A34})$$

$$g : \delta\Phi = \delta\Phi^1 + \delta\Phi^2 = \frac{2\pi}{N_1} n_1 \mathbf{q}^1 + \theta_2 \mathbf{q}^2 \quad (\text{A35})$$

$$\text{with } \mathbf{q}^1 = \begin{pmatrix} 1 \\ -p_1 \\ 0 \\ -p_{12} \end{pmatrix}, \quad \mathbf{q}^2 = \begin{pmatrix} 0 \\ -p_{21} \\ 1 \\ -p_2 \end{pmatrix} \quad (\text{A36})$$

with $n_1 \in \mathbb{Z}_{N_1}$ and $\theta_2 \in \text{U}(1)$.

While the K_G matrix from the response theory is derived in the same manner as in Secs. A 3 and A 4, we skip details and directly list the result

$$K_G = \begin{pmatrix} 2p_1 & N_1 & p_{12} & 0 \\ N_1 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (\text{A37})$$

The classification follows the logic in Secs. A 3 and A 4: we have that p_1, p_2, p_{12} each labels $\mathbb{Z}_{N_1}, \mathbb{Z}, \mathbb{Z}_{N_1}$.

APPENDIX B: SURFACE OF (3 + 1)D BF THEORY IS NOT GAUGE INVARIANT

There are some discussions on the surface theory of BF theory in the literature in different contexts. For example, in Appendix A of Ref. 41, the surface theory is interpreted as a (2 + 1)D electromagnetism, i.e., free photon theory. We comment that the theory is not a theory of photons claimed in the reference. In other words, it is not a usual electromagnetism. The reason is the following. If we use the definitions of ‘‘electric field’’ and ‘‘magnetic field’’ in the reference, the Lagrangian in Eq. (A6) can be reexpressed in the form of $\mathcal{L} = \dots + E^2 + B^2$ (in Euclidean metric), where ‘‘...’’ is an extra term that is not a gauge-invariant term. *Gauge invariance* is a fundamental requirement of electromagnetism. If one insists on the term ‘‘electromagnetism,’’ it is more appropriate to call it ‘‘anomalous electromagnetism’’ which can not exist alone in (2 + 1)D. It can exist as a surface of a (3 + 1)D bulk.

We emphasize that both Lagrangians in their Eq. (A6) and Lagrangians of the chiral boson theory in our paper are not gauge invariant *alone* on the surface. A gauge-invariant theory must be composed by the bulk BF theory and the surface theory *as a whole*. Thus, a full quantum effect of anomaly

of leaking currents to the bulk [here relation between chiral boson and BF theory is like the relation between chiral boson and Chern-Simons theory in (2 + 1)D quantum Hall effects] does happen.

APPENDIX C: DERIVATION OF SURFACE CHIRAL BOSON THEORY WITH QUANTUM ANOMALY

1. $\text{U}_c(1) \times [\mathbb{Z}_N \rtimes \mathbb{Z}_2]$ and $\mathbb{Z}_N \times [\text{U}_s(1) \rtimes \mathbb{Z}_2]$ symmetry groups

Physically, a SPT state with $\text{U}_c(1) \times [\mathbb{Z}_N \rtimes \mathbb{Z}_2]$ in Σ^3 can be viewed as a three-dimensional interacting bosonic ground state with charge-1 and spin-1 of discrete spin symmetry $\mathbb{Z}_N \rtimes \mathbb{Z}_2$ where \mathbb{Z}_N is $2\pi/N$ rotation about spin- z direction and \mathbb{Z}_2 is π rotation about spin- y direction. By collecting Eq. (85) and the θ_0 term in Eq. (3), we obtain the following dynamical gauge theory with path-integral measure $\mathcal{D}A^c \mathcal{D}A^s \mathcal{D}B^s$:

$$\mathcal{L} = \frac{\theta_0}{4\pi^2} \partial_\mu A_\nu^c \partial_\lambda A_\rho^s \epsilon^{\mu\nu\lambda\rho} + \frac{N}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^s \partial_\lambda A_\rho^s, \quad (\text{C1})$$

where $\theta_0 = \pi + 2\pi k$ ($k \in \mathbb{Z}$).

Similar to the previous Sec. III B, the (3 + 1)D bulk topological BF theory requires the (2 + 1)D edge theory of chiral (vector and scalar) bosons to preserve the gauge invariance on the manifold with boundary.^{79,81} We again first choose a temporal gauge choice $A_0^s = 0, B_{0i}^s = 0$. The gauge choice itself should not affect the overall physics; this choice should based only on the convenience. The EOM of A_0^s and B_{0i}^s impose the following constraints: $\epsilon^{0ijk} \partial_i B_{jk}^s = 0$ and $\epsilon^{0ijk} \partial_j A_k^s = 0$ which imply $B_{jk}^s = \partial_j \lambda_k - \partial_k \lambda_j$ and $A_k^s = \partial_k \phi$ as pure gauge forms. One interprets λ_k as vector bosons and ϕ as a scalar boson. Let us consider a $\partial\Sigma^3$ formed by x_1 - x_2 plane at $x_3 = 0$ and then collect the term on $\partial\Sigma^3$ to be

$$\frac{N}{2\pi} \int d^3x (\partial_1 \lambda_2 - \partial_2 \lambda_1) \partial_0 \phi. \quad (\text{C2})$$

By choosing a light-cone gauge $A_0^c + v_1 A_1^c + v_2 A_2^c = 0$, the action becomes

$$\frac{1}{2\pi} \int d^3x \epsilon^{ij} \partial_i \lambda_j (N \partial_0 \phi - v_1 \partial_1 \phi - v_2 \partial_2 \phi) \quad (\text{C3})$$

with i, j running in 1, 2. The pure gauge forms also affect the θ_0 term on $\partial\Sigma^3$, with $\frac{\theta_0}{4\pi^2} \epsilon^{\nu\lambda\rho} A_\nu^c \partial_\lambda A_\rho^s = 0$ because of $A_k^s = \partial_k \phi$. So, the θ_0 term becomes strictly zero on the surface. In this sense, Eq. (C3) is the only leftover term, which is required to cancel the anomaly from the bulk BF theory in Σ^3 .

On \mathbb{Z}_2 broken Σ^2 , collecting Eqs. (71) and (86) leads to the dynamical gauge theory

$$\mathcal{L} = \frac{1}{4\pi} (A_\mu^c, \bar{A}_\mu^c, A_\mu^s, \bar{A}_\mu^s) \begin{pmatrix} 2p_1 & 0 & p_{12} & 0 \\ 0 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & N \\ 0 & 0 & N & 0 \end{pmatrix} \times \partial_\nu \begin{pmatrix} A_\lambda^c \\ \bar{A}_\lambda^c \\ A_\lambda^s \\ \bar{A}_\lambda^s \end{pmatrix} \epsilon^{\mu\nu\lambda} \quad (\text{C4})$$

with path-integral measure $\mathcal{D}A^c \overline{\mathcal{D}A}^c$ and $p_1, p_{12}, p_2 \in \mathbb{Z}$. $Z_N \times [U_s(1) \times Z_2]$ symmetry group is similar and the results are shown in Table III.

2. $Z_{N_1} \times [Z_{N_2} \times Z_2]$ symmetry group

Physically, a SET state with $Z_{N_1} \times [Z_{N_2} \times Z_2]$ in Σ^3 can be viewed as a three-dimensional bosonic superconductor with charge- N_1 condensate and spin-1 of discrete spin symmetry $Z_N \times Z_2$ where Z_2 is π rotation about S^y . By collecting Eq. (85) and the θ_0 term in Eq. (3), we obtain the following dynamical gauge theory with path-integral measure $\mathcal{D}A^c \mathcal{D}A^s \mathcal{D}B^c \mathcal{D}B^s$:

$$\mathcal{L} = \frac{\theta_0}{4\pi^2} \partial_\mu A_\nu^c \partial_\lambda A_\rho^s \epsilon^{\mu\nu\lambda\rho} + \frac{N_1}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^c \partial_\lambda A_\rho^c + \frac{N_2}{4\pi} \epsilon^{\mu\nu\lambda\rho} B_{\mu\nu}^s \partial_\lambda A_\rho^s, \quad (\text{C5})$$

where $\theta_0 = \pi + 2\pi k$ ($k \in \mathbb{Z}$).

Similar to the previous Secs. III B and C 1, the (3+1)D bulk topological BF theory requires the (2+1)D edge theory of chiral (vector and scalar) bosons to preserve the gauge invariance on the manifold with boundary.^{61,79,81} We again first choose a temporal gauge choice^{41,61,79,81} $A_0^c = A_0^s = 0$, $B_{0i}^c = B_{0i}^s = 0$. The gauge choice itself should not affect the overall physics; this choice should be based only on convenience. The EOM of $A_0^c, A_0^s, B_{0i}^c, B_{0i}^s$ impose the following constraints: $\epsilon^{0ijk} \partial_i B_{jk}^c = \epsilon^{0ijk} \partial_i B_{jk}^s = 0$ and $\epsilon^{0ijk} \partial_j A_k^c = \epsilon^{0ijk} \partial_j A_k^s = 0$. These imply $B_{jk}^c = \partial_j \lambda_k^c - \partial_k \lambda_j^c$, $B_{jk}^s = \partial_j \lambda_k^s - \partial_k \lambda_j^s$, $A_k^c = \partial_k \phi^c$, and $A_k^s = \partial_k \phi^s$ as pure gauge forms. One interprets λ_k^c, λ_k^s as vector bosons and ϕ^c, ϕ^s as scalar bosons. Let us consider $\partial\Sigma^3$ formed by the x_1 - x_2 plane at $x_3 = 0$ and then

collect the term on $\partial\Sigma^3$ to be

$$\int d^3x \left(\frac{N_1}{2\pi} (\partial_1 \lambda_2^c - \partial_2 \lambda_1^c) \partial_0 \phi^c + \frac{N_2}{2\pi} (\partial_1 \lambda_2^s - \partial_2 \lambda_1^s) \partial_0 \phi^s \right). \quad (\text{C6})$$

By choosing light-cone gauges $A_0^c + v_1 A_1^c + v_2 A_2^c = 0$ and $A_0^s + v_1 A_1^s + v_2 A_2^s = 0$, the action becomes

$$\frac{1}{2\pi} \int d^3x \epsilon^{ij} (\partial_i \lambda_j^c (N_1 \partial_0 \phi^c - v_1 \partial_1 \phi^c - v_2 \partial_2 \phi^c) + \partial_i \lambda_j^s (N_2 \partial_0 \phi^s - v_1 \partial_1 \phi^s - v_2 \partial_2 \phi^s)) \quad (\text{C7})$$

with i, j running in 1, 2. The pure gauge forms also affect the θ_0 term on $\partial\Sigma^3$, with $\frac{\theta_0}{4\pi^2} \epsilon^{\nu\lambda\rho} A_\nu^c \partial_\lambda A_\rho^s = 0$ because of $A_k^c = \partial_k \phi^c$, $A_k^s = \partial_k \phi^s$. So, the θ_0 term becomes strictly zero on the surface. In this sense, Eq. (C7) is the only leftover term, which is required to cancel the anomaly from the bulk BF theory in Σ^3 .

On Z_2 broken Σ^2 , collecting Eqs. (71) and (86) leads to the dynamical gauge theory

$$\mathcal{L} = \frac{1}{4\pi} (A_\mu^c, \overline{A}_\mu^c, A_\mu^s, \overline{A}_\mu^s) \begin{pmatrix} 2p_1 & N_1 & p_{12} & 0 \\ N_1 & 0 & 0 & 0 \\ p_{12} & 0 & 2p_2 & N_2 \\ 0 & 0 & N_2 & 0 \end{pmatrix} \times \partial_\nu \begin{pmatrix} A_\lambda^c \\ \overline{A}_\lambda^c \\ A_\lambda^s \\ \overline{A}_\lambda^s \end{pmatrix} \epsilon^{\mu\nu\lambda} \quad (\text{C8})$$

with path-integral measure $\mathcal{D}A^c \overline{\mathcal{D}A}^c \mathcal{D}A^s \overline{\mathcal{D}A}^s$ and $p_1, p_{12}, p_2 \in \mathbb{Z}$.

¹M. Levin and X.-G. Wen, *Phys. Rev. Lett.* **96**, 110405 (2006).

²A. Kitaev and J. Preskill, *Phys. Rev. Lett.* **96**, 110404 (2006).

³Xie Chen, Z. C. Gu, Z. X. Liu, and X. G. Wen, *Phys. Rev. B* **87**, 155114 (2013).

⁴Xie Chen, Z. C. Gu, and X. G. Wen, *Phys. Rev. B* **82**, 155138 (2010).

⁵X. Chen, Z.-C. Gu, Z.-X. Liu, X.-G. Wen, *Science* **338**, 1604 (2012); X. L. Qi, *ibid.* **338**, 1550 (2012).

⁶D. C. Tsui, H. L. Stormer, and A. C. Gossard, *Phys. Rev. Lett.* **48**, 1559 (1982).

⁷X.-G. Wen, *Phys. Rev. B* **40**, 7387 (1989).

⁸X.-G. Wen and Q. Niu, *Phys. Rev. B* **41**, 9377 (1990).

⁹X.-G. Wen, *Int. J. Mod. Phys. B* **4**, 239 (1990).

¹⁰Juven Wang and X.-G. Wen, arXiv:1212.4863.

¹¹B. I. Halperin, *Phys. Rev. Lett.* **52**, 1583 (1984).

¹²D. Arovav, J. R. Schrieffer, and F. Wilczek, *Phys. Rev. Lett.* **53**, 722 (1984).

¹³B. I. Halperin, *Phys. Rev. B* **25**, 2185 (1982).

¹⁴X.-G. Wen, *Int. J. Mod. Phys. B* **6**, 1711 (1992).

¹⁵J. E. Moore and L. Balents, *Phys. Rev. B* **75**, 121306 (2007).

¹⁶L. Fu, C. L. Kane, and E. J. Mele, *Phys. Rev. Lett.* **98**, 106803 (2007).

¹⁷R. Roy, *Phys. Rev. B* **79**, 195322 (2009).

¹⁸M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).

¹⁹J. E. Moore, *Nature (London)* **464**, 194 (2010).

²⁰X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).

²¹X. L. Qi, T. L. Hughes, and S.-C. Zhang, *Phys. Rev. B* **78**, 195424 (2008).

²²F. D. M. Haldane, *Phys. Rev. Lett.* **50**, 1153 (1983); *Phys. Lett. A* **93**, 464 (1983); I. Affleck and F. D. M. Haldane, *Phys. Rev. B* **36**, 5291 (1987); I. Affleck, *J. Phys.: Condens. Matter.* **1**, 3047 (1989).

²³M. Levin and Z. C. Gu, *Phys. Rev. B* **86**, 115109 (2012).

²⁴F. J. Burnell, Xie Chen, L. Fidkowski, and A. Vishwanath, arXiv:1302.7072.

²⁵X. Chen, Y.-M. Lu, and A. Vishwanath, arXiv:1303.4301.

²⁶L. Fidkowski, X. Chen, and A. Vishwanath, *Phys. Rev. X* **3**, 041016 (2013).

²⁷K. Walker and Zhenghan Wang, *Frontiers Phys.* **7**, 150 (2012).

²⁸X. Chen and X.-G. Wen, *Phys. Rev. B* **86**, 235135 (2012).

²⁹L. H. Santos and J. Wang, arXiv:1310.8291.

³⁰L. -Y. Hung and Y. Wan, *Phys. Rev. B* **86**, 235132 (2012).

³¹M. Freedman, C. Nayak, K. Shtengel, K. Walker, and Z. Wang, *Ann. Phys. (NY)* **310**, 428 (2004).

³²M. A. Levin and X. G. Wen, *Phys. Rev. B* **71**, 045110 (2005).

- ³³Peng Ye and X.-G. Wen, *Phys. Rev. B* **87**, 195128 (2013).
- ³⁴Peng Ye and X.-G. Wen, arXiv:1303.3572.
- ³⁵M. A. Metlitski, C. L. Kane, and M. P. A. Fisher, *Phys. Rev. B* **88**, 035131 (2013).
- ³⁶Y. M. Lu and A. Vishwanath, *Phys. Rev. B* **86**, 125119 (2012).
- ³⁷Y.-M. Lu and A. Vishwanath, arXiv:1302.2634.
- ³⁸L.-Y. Hung and X.-G. Wen, *Phys. Rev. B* **87**, 165107 (2013).
- ³⁹Meng Cheng and Zheng-Cheng Gu, arXiv:1302.4803.
- ⁴⁰Ling-Yan Hung and Yidun Wan, *Phys. Rev. B* **87**, 195103 (2013).
- ⁴¹Ashvin Vishwanath and T. Senthil, *Phys. Rev. X* **3**, 011016 (2013).
- ⁴²Kenke Xu, *Phys. Rev. B* **87**, 144421 (2013).
- ⁴³J. Oon, G. Y. Cho, and Kenke Xu, *Phys. Rev. B* **88**, 014425 (2013).
- ⁴⁴Z. Bi, A. Rasmussen, and Kenke Xu, arXiv:1304.7272.
- ⁴⁵Z.-X. Liu and X.-G. Wen, *Phys. Rev. Lett.* **110**, 067205 (2013).
- ⁴⁶Olabode M. Sule, Xiao Chen, and Shinsei Ryu, *Phys. Rev. B* **88**, 075125 (2013).
- ⁴⁷Kenke Xu and T. Senthil, *Phys. Rev. B* **87**, 174412 (2013).
- ⁴⁸C. Wang and T. Senthil, *Phys. Rev. B* **87**, 235122 (2013).
- ⁴⁹R. Dijkgraaf and E. Witten, *Commun. Math. Phys.* **129**, 393 (1990).
- ⁵⁰T. Senthil and M. Levin, *Phys. Rev. Lett.* **110**, 046801 (2013).
- ⁵¹T. Grover and A. Vishwanath, *Phys. Rev. B* **87**, 045129 (2013).
- ⁵²Y.-M. Lu and D.-H. Lee, arXiv:1210.0909.
- ⁵³Y.-M. Lu and D.-H. Lee, arXiv:1212.0863.
- ⁵⁴X.-G. Wen, arXiv:1301.7675.
- ⁵⁵A. Mesaros and Y. Ran, *Phys. Rev. B* **87**, 155115 (2013).
- ⁵⁶A. M. Essin and M. Hermele, *Phys. Rev. B* **87**, 104406 (2013).
- ⁵⁷E. Witten, *Phys. Lett. B* **86**, 283 (1979).
- ⁵⁸G. Rosenberg and M. Franz, *Phys. Rev. B* **82**, 035105 (2010).
- ⁵⁹J. S. Schwinger, *Science* **165**, 757 (1969).
- ⁶⁰D. Zwanziger, *Phys. Rev.* **176**, 1489 (1968).
- ⁶¹X.-G. Wen, *Quantum Field Theory of Many-Body Systems: From the Origin of Sound to an Origin of Light and Electrons* (Oxford University Press, Oxford, 2004).
- ⁶²A. Zee, *Quantum Field Theory in a Nutshell* (Princeton University Press, Princeton, NJ, 2010), and references therein.
- ⁶³L.-Y. Hung and X.-G. Wen, arXiv:1211.2767.
- ⁶⁴G. W. Moore and N. Seiberg, *Phys. Lett. B* **220**, 422 (1989).
- ⁶⁵S. Adler, *Phys. Rev.* **177**, 2426 (1969).
- ⁶⁶J. S. Bell and R. Jackiw, *II Nuovo Cimento A* **60**, 47 (1969).
- ⁶⁷Kenke Xu, *Phys. Rev. B* **83**, 024408 (2011).
- ⁶⁸Victor M. Galitski, G. Refael, Matthew P. A. Fisher, and T. Senthil, *Phys. Rev. Lett.* **95**, 077002 (2005).
- ⁶⁹M. C. Diamantini, P. Sodano, and C. A. Trugenberger, *Nucl. Phys. B* **474**, 641 (1996).
- ⁷⁰Peng Ye, C. S. Tian, X.-L. Qi, and Zheng-Yu Weng, *Phys. Rev. Lett.* **106**, 147002 (2011).
- ⁷¹Peng Ye, C. S. Tian, X.-L. Qi, and Zheng-Yu Weng, *Nucl. Phys. B* **854**, 815 (2012).
- ⁷²Maissam Barkeshli and Xiao-Gang Wen, *Phys. Rev. B* **86**, 085114 (2012).
- ⁷³X. L. Qi, R. Li, J. Zhang, and S.-C. Zhang, *Science* **323**, 1184 (2009).
- ⁷⁴J. D. Jackson, *Classical Electromagnetics* (Wiley, New York, 1999).
- ⁷⁵T. Banks, *Nucl. Phys. B* **323**, 90 (1989).
- ⁷⁶A. Y. Kitaev, *Ann. Phys. (NY)* **303**, 2 (2003).
- ⁷⁷T. H. Hansson, V. Oganesyan, and S. L. Sondhi, *Ann. Phys. (NY)* **313**, 497 (2004).
- ⁷⁸G. Y. Cho and J. E. Moore, *Ann. Phys. (NY)* **326**, 1515 (2011).
- ⁷⁹S. Elitzur, G. W. Moore, A. Schwimmer, and N. Seiberg, *Nucl. Phys. B* **326**, 108 (1989).
- ⁸⁰S. B. Treiman, E. Witten, R. Jackiw, and B. Zumino, *Current Algebra and Anomalies* (World Scientific, Singapore, 1985), p. 537.
- ⁸¹X.-G. Wen, *Adv. Phys.* **44**, 405 (1995).
- ⁸²Y. C. Kao and D. H. Lee, *Phys. Rev. B* **54**, 16903 (1996).
- ⁸³X.-G. Wen, *Phys. Rev. D* **88**, 045013 (2013).
- ⁸⁴Juven Wang and X.-G. Wen, arXiv:1307.7480.
- ⁸⁵Y. Hu, Y. Wan, and Y.-S. Wu, *Phys. Rev. B* **87**, 125114 (2013).
- ⁸⁶C.-Y. Huang, X. Chen, and F.-L. Lin, *Phys. Rev. B* **88**, 205124 (2013).
- ⁸⁷G. Vidal, arXiv:0912.1651.
- ⁸⁸M. Levin and C. P. Nave, *Phys. Rev. Lett.* **99**, 120601 (2007).
- ⁸⁹Z. C. Gu, M. Levin, and X.-G. Wen, *Phys. Rev. B* **78**, 205116 (2008).
- ⁹⁰J. I. Cirac and F. Verstraete, *J. Phys. A: Math. Gen.* **42**, 504004 (2009).
- ⁹¹F. Verstraete, V. Murg, and J. I. Cirac, *Adv. Phys.*, **57**, 143 (2008).
- ⁹²B. Swingle, arXiv:1209.0776.
- ⁹³X. G. Wen and A. Zee, *Phys. Rev. B* **46**, 2290 (1992).
- ⁹⁴<http://mathoverflow.net/questions/97448>