

## Symmetry-protected many-body Aharonov-Bohm effect

Luiz H. Santos<sup>1</sup> and Juven Wang<sup>1,2</sup>

<sup>1</sup>*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada N2L 2Y5*

<sup>2</sup>*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

(Received 5 November 2013; revised manuscript received 14 April 2014; published 16 May 2014)

It is known as a purely quantum effect that a magnetic flux affects the real physics of a particle, such as the energy spectrum, even if the flux does not interfere with the particle's path—the Aharonov-Bohm effect. Here we examine an Aharonov-Bohm effect on a many-body wave function. Specifically, we study this many-body effect on the gapless edge states of a bulk gapped phase protected by a global symmetry (such as  $\mathbb{Z}_N$ )—the symmetry-protected topological (SPT) states. The many-body analog of spectral shifts, the twisted wave function, and the twisted boundary realization are identified in this SPT state. An explicit lattice construction of SPT edge states is derived, and a challenge of gauging its non-onsite symmetry is overcome. Agreement is found in the twisted spectrum between a numerical lattice calculation and a conformal field theory prediction.

DOI: [10.1103/PhysRevB.89.195122](https://doi.org/10.1103/PhysRevB.89.195122)

PACS number(s): 03.65.Vf, 67.10.-j, 75.10.Pq

Mysteriously, an external magnetic flux can affect the physical properties of particles even without interfering directly on their paths. This is known as the Aharonov-Bohm (AB) effect [1]. For instance, a particle of charge  $q$  and mass  $m$  confined in a ring (parametrized by  $0 \leq \theta < 2\pi$ ) of radius  $a$  threaded with a flux  $\Phi_B$  [see Fig. 1(a)] would have its energy spectrum shifted as

$$E_n = \frac{1}{2ma^2} \left( n + \frac{\Phi_B}{\Phi_0} \right)^2, \quad n = 0, \pm 1, \dots, \quad (1)$$

where  $\Phi_0 = 2\pi/q$  is the quantum of magnetic flux and we adopt  $e = \hbar = c = 1$  units. One can dispose of the gauge potential in Schrödinger's equation of the wave function  $\psi(\theta)$  by a gauge transformation that changes the wave function to  $\tilde{\psi}(\theta) = \psi(\theta) \exp[iq \int^\theta A(\theta') d\theta']$ . So, the effect of the external flux can be enforced by the condition that the phase  $\tilde{\varphi}(\theta)$  of the new wave function satisfies a twisted boundary condition,

$$(1/2\pi) \oint d\theta \frac{\partial \tilde{\varphi}(\theta)}{\partial \theta} = \Phi_B / \Phi_0, \quad (2)$$

as the particle trajectory encloses the ring; thus, this twisted boundary condition implies a “branch cut” [see Fig. 1(b)]. We may refer to this twist effect as an “*Aharonov-Bohm twist*.” For electrons confined on a mesoscopic ring, for example, even though interactions are not negligible, the sensitivity of the system to the presence of the external flux can be rationalized as a single-particle phenomenon [2].

It is then opportune, as matter of principle, to ask whether such an AB effect can take place as an *intrinsically* interacting many-body phenomenon. More concretely, we ask whether the low-energy properties of such interacting systems display a response analogous to Eq. (1) when subject to a gauge perturbation and, in turn, how this effect is encoded in the “topology” (or boundary conditions) of the the wave-functional  $\Psi[\phi(x)]$ ; see Figs. 1(c) and 1(d). We shall refer to this as a many-body AB effect or twist.

In this paper, we show that two-dimensional (2D) symmetry-protected topological (SPT) states [3–5] offer a natural platform for observing the many-body AB effect. SPT states are quantum many-body states of matter with a finite gap to bulk excitations and no fractionalized degrees

of freedom. Due to a global symmetry, the system has the property that its edge states can only be gapped if a symmetry breaking occurs, either explicitly or spontaneously. So, in the absence of any symmetry breaking, the edge is described by robust edge excitations that cannot be localized due to weak symmetry-preserving disorder, in contrast to purely one-dimensional systems [6]. Assuming then that the edge states are in this gapless phase (an assumption that we will take throughout the paper), we shall demonstrate that the system will respond to the insertion of a gauge flux in a nontrivial way, whereas if the edge degrees of freedom were to become gapped, then they would be insensitive to the flux. We note that in 2D systems displaying the integer quantum Hall effect, the insertion of a flux also induces a nontrivial response of the chiral edge states [7]. In contrast to this situation, here we shall be concerned with 2D nonchiral SPT states for which gapless edge excitations, such as the single-particle modes on a ring, propagate in both directions. The spectrum of these gapless modes characterizes the low-energy properties of the system.

We approach this problem from two directions: (i) First, we study the response of the SPT state to the insertion of a gauge flux by means of a low-energy effective theory for the edge states, and we derive the change in the spectrum of edge states akin to Eq. (1). (ii) Complementarily, we show that the many-body AB effect derived in (i) can also be captured by formulating a lattice model describing the edge states. Twisted boundary conditions defined for these models are shown to account for the presence of a gauge flux, which we confirm numerically.

### I. MANY-BODY AHARONOV-BOHM EFFECT

To capture the essence of the AB effect on a symmetry-protected many-body wave function, we imagine threading a gauge flux through an effective 1D edge on one side of a 2D bulk SPT annulus (or cylinder). This many-body wave function on the 1D edge (parametrized by  $0 \leq x < L$ ) of SPT states is the analog of a single-body wave function of a particle in a ring. Since the bulk degrees of freedom are gapped, we concentrate on the low-energy properties on the edge described by a nonchiral Luttinger liquid action  $I_{\text{edge}}[\phi_I]$  [8,9]. To capture the gauge flux effect on a many-body wave




FIG. 1. (Color online) (a) and (c) Single- and many-body wave functions upon flux insertion, respectively. (b) and (d) Flux effect captured by twisted boundary conditions showing the associated branch cut.

function  $|\Psi\rangle$ , we formulate it in the path integral,

$$\begin{aligned} |\Psi(t_f)\rangle &= \sum_n |\Psi_n(t_f)\rangle \langle \Psi_n(t_f) | e^{-i \int_{t_i}^{t_f} H(t) dt} | \Psi(t_i)\rangle \\ &= \sum_n |\Psi_n(t_f)\rangle \int_{\phi_I(t_i)}^{\phi_I(t_f)} \mathcal{D}\phi_I e^{i \int_{t_i}^{t_f} (\mathcal{L}_{\text{edge}}[\phi_I] + (1/2\pi) \int q^I A \wedge d\phi_I)}, \end{aligned} \quad (3)$$

with  $\phi_I$  the intrinsic field on the edge. Our goal is to interpret this many-body AB twist  $(1/2\pi) \int q^I A \wedge d\phi_I$ . We anticipate the energy spectrum under the flux would be adjusted, and we aim to capture this “twist” effect on the energy spectrum. Below we focus on bosonic SPT states with  $\mathbb{Z}_N$  symmetry [8–12], with global symmetry transformation on the edge (see Appendix A for details on the field-theoretic input),

$$\mathcal{S}_N^{(p)} = e^{\frac{i}{N} \int_0^L dx \partial_x \phi_2 + p \int_0^L dx \partial_x \phi_1}, \quad (4)$$

where  $p \in \{0, \dots, N-1\}$  and  $(1/2\pi) \partial_x \phi_2(x)$  is the canonical momentum associated with  $\phi_1(x)$  [13].

The Lagrangian density associated with Eq. (3) reads

$$\begin{aligned} \mathcal{L}_{\text{edge}}[A] &= \frac{1}{4\pi} K_{IJ} \partial_t \phi_I \partial_x \phi_J - \mathcal{H}_f[\phi_I] \\ &\quad + \frac{1}{2\pi} q^I A_\mu \varepsilon^{\mu\nu} \partial_\nu \phi_I, \end{aligned} \quad (5)$$

where indices  $\mu, \nu \in \{0, 1\}$ ,  $I, J \in \{1, 2\}$ ,  $K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $\mathcal{H}_f[\phi_I]$  is the Hamiltonian density describing a free boson, and  $q^I = (q^1, q^2) = (1, p)$  specify the charges carried by the currents  $J_I^\mu = (1/2\pi) \varepsilon^{\mu\nu} \partial_\nu \phi_I$ . The right (left) -moving modes are described by  $\phi_{R,L} \propto \phi_1 \pm \phi_2$ .

Integrating the equations of motion of (5), with respect to  $\phi_I$ , along the boundary coordinate  $x$  in the presence of a static background  $\mathbb{Z}_N$  gauge flux configuration  $\oint_0^L dx A_1(x) = \frac{2\pi}{N}$  yields

$$(1/2\pi) \oint_0^L dx \partial_x \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = \frac{1}{N} \begin{pmatrix} p \\ 1 \end{pmatrix}. \quad (6)$$

(See Appendix B for an alternative derivation from a bulk-edge Chern-Simons approach.) Equation (6) represents the shift in winding modes of the edge boson fields, and it plays a role

analogous to the single-particle twisted boundary condition, Eq. (2). The spectrum of the central charge  $c = 1$  free boson at compactification radius  $R$  is labeled by the primary states  $|n, m\rangle$  ( $n, m \in \mathbb{Z}$ ) with scaling dimension

$$\Delta(n, m; R) = \frac{n^2}{R^2} + \frac{R^2 m^2}{4} \quad (7)$$

and momentum  $\mathcal{P}(n, m) = nm$  [14]. Then, according to Eq. (6), after the flux insertion, we derive the new spectrum (also see another related setting [15])

$$\tilde{\Delta}_N^{(p)}(n, m; R) = \frac{1}{R^2} \left( n + \frac{p}{N} \right)^2 + \frac{R^2}{4} \left( m + \frac{1}{N} \right)^2 \quad (8)$$

and momenta  $\tilde{\mathcal{P}}_N^{(p)}(n, m) = (n + \frac{p}{N})(m + \frac{1}{N})$  for each SPT state  $p \in \{0, \dots, N-1\}$ . Equations (6) and (8) capture the essence of the many-body AB effect analogous to Eqs. (1) and (2).

## II. EFFECTIVE LATTICE MODEL FOR THE EDGE OF SPT STATES

### A. Symmetry transformation and domain wall

The twist effect encoded in Eq. (8) comes from an effective low-energy description of the edge. We aim, as a complementary and perhaps more fundamental point of view, to capture this twist effect from a lattice model. As a first step in this program, we shall construct a global  $\mathbb{Z}_N$  symmetry transformation in terms of discrete degrees of freedom on the edge whose action reduces to Eq. (4) at long wavelengths. The hallmark of a nontrivial SPT state is that the symmetry transformation on the boundary cannot be in a tensor product form on each single site, i.e., it acts as a non-onsite symmetry transformation [3,4,16]. We propose the following ansatz for the symmetry transformation:

$$\begin{aligned} \mathcal{S}_N^{(p)} &\equiv \prod_{j=1}^M \tau_j \prod_{j=1}^M \exp \left\{ i \frac{p}{N} \left[ \frac{2\pi}{N} (\delta N_{\text{DW}})_{j,j+1} \right] \right\} \\ &\equiv \prod_{j=1}^M \tau_j \prod_{j=1}^M e^{\frac{i}{N} \mathcal{Q}_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})}, \end{aligned} \quad (9)$$

acting on a ring with  $M$  sites that we take to describe the 1D edge, with  $\sigma_{M+1} \equiv \sigma_1$ . At every site of the ring we consider  $\mathbb{Z}_N$  operators  $(\tau_j, \sigma_j)$ ,  $j = 1, \dots, M$ , satisfying  $\tau_j^N = \sigma_j^N = \mathbb{1}$  and  $\tau_j^\dagger \sigma_j \tau_j = \omega \sigma_j$ , where  $\omega \equiv e^{i2\pi/N}$ . We shall use the following representation:

$$\begin{aligned} \sigma_j &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \omega & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \omega^{N-1} \end{pmatrix}_j, \\ \tau_j &= \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & 0 & 0 & \cdots & 1 & 0 \end{pmatrix}_j. \end{aligned} \quad (10)$$

The overall symmetry transformation contains the onsite transformation part generated by the string of  $\tau$ 's and the "non-onsite domain wall (DW)" part  $(\delta N_{\text{DW}})_{j,j+1}$  between sites  $j$  and  $(j+1)$ . The ansatz form Eq. (9) has the property that  $\prod_{j=1}^M \tau_j$  and  $\prod_{j=1}^M e^{\frac{i}{N} Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})}$  commute, and the unitarity of  $S_N^{(p)}$  implies  $[Q_N^{(p)}]^\dagger = Q_N^{(p)}$ . It follows that

$$(S_N^{(p)})^N = \prod_{j=1}^M e^{i Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})}. \quad (11)$$

The construction above then naturally yields  $N$  distinct classes of  $\mathbb{Z}_N$  symmetry transformations, labeled by  $p \in \mathbb{Z}_N$ , upon imposing the following condition on the  $(N-1)$ th-order polynomial operator  $Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})$ :

$$e^{i Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})} = (\sigma_j^\dagger \sigma_{j+1})^p, \quad p = 0, \dots, N-1, \quad (12)$$

which guarantees (due to periodic boundary conditions) that  $(S_N^{(p)})^N = \mathbb{1}$ . The symmetry transformation in the trivial case corresponds to  $p = 0 \pmod{N}$ , for which  $\prod_j e^{\frac{i}{N} Q_N^{(p=0)}(\sigma_j^\dagger \sigma_{j+1})} = \mathbb{1}$ , while  $p \neq 0 \pmod{N}$  describe the other  $N-1$  nontrivial SPT classes. Identifying  $\sigma_j \sim e^{i \phi_1(j)}$ , then the domain-wall variable  $(\delta N_{\text{DW}})_{j,j+1}$  counts the number of units of the  $\mathbb{Z}_N$  angle between sites  $j$  and  $j+1$ , so  $(2\pi/N)(\delta N_{\text{DW}})_{j,j+1} = \phi_{1,j+1} - \phi_{1,j}$ , which produces the expected long-distance behavior of the symmetry transformation Eq. (4). Our ansatz nicely embodies two interpretations together, on both a continuum field theory and a discrete lattice model. The  $\mathbb{Z}_N$  symmetry transformations Eq. (9) that satisfy Eq. (12) can be explicitly written as

$$S_N^{(p)} = \prod_{j=1}^M \tau_j \prod_{j=1}^M e^{-i \frac{2\pi}{N^2} p \{ (\frac{N-1}{2}) \mathbb{1} + \sum_{k=1}^{N-1} \frac{(\sigma_j^\dagger \sigma_{j+1})^k}{(\omega^{k-1})} \}}. \quad (13)$$

In Ref. [16], the edge symmetry for  $\mathbb{Z}_N$  SPT states was proposed in terms of effective long-wavelength rotor variables. We emphasize that the construction of the edge symmetry transformations Eq. (13) described here does not rely on a long-wavelength description; rather, it can be viewed as a fully regularized symmetry transformation. In Appendixes C and D, we give explicit formulas for the  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  symmetry transformations, and we draw a connection between the lattice operators  $(\tau_j, \sigma_j)$  and quantum rotor variables.

### B. Lattice model

Having constructed all the classes of  $\mathbb{Z}_N$  symmetry transformations, Eq. (13), we now propose our translation invariant and  $\mathbb{Z}_N$ -symmetric lattice model Hamiltonians  $H_N^{(p)}$  on the edge of  $\mathbb{Z}_N$  SPT states, i.e.,

$$[H_N^{(p)}, T] = 0, \quad [H_N^{(p)}, S_N^{(p)}] = 0, \quad (14)$$

where  $T$  performs a translation by one lattice site. Our model Hamiltonian is (with  $\lambda_N^{(p)}$  a constant),

$$H_N^{(p)} = \lambda_N^{(p)} \sum_{j=1}^M h_{N,j}^{(p)} \equiv -\lambda_N^{(p)} \sum_{j=1}^M \sum_{\ell=0}^{N-1} (S_N^{(p)})^{-\ell} (\tau_j + \tau_j^\dagger) (S_N^{(p)})^\ell. \quad (15)$$




FIG. 2. (Color online) Spectrum of the SPT Hamiltonian Eq. (15) with respect to the lowest energy  $E_{N,0}^{(p)}$ , on a ring as a function of the lattice momentum  $k \in \mathbb{Z}$ . The first few primary states are labeled by  $(n, m)$ . (a) Spectrum of  $H_2^{(p=1)}$  with  $\lambda_2^{(p=1)} = 0.82$  and  $M = 20$  sites. (b) Spectrum of  $H_3^{(p=1,2)}$  with  $\lambda_3^{(p=1,2)} = 0.26$  and  $M = 12$  sites. The values of  $\lambda_N^{(p)}$  above guarantee a proper normalization so that states in the same conformal tower separated by  $\delta k = \pm 1$  are integer-spaced (up to finite size effects) (see Ref. [17]).

Notice that  $H_N^{(p)}$  is manifestly  $\mathbb{Z}_N$  symmetric since it is constructed from the superposition of  $\tau_j$  conjugated to all powers of  $S_N^{(p)}$ . In the trivial SPT case for which  $H_N^{(p=0)} \propto -\sum_{j=1}^M (\tau_j + \tau_j^\dagger)$ , the model gives a gapped and symmetry-preserving ground state. In Appendix C, we provide explicit forms of the nontrivial classes of SPT Hamiltonians for the  $N = 2$  and  $N = 3$  cases. We note that for the  $\mathbb{Z}_2$  case, our symmetry transformation and edge Hamiltonian are the same as that obtained in Ref. [9] (where the low-energy theory in terms of a nonchiral Luttinger liquid has been discussed), despite the fact that our method of constructing the symmetry is independent of that in Ref. [9] and provides a generalization for all  $\mathbb{Z}_N$  groups. It is noteworthy to mention that the authors of Ref. [9] argue that the edge of the  $\mathbb{Z}_2$  bosonic SPT state is generically unstable to symmetry-preserving perturbations. Nevertheless, we shall still study the model Hamiltonian (15) for the  $\mathbb{Z}_2$  as a means to address our numerical methods. A common feature of these Hamiltonian classes is the existence of combinations of terms such as  $\sigma_{j-1} \tau_j \sigma_{j+1}$  due to the non-onsite global symmetry. Their effect, as we shall see, is to give rise to a gapless spectrum. To understand their effect on the low-energy properties, we perform an exact diagonalization study of the nontrivial Hamiltonian classes Eq. (15) on finite systems.

In Fig. 2, we plot the lowest energy eigenvalues for the  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  nontrivial SPT states as a function of the lattice momentum  $k \in \mathbb{Z}$  defined by  $T = e^{i \frac{2\pi}{M} k}$ . The spectrum of  $H_2^{(1)}$  with  $M = 20$  sites shows very good agreement with the bosonic spectrum Eq. (7) at  $R = 2$ , with states being labeled by  $|n, m\rangle$ . The global  $\mathbb{Z}_2$  charges relative to the ground state were found to be  $e^{i\pi(n+m)}$  in accordance with Eq. (4) (we note that similar results have been obtained for the  $\mathbb{Z}_2$  case in Ref. [16]). For the  $\mathbb{Z}_3$  SPT states, which have not been investigated before, with  $M = 12$  sites, the spectra of  $H_3^{(1)}$  and  $H_3^{(2)}$  are identical [18]. Finite-size effects are more prominent than in the  $\mathbb{Z}_2$  case, but the overall structure of the spectrum is very similar, with the second and third states being degenerate with energy close to  $1/4$  and global  $\mathbb{Z}_3$  charges  $e^{\pm 2\pi i/3}$  (which we identify as the  $|n = \pm 1, m = 0\rangle$  states), suggesting the same spectrum Eq. (7) at  $R = 2$ .

In Appendix D, following the methods of Refs. [3,4,16], we show that the symmetry classes defined in Eq. (9) subject to condition Eq. (12) are related to all  $\mathbb{Z}_N$  3-cocycles of the group cohomology classification of 2D SPT states [3]. Thus, our lattice model completely realizes all  $N$  classes of  $\mathcal{H}^3(\mathbb{Z}_N, U(1)) = \mathbb{Z}_N$ , where  $p$  stands for the  $p$ th class in the third cohomology group.

### III. TWISTED BOUNDARY CONDITIONS AND TWISTED HAMILTONIAN ON THE LATTICE

We now seek to build a lattice model with twisted boundary conditions to capture the edge state spectral shift in the presence of a unit of  $\mathbb{Z}_N$  flux insertion. It is instructive to revisit the case of twisted boundary conditions where the symmetry transformation acts as an on-site symmetry. For the sake of concreteness, let us consider the one-dimensional quantum Ising model  $H_{\text{Ising}} = \sum_{j=1}^M (J\sigma_j^z\sigma_{j+1}^z + h\sigma_j^x)$  with global  $\mathbb{Z}_2$  symmetry  $\prod_{j=1}^M \sigma_j^x$ . The  $\mathbb{Z}_2$  twisted sector (or equivalently, in this case, the antiperiodic boundary condition sector) of the model is realized by flipping the sign of a pair interaction  $\sigma_k^z\sigma_{k+1}^z \rightarrow -\sigma_k^z\sigma_{k+1}^z$ , for some site  $k$ , while leaving all the other terms unchanged. If the Ising model is defined on an open line, the twist effect is implemented by conjugating the  $H_{\text{Ising}}$  with the operator  $\prod_{\ell \leq k} \sigma_\ell^x$ . When the model is defined on a ring, the same effect is obtained by defining a new translation operator  $\tilde{T} = T\sigma_k^x$  and demanding that the twisted Hamiltonian  $\tilde{H}_{\text{Ising}}$  commutes with  $\tilde{T}$ . It is straightforward to see that the twisted Ising Hamiltonian on a ring that commutes with  $\tilde{T}$  indeed has  $\sigma_k^z\sigma_{k+1}^z \rightarrow -\sigma_k^z\sigma_{k+1}^z$ . We also note that  $(\tilde{T})^M = \prod_{j=1}^M \sigma_j^x$  generates the  $\mathbb{Z}_2$  symmetry of  $H_{\text{Ising}}$ , which is also a symmetry of  $\tilde{H}_{\text{Ising}}$ .

We now generalize the construction above for the SPT edge Hamiltonians on a ring with a non-onsite symmetry by defining the unitary twisted lattice translation operator [19]

$$\tilde{T}^{(p)} = T e^{\frac{i}{N} Q_N^{(p)}(\sigma_M^\dagger \sigma_1)} \tau_1 \quad (16)$$

for each  $p \in \mathbb{Z}_N$  class, which incorporates the effect of the branch cut as in Fig. 1(d). The twisted Hamiltonian  $\tilde{H}_N^{(p)}$ , constructed from  $H_N^{(p)}$  of Eq. (15) and satisfying

$$[\tilde{H}_N^{(p)}, \tilde{T}^{(p)}] = 0, \quad (17)$$

reads (see Appendix C 2 for explicit results)

$$\tilde{H}_N^{(p)} = \lambda_N^{(p)} \sum_{j=1}^M \tilde{h}_{N,j}^{(p)}, \quad (18a)$$

$$\tilde{h}_{N,1}^{(p)} = \tau_1^\dagger \tau_2^\dagger h_{N,1}^{(p)} \tau_1 \tau_2,$$

$$\tilde{h}_{N,j}^{(p)} = h_{N,j}^{(p)} (2 \leq j \leq M-1), \quad (18b)$$

$$\tilde{h}_{N,M}^{(p)} = \tau_1^\dagger e^{-\frac{i}{N} Q_N^{(p)}(\sigma_M^\dagger \sigma_1)} h_{N,M}^{(p)} e^{\frac{i}{N} Q_N^{(p)}(\sigma_M^\dagger \sigma_1)} \tau_1.$$

Notice that, due to the intrinsic non-onsite term in the symmetry transformation,

$$\tilde{S}_N^{(p)} \equiv (\tilde{T}^{(p)})^M = e^{\frac{i}{N} [Q_N^{(p)}(\omega \sigma_M^\dagger \sigma_1) - Q_N^{(p)}(\sigma_M^\dagger \sigma_1)]} S_N^{(p)}, \quad (19)$$

the twisted nontrivial Hamiltonian breaks the SPT global symmetry [i.e.,  $[\tilde{H}_N^{(p)}, S_N^{(p)}] \neq 0$  if  $p \neq 0 \pmod{N}$ ], signaling




FIG. 3. (Color online) Spectrum of the twisted SPT Hamiltonian with respect to the lowest energy  $E_{N,0}^{(p)}$  on a ring as a function of the lattice momentum  $\tilde{k}$ , with the same values of  $\lambda_N^{(p)}$  as in Fig. 2. The first few primary states are labeled by  $(n,m)$ . (a) Spectrum of  $\tilde{H}_2^{(1)}$  with  $M = 20$  sites. (b) Spectrum of  $\tilde{H}_3^{(1)}$  (+) and  $\tilde{H}_3^{(2)}$  (x) with  $M = 12$  sites. (c) Comparison between  $\tilde{\Delta}_2^{(1)}$  (circles) and numerical results (+) plotted as a function of the momentum  $\tilde{P}_2^{(1)}$ . All points are twofold-degenerate. Red circles represent primary states, while the remaining points account for descendant states in the CFT spectrum. (d) Comparison between  $\tilde{\Delta}_3^{(1)}$  (circles) and data points (+) plotted in terms of the momentum  $\tilde{P}_3^{(1)}$ . Same for  $\tilde{\Delta}_3^{(2)}$  (squares) and data points (x) plotted in terms of the momentum  $\tilde{P}_3^{(2)}$ .

an anomaly effect [20,21]. (For a more systematic discussion of bosonic anomalies in the context of 2D SPT states, see Ref. [21].) However, in the trivial state, Eq. (19) yields  $\tilde{S}_N^{(p=0)} = S_N^{(p=0)} = \prod_{j=1}^M \tau_j$ , so that the twisted trivial Hamiltonian still commutes with the global  $\mathbb{Z}_N$  onsite symmetry, and the twisted effect is equivalent to the usual toroidal boundary conditions [17], as exemplified before for the antiperiodic boundary condition of the Ising model.

In Figs. 3(a) and 3(b), we display the low-energy spectrum of the twisted  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  SPT Hamiltonians with a  $\pi$ -flux and  $2\pi/3$ -flux, respectively, as a function of twisted lattice momentum  $\tilde{k}$  defined as  $\tilde{T} = e^{i \frac{2\pi}{M} \tilde{k}}$ . The eigenvalues of the primary states show very good agreement with  $\tilde{\Delta}_2^{(1)}(n,m; R=2)$  and  $\tilde{\Delta}_3^{(1,2)}(n,m; R=2)$  in Eq. (8), which we compare, in Figs. 3(c) and 3(d), by folding the spectrum so that the primary states are plotted as a function of the continuum momenta  $\tilde{P}_2^{(1)}(n,m)$  and  $\tilde{P}_3^{(1,2)}(n,m)$ . Our findings thus establish a relationship between the many-body AB effect in terms of both a long-wavelength description in the field theory as well as twisted boundary conditions in a lattice model.

### IV. SUMMARY

We have demonstrated that an intrinsically many-body realization of the Aharonov-Bohm phenomenon takes place on the edge of a 2D symmetry-protected many-body system in the presence of a background gauge flux. In our construction, we have assumed that the edge state is in a gapless phase and is described by a simple nonchiral Luttinger liquid action with one right- and one left-moving propagating mode carrying

different  $\mathbb{Z}_N$  charges [22], in which case the spectrum in the presence of a gauge flux displays quantization as Eq. (8) due to global symmetry protection ( $\mathbb{Z}_N$  symmetry in our work), analogous to the quantization of the energy spectrum of a superconducting ring due to the  $\mathbb{Z}_2$  symmetry inherent to superconductors [23]. The universal information carried by the counterpropagating edge modes is that they carry different  $\mathbb{Z}_N$  charges, which has been numerically verified for the  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  SPT classes in Fig. 2, where this difference is parametrized by the integer  $p \in \{1, \dots, N-1\}$  that characterizes the SPT class. This quantum number should remain invariant as long as the SPT order is not destroyed in the bulk. The offset in the charges carried by the right- and left-moving modes has then been shown to reflect itself in the edge spectrum according to Eq. (8) (where  $R$  is a nonuniversal parameter), which we have confirmed numerically in our model Hamiltonians for the  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  SPT classes in Fig. 3.

We have proposed general principles guiding the construction of the lattice Hamiltonians, Eqs. (15) and (18), of the bosonic  $\mathbb{Z}_N$ -symmetric SPT edge states for both the untwisted and twisted (without and with gauge fluxes) cases. The twisted spectra (i.e., with gauge flux) characterize all types of  $\mathbb{Z}_N$  bosonic anomalies [20,21], which naturally serve as ‘‘SPT invariants [5]’’ to detect and distinguish all  $\mathbb{Z}_N$  classes of SPT states numerically and experimentally. (See also the recent work in Refs. [21,24].)

Gauging a non-onsite symmetry of SPT has been noticed relating to the Ginsparg-Wilson (GW) fermion [25] approach of a lattice field theory problem [26]. We remark that our current work achieves gauging a non-onsite symmetry for a bosonic system, thus providing an important step in this direction. Whether our work can be extended to more general symmetry classes and to fermionic systems [such as U(1) symmetry in the GW fermion approach] is an open question, which we leave for future work.

## ACKNOWLEDGMENTS

We acknowledge useful discussions with X. Chen, L. Cincio, D. Gaiotto, T. Senthil, G. Vidal, A. Vishwanath, and X.-G. Wen. We are particularly grateful to Xie Chen on clarifying her work, and to Lukasz Cincio for introducing us to some of the numerical methods used here. We thank an anonymous referee for pointing out the possibility of edge reconstructions. L.H.S. would like to thank the ICPT South American Institute for Fundamental Research (ICTP-SAIFR), where part of this work was carried out via a partnership with the Perimeter Institute, for their hospitality. This research is supported by NSF Grants No. DMR-1005541, No. NSFC 11074140, and No. NSFC 11274192 (J.W.). Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development & Innovation (L.H.S. and J.W.).

## APPENDIX A: FIELD THEORY REALIZATION OF $\mathbb{Z}_N$ SPT STATES

In this appendix, we briefly review the field theory tool for topological states, especially symmetry-protected topological

(SPT) states, but with an emphasis on canonical quantization and how the global symmetry transformation  $\mathcal{S}_N^{(p)}$  on the edge is encoded in the canonical quantization.

### 1. Bulk and boundary actions

A general framework of categorizing and classifying Abelian topological orders, especially the SPT ones, in  $2+1$ D, makes use of Abelian  $K$ -matrix Chern-Simons theory [27]. We now derive the  $K$ -matrix construction for the SPT order, following the pioneering work of Refs. [8–13].

The intrinsic field theory description of SPT states, on a 2D spatial surface  $\mathcal{M}^2$ , is the Chern-Simons action

$$I_{\text{SPT}, \mathcal{M}^2} = \frac{1}{4\pi} \int dt d^2x K_{IJ} \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J, \quad (\text{A1})$$

where  $a$  is the intrinsic (or statistical) gauge field, and  $K$  is the  $K$  matrix, which categorizes the SPT orders. An SPT state is not intrinsically topologically ordered [3], so it has no topological degeneracy [13,27]. Ground-state degeneracy (GSD) of SPT on the torus is  $\text{GSD} = |\det K| = 1$  [8,13,27]; this suggests a constrained canonical form of  $K$  [8,12,13].

The SPT order is symmetry-protected, so tautologically its order is protected by a global symmetry. The novel features of SPT distinct from a trivial insulator are its symmetry-protected edge states on the boundary. The effective degree of freedom of its 1D edge,  $\partial\mathcal{M}^2$ , is the chiral bosonic field  $\phi$ , where  $\phi$  is meant to preserve gauge invariance on the bulk edge under gauge transformation of the field  $a$  [27]. The boundary action is

$$I_{\text{SPT}, \partial\mathcal{M}^2} = \frac{1}{4\pi} \int dt dx (K_{IJ} \partial_t \phi_I \partial_x \phi_J - V_{IJ} \partial_x \phi_I \partial_x \phi_J). \quad (\text{A2})$$

### 2. $\mathbb{Z}_N$ symmetry transformation

The  $\mathbb{Z}_N$  symmetry simply requires a rank-2  $K$  matrix, which exhausts all the group cohomology class,  $\mathcal{H}^3(\mathbb{Z}_N, U(1)) = \mathbb{Z}_N$ ,

$$K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (\text{A3})$$

The  $\mathbb{Z}_N$  symmetry transformation with a  $\mathbb{Z}_N$  angle specifies the group element  $g$  [8],

$$g_n : \delta\phi_{g_n} = \frac{2\pi}{N} n \begin{pmatrix} 1 \\ p \end{pmatrix}, \quad (\text{A4})$$

where  $p$  labels the  $\mathbb{Z}_N$  class of the cohomology group  $\mathcal{H}^3(\mathbb{Z}_N, U(1)) = \mathbb{Z}_N$ . Both  $n$  and  $p$  are module  $N$  as elements in  $\mathbb{Z}_N$ . It can be shown that under  $\phi_{g_n} \rightarrow \phi_{g_n} + \delta\phi_{g_n}$ , the action Eq. (A2) is invariant, and the  $\mathbb{Z}_N$  group structure is realized through  $g_n^N = \mathbb{1}$ . The construction of more general symmetry classes can be found in Refs. [8,12].

### 3. Canonical quantization

Here we go through the canonical quantization of the boson field  $\phi_I$ . For canonical quantization, we mean imposing a commutation relation between  $\phi_I$  and its conjugate momentum field  $\Pi_I(x) = \frac{\delta L}{\delta(\partial_t \phi_I)} = \frac{1}{2\pi} K_{IJ} \partial_x \phi_J$ . Because  $\phi_I$  is a compact

phase of a matter field, its bosonization contains both zero mode  $\phi_{0I}$  and winding momentum  $P_{\phi_I}$ , in addition to nonzero modes [13]:

$$\phi_I(x) = \phi_{0I} + K_{IJ}^{-1} P_{\phi_J} \frac{2\pi}{L} x + i \sum_{n \neq 0} \frac{1}{n} \alpha_{I,n} e^{-inx \frac{2\pi}{L}}. \quad (\text{A5})$$

The periodic boundary has size  $0 \leq x < L$ . First, we impose the commutation relation for zero mode and winding modes, and we generalize Kac-Moody algebra for nonzero modes:

$$[\phi_{0I}, P_{\phi_J}] = i \delta_{IJ}, \quad [\alpha_{I,n}, \alpha_{J,m}] = n K_{IJ}^{-1} \delta_{n,-m}. \quad (\text{A6})$$

We thus derive canonical quantized fields with the commutation relation,

$$[\phi_I(x_1), K_{IJ} \partial_x \phi_J(x_2)] = 2\pi i \delta_{IJ} \delta(x_1 - x_2), \quad (\text{A7})$$

$$[\phi_I(x_1), \Pi_J(x_2)] = i \delta_{IJ} \delta(x_1 - x_2). \quad (\text{A8})$$

The symmetry transformation of Eq. (A4) implies  $\phi_{g_n} \rightarrow \phi_{g_n} + \delta\phi_{g_n}$ :

$$\begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix} \rightarrow \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix} + \frac{2\pi}{N} \begin{pmatrix} 1 \\ p \end{pmatrix}. \quad (\text{A9})$$

It can be easily checked, using Eq. (A7), that

$$\mathcal{S}_N^{(p)} = e^{\frac{i}{N} (\int_0^L dx \partial_x \phi_2 + p \int_0^L dx \partial_x \phi_1)} \quad (\text{A10})$$

implements the global symmetry transformation

$$\mathcal{S}_N^{(p)} \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix} (\mathcal{S}_N^{(p)})^{-1} = \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix} + \frac{2\pi}{N} \begin{pmatrix} 1 \\ p \end{pmatrix}. \quad (\text{A11})$$

## APPENDIX B: TWISTED BOUNDARY CONDITION FROM A GAUGE FLUX INSERTION

Using the same formalism as in Appendix A, in this appendix we derive the twisted boundary condition due to a gauge flux insertion. Here we apply the canonical quantization method to formulate the effect of a gauge flux insertion through a cylinder (an analog of Laughlin thought experiments [7]) in terms of a twisted boundary condition effect. The *canonical quantization approach* here can be compared with the alternate *path-integral approach* motivated in the main text. The canonical quantization offers a solid view as to why the twisted boundary condition resulting from a gauge flux is a quantum effect. We will first present the bulk theory viewpoint, then the edge theory viewpoint.

### 1. Bulk theory

Our setting is an external adiabatic gauge flux insertion through a cylinder or annulus. Here the gauge field (such as the electromagnetic field) couples to (SPT or intrinsic) topologically ordered states, by a coupling charge vector  $q_I$ . The bulk term (here we recover the right dimension, while one

can set these to be  $e = \hbar = c = 1$  in the end)

$$I_{\text{bulk}} = \int_{\mathcal{M}} (c dt) d^2x \left[ \left( \frac{e^2}{\hbar} \right) \frac{K_{IJ}}{4\pi} \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J + e q^I A_\mu J_I^\mu \right], \quad (\text{B1})$$

where  $J_I^\mu$  is in a conserved current form

$$J_I^\mu = \left( \frac{e}{\hbar} \right) \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \partial_\nu a_{\rho,I}. \quad (\text{B2})$$

From the action, we derive the EOM,

$$J_J^\mu = -q_I \frac{e}{2\pi} K_{IJ}^{-1} \frac{c}{\hbar} \epsilon^{\mu\nu\rho} \partial_\nu a_{\rho}. \quad (\text{B3})$$

From the bulk theory side, an adiabatic flux  $\Delta\Phi_B$  induces an electric field  $E_x$  by the Faraday effect, causing a perpendicular current  $J_y$  flow to the boundary edge states. We can explicitly derive the flux effect from the Faraday-Maxwell equation in the 2 + 1D bulk,

$$\begin{aligned} q_I \Delta\Phi_B &= -q_I \int dt \int \vec{E} \cdot d\vec{l} = q_I \int dt dl_\mu c \epsilon^{\mu\nu\rho} \partial_\nu a_\rho \\ &= -\frac{2\pi}{e} K_{IJ} \hbar \int J_{y,J} dt dx = -\frac{2\pi}{e} K_{IJ} \frac{\hbar}{e} Q_J, \end{aligned} \quad (\text{B4})$$

which relates to the induced charge transported through the bulk, via the Hall effect mechanism. This is a derivation of the Laughlin flux insertion argument.  $Q$  is the total charge transported through the bulk, which should condense on the edge of the cylinder.

### 2. Edge theory

On the other hand, from the boundary theory side, the induced charge  $Q_I$  on the edge can be derived from the edge state dynamics affecting winding modes [see Eq. (A5)] by

$$Q_I = \int J_{\partial,I}^0 dx = -\oint_0^L \frac{e}{2\pi} \partial_x \phi_I dx = -e K_{IJ}^{-1} P_{\phi,J}. \quad (\text{B5})$$

Combine Eqs. (B4) and (B5),

$$q_I \Delta\Phi_B \Big/ \left( 2\pi \frac{\hbar}{e} \right) = \Delta P_{\phi,I}. \quad (\text{B6})$$

An equivalent interpretation is that the flux insertion twists the boundary conditions of the  $\phi_I$  field,

$$\begin{aligned} \frac{1}{2\pi} [\phi_I(L) - \phi_I(0)] &= \oint_0^L \frac{1}{2\pi} \partial_x \phi_I dx = K_{IJ}^{-1} \Delta P_{\phi,J} \quad (\text{B7}) \\ &= K_{IJ}^{-1} q_J \left[ \Delta\Phi_B \Big/ \left( 2\pi \frac{\hbar}{e} \right) \right]. \end{aligned} \quad (\text{B8})$$

In the  $\mathbb{Z}_N$  symmetry SPT case at hand, we should replace  $e$  to the condensate (order parameter) charge  $e^* = Ne$ . This affects the unit of  $\Delta\Phi_B$  as  $2\pi \frac{\hbar}{e^*}$ , so  $\Delta\Phi_B = 2\pi n \frac{\hbar}{Ne}$ , and the twisted boundary condition is

$$\frac{1}{2\pi} [\phi_I(L) - \phi_I(0)] = K_{IJ}^{-1} q_J (n/N). \quad (\text{B9})$$

Notice  $q_J$  is the crucial coupling in the global symmetry transformation, where we gauge it by minimal coupling to a gauge field  $A$  with a term  $q^I A_\mu J_I^\mu$ . Here  $q_J$  is realized by  $(1, p)$  from Eq. (A4), so inserting a unit  $\mathbb{Z}_N$  flux produces

$$[\phi_I(L) - \phi_I(0)] = \frac{2\pi}{N} \begin{pmatrix} p \\ 1 \end{pmatrix}. \quad (\text{B10})$$

In other words, while the global  $\mathbb{Z}_N$  symmetry transformation is realized by

$$\mathcal{S}_N^{(p)} \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix} (\mathcal{S}_N^{(p)})^{-1} = \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix} + \frac{2\pi}{N} \begin{pmatrix} 1 \\ p \end{pmatrix}, \quad (\text{B11})$$

the insertion of a unit  $\mathbb{Z}_N$  gauge flux implies the twisted boundary condition

$$\begin{pmatrix} \phi_1(L) \\ \phi_2(L) \end{pmatrix} = \begin{pmatrix} \phi_1(0) \\ \phi_2(0) \end{pmatrix} + \frac{2\pi}{N} \begin{pmatrix} p \\ 1 \end{pmatrix}. \quad (\text{B12})$$

Here  $\phi_1(x)$  is realized as the long-wavelength description of the rotor angle variable introduced in the main text, while its conjugate momentum is the angular momentum,

$$L_{\phi_1}(x) = \frac{1}{2\pi} \partial_x \phi_2(x), \quad (\text{B13})$$

where

$$[\phi_1(x_1), L_{\phi_1}(x_2)] = i\delta(x_1 - x_2). \quad (\text{B14})$$

We stress that our result is very different from a seemingly similar study in Ref. [5], where ‘‘the gauging process’’ is done by coupling the bulk state to an external gauge field  $A$ , and integrating out the intrinsic field  $a$ , to get an effective response theory description. However, the twisted boundary condition derived in [5] does not capture the dynamical effect on the edge under gauge flux insertion. Instead, in our case, we can capture this effect in Eq. (B12).

### APPENDIX C: FROM FIELD THEORY TO THE LATTICE MODEL

In this appendix, we provide our detailed lattice construction (with  $\mathbb{Z}_N$  symmetry) for both the untwisted and twisted (without and with gauge flux) cases. Here we motivate the construction of our lattice model from the field theory. Our lattice model uses the rotor eigenstate  $|\phi\rangle$  as a basis, where in  $\mathbb{Z}_N$  symmetry,  $\phi = n(2\pi/N)$ , where  $n$  is a  $\mathbb{Z}_N$  variable. The conjugate variable of  $\phi$  is the angular momentum  $L$ , which again is a  $\mathbb{Z}_N$  variable. The  $|\phi\rangle$  and  $|L\rangle$  eigenstates are related by a Fourier transformation,  $|\phi\rangle = \sum_{L=0}^{N-1} \frac{1}{\sqrt{N}} e^{iL\phi} |L\rangle$ .

#### 1. General Hamiltonian construction

The  $\mathbb{Z}_N$  class Hamiltonian may be realized by  $H_N^{(p)}$ , with  $p \in \mathbb{Z}_N$ ,

$$\begin{aligned} H_N^{(p)} &\equiv \lambda_N^{(p)} \sum_{j=1}^M h_{N,j}^{(p)} \\ &= -\lambda_N^{(p)} \sum_{j=1}^M \sum_{\ell=0}^{N-1} (\mathcal{S}_N^{(p)})^{-\ell} (\tau_j + \tau_j^\dagger) (\mathcal{S}_N^{(p)})^\ell, \quad (\text{C1}) \end{aligned}$$

with the parametrization

$$\tau_j = e^{i2\pi L_j/N}. \quad (\text{C2})$$

$\mathcal{S}_N^{(p)}$  is the  $\mathbb{Z}_N$  class of symmetry transformation

$$\begin{aligned} \mathcal{S}_N^{(p)} &\equiv \prod_{j=1}^M \tau_j \prod_{j=1}^M \exp \left\{ i \frac{p}{N} \left[ \frac{2\pi}{N} (\delta N_{\text{DW}})_{j,j+1} \right] \right\} \\ &\equiv \prod_{j=1}^M \tau_j \prod_{j=1}^M e^{\frac{i}{N} Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})}, \quad (\text{C3}) \end{aligned}$$

where

$$Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1}) = \sum_{a=0}^{N-1} q_{N,a}^{(p)} (\sigma_j^\dagger \sigma_{j+1})^a. \quad (\text{C4})$$

The hermiticity of  $Q_N^{(p)}$  combined with  $\sigma_j^\dagger \sigma_{j+1} \in \mathbb{Z}_N$  imply the constraint on the complex coefficients  $q_a$  (we drop indices  $p, N$  to simplify notation, and, in the following, an overbar denotes complex conjugation):

$$q_0 \in \mathbb{R}; q_a = \bar{q}_{N-a}, a = 1, \dots, (N-1)/2 \quad (\text{C5})$$

for odd  $N$ , while

$$q_0 \in \mathbb{R}; q_a = \bar{q}_{N-a}, a = 1, \dots, N/2 - 1; q_{N/2} \in \mathbb{R} \quad (\text{C6})$$

for even  $N$ . The coefficients of the  $(N-1)$ th-order polynomial operator  $Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})$  are determined, up to unimportant phases, by the condition

$$e^i Q_N^{(p)}(\sigma_j^\dagger \sigma_{j+1}) = (\sigma_j^\dagger \sigma_{j+1})^p, \quad p = 0, \dots, N-1. \quad (\text{C7})$$

The solution of Eq. (C7) can be systematically found for each value of  $p \in \mathbb{Z}_N$  giving rise to different symmetry classes. Below, for the sake of concreteness, we give explicit forms of the symmetry transformations and Hamiltonians for  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$  groups.

#### a. $\mathbb{Z}_2$ lattice model

For the  $N=2$  lattice model, in the  $|\phi\rangle$  basis, we have  $|\phi=0\rangle$ ,  $|\phi=\pi\rangle$ , and  $\omega = e^{i\pi} = -1$ ,

$$\langle \phi_a | e^{i\phi_j} | \phi_b \rangle = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}_{ab,j} = \sigma_{ab,j} = (\sigma_z)_{ab,j}, \quad (\text{C8})$$

$$\begin{aligned} \langle \phi_a | \tau_j | \phi_b \rangle &= \langle \phi_a | e^{i2\pi L_j/N} | \phi_b \rangle \\ &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}_{ab,j} = \tau_{ab,j} = (\sigma_x)_{ab,j}. \quad (\text{C9}) \end{aligned}$$

The symmetry transformation reads

$$\mathcal{S}_2^{(p)} = \prod_{j=1}^M \tau_j \prod_{j=1}^M e^{\frac{i}{2} Q_2^{(p)}(\sigma_j^z \sigma_{j+1}^z)}, \quad (\text{C10})$$

where we find, by imposing condition (C7),

$$Q_2^{(p)}(\sigma_j^z \sigma_{j+1}^z) = p \frac{\pi}{2} (1 - \sigma_j^z \sigma_{j+1}^z), \quad p = 0, 1. \quad (\text{C11})$$

With that, we obtain the Hamiltonian in the trivial class as

$$H_2^{(0)} = -2\lambda_2^{(0)} \sum_{j=1}^M \sigma_j^x, \quad (\text{C12})$$

and in the nontrivial SPT class as

$$H_2^{(1)} = -\lambda_2^{(1)} \sum_{j=1}^M (\sigma_j^x - \sigma_{j-1}^z \sigma_j^x \sigma_{j+1}^z). \quad (\text{C13})$$

### b. $\mathbb{Z}_3$ lattice model

For the  $N = 3$  lattice model, in the  $|\phi\rangle$  basis, we have  $|\phi = 0\rangle$ ,  $|\phi = 2\pi/3\rangle$ ,  $|\phi = 4\pi/3\rangle$ , and  $\omega = e^{i2\pi/3}$ ,

$$e^{i\phi_j} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}_j = \sigma_j, \quad (\text{C14})$$

$$e^{i2\pi L_j/N} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}_j = \tau_j. \quad (\text{C15})$$

The symmetry transformation reads

$$S_3^{(p)} = \prod_{j=1}^M \tau_j \prod_{j=1}^M e^{\frac{i}{3} Q_3^{(p)}(\sigma_j^\dagger \sigma_{j+1})}, \quad (\text{C16})$$

where we find, by imposing condition Eq. (C7),

$$\begin{aligned} Q_3^{(p)}(\sigma_j^\dagger \sigma_{j+1}) &= q_0^{(p)} + q_1^{(p)}(\sigma_j^\dagger \sigma_{j+1}) + \bar{q}_1^{(p)}(\sigma_j^\dagger \sigma_{j+1})^2, \\ q_0^{(p)} &= -p \frac{2\pi}{3}, \quad q_1^{(p)} = p \frac{\pi}{3}(1 + i/\sqrt{3}), \quad p = 0, 1, 2. \end{aligned} \quad (\text{C17})$$

With that, we obtain the Hamiltonian in the trivial class as

$$H_3^{(0)} = -3\lambda_3^{(0)} \sum_{j=1}^M (\tau_j + \tau_j^\dagger), \quad (\text{C18})$$

and in the nontrivial SPT classes  $p = 1, 2$  as

$$\begin{aligned} H_3^{(p)} &= -\lambda_3^{(p)} \sum_{j=1}^M \left\{ \tau_j \left[ \frac{5}{3} + \frac{\omega + \bar{\omega}}{3} (\sigma_{j-1}^\dagger \sigma_j + \sigma_{j-1} \sigma_j^\dagger) \right. \right. \\ &\quad \left. \left. + \left( \frac{(1+\omega)}{3} \sigma_j^\dagger \sigma_{j+1} + \frac{2\bar{\omega}}{3} \sigma_{j-1}^\dagger \sigma_{j+1} + \frac{2\omega}{3} \sigma_{j-1} \sigma_j^\dagger \sigma_{j+1} \right. \right. \right. \\ &\quad \left. \left. \left. + \text{H.c.} \right) \right] + \text{H.c.} \right\}. \end{aligned} \quad (\text{C19})$$

### c. $\mathbb{Z}_N$ lattice model

For a generic  $\mathbb{Z}_N$  lattice model, we have  $|\phi = 0\rangle, |\phi = 2\pi/N\rangle, \dots, |\phi = 2\pi(N-1)/N\rangle$ , and  $\omega = e^{i2\pi/N}$ . Applying the Fourier transformation,  $|\phi\rangle = \sum_{L=0}^{N-1} \frac{1}{\sqrt{N}} e^{iL\phi} |L\rangle$ , in the  $|\phi\rangle$  basis, we derive

$$e^{i\phi_j} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \omega & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & \omega^{N-1} \end{pmatrix}_j = \sigma_j, \quad (\text{C20})$$

$$e^{i2\pi L_j/N} = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & 0 & 0 & \cdots & 1 & 0 \end{pmatrix}_j = \tau_j. \quad (\text{C21})$$

Explicit forms of  $S_N^{(p)}$  can systematically be found by imposing condition (C7) for all  $p \in \mathbb{Z}_N$ . The explicit form of the symmetry transformation reads

$$S_N^{(p)} = \prod_{j=1}^M \tau_j \prod_{j=1}^M e^{-i \frac{2\pi}{N^2} p \left( \left( \frac{N-1}{2} \right) \mathbb{1} + \sum_{k=1}^{N-1} \frac{(\sigma_j^\dagger \sigma_{j+1})^k}{(\omega^k - 1)} \right)}. \quad (\text{C22})$$

## 2. Twisted boundary conditions on the lattice model

We clarify some of the steps leading to an edge Hamiltonian satisfying twisted boundary conditions accounting for the presence of one unit of background  $\mathbb{Z}_N$  gauge flux. The case with a general number of flux quanta can be equally worked out.

Let  $T$  be the lattice translation operator satisfying

$$T^\dagger X_j T = X_{j+1}, \quad j = 1, \dots, M \quad (\text{C23})$$

for any operator  $X_j$  on a ring such that  $X_{M+1} \equiv X_1$ . It satisfies  $T^M = \mathbb{1}$ . One can then immediately verify from Eqs. (C1) and (C3) that  $[S_N^{(p)}, T] = 0$  and

$$T^\dagger h_{N,j}^{(p)} T = h_{N,j+1}^{(p)}, \quad (\text{C24})$$

from which it follows that  $H_N^{(p)}$  in Eq. (C1) is translational invariant, i.e.,

$$[H_N^{(p)}, T] = 0. \quad (\text{C25})$$

Twisted boundary conditions are implemented by defining a modified translation operator

$$\tilde{T}^{(p)} = T e^{\frac{i}{N} Q_N^{(p)}(\sigma_M^\dagger \sigma_1)} \tau_1 \quad (\text{C26})$$

and seeking a twisted Hamiltonian

$$\tilde{H}_N^{(p)} \equiv \lambda_N^{(p)} \sum_{j=1}^M \tilde{h}_{N,j}^{(p)} \quad (\text{C27})$$

under the condition that

$$(\tilde{T}^{(p)})^\dagger \tilde{h}_{N,j}^{(p)} (\tilde{T}^{(p)}) = \tilde{h}_{N,j+1}^{(p)}, \quad (\text{C28})$$

which then yields

$$[\tilde{H}_N^{(p)}, \tilde{T}^{(p)}] = 0. \quad (\text{C29})$$



We now compute, iteratively,  $(\tilde{T}^{(p)})^M$  [where we use  $U_{M,1}^{(p)} = e^{\frac{i}{N} Q_N^{(p)}(\sigma_M^\dagger \sigma_1)}$ ],

$$\begin{aligned}
 (\tilde{T}^{(p)})^2 &= T U_{M,1}^{(p)} \tau_1 T U_{M,1}^{(p)} \tau_1 = T^2 U_{1,2}^{(p)} \tau_2 U_{M,1}^{(p)} \tau_1, \\
 &\vdots \\
 (\tilde{T}^{(p)})^M &= \underbrace{T^M}_{=1} (U_{M-1,M}^{(p)} \tau_M) (U_{M-2,M-1}^{(p)} \tau_{M-1}) \cdots (U_{1,2}^{(p)} \tau_2) (U_{M,1}^{(p)} \tau_1) \\
 &= (U_{M-1,M}^{(p)} U_{M-2,M-1}^{(p)} \cdots U_{1,2}^{(p)}) \tau_M U_{M,1}^{(p)} (\tau_{M-1} \tau_{M-2} \cdots \tau_1) \\
 &= \left( \prod_{j=1}^M U_{j,j+1}^{(p)} \right) (U_{M,1}^{(p)})^{-1} \tau_M U_{M,1}^{(p)} \tau_M^\dagger \left( \prod_{j=1}^M \tau_j \right) \\
 &= \left( \prod_{j=1}^M U_{j,j+1}^{(p)} \right) e^{-\frac{i}{N} Q_N^{(p)}(\sigma_M^\dagger \sigma_1)} e^{\frac{i}{N} Q_N^{(p)}(\omega \sigma_M^\dagger \sigma_1)} \left( \prod_{j=1}^M \tau_j \right). \tag{C30}
 \end{aligned}$$

Thus we obtain

$$\tilde{S}_N^{(p)} \equiv (\tilde{T}^{(p)})^M = e^{\frac{i}{N} [Q_N^{(p)}(\omega \sigma_M^\dagger \sigma_1) - Q_N^{(p)}(\sigma_M^\dagger \sigma_1)]} S_N^{(p)}. \tag{C31}$$

Notice that in a trivial case ( $p = 0$ ), the relation

$$\tilde{S}_N^{(p=0)} = (\tilde{T}^{(p=0)})^M = \prod_{j=1}^N \tau_j = S_N^{(p=0)} \tag{C32}$$

reduces to to the global *onsite* symmetry  $S_N^{(p=0)}$ . In this case, the twisted Hamiltonian commutes with the onsite symmetry since  $0 = [\tilde{H}_N^{(p=0)}, (\tilde{T}^{(p=0)})^M] = [\tilde{H}_N^{(p=0)}, S_N^{(p=0)}]$ , and the states in the twisted sector are still labeled by the global trivial  $\mathbb{Z}_N$  charges, corresponding to usual toroidal boundary conditions. In a nontrivial SPT state ( $p \neq 0$ ), however, we find  $0 = [\tilde{H}_N^{(p)}, (\tilde{T}^{(p)})^M] \neq [\tilde{H}_N^{(p)}, S_N^{(p)}]$ , so that the twisted Hamiltonian breaks the nontrivial  $\mathbb{Z}_N$  SPT global symmetry. We should regard  $(\tilde{T}^{(p)})^M \equiv \tilde{S}_N^{(p)}$  as a new *twisted symmetry transformation* incorporating the gauge flux effect on the branch cut.

#### a. Twisted boundary conditions for the $\mathbb{Z}_2$ SPT state

We now explicitly work out the twisted Hamiltonian for the nontrivial  $\mathbb{Z}_2$  SPT state and later mention the general  $\mathbb{Z}_N$  case. The global SPT symmetry reads

$$S_2^{(1)} = \prod_{j=1}^M \sigma_j^x \prod_{j=1}^M e^{\frac{i}{2} Q_2^{(1)}(\sigma_j^z \sigma_{j+1}^z)} = \prod_{j=1}^M \sigma_j^x \prod_{j=1}^M e^{\frac{i\pi}{4} [1 - \sigma_j^z \sigma_{j+1}^z]}. \tag{C33}$$

Define  $U_{j,j+1} \equiv e^{\frac{i\pi}{4} [1 - \sigma_j^z \sigma_{j+1}^z]}$ . Then the nontrivial SPT Hamiltonian  $H = \sum_{j=1}^M h_j$  (we drop overall constants for simplicity) is

$$\begin{aligned}
 h_j &= \sigma_j^x + S^{-1} \sigma_j^x S \\
 &= \sigma_j^x + U_{j-1,j}^{-1} U_{j,j+1}^{-1} \sigma_j^x U_{j-1,j} U_{j,j+1} \\
 &= \sigma_j^x - \sigma_{j-1}^z \sigma_j^x \sigma_{j+1}^z \tag{C34}
 \end{aligned}$$

for  $j = 1, \dots, M$ . The modified translation operator reads

$$\tilde{T} = T U_{M,1} \sigma_1^x = T e^{\frac{i\pi}{4} [1 - \sigma_M^z \sigma_1^z]} \sigma_1^x. \tag{C35}$$

We seek a twisted Hamiltonian  $\tilde{H} \equiv \sum_{j=1}^M \tilde{h}_j$  that commutes with  $\tilde{T}$ . It is a simple exercise to check that

$$\begin{aligned}
 \tilde{T}^\dagger h_2 \tilde{T} &= h_3, \\
 \tilde{T}^\dagger h_3 \tilde{T} &= h_4, \\
 &\vdots \\
 \tilde{T}^\dagger h_{M-2} \tilde{T} &= h_{M-1}. \tag{C36}
 \end{aligned}$$

We are then led to identify

$$\tilde{h}_j \equiv h_j, \quad j = 2, \dots, M-1. \tag{C37}$$

We now consider

$$\tilde{h}_M \equiv \tilde{T}^\dagger h_{M-1} \tilde{T} = \sigma_1^x U_{M,1}^{-1} h_M U_{M,1} \sigma_1^x \tag{C38}$$

and

$$\begin{aligned}
 \tilde{h}_1 &\equiv \tilde{T}^\dagger \tilde{h}_M \tilde{T} \\
 &= \sigma_1^x U_{M,1}^{-1} (\sigma_2^x U_{1,2}^{-1} h_1 U_{1,2} \sigma_2^x) U_{M,1} \sigma_1^x \\
 &= \sigma_1^x \sigma_2^x \underbrace{(U_{M,1}^{-1} U_{1,2}^{-1} h_1 U_{1,2} U_{M,1})}_{=h_1} \sigma_1^x \sigma_2^x \\
 &= \sigma_1^x \sigma_2^x h_1 \sigma_1^x \sigma_2^x. \tag{C39}
 \end{aligned}$$

Now it remains to be shown that  $\tilde{T}^\dagger \tilde{h}_1 \tilde{T} = \tilde{h}_2 = h_2$ . And indeed

$$\begin{aligned}
 \tilde{T}^\dagger \tilde{h}_1 \tilde{T} &= \sigma_1^x U_{M,1}^{-1} (\sigma_2^x \sigma_3^x h_2 \sigma_2^x \sigma_3^x) U_{M,1} \sigma_1^x \\
 &= \sigma_1^x \sigma_2^x \sigma_3^x h_2 \sigma_1^x \sigma_2^x \sigma_3^x \\
 &= h_2. \tag{C40}
 \end{aligned}$$

So we have found new terms  $\tilde{h}_j$  such that  $\tilde{T}^\dagger \tilde{h}_j \tilde{T} = \tilde{h}_{j+1}$ , thus implying that  $[\tilde{T}, \tilde{H}] = 0$ .

Explicitly, the twisted Hamiltonian for the  $\mathbb{Z}_2$  nontrivial SPT state reads

$$\tilde{H} = \sum_{j=1}^M \tilde{h}_j, \tag{C41a}$$

where

$$\begin{aligned} \tilde{h}_1 &= \sigma_1^x \sigma_2^x h_1 \sigma_1^x \sigma_2^x = \sigma_1^x + \sigma_M^z \sigma_1^x \sigma_2^z, \\ \tilde{h}_2 &= h_2 = \sigma_2^x - \sigma_1^z \sigma_2^x \sigma_3^z, \\ &\vdots \\ \tilde{h}_{M-1} &= h_{M-1} = \sigma_{M-1}^x - \sigma_{M-2}^z \sigma_{M-1}^x \sigma_M^z, \\ \tilde{h}_M &= \sigma_1^x U_{M,1}^{-1} h_M U_{M,1} \sigma_1^x = \sigma_M^y \sigma_1^z + \sigma_{M-1}^z \sigma_M^y. \end{aligned} \quad (\text{C41b})$$

**b. Twisted boundary conditions for the  $\mathbb{Z}_N$  SPT state**

Generalization to the  $\mathbb{Z}_N$  case follows very similar lines to the  $\mathbb{Z}_2$  case above. We have for the twisted Hamiltonian (again we drop overall constants)

$$\tilde{H}_N^{(p)} = \sum_{j=1}^M \tilde{h}_{N,j}^{(p)}, \quad (\text{C42a})$$

where

$$\begin{aligned} \tilde{h}_{N,1}^{(p)} &= \tau_1^\dagger \tau_2^\dagger h_{N,1}^{(p)} \tau_1 \tau_2, \\ \tilde{h}_{N,2}^{(p)} &= h_{N,2}^{(p)}, \\ &\vdots \\ \tilde{h}_{N,M-1}^{(p)} &= h_{N,M-1}^{(p)}, \\ \tilde{h}_{N,M}^{(p)} &= \tau_1^\dagger (U_{M,1}^{(p)})^{-1} h_{N,M}^{(p)} U_{M,1}^{(p)} \tau_1, \end{aligned} \quad (\text{C42b})$$

where  $U_{M,1}^{(p)} = e^{\frac{i}{N} \mathcal{Q}_N^{(p)}(\sigma_M^\dagger \sigma_1)}$ . One can easily verify that Eqs. (C28) and (C29) are satisfied.

**APPENDIX D: CORRESPONDENCE IN GROUP COHOMOLOGY AND NONTRIVIAL 3-COCYCLES FROM A MPS PROJECTIVE REPRESENTATION**

In this appendix, we match each SPT class of our lattice construction to the 3-cocycles in the group cohomology classification. Importantly, we notice that the non-onsite piece in  $S_N^{(p)}$  is

$$U_{j,j+1} \equiv e^{i \mathcal{Q}_N^{(p)}(\sigma_j^\dagger \sigma_{j+1})} = \exp \left[ \frac{i}{N} \sum_{a=0}^{N-1} q_a (\sigma_j^\dagger \sigma_{j+1})^a \right] \quad (\text{D1})$$

$$\equiv \exp \left\{ i \frac{p}{N} \left[ \frac{2\pi}{N} (\delta N_{\text{DW}})_{j,j+1} \right] \right\}. \quad (\text{D2})$$

We seek a quantum rotor description of the above form. We claim that

$$U_{j,j+1} = \exp \left[ i \frac{p}{N} (\phi_{1,j+1} - \phi_{1,j})_r \right], \quad (\text{D3})$$

which is equivalent to (i) the domain-wall picture using rotor angle variables [here  $(\phi_{1,j+1} - \phi_{1,j})_r$ , where the subscript  $r$  means that we take the module  $2\pi$  on the angle [16]], and (ii) the field theory formalism in Eq. (A10).

The reason is as follows: as we mention in the  $p$ th case of  $\mathbb{Z}_N$  class, we impose the constraint

$$U_{j,j+1}^N = (\sigma_j^\dagger \sigma_{j+1})^p \quad (\text{D4})$$

to solve the polynomial ansatz  $\sum_{a=0}^{N-1} q_a (\sigma_j^\dagger \sigma_{j+1})^a$ . This is equivalent to the fact that

$$\begin{aligned} U_{j,j+1}^N &= (\sigma_j^\dagger \sigma_{j+1})^p = (\exp[i\phi_{1,j}]^\dagger \exp[i\phi_{1,j+1}])^p \\ &= \exp[ip(\phi_{1,j+1} - \phi_{1,j})_r], \end{aligned} \quad (\text{D5})$$

since  $\exp[i\phi_{1,j}]_{ab} = \langle \phi_a | e^{i\phi_j} | \phi_b \rangle = \sigma_{ab,j}$ . Therefore, the domain-wall variable  $(\delta N_{\text{DW}})_{j,j+1}$  indeed counts the number of units of  $\mathbb{Z}_N$  angle between sites  $j$  and  $j+1$ , so  $(2\pi/N)(\delta N_{\text{DW}})_{j,j+1} = \phi_{1,j+1} - \phi_{1,j}$ . We thus have shown Eq. (D3), and we have confirmed that our approach of lattice regularization is indeed a rotor realization in Ref. [16] with the same symmetry transformation  $S_N^{(p)}$ , but it captures much more than the low-energy rotor model there.

The argument on nontrivial 3-cocycles from matrix product states (MPSs) projective representation follows closely Ref. [16]. We start by writing the symmetry transformation  $S_N^{(p)}$  in terms of the rotor variable; this is achieved based on the mapping derived above. So

$$\begin{aligned} S_N^{(p)} &\equiv \prod_{j=1}^M \tau_j \prod_{j=1}^M U_{j,j+1}^{(p)} \\ &= \prod_j e^{i2\pi L_j/N} \exp \left[ i \frac{p}{N} (\phi_{1,j+1} - \phi_{1,j})_r \right]. \end{aligned} \quad (\text{D7})$$

We then formulate  $S_N^{(p)}$  as the MPS with the form

$$S_N^{(p)} = \sum_{\{j,j'\}} \text{tr} [T_{\alpha_1 \alpha_2}^{j_1 j'_1} T_{\alpha_2 \alpha_3}^{j_2 j'_2} \cdots T_{\alpha_M \alpha_1}^{j_M j'_M}] |j'_1, \dots, j'_M\rangle \langle j_1, \dots, j_M|. \quad (\text{D8})$$

Here  $j_1, j_2, \dots, j_M$  and  $j'_1, j'_2, \dots, j'_M$  are labeled by input or output physical eigenvalues (here  $\mathbb{Z}_N$  angle), and the subscripts  $1, 2, \dots, M$  are the physical site indices. There are also inner indices  $\alpha_1, \alpha_2, \dots, \alpha_M$  that are traced in the end. Summing over the entire operation from  $\{j, j'\}$  indices is supposed to reproduce the symmetry transformation operator  $S_N^{(p)}$ . This tensor  $T$  is suggested [16] to be (with the  $\mathbb{Z}_N$  angle element  $\frac{2\pi k}{N}$ )

$$\begin{aligned} (T^{\phi_{\text{in}}, \phi_{\text{out}}})_{\varphi_\alpha, \varphi_\beta, N}^{(p)} &\left( \frac{2\pi k}{N} \right) \\ &= \delta \left( \phi_{\text{out}} - \phi_{\text{in}} - \frac{2\pi k}{N} \right) \\ &\times \int d\varphi_\alpha d\varphi_\beta |\varphi_\beta\rangle \langle \varphi_\alpha | \delta(\varphi_\beta - \phi_{\text{in}}) e^{ipk(\varphi_\alpha - \phi_{\text{in}})_r/N}. \end{aligned} \quad (\text{D9})$$

We verify the tensor  $T$  by computing  $S_N^{(p)}$ ,

$$S_N^{(p)} = \sum_{\{j,j'\}} \text{tr} [T_{\varphi_{\alpha_1} \varphi_{\alpha_2}}^{\phi_{\text{in}}^1, \phi_{\text{out}}^1} T_{\varphi_{\alpha_2} \varphi_{\alpha_3}}^{\phi_{\text{in}}^2, \phi_{\text{out}}^2} \cdots T_{\varphi_{\alpha_M} \varphi_{\alpha_1}}^{\phi_{\text{in}}^M, \phi_{\text{out}}^M}] |\phi_{\text{out}}^1, \phi_{\text{out}}^2, \dots, \phi_{\text{out}}^M\rangle \langle \phi_{\text{in}}^1, \phi_{\text{in}}^2, \dots, \phi_{\text{in}}^M| \quad (\text{D10})$$

$$= e^{i \frac{p}{N} [(\phi_{\text{in}}^2 - \phi_{\text{in}}^1)_r + (\phi_{\text{in}}^3 - \phi_{\text{in}}^2)_r + \dots + (\phi_{\text{in}}^M - \phi_{\text{in}}^{M-1})_r]} \left| \phi_{\text{in}}^1 + \frac{2\pi}{N}, \phi_{\text{in}}^2 + \frac{2\pi}{N}, \dots, \phi_{\text{in}}^M + \frac{2\pi}{N} \right\rangle \langle \phi_{\text{in}}^1, \phi_{\text{in}}^2, \dots, \phi_{\text{in}}^M | \quad (\text{D11})$$

$$= e^{i \frac{p}{N} [\sum_{j=1}^M (\phi_{\text{in}}^{j+1} - \phi_{\text{in}}^j)_r]} \left| \dots, \phi_{\text{in}}^j + \frac{2\pi}{N}, \dots \right\rangle \langle \dots, \phi_{\text{in}}^j, \dots | \quad (\text{D12})$$

$$= \prod_j \exp \left[ i \frac{p}{N} (\phi_{1,j+1} - \phi_{1,j})_r \right] \prod_j e^{i 2\pi L_j / N}, \quad (\text{D13})$$

which justifies the claim for MPSs of  $S_N^{(p)}$ .

To find out the projective representation  $e^{i\theta(g_1, g_2, g_3)}$  of these tensors  $T(g_1), T(g_2), T(g_3)$  acting on three neighbored sites, we follow the fact that

$$P_{g_1, g_2}^\dagger T(g_1) T(g_2) P_{g_1, g_2} = T(g_1 g_2) \quad (\text{D14})$$

and contracting the three neighbored-site tensors in two different orders,

$$(P_{g_1, g_2} \otimes I_3) P_{g_1, g_2, g_3} \simeq e^{i\theta(g_1, g_2, g_3)} (I_1 \otimes P_{g_2, g_3}) P_{g_1, g_2, g_3}. \quad (\text{D15})$$

Here  $\simeq$  means the equivalence is up to a projection out of an unparallel state transformation.

To derive  $P_{g_1, g_2}$ , notice that  $P_{g_1, g_2}$  inputs one state and output two states. This has the expected form

$$P_{N, m_1, m_2}^{(p)} = \int d\phi_{\text{in}} \left| \phi_{\text{in}} + \frac{2\pi}{N} m_2 \right\rangle |\phi_{\text{in}}\rangle \langle \phi_{\text{in}}| \times e^{-ip\phi_{\text{in}}[m_1 + m_2 - (m_1 + m_2)_N]/N}, \quad (\text{D16})$$

where  $(m_1 + m_2)_N$  with subscript  $N$  means taking the value module  $N$ .

To derive  $\theta(g_1, g_2, g_3)$ , we start by contracting  $T_N^{(p)}(m_1)$  and  $T_N^{(p)}(m_2)$  first, and then the combined tensor contracts with  $T_N^{(p)}(m_3)$  give

$$(P_{g_1, g_2} \otimes I_3) P_{g_1, g_2, g_3} = \int d\phi_{\text{in}} \left| \phi_{\text{in}} + \frac{2\pi}{N} (m_2 + m_3) \right\rangle \left| \phi_{\text{in}} + \frac{2\pi}{N} m_3 \right\rangle |\phi_{\text{in}}\rangle \langle \phi_{\text{in}}|$$

$$\times e^{-ip\phi_{\text{in}}[m_1 + m_2 + m_3 - (m_1 + m_2 + m_3)_N]} \times e^{-ip \frac{2\pi}{N} m_3 \frac{m_1 + m_2 - (m_1 + m_2)_N}{N}}, \quad (\text{D17})$$

the form of which inputs one state  $\langle \phi_{\text{in}} |$  and outputs three states  $|\phi_{\text{in}} + \frac{2\pi}{N} (m_2 + m_3)\rangle$ ,  $|\phi_{\text{in}} + \frac{2\pi}{N} m_3\rangle$ , and  $|\phi_{\text{in}}\rangle$ .

On the other hand, one can contract  $T_N^{(p)}(m_2)$  and  $T_N^{(p)}(m_3)$  first, and then the combined tensor contracted with  $T_N^{(p)}(m_1)$  gives

$$(I_1 \otimes P_{g_2, g_3}) P_{g_1, g_2, g_3} = \int d\phi_{\text{in}} \left| \phi_{\text{in}} + \frac{2\pi}{N} (m_2 + m_3) \right\rangle \left| \phi_{\text{in}} + \frac{2\pi}{N} m_3 \right\rangle |\phi_{\text{in}}\rangle \langle \phi_{\text{in}}| \times e^{-ip\phi_{\text{in}}[m_1 + m_2 + m_3 - (m_1 + m_2 + m_3)_N]}, \quad (\text{D18})$$

again the form of which inputs one state  $\langle \phi_{\text{in}} |$  and outputs three states  $|\phi_{\text{in}} + \frac{2\pi}{N} (m_2 + m_3)\rangle$ ,  $|\phi_{\text{in}} + \frac{2\pi}{N} m_3\rangle$ , and  $|\phi_{\text{in}}\rangle$ . From Eqs. (D15), (D17), and (D18), we derive

$$e^{i\theta(g_1, g_2, g_3)} = e^{-ip \frac{2\pi}{N} m_3 \frac{m_1 + m_2 - (m_1 + m_2)_N}{N}}, \quad (\text{D19})$$

which indeed is the 3-cocycle in the third cohomology group  $\mathcal{H}^3(\mathbb{Z}_N, U(1)) = \mathbb{Z}_N$ . We thus verify that the projective representation  $e^{i\theta(g_1, g_2, g_3)}$  from MPS tensors corresponds to the group cohomology approach [3]. This demonstrates that our lattice model construction completely maps to all classes of SPT, which is what we aimed for.

[1] W. Ehrenberg and R. E. Siday, *Proc. Phys. Soc. London, Sect. B* **62**, 8 (1949); Y. Aharonov and D. Bohm, *Phys. Rev.* **115**, 485 (1959).  
 [2] A. G. Aronov and Y. V. Sharvin, *Rev. Mod. Phys.* **59**, 755 (1987).  
 [3] X. Chen, Z. C. Gu, Z. X. Liu and X. G. Wen, *Phys. Rev. B* **87**, 155114 (2013).  
 [4] X. Chen, Z.-X. Liu, and X.-G. Wen, *Phys. Rev. B* **84**, 235141 (2011).  
 [5] X. G. Wen, *Phys. Rev. B* **89**, 035147 (2014).  
 [6] P. A. Lee and T. V. Ramakrishnan, *Rev. Mod. Phys.* **57**, 287 (1985).  
 [7] R. B. Laughlin, *Phys. Rev. B* **23**, 5632(R) (1981).  
 [8] Y. M. Lu and A. Vishwanath, *Phys. Rev. B* **86**, 125119 (2012).  
 [9] M. Levin and Z. C. Gu, *Phys. Rev. B* **86**, 115109 (2012).  
 [10] L. Y. Hung and Y. Wan, *Phys. Rev. B* **87**, 195103 (2013).  
 [11] M. Cheng and Z.-C. Gu, *Phys. Rev. Lett.* **112**, 141602 (2014).

[12] P. Ye and J. Wang, *Phys. Rev. B* **88**, 235109 (2013).  
 [13] J. Wang and X.-G. Wen, [arXiv:1212.4863](https://arxiv.org/abs/1212.4863).  
 [14] P. Di Francesco, P. Mathieu, and D. Senechal, *Conformal Field Theory, Graduate Texts in Contemporary Physics* (Springer-Verlag, New York, 1996).  
 [15] O. M. Sule, X. Chen, and S. Ryu, *Phys. Rev. B* **88**, 075125 (2013).  
 [16] X. Chen and X. G. Wen, *Phys. Rev. B* **86**, 235135 (2012).  
 [17] M. Henkel, *Conformal Invariance and Critical Phenomena* (Springer-Verlag, Berlin, Heidelberg, 1999).  
 [18] A side comment is that our Hamiltonian Eq. (15) is the same for all  $\mathbb{Z}_N$  classes, although the symmetry transformation Eq. (9) is realized differently. It is analogous to the field theory approach in [8, 12] choosing the same canonical form of the Lagrangian but realizing the symmetry transformation differently.  
 [19] For  $s$  units of flux, the generalization follows  $\tilde{T}^{(p)} = T(e^{i \frac{p}{N} \mathcal{Q}_N^{(p)} (\sigma_M^\dagger \sigma_1)})^s \tau_1^s$ .

- [20] X. G. Wen, *Phys. Rev. D* **88**, 045013 (2013).
- [21] J. Wang, L. H. Santos, and X.-G. Wen, [arXiv:1403.5256](https://arxiv.org/abs/1403.5256) (cond-mat.str-el).
- [22] When edge reconstructions give rise to new gapless degrees of freedom, the analysis presented here for a single pair of modes should be extended to all the pairs of counterpropagating modes, in which case the form of the spectrum would depend on the details of the reconstruction itself, a problem that is beyond the scope of this paper.
- [23] B. S. Deaver and W. M. Fairbank, *Phys. Rev. Lett.* **7**, 43 (1961); N. Byers and C. N. Yang, *ibid.* **7**, 46 (1961); R. Doll and M. Nbauer, *ibid.* **7**, 51 (1961).
- [24] M. P. Zaletel, [arXiv:1309.7387](https://arxiv.org/abs/1309.7387).
- [25] P. H. Ginsparg and K. G. Wilson, *Phys. Rev. D* **25**, 2649 (1982).
- [26] Juven Wang and X.-G. Wen, [arXiv:1307.7480](https://arxiv.org/abs/1307.7480).
- [27] X.-G. Wen, *Quantum Field Theory of Many-Body Systems—From the Origin of Sound to an Origin of Light and Electrons* (Oxford University Press, Oxford, 2004).