

Quantum fluctuations beyond the Gutzwiller approximation

Michele Fabrizio

International School for Advanced Studies (SISSA), Via Bonomea 265, I-34136 Trieste, Italy

(Received 12 December 2016; revised manuscript received 10 February 2017; published 28 February 2017)

We present a simple scheme to evaluate linear response functions including quantum fluctuation corrections on top of the Gutzwiller approximation. The method is derived for a generic multiband lattice Hamiltonian without any assumption about the dynamics of the variational correlation parameters that define the Gutzwiller wave function, and which thus behave as genuine dynamical degrees of freedom that add on those of the variational uncorrelated Slater determinant. We apply the method to the standard half-filled single-band Hubbard model. We are able to recover known results, but, as a by-product, we also obtain a few other results. In particular, we show that quantum fluctuations can reproduce, almost quantitatively, the behavior of the uniform magnetic susceptibility uncovered by dynamical mean-field theory, which, though enhanced by correlations, is found to be smooth across the paramagnetic Mott transition. By contrast, the simple Gutzwiller approximation predicts that susceptibility to diverge at the transition.

DOI: [10.1103/PhysRevB.95.075156](https://doi.org/10.1103/PhysRevB.95.075156)

I. INTRODUCTION

The Gutzwiller approximation [1,2] is likely the simplest tool to deal with strong correlations in lattice models of interacting electrons. It consists of a recipe for approximate analytical expressions of expectation values in a class of wave functions, named Gutzwiller wave functions, of the form

$$|\Psi\rangle = \prod_i \mathcal{P}(i)|\Psi_0\rangle, \quad (1)$$

where $|\Psi_0\rangle$ is a variational Slater determinant, and $\mathcal{P}(i)$ a linear operator that acts on the local Hilbert space at site i and depends on a set of variational parameters.

Curiously, the Gutzwiller approximation often provides physically more sound results than a direct evaluation of expectation values in wave functions such as Eq. (1). For instance, the numerical optimization on a finite-dimensional lattice of a variational Gutzwiller wave function for a single-band half-filled Hubbard model never stabilizes a genuine Mott insulating phase [3,4], i.e., an insulator that does not break any symmetry, which intuitively is to be expected beyond a critical strength of the on-site repulsion. By contrast, the Gutzwiller approximation is instead able to describe such a genuine Mott transition [5]. The explanation of this strange outcome relies on the following observations. The first is that in order to describe a genuine Mott insulator, one needs to add to the Gutzwiller wave function, given by Eq. (1), long-range density-density Jastrow factors [4]. However, the effect of such Jastrow factors disappears in lattices with coordination number $z \rightarrow \infty$; therefore, only in that limit, wave functions such as Eq. (1) can faithfully describe Mott insulators. Moreover, right in that limit of $z \rightarrow \infty$, the Gutzwiller approximation provides the exact expression of expectation values [6,7]. Therefore, the Gutzwiller approximation is better regarded as a recipe to evaluate approximate expectation values in Gutzwiller-Jastrow wave functions, which becomes exact when the coordination number tends to infinity, rather than in Gutzwiller-only wave functions. In other words, the Gutzwiller approximation applied on a lattice with finite z is just the variational counterpart of dynamical mean-field theory (DMFT) [8] applied on that same lattice.

Recently, several attempts to include the Gutzwiller approximation inside density functional theory (DFT) electronic structure codes have been performed with quite encouraging outcomes [9–20]. In this perspective, it might be useful to have at one's disposal a simple and flexible method to calculate linear response functions within the Gutzwiller approximation, in view of an extension of the so-called linear response time-dependent DFT (TDDFT) [21,22] to the case when DFT is combined with the Gutzwiller approximation.

There are already several works dealing with linear response in the Gutzwiller approximation, most of which are limited to the single-band Hubbard model [23–28]. Extensions to multiband models have been attempted [29,30], though under an assumption about the dynamics of the variational parameters that determine the linear operators $\mathcal{P}(i)$ in Eq. (1). Here we shall instead present a very simple and general method to evaluate linear response functions within the Gutzwiller approximation without any preliminary assumption. The method is essentially an extension of the time-dependent Gutzwiller approximation of Ref. [31] to a generic multiband Hamiltonian, where the dynamics of the linear operators $\mathcal{P}(i)$ and of the Slater determinant $|\Psi_0\rangle$ [see Eq. (1)] are treated on equal footing. Linearization of the equations of motion around the stationary solution, which is the equilibrium state, thus allows the calculation of linear response functions.

We note that the results of the Gutzwiller approximation at equilibrium coincide with the saddle-point solution of the slave-boson theory in the path-integral formulation [32], which, in multiband models, corresponds to the so-called rotationally invariant slave-boson (RISB) formalism [33]. Our present results in the linear response regime can therefore be considered equivalent to the quantum fluctuation corrections above the RISB saddle-point solution. We preferred here to derive such corrections to the action directly from the time-dependent Gutzwiller approximation rather than from the RISB theory, since the former is at least a well-controlled variational scheme in lattices with infinite coordination number. However, both the notations as well as the language we shall use are actually closely related to RISB theory.

The paper is organized as follows. In Sec. II, we briefly present the time-dependent Gutzwiller approximation, with

some additional technical details postponed to the Appendix. In Sec. III, we linearize the equations of motion around the stationary solution and derive an effective action for the fluctuations in the harmonic approximation. In Sec. IV, we apply the method to the single-band half-filled Hubbard model, which allows a comparison with already existing results. Section V is devoted to concluding remarks.

II. THE GUTZWILLER APPROXIMATION IN BRIEF

Besides the original works [1,2] where Gutzwiller introduced a novel class of variational wave functions as well as an approximate scheme to compute expectation values, named Gutzwiller wave functions and approximation, and the subsequent demonstration that such an approximation becomes exact in the limit of infinite-coordination lattices [6,7], there are by now many articles where the Gutzwiller approximation is described in detail. Here we shall follow Ref. [34] and use the same notations.

The time-dependent Gutzwiller wave function is defined through [24,31,34]

$$|\Psi(t)\rangle = \prod_i \mathcal{P}(i,t) |\Psi_0(t)\rangle, \quad (2)$$

which is analogous to Eq. (1), but now $|\Psi_0(t)\rangle$ is a time-dependent variational Slater determinant and $\mathcal{P}(i,t)$ are linear operators on the local Hilbert space that depend on time-dependent variational parameters. For the sake of simplicity, we shall not include in our analysis BCS wave functions or operators $\mathcal{P}(i,t)$ that are charge nonconserving. The extension to those cases is simple, though notations get more involved.

Suppose that the Hamiltonian is written in terms of fermionic operators $c_{i\alpha}$ and $c_{i\alpha}^\dagger$, $\alpha = 1, \dots, 2M$, where M is the number of orbitals and 2 is the spin degeneracy, which correspond to annihilating or creating a fermion at site i in a chosen basis of Wannier functions $\phi_{i\alpha}(\mathbf{x}, t)$, where α indicates both spin and orbital indices. Let us imagine a $U(2M)$ unitary transformation,

$$\mathcal{W}(i,t) = \exp \left[i \sum_{\alpha\beta} K_{\alpha\beta}(i,t) c_{i\alpha}^\dagger c_{i\beta} \right], \quad (3)$$

with $K_{\alpha\beta}(i,t) = K_{\beta\alpha}(i,t)^*$ the matrix elements of a Hermitian matrix $\hat{K}(i,t)$, which maps $c_{i\alpha}$ into a new basis set $d_{i\alpha}$ of single-particle operators,

$$d_{i\alpha} = \mathcal{W}(i,t)^\dagger c_{i\alpha} \mathcal{W}(i,t) = \sum_{\beta} U_{\alpha\beta}(i,t) c_{i\beta}, \quad (4)$$

where $U_{\alpha\beta}(i,t)$ are the matrix elements of

$$\hat{U}(i,t) = e^{i\hat{K}(i,t)}. \quad (5)$$

Evidently, if we consider the *gauge* transformation

$$\mathcal{P}(i,t) \rightarrow \mathcal{P}(i,t) \mathcal{W}(i,t)^\dagger, \quad (6)$$

$$|\Psi_0(t)\rangle \rightarrow \mathcal{W}(t) |\Psi_0(t)\rangle = \prod_i \mathcal{W}(i,t) |\Psi_0(t)\rangle, \quad (7)$$

the Gutzwiller wave function $|\Psi(t)\rangle$ in (2) stays invariant and the transformed $|\Psi_0(t)\rangle$ remains a Slater determinant. Such

gauge invariance, analogous to that of the RISB theory [33], repeatedly appears in the calculations that follow.

The most general $\mathcal{P}(i,t)$ can be written [34,35] as

$$\begin{aligned} \mathcal{P}(i,t) &= \sum_{nm} \lambda_{nm}(i,t) |n;i\rangle \langle m;i| \\ &= \sum_{\bar{n}\bar{m}} \lambda_{\bar{n}\bar{m}}(i,t) |n;i\rangle \langle \bar{m};i|, \end{aligned} \quad (8)$$

where the *ket* $|n;i\rangle$ is a local Fock state of the operators $c_{i\alpha}$, while the *bra* is a Fock state of the same operators, $\langle m;i|$ in the first equation, or different ones, $\langle \bar{m};i|$ in the second equation, e.g., the basis of the operators $d_{i\alpha}$ in Eq. (4). The latter option, which we denote as the *mixed-basis* representation [36], is evidently better when the Slater determinant $|\Psi_0(t)\rangle$ is, for convenience, written in terms of single-particle operators different from those instead used to write the Hamiltonian, which we shall hereafter assume are the $c_{i\alpha}$. We define the uncorrelated local probability distribution $\hat{P}_0(i,t)$, which is positive definite, by its matrix elements,

$$P_{0\bar{n}\bar{m}}(i,t) = \langle \Psi_0(t) | \bar{m};i\rangle \langle \bar{n};i | \Psi_0(t)\rangle, \quad (9)$$

as well as the Gutzwiller variational matrix,

$$\hat{\Phi}(i,t) \equiv \hat{\lambda}(i,t) \sqrt{\hat{P}_0(i,t)}, \quad (10)$$

with matrix elements $\Phi_{\bar{n}\bar{m}}(i,t)$, which changes under the gauge transformation (6) as

$$\hat{\Phi}(i,t) \rightarrow \hat{\Phi}(i,t) \hat{W}(i,t)^\dagger, \quad (11)$$

where $\hat{W}(i,t)^\dagger$ is the matrix representation of $\mathcal{W}(i,t)^\dagger$, given by Eq. (3), in the local Fock basis. Expectation values of local and nonlocal operators in the Gutzwiller wave function (2) can be calculated explicitly in infinite-coordination lattices if one imposes the following two constraints at any time [7,34]:

$$\text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t)] = 1, \quad (12)$$

$$\begin{aligned} \text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \hat{c}_{i\alpha}^\dagger \hat{c}_{i\beta}] &\equiv n_{\alpha\beta}(i,t) \\ &= \langle \Psi_0(t) | c_{i\alpha}^\dagger c_{i\beta} | \Psi_0(t)\rangle, \end{aligned} \quad (13)$$

where the fermionic operators within the spur must be regarded as their matrix representation in the local Fock space. The first constraint, given by Eq. (12), is a normalization condition that does not reduce the variational freedom, unlike the second constraint, given by Eq. (13), that plays the same role as the projector onto the physical states in the RISB technique [33].

Another important ingredient is the wave-function renormalization matrix $\hat{R}(i,t)$ with elements $R_{\alpha\beta}(i,t)$, defined by solving the set of equations

$$\begin{aligned} \langle \Psi_0(t) | c_{i\gamma}^\dagger \mathcal{P}(i,t)^\dagger c_{i\alpha} \mathcal{P}(i,t) | \Psi_0(t)\rangle \\ = \sum_{\beta} n_{\gamma\beta}(i,t) R_{\alpha\beta}(i,t), \end{aligned} \quad (14)$$

where the left-hand side can be straightforwardly evaluated by the Wick's theorem. As shown in the Appendix, the solution of the above equation reads

$$\hat{R}(i,t) = \hat{Q}(i,t) \hat{S}(i,t), \quad (15)$$

where $\hat{Q}(i,t)$ has matrix elements

$$Q_{\alpha\beta}(i,t) = \text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{c}_{i\alpha} \hat{\Phi}(i,t) \hat{c}_{i\beta}^\dagger], \quad (16)$$

and the Hermitian matrix $\hat{S}(i,t)$ is defined through

$$4\hat{S}(i,t)^{-2} = 1 - \hat{\Delta}(i,t)^2, \quad (17)$$

where the matrix elements of $\hat{\Delta}(i,t)$ are

$$\Delta_{\alpha\beta}(i,t) = \text{Tr}\{\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) [\hat{c}_{i\alpha}, \hat{c}_{i\beta}^\dagger]\}. \quad (18)$$

The meaning of $\hat{R}(i,t)$ is that the action of the annihilation operator $c_{i\alpha}$ on the Gutzwiller wave function is equivalent to the action of the operator,

$$\mathcal{P}(i,t)^\dagger \mathbf{c}_i \mathcal{P}(i,t) \rightarrow \hat{R}(i,t) \mathbf{c}_i, \quad (19)$$

on the Slater determinant $|\Psi_0(t)\rangle$, where \mathbf{c}_i is a spinor with components $c_{i\alpha}$. One can readily show that under the gauge transformation (6),

$$\hat{R}(i,t) \rightarrow \hat{R}(i,t)^W = \hat{R}(i,t) \hat{U}(i,t)^\dagger, \quad (20)$$

where $\hat{U}(i,t)$ has been defined in Eq. (5), so that Eq. (19) transforms into

$$\mathcal{W}(i,t) \mathcal{P}(i,t)^\dagger \mathbf{c}_i \mathcal{P}(i,t) \mathcal{W}(i,t)^\dagger \rightarrow \hat{R}(i,t)^W \mathbf{d}_i.$$

Since we have complete freedom in choosing $\mathcal{W}(i,t)$, a convenient choice is the unitary transformation that diagonalizes the local single-particle density matrix, in which case the operators $d_{i\alpha}$ are associated to the natural orbitals and satisfy

$$\text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta}] = \delta_{\alpha\beta} n_\alpha(i,t), \quad (21)$$

while the matrix elements of $\hat{R}(i,t)^W$ acquire the simple expression

$$R_{\alpha\beta}(i,t)^W = \frac{\text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{c}_{i\alpha} \hat{\Phi}(i,t) \hat{d}_{i\beta}^\dagger]}{\sqrt{n_\beta(i,t)[1 - n_\beta(i,t)]}}. \quad (22)$$

The matrix $\hat{\Phi}(i,t)$ is in this case conveniently defined in the mixed-basis representation, where n in $\Phi_{n\bar{m}}(i,t)$ refers to a Fock state in the original basis, and \bar{m} to a Fock state in the natural one. Such a mixed-basis representation is useful since, throughout all calculations, one does not actually need to know what the natural basis is in terms of the original one [36]. Such a nice property is linked to the gauge invariance, given by Eqs. (6) and (7), of the theory [33].

A. The model

We shall assume the generic Hamiltonian

$$\mathcal{H} = \sum_{i \neq j} \mathbf{c}_i^\dagger \hat{t}_{ij} \mathbf{c}_j^\dagger + \sum_i \mathcal{H}_i, \quad (23)$$

where \mathcal{H}_i includes all on-site terms. If the constraints Eq. (12) and Eq. (13) are satisfied at any time t , then, in infinite

coordination lattices, it holds that [7,34]

$$\begin{aligned} E(t) &= \langle \Psi(t) | \mathcal{H} | \Psi(t) \rangle = \langle \Psi_0(t) | \mathcal{H}_*(t) | \Psi_0(t) \rangle \\ &+ \sum_i \text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{H}_i \hat{\Phi}(i,t)] \\ &\equiv E_*(t) + \sum_i \text{Tr}[\hat{\Phi}(i,t)^\dagger \hat{H}_i \hat{\Phi}(i,t)], \end{aligned} \quad (24)$$

where

$$\mathcal{H}_*(t) = \sum_{i \neq j} \mathbf{c}_i^\dagger \hat{R}(i,t)^\dagger \hat{t}_{ij} \hat{R}(j,t) \mathbf{c}_i \quad (25)$$

may be interpreted as the Hamiltonian of the quasiparticles. Evidently, all expectation values can be straightforwardly evaluated since the uncorrelated wave function $|\Psi_0(t)\rangle$ allows use of Wick's theorem.

B. The action

In the time domain, the variational principle corresponds to searching for the saddle point of the action [31],

$$\begin{aligned} \mathcal{S} &= \int dt [i \langle \Psi(t) | \dot{\Psi}(t) \rangle - E(t)] \\ &\equiv \int dt \left\{ i \sum_i \text{Tr} \left[\hat{\Phi}(i,t)^\dagger \frac{\partial \hat{\Phi}(i,t)}{\partial t} \right] \right. \\ &\quad \left. + i \langle \Psi_0(t) | \dot{\Psi}_0(t) \rangle - E(t) \right\}, \end{aligned} \quad (26)$$

where the equivalence holds on provision that the constraints (12) and (13) are fulfilled at any time. The saddle-point equations are readily obtained:

$$i \frac{\partial \hat{\Phi}(i,t)}{\partial t} = \hat{H}_i \hat{\Phi}(i,t) + \frac{\partial E_*(t)}{\partial \hat{\Phi}(i,t)^\dagger}, \quad (27)$$

$$i |\dot{\Psi}_0(t)\rangle = \mathcal{H}_*(t) |\Psi_0(t)\rangle, \quad (28)$$

where

$$\begin{aligned} \frac{\partial E_*(t)}{\partial \hat{\Phi}(i,t)^\dagger} &= \langle \Psi_0(t) | \frac{\partial \mathcal{H}_*(t)}{\partial \hat{\Phi}(i,t)^\dagger} | \Psi_0(t) \rangle \\ &\equiv \hat{T}(i,t) \hat{\Phi}(i,t). \end{aligned} \quad (29)$$

$\hat{T}(i,t)$ is a tensor with components $T_{nm;n'm'}(i,t)$, which is still a functional of the matrices $\hat{\Phi}$ and $\hat{\Phi}^\dagger$ at site i as well as at all sites connected to i by the hopping. One can show that this tensor is Hermitian, $\hat{T}(i,t) = \hat{T}(i,t)^\dagger$, which implies that the normalization (12) is conserved by the time evolution.

C. Fate of the constraint

Concerning the second constraint, given by Eq. (13), we now prove that if it is satisfied at the initial time, it will remain so at the saddle-point solutions of Eqs. (27) and (28). Suppose we have indeed found the saddle point $\hat{\Phi}(i,t)$ and $|\Psi_0(t)\rangle$. By definition, any small variation with respect to that solution must lead to a vanishing variation of the action. Let us therefore consider the gauge transformation (6) and (11), with

infinitesimal $K_{\alpha\beta}(i,t)$ [see Eq. (3)]. Under this transformation,

$$\hat{\Phi}(i,t) \hat{W}(i,t)^\dagger \simeq \hat{\Phi}(i,t) [1 - i \delta \hat{W}(i,t)], \quad \mathcal{W}(t)|\Psi_0(t)\rangle \simeq \left[1 + i \sum_i \delta \mathcal{W}(i,t) \right] |\Psi_0(t)\rangle,$$

where the operator

$$\delta \mathcal{W}(i,t) = \sum_{\alpha\beta} K_{\alpha\beta}(i,t) c_{i\alpha}^\dagger c_{i\beta}, \quad (30)$$

with $\delta \hat{W}(i,t)$ its matrix representation in the local Fock space.

We already mentioned that the energy $E(t)$ is gauge invariant so that the variation of the action, $\delta \mathcal{S} = \mathcal{S}^\mathcal{W} - \mathcal{S}$, simply reads

$$\begin{aligned} \delta \mathcal{S} &= \int dt \left\{ i \sum_i \text{Tr} \left[\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \frac{\partial \hat{W}(i,t)^\dagger}{\partial t} \hat{W}(i,t) \right] + i \sum_i \langle \Psi_0(t) | \mathcal{W}(i,t)^\dagger \dot{\mathcal{W}}(i,t) | \Psi_0(t) \rangle \right\} \\ &\simeq \sum_i \int dt \left\{ \text{Tr} \left[\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \frac{\partial \delta \hat{W}(i,t)}{\partial t} \right] - \langle \Psi_0(t) | \delta \dot{\mathcal{W}}(i,t) | \Psi_0(t) \rangle \right\} \\ &= \sum_i \sum_{\alpha\beta} \int dt \dot{K}_{\alpha\beta}(i,t) \{ \text{Tr} [\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \hat{c}_{i\alpha}^\dagger \hat{c}_{i\beta}^\dagger] - \langle \Psi_0(t) | c_{i\alpha}^\dagger c_{i\beta} | \Psi_0(t) \rangle \} \\ &= - \sum_i \sum_{\alpha\beta} \int dt K_{\alpha\beta}(i,t) \frac{\partial}{\partial t} \{ \text{Tr} [\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \hat{c}_{i\alpha}^\dagger \hat{c}_{i\beta}^\dagger] - \langle \Psi_0(t) | c_{i\alpha}^\dagger c_{i\beta} | \Psi_0(t) \rangle \}. \end{aligned}$$

Since $\hat{\Phi}(i,t)$ and $|\Psi_0(t)\rangle$ are solutions of the saddle-point equations, it follows that $\delta \mathcal{S}$ must strictly vanish for any choice of the infinitesimally small matrix elements $K_{\alpha\beta}(t)$, which implies

$$\frac{\partial}{\partial t} \{ \text{Tr} [\hat{\Phi}(i,t)^\dagger \hat{\Phi}(i,t) \hat{c}_{i\alpha}^\dagger \hat{c}_{i\beta}^\dagger] - \langle \Psi_0(t) | c_{i\alpha}^\dagger c_{i\beta} | \Psi_0(t) \rangle \} = 0,$$

thus just the desired result. It actually means that the term in parentheses is conserved in the evolution. Therefore, if it is initially vanishing, it will remain so at any time, which thus implies that the constraint given by Eq. (13) is fulfilled during the whole time evolution.

D. Stationary problem

At equilibrium, one needs to find the minimum of the energy with the two constraints (12) and (13), which can be enforced, e.g., by Lagrange multipliers, leading to the set of equations

$$\Lambda(i) \hat{\Phi}(i) = [\hat{H}_i + \hat{T}(i)] \hat{\Phi}(i) + \sum_{\alpha\beta} \mu_{\alpha\beta}(i) \hat{\Phi}(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta}, \quad (31)$$

$$E_* |\Psi_0\rangle = \left[\mathcal{H}_* - \sum_i \mu_{\alpha\beta}(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta} \right] |\Psi_0\rangle, \quad (32)$$

where $\Lambda(i)$ enforces Eq. (12), and the Hermitian matrix $\hat{\mu}(i)$ with components $\mu_{\alpha\beta}(i)$ enforces Eq. (13). In whatever follows, we shall assume to work in a mixed-basis representation where the operators $\hat{d}_{i\alpha}$ are associated to the natural orbitals, so that we must also ensure that

$$\text{Tr} [\hat{\Phi}^\dagger(i) \hat{\Phi}(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta}] = \langle \Psi_0 | \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta} | \Psi_0 \rangle = \delta_{\alpha\beta} n_\alpha(i).$$

The quasiparticle Hamiltonian in the natural basis, including explicitly the Lagrange multipliers, is therefore

$$\mathcal{H}_* \rightarrow \sum_{i \neq j} \mathbf{d}_i^\dagger \hat{R}(i)^\dagger \hat{t}_{ij} \hat{R}(j) \mathbf{d}_i - \sum_i \mathbf{d}_i^\dagger \hat{\mu}(i) \mathbf{d}_i, \quad (33)$$

with \hat{R} defined in Eq. (22). Working in the mixed-basis representation with the natural orbitals considerably simplifies all calculations.

Recalling that $\hat{T}(i)$ is still a functional of $\hat{\Phi}$, Eq. (31) looks like a stationary nonlinear Schrödinger equation [20,37]. One can, for instance, solve it as in any Hartree-Fock calculation. Namely, one can find the eigenstates and eigenvalues of Eq. (31) assuming $\hat{T}(i)$ fixed, and impose that when $\hat{T}(i)$ is calculated by substituting the actual expression of the lowest-energy solution $\hat{\Phi}_0(i)$, the two values coincide. The Lagrange multiplier $\hat{\mu}$ is fixed by imposing Eqs. (13) and (21). In this way, one finally gets the self-consistent $\hat{T}(i)$, which we shall hereafter denote as

$$\hat{T}^{(0)}(i) \equiv \hat{T}[\hat{\Phi}_0, \hat{\Phi}_0^\dagger]. \quad (34)$$

Once the latter is known, as well as the value of $\hat{\mu}$, one can also solve (31) for all eigenvectors $\hat{\Phi}_n(i)$ and corresponding eigenvalues $E_n(i)$, with $E_0(i) = \Lambda(i)$, namely,

$$E_n(i) \hat{\Phi}_n(i) = [\hat{H}_i + \hat{T}^{(0)}(i)] \hat{\Phi}_n(i) + \sum_{\alpha\beta} \mu_{\alpha\beta}(i) \hat{\Phi}_n(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta}. \quad (35)$$

We shall denote \mathcal{H}_* , \hat{R} , \hat{Q} , \hat{n} , and \hat{S} calculated with $\hat{\Phi}_0$ as $\mathcal{H}_*^{(0)}$, $\hat{R}^{(0)}$, $\hat{Q}^{(0)}$, $\hat{n}^{(0)}$, and $\hat{S}^{(0)}$, respectively, with the latter two matrices diagonal in the natural basis,

$$n_{\alpha\beta}^{(0)} = \delta_{\alpha\beta} n_\alpha^{(0)}, \quad (36)$$

$$S_{\alpha\beta}^{(0)} = \delta_{\alpha\beta} S_\alpha^{(0)} = \delta_{\alpha\beta} [n_\alpha^{(0)} (1 - n_\alpha^{(0)})]^{-1/2}. \quad (37)$$

We conclude by noting that the saddle-point Hamiltonian (33) with the inclusion of the Lagrange multipliers is no longer invariant under the most general $U(2M)$ gauge transformation, but only under a subgroup G with generators \hat{T}^a that commute with $\hat{\mu}$. This is common in theories where the gauge invariance implements constraints about physical states. In the natural basis representation, $\mu_{i,\alpha\beta} = \delta_{\alpha\beta} \mu_{i\alpha}$ is diagonal, so that the matrix elements of \hat{T}^a must satisfy

$$T_{i,\alpha\beta}^a (\mu_{i\alpha} - \mu_{i\beta}) = 0, \quad (38)$$

whose solution is straightforward. For any nondegenerate α , i.e., such that $\mu_{i\alpha} \neq \mu_{i\beta}, \forall \beta \neq \alpha$, we associate the generators $T_{i,\gamma\beta}^\alpha = \delta_{\alpha\beta} \delta_{\gamma\beta}$ of $U(1)$ Abelian groups. On the contrary, for any set of $\alpha_i, i = 1, \dots, k$, such that $\mu_{i\alpha_i} = \mu_{i\alpha_j} \neq \mu_{i\beta}, \forall \beta \neq \alpha_1, \dots, \alpha_k$, we can associate generators of a $U(k)$ Lie algebra.

III. FLUCTUATIONS ABOVE THE SADDLE-POINT SOLUTION

Our goal is to determine the action of the fluctuations beyond the saddle point within the harmonic approximation. To that purpose, we assume that

$$\hat{\Phi}(i,t) = e^{-iE_0 t} \sum_n \phi_n(i,t) \hat{\Phi}_n(i) \hat{W}(i,t)^\dagger, \quad (39)$$

where $\phi_n(i,t)$ for $n > 0$ is regarded as a first-order fluctuation, while to enforce normalization,

$$\phi_0(i,t) = 1 - \frac{1}{2} \sum_{n>0} |\phi_n(i,t)|^2. \quad (40)$$

The physical meaning of Eq. (39) is that we expand the fluctuations around the stationary solution in terms of the ground and excited states of the local eigenvalue problem described by Eq. (35), with each eigenstate weighed by the probability amplitude $\phi_n(i,t)$. In addition, the Slater determinant is defined through

$$|\Psi_0(t)\rangle \rightarrow e^{-iE_* t} \mathcal{W}(t) |\Psi_0(t)\rangle, \quad (41)$$

where $|\Psi_0(t)\rangle$ is properly normalized and includes the zeroth-order $|\Psi_0^{(0)}\rangle$ solution of the saddle point, as well as a fluctuation correction $|\delta\Psi_0(t)\rangle$. The unitary operator $\mathcal{W}(i,t)$ is chosen as

$$\mathcal{W}(i,t) = \exp[-i t \mathbf{d}_i^\dagger \hat{\mu}(i) \mathbf{d}_i], \quad (42)$$

where $\hat{\mu}(i)$ is the equilibrium Lagrange multiplier and $\hat{W}(i,t)$ is the matrix representation of $\mathcal{W}(i,t)$.

Through the above definitions, the action becomes

$$\begin{aligned} \mathcal{S} = \int dt \left\{ i \sum_i \sum_{n>0} \phi_n(i,t)^* \dot{\phi}_n(i,t) + i \langle \Psi_0(t) | \dot{\Psi}_0(t) \rangle \right. \\ \left. - \sum_i \sum_{nm} \phi_n(i,t)^* V_{nm}(i) \phi_m(i,t) + E_0 + E_* - E_*(t) \right\}, \end{aligned} \quad (43)$$

where $E_*(t) = \langle \Psi_0(t) | \mathcal{H}_*(t) | \Psi_0(t) \rangle$, being now

$$\mathcal{H}_*(t) = \sum_{i \neq j} \mathbf{d}_i^\dagger \hat{R}(i,t)^\dagger \hat{t}_{ij} \hat{R}(j,t) \mathbf{d}_i - \sum_i \mathbf{d}_i^\dagger \hat{\mu}(i) \mathbf{d}_i \quad (44)$$

and

$$\begin{aligned} V_{nm}(i) = \text{Tr}[\hat{\Phi}_n(i)^\dagger \hat{H}_i \hat{\Phi}_m(i)] \\ + \text{Tr}[\hat{\Phi}_n(i)^\dagger \hat{\Phi}_m(i) \hat{\mathbf{d}}_i^\dagger \hat{\mu}(i) \hat{\mathbf{d}}_i]. \end{aligned} \quad (45)$$

We expand $\mathcal{H}_*(t)$ up to second order in the fluctuations. The zeroth order is just $\mathcal{H}_*^{(0)}$. Since the stationary solution is the saddle point of the action, the expectation value of the first-order expansion $\mathcal{H}^{(1)}(t)$ over the saddle-point Slater determinant $|\Psi_0^{(0)}\rangle$ cancels with the first-order expansion of the local energy $\sum_i \sum_{nm} \phi_n(i,t)^* V_{nm}(i) \phi_m(i,t)$. Therefore, $\mathcal{H}^{(1)}(t)$ contributes to $E_*(t)$ with a second-order term that, by linear response theory, reads

$$\begin{aligned} \delta_1 E_*(t) = \langle \delta\Psi_0(t) | \mathcal{H}^{(1)}(t) | \Psi_0^{(0)} \rangle + \text{c.c.} \\ = -i \int_0^t d\tau \langle [\mathcal{H}^{(1)}(t), \mathcal{H}^{(1)}(\tau)] \rangle_0, \end{aligned} \quad (46)$$

where, hereafter, $\langle \cdot \rangle_0$ will denote the average over $|\Psi_0^{(0)}\rangle$, and the operators in Eq. (46) have an additional time dependence since they are evolved with the saddle-point Hamiltonian $\mathcal{H}_*^{(0)}$. The explicit expression of $\mathcal{H}^{(1)}(t)$ is

$$\mathcal{H}^{(1)}(t) = \sum_{i \neq j} [\mathbf{d}_j^\dagger \hat{R}^{(0)}(j)^\dagger \hat{t}_{ji} \hat{R}^{(1)}(i,t) \mathbf{d}_i + \text{H.c.}], \quad (47)$$

where $\hat{R}^{(0)}(i)$ is the stationary value, while the explicit expression of the first-order Taylor expansion $\hat{R}^{(1)}(i,t)$ is given in the Appendix; see Eq. (A17).

There are several second-order terms upon expanding $\mathcal{H}_*(t)$, which we shall consider separately. The first is simply

$$\mathcal{H}_1^{(2)}(t) = \sum_{i \neq j} \mathbf{d}_i^\dagger \hat{R}^{(1)}(i,t)^\dagger \hat{t}_{ij} \hat{R}^{(1)}(j,t) \mathbf{d}_j, \quad (48)$$

whose expectation value over $|\Psi_0^{(0)}\rangle$ is an additional second-order contribution,

$$\delta_2 E_*(t) = \langle \mathcal{H}_1^{(2)}(t) \rangle_0, \quad (49)$$

which, together with $\delta_1 E_*(t)$ in Eq. (46), endow the action with spatial correlations among the $\phi_n(i,t)$ at different sites.

The next second-order corrections to $\mathcal{H}_*(t)$ derive from the second-order expansion of $\hat{R}(i,t)$,

$$\hat{R}^{(2)}(i,t) = \hat{R}_1^{(2)}(i,t) + \hat{R}_2^{(2)}(i,t), \quad (50)$$

where we distinguish two different contributions; see Eqs. (A19) and (A20) in the Appendix. The reason for this distinction is that

$$\begin{aligned} \delta_3 E_*(t) = \sum_i \sum_{nm} \phi_n(i,t)^* V_{nm}(i) \phi_m(i,t) \\ + \left\langle \sum_{i \neq j} [\mathbf{d}_j^\dagger \hat{R}^{(0)}(j)^\dagger \hat{t}_{ji} \hat{R}_1^{(2)}(i,t) \mathbf{d}_i + \text{H.c.}] \right\rangle_0 \\ = \sum_{n>0} (E_n - E_0) \phi_n(i,t)^* \phi_n(i,t) \end{aligned} \quad (51)$$

reproduces the bare excitation energy of the fluctuations. The last contribution to the energy of the fluctuations is therefore

$$\delta_4 E_*(t) = \left\langle \sum_{i \neq j} [d_j^\dagger \hat{R}^{(0)}(j) \hat{r}_{ji} \hat{R}_2^{(2)}(i, t) d_i + \text{H.c.}] \right\rangle. \quad (52)$$

If we define new variables

$$x_n(i, t) = \frac{1}{\sqrt{2}} [\phi_n(i, t) + \phi_n(i, t)^*], \quad (53)$$

$$p_n(i, t) = -\frac{i}{\sqrt{2}} [\phi_n(i, t) - \phi_n(i, t)^*], \quad (54)$$

and the quadratic potential

$$U(t, \{x, p\}) = \delta_1 E_*(t) + \delta_2 E_*(t) + \delta_4 E_*(t), \quad (55)$$

which has a retarded component $\delta_1 E_*(t)$ [see Eq. (46)], then the action of the fluctuations, upon defining $\omega_n = E_n - E_0$, reads

$$\delta S = \int dt \left(\sum_i \sum_{n>0} \left\{ p_n(i, t) \dot{x}_n(i, t) - \frac{\omega_n}{2} [x_n(i, t)^2 + p_n(i, t)^2] \right\} - U(t, \{x, p\}) \right), \quad (56)$$

which is just the action of coupled harmonic oscillators.

δS in Eq. (56) can, for instance, be used to evaluate the fluctuation corrections to linear response functions of local operators. For any local observable $\hat{O}(i)$, let us define the matrix element

$$O_n(i) \equiv \text{Tr}[\hat{\Phi}_n(i)^\dagger \hat{O}(i) \hat{\Phi}_0(i)]. \quad (57)$$

Suppose we add a perturbation that couples to the local density matrix,

$$\delta \mathcal{H}(t) = \sum_i c_i^\dagger \hat{V}(i, t) c_i, \quad (58)$$

where the matrix $\hat{V}(i, t)$ with elements $V_{\alpha\beta}(i, t)$ represents the external field. Without loss of generality, we can assume that the expectation value of $\delta \mathcal{H}(t)$ in Eq. (58) vanishes at the stationary solution. Since by assumption the external field is first order, the perturbation adds a second-order correction to the action (56),

$$\begin{aligned} V(t) &= \sum_i \sum_n \{ \phi_n(i, t)^* \text{Tr}[\hat{\Phi}_n(i)^\dagger \hat{c}_i^\dagger \hat{V}(i, t) \hat{c}_i \hat{\Phi}_0(i)] \\ &\quad + \phi_n(i, t) \text{Tr}[\hat{\Phi}_0(i)^\dagger \hat{c}_i^\dagger \hat{V}(i, t) \hat{c}_i \hat{\Phi}_n(i)] \} \\ &\equiv \sum_i \sum_n [\phi_n(i, t)^* V_n(i, t) + \phi_n(i, t) V_n(i, t)^*] \\ &= \sqrt{2} \sum_{in} [\text{Re} V_n(i, t) x_n(i, t) + \text{Im} V_n(i, t) p_n(i, t)]. \end{aligned} \quad (59)$$

In the presence of $V(t)$, the action transforms into that of forced harmonic oscillators, whose solution allows calculation of the

expectation value of any local operator $\hat{O}(i)$ [see Eq. (57)],

$$\begin{aligned} O(i, t) &= \text{Tr}[\hat{\Phi}(i, t)^\dagger \hat{O}(i) \hat{\Phi}(i, t)] \\ &\simeq \sqrt{2} \sum_n [\text{Re} O_n(i) x_n(i, t) + \text{Im} O_n(i) p_n(i, t)], \end{aligned}$$

at linear order in the external field.

Residual gauge invariance and would-be Goldstone modes

As we mentioned, the action given by Eq. (43), with the time-dependent quasiparticle Hamiltonian defined in Eq. (44), is invariant under a subgroup G of the initial $U(2M)$ gauge symmetry. This implies the existence of massless modes with singular propagators that diverge as $1/\omega^2$ at low frequency, which are the would-be Goldstone modes related to the fact that the saddle point $\hat{\Phi}_0(i)$ is not invariant under G . Since the gauge invariance breakdown is fake, a $U(2M)$ gauge symmetry cannot be spontaneously broken; such singular behavior is unphysical and simply signals that gauge symmetry is to be restored beyond the saddle-point approximation [38].

Let us consider, for instance, a $U(1)$ subgroup of G related to the nondegenerate state α in the natural basis. The associated adjoint charge is

$$n_\alpha(i, t) \simeq \sum_{n>0} \{ \phi_n(i, t)^* \text{Tr}[\hat{\Phi}_n(i)^\dagger \hat{\Phi}_0(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\alpha}] + \text{c.c.} \},$$

and its conjugate variable is readily found to be

$$\begin{aligned} \varphi_\alpha(i, t) &\simeq \frac{i}{2n_\alpha^{(0)}} \sum_{n>0} \{ \phi_n(i, t)^* \text{Tr}[\hat{\Phi}_n(i)^\dagger \hat{\Phi}_0(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\alpha}] \\ &\quad - \phi_n(i, t) \text{Tr}[\hat{\Phi}_0(i)^\dagger \hat{\Phi}_n(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\alpha}] \}. \end{aligned}$$

The role of $\varphi_\alpha(i, t)$ is just to enforce the constraint (13), i.e.,

$$n_\alpha(i, t) = \langle \Psi_0(t) | c_{i\alpha}^\dagger c_{i\alpha} | \Psi_0(t) \rangle \equiv \langle c_{i\alpha}^\dagger c_{i\alpha} \rangle_t.$$

Indeed we can always perform a gauge transformation on the fermions,

$$c_{i\alpha} \rightarrow e^{-i\varphi_\alpha(i, t)} c_{i\alpha},$$

which makes $\varphi_\alpha(i, t)$ disappear from the energy, leaving just the time derivative term in the action,

$$\delta S = - \int dt \dot{\varphi}_\alpha(i, t) [n_\alpha(i, t) - \langle c_{i\alpha}^\dagger c_{i\alpha} \rangle_t].$$

The condition of vanishing derivative with respect to $\varphi_\alpha(i, t)$ is therefore just the condition that the constraint is conserved.

It follows that we can always drop from the action all terms that contain the variables conjugate to the adjoint charges associated with the gauge symmetry G , with the provision that wherever $n_\alpha(i, t)$ appears, we replace it with $\langle c_{i\alpha}^\dagger c_{i\alpha} \rangle_t$.

However, the above procedure does not involve all the coefficients $\phi_n(i, t)$; some of their linear combinations are untouched by *gauge fixing* and remain genuine independent dynamical degrees of freedom [39]. This fact, rather than being a limitation, endows the theory with a richer dynamics.

IV. APPLICATION TO THE HALF-FILLED HUBBARD MODEL

We now apply the above formalism to the simple case of a single-band Hubbard model at half filling, where all calculations can be worked out analytically and which also allows for a direct comparison with previous works [23–28,39–41]. We will show that we can indeed recover known results, but also find a few others.

The Hamiltonian in this case is

$$\mathcal{H} = -\frac{t}{\sqrt{z}} \sum_{(ij)\sigma} (c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.}) + \frac{U}{4} \sum_i [2(n_i - 1)^2 - 1], \quad (60)$$

where (ij) means nearest-neighbor bonds on a d -dimensional hypercubic lattice, and $z = 2d$ is the lattice coordination number that must be sent to $+\infty$ for the calculation to be really variational.

The local basis comprises four states which we choose to be, in order, empty configuration $|0\rangle$, doubly occupied $|2\rangle$, singly occupied by a spin-up electron $|\uparrow\rangle$, and occupied by a spin-down electron $|\downarrow\rangle$. The most general charge-conserving $\hat{\Phi}$ has the following form, dropping for now the site index:

$$\hat{\Phi} = \frac{1}{\sqrt{2}} \begin{pmatrix} \hat{\Phi}_c & 0 \\ 0 & \hat{\Phi}_s \end{pmatrix}, \quad (61)$$

where the charge component, i.e., the matrix elements in the subspace ($|0\rangle, |2\rangle$), is

$$\hat{\Phi}_c = \begin{pmatrix} \phi_{c0} + \phi_{c3} & 0 \\ 0 & \phi_{c0} - \phi_{c3} \end{pmatrix} = \phi_{c0} \sigma_0 + \phi_{c3} \sigma_3, \quad (62)$$

with σ_0 the 2×2 identity matrix, and $\sigma_i, i = 1, \dots, 3$ the Pauli matrices, whereas the spin component, namely, the matrix elements in the subspace ($|\uparrow\rangle, |\downarrow\rangle$), is instead

$$\hat{\Phi}_s = \sum_{i=0}^3 \phi_{si} \sigma_i = \phi_{s0} \sigma_0 + \boldsymbol{\phi}_s \cdot \boldsymbol{\sigma}, \quad (63)$$

which allows a full spin-SU(2) invariant analysis [26,42]. Normalization implies that

$$1 = |\phi_{c0}|^2 + |\phi_{c3}|^2 + |\phi_{s0}|^2 + \boldsymbol{\phi}_s^* \cdot \boldsymbol{\phi}_s.$$

One can readily verify that the matrix \hat{Q} with components,

$$Q_{\sigma\sigma'} = \text{Tr}(\hat{\Phi}^\dagger c_\sigma \hat{\Phi} c_{\sigma'}^\dagger), \quad (64)$$

can be written as

$$\hat{Q} = Q_0 \sigma_0 + \mathbf{Q} \cdot \boldsymbol{\sigma}, \quad (65)$$

where

$$2Q_0 = (\phi_{c0}^* \phi_{s0} + \phi_{s0}^* \phi_{c0}) + (\phi_{c3}^* \phi_{s0} - \phi_{s0}^* \phi_{c3}), \quad (66)$$

$$2Q_i = (\phi_{c0}^* \phi_{si} - \phi_{si}^* \phi_{c0}) + (\phi_{c3}^* \phi_{si} + \phi_{si}^* \phi_{c3}), \quad (67)$$

with $i = 1, \dots, 3$. Seemingly,

$$\begin{aligned} \hat{\Delta} &\equiv (\phi_{c0}^* \phi_{c3} + \phi_{c3}^* \phi_{c0}) \sigma_0 - (\phi_{s0}^* \boldsymbol{\phi}_s + \phi_{s0} \boldsymbol{\phi}_s^* + i \boldsymbol{\phi}_s^* \wedge \boldsymbol{\phi}_s) \cdot \boldsymbol{\sigma} \\ &\equiv \Delta_0 \sigma_0 + \boldsymbol{\Delta} \cdot \boldsymbol{\sigma}. \end{aligned} \quad (68)$$

A. Stationary solution

As is common when discussing the Mott transition in the single-band Hubbard model, we shall be interested in the stationary solution within the paramagnetic sector, i.e., neglecting spontaneous breakdown of spin-SU(2) symmetry. Such solution at half filling is characterized by a site-independent

$$\hat{\Phi}_0 = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_{c0}^{(0)} \sigma_0 & 0 \\ 0 & \phi_{s0}^{(0)} \sigma_0 \end{pmatrix},$$

with

$$1 = |\phi_{c0}^{(0)}|^2 + |\phi_{s0}^{(0)}|^2.$$

Under this assumption,

$$\hat{R}(i) = (\phi_{c0}^{(0)*} \phi_{s0}^{(0)} + \phi_{s0}^{(0)*} \phi_{c0}^{(0)}) \sigma_0 = R^{(0)} \sigma_0 \forall i, \quad (69)$$

so that the quasiparticle Hamiltonian is just a tight-binding model with renormalized hopping, i.e.,

$$\mathcal{H}_*^{(0)} = -\frac{t}{\sqrt{z}} R^{(0)2} \sum_{(ij)} (c_i^\dagger c_j + \text{H.c.}), \quad (70)$$

and natural and original orbitals coincide. It follows that the stationary Slater determinant is the noninteracting Fermi sea. We define

$$-\sum_i T_0 \equiv -\frac{t}{\sqrt{z}} \sum_{(ij)\sigma} \langle c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.} \rangle_0,$$

where $\langle \cdot \rangle_0$ is the average over the Fermi sea. Therefore, $-T_0$ is the hopping energy per site and $-2T_0/z$ is the hopping energy per bond of the Fermi sea.

The saddle-point equations for $\hat{\Phi}_0$ can be readily found,

$$E \phi_{c0}^{(0)} = -2T_0 R^{(0)} \phi_{s0} + \frac{U}{4} \phi_{c0},$$

$$E \phi_{s0}^{(0)} = -2T_0 R^{(0)} \phi_{c0} - \frac{U}{4} \phi_{s0}.$$

The lowest-energy eigenvalue is

$$E_0 = -\frac{1}{2} \sqrt{U^2 + (8T_0 R^{(0)})^2}, \quad (71)$$

and is characterized by

$$\phi_{c0}^{(0)} = \sin \frac{\theta}{2}, \quad \phi_{s0}^{(0)} = \cos \frac{\theta}{2},$$

with $\tan \theta = 8T_0 R^{(0)}/U$. Since through Eq. (69) $R^{(0)} = \sin \theta$, the self-consistency condition implies

$$\tan \theta = \frac{8T_0 R^{(0)}}{U} = \frac{8T_0}{U} \sin \theta, \quad (72)$$

namely,

$$\cos \theta = \begin{cases} U/U_c, & U \leq U_c = 8T_0 \\ 1, & U > U_c. \end{cases} \quad (73)$$

U_c is the well-known value of the Brinkman-Rice [5] metal-insulator transition within the Gutzwiller approximation. Indeed, for any $U \geq U_c$, $R^{(0)} = \sin \theta = 0$, so that the renormalized hopping in Eq. (70) vanishes; the quasiparticle motion is completely suppressed beyond U_c .

In conclusion, the lowest-energy eigenstate is

$$\hat{\Phi}_0 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sin \frac{\theta}{2} \sigma_0 & 0 \\ 0 & \cos \frac{\theta}{2} \sigma_0 \end{pmatrix}, \quad (74)$$

where $\cos \theta = \min(1, U/U_c)$, and has eigenvalue

$$E_0 = -\frac{U}{4 \cos \theta} = -\frac{\text{Max}(U, U_c)}{4}. \quad (75)$$

We can now find all other eigenvalues and eigenvectors. The highest-energy one is

$$\hat{\Phi}_3 = \frac{1}{\sqrt{2}} \begin{pmatrix} \cos \frac{\theta}{2} \sigma_0 & 0 \\ 0 & -\sin \frac{\theta}{2} \sigma_0 \end{pmatrix}, \quad (76)$$

with eigenvalue

$$E_3 = -E_0. \quad (77)$$

This eigenstate actually corresponds to the high-energy Hubbard bands. Indeed, above U_c , its excitation energy $E_3 - E_0 = U/2$ is just the cost of creating an empty or doubly occupied site in the atomic limit described by the Hamiltonian (60) with $t = 0$.

The lowest excited eigenstate is threefold degenerate ($i = 1, 2, 3$),

$$\hat{\Phi}_{1i} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 \\ 0 & \sigma_i \end{pmatrix}, \quad (78)$$

with eigenvalue

$$E_1 = -\frac{U}{4}, \quad (79)$$

and describes spin fluctuations. We note that above the Brinkmann-Rice transition, $U > U_c$, this magnetic state becomes degenerate with the ground state, as both imply strict single occupancy. In what follows, we shall nevertheless always expand around $\hat{\Phi}_0$ and, to avoid problems, we will mostly consider the metal phase at $U \leq U_c$.

Finally, the last eigenstate is

$$\hat{\Phi}_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sigma_3 & 0 \\ 0 & 0 \end{pmatrix}, \quad (80)$$

with eigenvalue

$$E_2 = +\frac{U}{4}, \quad (81)$$

and instead describes charge fluctuations, i.e., deviations of the local density from its average value of one. This mode becomes evidently degenerate with $\hat{\Phi}_3$ above the transition.

B. Action of the fluctuations

Following Sec. III, we write

$$\hat{\Phi}(i, t) = \phi_0(i, t) \hat{\Phi}_0 + \sum_{i=1}^3 \phi_{1i}(i, t) \hat{\Phi}_{1i} + \sum_{n=2}^3 \phi_n(i, t) \hat{\Phi}_n, \quad (82)$$

with $\phi_0(i, t)$ fixed by normalization. Through Eqs. (65)–(67), we find that

$$\begin{aligned} \hat{R}^{(1)}(i, t) &= \sin \frac{\theta}{2} [\phi_1(i, t) - \phi_1(i, t)^*] \cdot \sigma \\ &\quad - \cos \frac{\theta}{2} [\phi_2(i, t) - \phi_2(i, t)^*] \\ &\quad + \cos \theta [\phi_3(i, t) + \phi_3(i, t)^*] \\ &\equiv i \sqrt{2} \sin \frac{\theta}{2} \mathbf{p}_1(i, t) \cdot \sigma - i \sqrt{2} \cos \frac{\theta}{2} p_2(i, t) \\ &\quad + \sqrt{2} \cos \theta x_3(i, t), \end{aligned} \quad (83)$$

where we have introduced the conjugate variables associated with ϕ_n and ϕ_n^* . Equation (47) explicitly reads

$$\begin{aligned} \mathcal{H}_*^{(1)} &= \sum_i \left\{ \sqrt{2} \left(2 \cos \frac{\theta}{2} \right)^{-1} \nabla \cdot \mathbf{J}_s(i) \cdot \mathbf{p}_1(i, t) \right. \\ &\quad - \sqrt{2} \left(2 \sin \frac{\theta}{2} \right)^{-1} \nabla \cdot \mathbf{J}_c(i) p_2(i, t) \\ &\quad \left. + 2\sqrt{2} \cot \theta h_*(i) x_3(i, t) \right\}, \end{aligned} \quad (84)$$

where ∇ is the lattice divergence, $\mathbf{J}_s(i)$ and $\mathbf{J}_c(i)$ are the spin and charge currents, respectively, defined through the continuity equations

$$i \frac{\partial}{\partial t} (c_i^\dagger \sigma_0 c_i) = [c_i^\dagger \sigma_0 c_i, \mathcal{H}_*^{(0)}] \equiv -i \nabla \cdot \mathbf{J}_c(i), \quad (85)$$

$$i \frac{\partial}{\partial t} (c_i^\dagger \sigma c_i) = [c_i^\dagger \sigma c_i, \mathcal{H}_*^{(0)}] \equiv -i \nabla \cdot \mathbf{J}_s(i), \quad (86)$$

and, finally, $h_*(i)$ is the Hamiltonian density,

$$h_*(i) = -\frac{t}{2\sqrt{z}} R^{(0)2} \sum_{j \text{ n.n. } i} (c_i^\dagger c_j + \text{H.c.}). \quad (87)$$

Therefore, $\delta_1 E_*(t)$ defined in Eq. (46) becomes, due to particle-hole and spin-SU(2) symmetry,

$$\begin{aligned} \delta_1 E_*(t) &= \sum_{i,j} \int d\tau \left\{ \frac{1}{1 + \cos \theta} \chi_{\nabla J \nabla J}(i - j, t - \tau) \mathbf{p}_1(i, t) \cdot \mathbf{p}_1(j, \tau) \right. \\ &\quad + \frac{1}{1 - \cos \theta} \chi_{\nabla J \nabla J}(i - j, t - \tau) p_2(i, t) p_2(j, \tau) \\ &\quad \left. + 8 \cot^2 \theta \chi_{h_* h_*}(i - j, t - \tau) x_3(i, t) x_3(j, \tau) \right\}, \end{aligned} \quad (88)$$

where $\chi_{\nabla J \nabla J}$ is the linear response function of $\nabla \mathbf{J}$ with the Hamiltonian $\mathcal{H}_*^{(0)}$, which is actually the same for charge and spin currents, and $\chi_{h_* h_*}$ is the response function of h_* . We observe that because of charge and spin continuity equations, in Fourier space the following equivalence holds:

$$2T_0 \sin^2 \theta (\gamma_0 - \gamma_{\mathbf{q}}) + \chi_{\nabla J \nabla J}(\mathbf{q}, \omega) = \omega^2 \chi(\mathbf{q}, \omega), \quad (89)$$

where $\chi(\mathbf{q}, \omega)$ is the density-density response function, which is the same both in the charge and spin channels, and,

by definition,

$$\gamma_{\mathbf{q}} = \frac{2}{z} \sum_{i=1}^d \cos q_i \in [-1, +1]. \quad (90)$$

Without going into further details, we find the following expressions for the remaining contributions $\delta_2 E_*(t)$ in Eq. (49) and $\delta_4 E_*(t)$ in Eq. (52):

$$\delta_2 E_*(t) = -\frac{4T_0}{z} \sum_{(ij)} \left\{ \sin^2 \frac{\theta}{2} \mathbf{p}_1(i,t) \cdot \mathbf{p}_1(j,t) + \cos^2 \frac{\theta}{2} p_2(i,t) p_2(j,t) + \cos^2 \theta x_3(i,t) x_3(j,t) \right\}, \quad (91)$$

$$\delta_4 E_*(t) = -2T_0 \sin^2 \theta \sum_i \left\{ \cos^2 \frac{\theta}{2} \mathbf{x}_1(i,t) \cdot \mathbf{x}_1(i,t) + \sin^2 \frac{\theta}{2} x_2(i,t)^2 \right\}. \quad (92)$$

We now have all of the ingredients required to evaluate linear response functions of local operators within the harmonic approximation for the fluctuations.

C. Hubbard-band dispersion mode

As we mentioned, the Hubbard bands may be associated with the excited state $\hat{\Phi}_3$, hence with the operators x_3 and p_3 . Their equations of motion in Fourier space are

$$\begin{aligned} -i\omega x_3(\mathbf{q}, \omega) &= \omega_3 p_3(\mathbf{q}, \omega), \quad (93) \\ -i\omega p_3(\mathbf{q}, \omega) &= -[\omega_3 - 4T_0 \gamma_{\mathbf{q}} + 8 \cot^2 \theta \chi_{h_* h_*}(\mathbf{q}, \omega)] \\ &\quad \times x_3(\mathbf{q}, \omega). \quad (94) \end{aligned}$$

Within the metal phase, $U < U_c$, $\omega_3 = E_3 - E_0 = 4T_0$, so that, upon defining $\cos \theta = U/U_c \equiv u$, and noting that for small $|\mathbf{q}|$, $\chi_{hh}(\mathbf{q}, \omega) = O(q^4)$, the eigenmode energy is the solution of the equation

$$\begin{aligned} \omega_{3\mathbf{q}}^2 &= 4T_0 \left[4T_0(1 - u^2) + 4T_0 u^2 (\gamma_0 - \gamma_{\mathbf{q}}) \right. \\ &\quad \left. + 8 \frac{u^2}{1 - u^2} \chi_{h_* h_*}(\mathbf{q}, \omega_{3\mathbf{q}}) \right] \\ &\simeq 16T_0^2 [(1 - u^2) + u^2 (\gamma_0 - \gamma_{\mathbf{q}})], \quad (95) \end{aligned}$$

and thus describes an optical mode that softens at the metal-insulator transition, $\omega_{30} = 4T_0 \sqrt{1 - u^2} \rightarrow 0$, when $u \rightarrow 1$. We observe that the continuum of quasiparticle-quasihole excitations extends up to an energy of the order of $T_0(1 - u^2)$, so that upon approaching the transition, $\omega_{3\mathbf{q}}$ must detach from the continuum and become a genuine coherent excitation. This coherent mode actually corresponds to the spin-wave excitations of the Ising field within the Z_2 slave-spin representation of the Hubbard model [27,43,44]. This is not surprising since, as shown in Ref. [27], the Gutzwiller wave function is just the mean-field variational state of the Z_2 slave-spin theory. At the mean-field level, the Mott transition in this representation translates into the order-disorder transition of a quantum Ising model. Therefore, the mode x_3 seems to be the real fingerprint of the Mott transition.

D. Dynamical charge susceptibility

We assume to perturb the system in the metal phase, $u \leq 1$, by an external potential that couples to the charge deviation

from half filling, namely,

$$\begin{aligned} \delta \mathcal{H}(t) &= \sum_i v(i,t) (n_i - 1) \\ &\simeq -\sqrt{2} \sin \frac{\theta}{2} \sum_i v(i,t) x_2(i,t). \quad (96) \end{aligned}$$

Since $\omega_2 = E_2 - E_0 = 2T_0(1 + u)$ and by means of Eq. (89), we find in the presence of the field the following equations of motion for the conjugate variables x_2 and p_2 :

$$\begin{aligned} -i\omega x_2(\mathbf{q}, \omega) &= \frac{\omega^2}{1 - u} \chi(\mathbf{q}, \omega) p_2(\mathbf{q}, \omega), \\ -i\omega p_2(\mathbf{q}, \omega) &= \sqrt{2} \sin \frac{\theta}{2} v(\mathbf{q}, \omega) \\ &\quad - 2T_0(1 + u)u(2 - u)x(\mathbf{q}, \omega), \end{aligned}$$

from which it follows that the dynamical charge susceptibility is

$$\begin{aligned} \chi_c(\mathbf{q}, \omega) &= \frac{(1 - u) \chi(\mathbf{q}, \omega)}{(1 - u) - 2T_0(1 + u)u(2 - u) \chi(\mathbf{q}, \omega)} \\ &\equiv \frac{\chi(\mathbf{q}, \omega)}{1 + \Gamma_c \chi(\mathbf{q}, \omega)}, \quad (97) \end{aligned}$$

where the analogy with conventional random-phase approximation (RPA) is evident, though with a renormalized coupling constant,

$$\Gamma_c = -\frac{U}{2} \frac{1 + u}{1 - u} \left(1 - \frac{u}{2} \right) < 0. \quad (98)$$

We note that

$$\chi(\mathbf{q} \rightarrow 0, \omega = 0) = -\mathcal{N}_*,$$

where

$$\mathcal{N}_* = \frac{\mathcal{N}_0}{1 - u^2} \quad (99)$$

is the quasiparticle density of states (DOS) at the chemical potential, as opposed to the bare DOS \mathcal{N}_0 , and diverges approaching the Mott transition. Therefore, through Eq. (97), the charge compressibility is readily obtained,

$$\kappa = \frac{\mathcal{N}_*}{1 - \Gamma_c \mathcal{N}_*} \equiv \frac{\mathcal{N}_*}{1 + F_0^S},$$

and defines the Landau F_0^S parameter,

$$F_0^S = -\mathcal{N}_* \Gamma_c. \quad (100)$$

Since approaching the transition, $u \rightarrow 1$, $F_0^S \sim (1-u)^{-2}$ diverges faster than $\mathcal{N}_* \sim (1-u)^{-1}$, we find that the charge compressibility correctly vanishes at the metal-insulator transition (MIT). The expression of F_0^S coincides with that originally obtained by Vollhardt [23].

In the opposite limit of small $|\mathbf{q}|$ with respect to frequency,

$$\chi(\mathbf{q}, \omega) \simeq \frac{2T_0(1-u^2)(\gamma_0 - \gamma_{\mathbf{q}})}{\omega^2},$$

which, inserted into Eq. (97), allows calculation of the poles of the dynamical charge susceptibility, which are

$$\omega_{c\mathbf{q}}^2 = 4T_0^2(1+u^2)u(2-u)(\gamma_0 - \gamma_{\mathbf{q}}). \quad (101)$$

This acoustic mode is above the quasiparticle-quasihole continuum and actually corresponds to the Landau's zero sound. Once again, this result is compatible with Vollhardt's description of the correlated metal within the Gutzwiller approximation in the framework of Landau-Fermi-liquid theory [23]. Indeed, the zero sound velocity has the expected Landau's expression, once one realizes that in a lattice with infinite coordination, $F_1^S = 0$ and it is unrelated to the enhancement of the effective mass.

We conclude, highlighting that the velocity of the zero sound stays constant approaching the Mott transition. In particular, for $\omega^2 \gg T_0(1-u^2)(\gamma_0 - \gamma_{\mathbf{q}})$, the dynamical charge susceptibility can be written as

$$\chi_c(\mathbf{q} \rightarrow 0, \omega) = \frac{2T_0(1-u^2)(\gamma_0 - \gamma_{\mathbf{q}})}{\omega^2 - \omega_{c\mathbf{q}}^2}, \quad (102)$$

and hence the pole at the zero sound has vanishing weight as the transition $u \rightarrow 1$ is approached, in agreement with the expectation that spectral weight is transferred at high energy.

We conclude by observing that the propagator $\Pi_2(\mathbf{q}, \omega)$ of $p_2(\mathbf{q}, \omega)$,

$$\Pi_2(\mathbf{q}, \omega) = -\frac{1}{\omega^2} \frac{(1-u)\Gamma_c}{1 + \Gamma_c \chi(\mathbf{q}, \omega)},$$

is singular at $\omega = 0$, although this singularity does not appear in the physical response function, which is proportional to the propagator of the conjugate variable $x_2(\mathbf{q}, \omega)$. Indeed, $p_2(\mathbf{q}, \omega)$ is one of the would-be Goldstone modes that we mentioned in Sec. III. The action of the single-band Hubbard model is $U(2) = U(1) \times SU(2)$ gauge invariant, and $p_2(\mathbf{q}, \omega)$ is just the would-be Goldstone mode associated with the Abelian $U(1)$, whereas we shall see that $\mathbf{p}_1(\mathbf{q}, \omega)$ are instead those associated with $SU(2)$. In fact, the RPA form of the charge susceptibility could be very easily obtained by the gauge-fixing prescription of Sec. III. If we drop all terms that contain $p_2(i, t)$ and replace

$$-\sqrt{2} \sin \frac{\theta}{2} x_2(i, t) \rightarrow \langle n_i - 1 \rangle_t,$$

we get an effective Hamiltonian of the quasiparticles, neglecting for convenience all other variables but $x_2(i, t)$,

$$\mathcal{H}_*(t) = \mathcal{H}_*^{(0)} + \sum_i v_*(i, t)(n_i - 1),$$

where

$$v_*(i, t) = v(i, t) - \Gamma_c \langle n_i - 1 \rangle_t, \quad (103)$$

which readily leads to Eq. (97).

E. Dynamical spin susceptibility

In order to study the spin response, we imagine to add an external field that couples to the spin density, e.g., to its z component, namely,

$$\begin{aligned} \delta\mathcal{H}(t) &= -\sum_i B_3(i, t)(n_{i\uparrow} - n_{i\downarrow}) \\ &= -\sqrt{2} \cos \frac{\theta}{2} \sum_i B_3(i, t) x_{1,3}(i, t). \end{aligned} \quad (104)$$

In the metal phase, $\omega_1 = E_1 - E_0 = 2T_0(1-u)$, and repeating all calculations done for the charge susceptibility, we finally obtain the dynamical spin susceptibility,

$$\chi_s(\mathbf{q}, \omega) = \frac{\chi(\mathbf{q}, \omega)}{1 + \Gamma_s \chi(\mathbf{q}, \omega)}, \quad (105)$$

where

$$\Gamma_s = \frac{U}{2} \frac{1-u}{1+u} \left(1 + \frac{u}{2}\right) > 0. \quad (106)$$

The above expression reproduces the small u Stoner's enhancement of the magnetic susceptibility. In addition, it satisfies the relationship $\Gamma_s(U) = \Gamma_c(-U)$ valid at particle-hole symmetry [23]. Since $\Gamma_s \sim (1-u)$ vanishes linearly approaching the transition, the Landau's parameter,

$$F_0^A = -\mathcal{N}_* \Gamma_s < 0, \quad (107)$$

is constant for $u \rightarrow 1$, which implies that the uniform static spin susceptibility diverges at the MIT. This result agrees with previous ones [23,26] also obtained within the Gutzwiller approximation, but contrasts DMFT, which instead finds a finite uniform spin susceptibility at the transition.

Such a negative outcome critically depends on the fact that the effective interaction Γ_s , given by Eq. (106), vanishes at the transition. We are going to show that beyond the harmonic approximation, this cancellation no longer occurs.

We note that $p_{1a}(i, t)$, $a = 1, \dots, 3$, are now the Goldstone modes associated with $SU(2)$ gauge invariance, and their propagators,

$$\Pi_{1a}(\mathbf{q}, \omega) = -\frac{1}{\omega^2} \frac{(1+u)\Gamma_s}{1 + \Gamma_s \chi(\mathbf{q}, \omega)},$$

diverge at $\omega = 0$. We can, as in Sec. IV D, drop $\mathbf{p}_1(i, t)$ from the action and replace

$$\sqrt{2} \cos \frac{\theta}{2} \mathbf{x}_1(i, t) \rightarrow \langle \mathbf{c}_i^\dagger \boldsymbol{\sigma} \mathbf{c}_i \rangle_t,$$

whose effect could be absorbed into an effective magnetic field,

$$B_{*a}(i, t) = \delta_{a3} B_3(i, t) - \Gamma_s \langle \mathbf{c}_i^\dagger \boldsymbol{\sigma}_a \mathbf{c}_i \rangle_t, \quad (108)$$

that straightforwardly leads to Eq. (105).

F. Beyond RPA in the x_3 mode

We observe that all of the above results in the metal phase correspond to expanding the action at second order in the fluctuations, but treating the linear coupling between the latter and the fermions just within RPA, i.e., not accounting for exchange processes. While this procedure is somehow forced by gauge invariance when it concerns charge and spin modes (see the ending parts of Secs. IV D and IV E), it is not really compulsory for the $x_3(i, t)$ mode that describes the Hubbard bands. We can therefore take a first step forward when dealing with $x_3(i, t)$ in the direction of the so-called RPA+Exchange. According to Eq. (84), promoting x_3 and p_3 to quantum conjugate variables, after defining $t_* = t \sin^2 \theta$ and

$$X(i) = 1 + \sqrt{2} \cot \theta x_3(i),$$

the Hamiltonian reads

$$\begin{aligned} \mathcal{H}_* = & -\frac{t_*}{\sqrt{z}} \sum_{(ij)} (c_i^\dagger c_j + \text{H.c.}) X(i) X(j) \\ & + \sum_i \{ [v_*(i, t) c_i^\dagger \sigma_0 c_i + \mathbf{B}_*(i, t) \cdot c_i^\dagger \boldsymbol{\sigma} c_i] \\ & + \frac{\omega_3}{2} [x_3(i)^2 + p_3(i)^2] + T_0 \sin 2\theta \sqrt{2} x_3(i) \}, \end{aligned} \quad (109)$$

where the effective fields are those in Eqs. (103) and (108). The last term in Eq. (109), linear in x_3 , derives from Eq. (45) and cancels the linear term of the hopping when the latter is averaged over the Fermi sea, which is just the saddle-point condition for x_3 .

Near the Mott transition from the metal side, $u \lesssim 1$, since t_* is small with respect to ω_3 , we can integrate out x_3 and neglect the frequency dependence of its propagator $D_3(\mathbf{q}, \omega)$, which, through Eqs. (93) and (94), implies that

$$D_3(\mathbf{q}, \omega) = \frac{E_3 - E_0}{\omega^2 - \omega_{3\mathbf{q}}^2} \simeq -\frac{E_3 - E_0}{\omega_{3\mathbf{q}}^2} \simeq -\frac{E_3 - E_0}{\omega_{30}^2},$$

where we have furthermore neglected the momentum dependence.

In this approximation, the mode x_3 simply induces a nonretarded electron-electron interaction, which, within RPA+Exchange, leads to a change of the charge and spin susceptibilities,

$$\chi_{c(s)}(\mathbf{q}, \omega) \rightarrow \frac{\chi(\mathbf{q}, \omega)}{1 + \Gamma_{c(s)}(\mathbf{q}) \chi(\mathbf{q}, \omega)}, \quad (110)$$

where

$$\Gamma_{c(s)} \rightarrow \Gamma_{c(s)}(\mathbf{q}) = \Gamma_{c(s)} - \frac{t^2 u^2}{4T_0} \gamma_{\mathbf{q}}, \quad (111)$$

which also implies that the Landau parameters change into

$$F_0^{S(A)} \rightarrow -\mathcal{N}_* \Gamma_{c(s)}(\mathbf{0}). \quad (112)$$

The charge $F_0^S > 0$ keeps its singularity $(1 - u)^{-2}$, so that the charge compressibility still vanishes. On the contrary,

$$F_0^A \xrightarrow{u \rightarrow 1} \frac{t^2}{4T_0} \mathcal{N}_*, \quad (113)$$

so that the uniform spin susceptibility,

$$\chi = -\chi_s(\mathbf{q} \rightarrow \mathbf{0}, 0) \xrightarrow{u \rightarrow 1} \frac{4T_0}{t^2} = \frac{U_c}{2t^2}, \quad (114)$$

is now finite. Remarkably, this expression agrees with that obtained by DMFT [8], although the numerical value of U_c in DMFT is smaller than in the Gutzwiller approximation.

The quantum Hamiltonian (109) also allows calculation of the optical conductivity. In the presence of a small transverse vector potential $A_{i \rightarrow j}(t) = -A_{j \rightarrow i}(t)$, the Hamiltonian acquires an additional term,

$$\begin{aligned} \delta \mathcal{H}_*(t) = & -i \frac{t_*}{\sqrt{z}} \sum_{(ij)} A_{i \rightarrow j}(t) (c_i^\dagger c_j - \text{H.c.}) X(i) X(j) \\ & + \frac{t_*}{2\sqrt{z}} \sum_{(ij)} A_{i \rightarrow j}(t)^2 (c_i^\dagger c_j + \text{H.c.}) X(i) X(j). \end{aligned}$$

The calculation of the optical conductivity is straightforward and follows exactly that obtained within slave bosons in Ref. [40]. Besides the Drude peak that is obtained taking $X(i) = 1$, and vanishes like $\sin^2 \theta = 1 - u^2$ at the transition, the optical conductivity gets high-frequency contributions from the absorption spectrum of the mode x_3 [40].

V. CONCLUSIONS

In this paper, we have presented a quite simple method to calculate linear response functions within the Gutzwiller approximation, including in a consistent way quantum fluctuations in the harmonic approximation. The calculation is straightforward and just requires a little more effort than the equilibrium one. In fact, besides the variational matrix $\hat{\Phi}_0$ that minimizes the energy at equilibrium, and which can be regarded as the lowest-energy eigenstate of a local Hamiltonian [20,37] [see Eq. (31)], one also needs all excited eigenstates and eigenvalues. In a model that involves M correlated orbitals in each unit cell, this local Hamiltonian is defined in a Hilbert space of dimension $\binom{4M}{2M}$, and can be conveniently recast into the problem of an impurity with M orbitals hybridized to a single bath site with the same number of orbitals, with the coupled system being at half filling [20].

As a check, we have applied the method to the single-band Hubbard model at half filling and recovered all known results about charge compressibility and spin susceptibility within the Gutzwiller approximation [23–27,39–41]. As a by-product, we also showed how to cure one flaw of the Gutzwiller approximation, i.e., the divergence of the uniform magnetic susceptibility approaching the Mott transition from the metal side.

ACKNOWLEDGMENT

This work has been supported by the European Union under H2020 Framework Programs, ERC Advanced Grant No. 692670 ‘‘FIRSTORM.’’

**APPENDIX: THE WAVE-FUNCTION
RENORMALIZATION MATRIX $\hat{R}(i)$**

At equilibrium and in the natural basis, constraint Eq. (13) reads

$$\begin{aligned} \text{Tr}[\hat{\Phi}_0(i)^\dagger \hat{\Phi}_0(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta}] &= \text{Tr}[\hat{P}_0^{(0)}(i) \hat{d}_{i\alpha}^\dagger \hat{d}_{i\beta}] \\ &= \delta_{\alpha\beta} n_\alpha^{(0)}(i), \end{aligned}$$

where $\hat{P}_0(i)$ is the local probability distribution of the Slater determinant. Hereafter, we shall drop for simplicity the site index i .

We can always write $\hat{P}_0^{(0)}$ as the Boltzmann distribution of a noninteracting Hamiltonian,

$$H = \sum_\alpha \epsilon_\alpha n_\alpha,$$

where $f(\epsilon_\alpha) = n_\alpha^{(0)}$ is the Fermi distribution function. If $\hat{\Phi}$ is varied, also the probability distribution must vary in such a way as to preserve the constraint. This change will generally correspond to

$$H \rightarrow H + \delta H.$$

Since H must still be a one-body Hamiltonian, it follows that

$$d_\alpha(\tau) = e^{\tau H} d_\alpha e^{-\tau H} = (e^{-\hat{H}\tau} \mathbf{d})_\alpha = \sum_\beta U_{\beta\alpha}(\tau) d_\beta,$$

where \hat{H} is the matrix representation of H in the single-particle basis, so that $d_\alpha(\tau)$ remains a combination of creation operators. Since $\hat{U}(\tau_1)\hat{U}(\tau_2) = \hat{U}(\tau_1 + \tau_2)$, it trivially holds that $\hat{U}(\tau)\hat{U}(-\tau) = 1$ and

$$\hat{U}(\beta/2)\hat{U}(\beta/2) = \hat{U}(\beta). \quad (\text{A1})$$

The local probability distribution,

$$\hat{P}_0 = \frac{e^{-\beta\hat{H}}}{\text{Tr}(e^{-\beta\hat{H}})},$$

so that

$$\begin{aligned} \text{Tr}(\hat{P}_0 \hat{d}_\beta(\beta) \hat{d}_\alpha^\dagger) &= \text{Tr}(\hat{P}_0 \hat{d}_\alpha^\dagger \hat{d}_\beta) \equiv n_{\alpha\beta} \\ &= \sum_\gamma U_{\gamma\beta}(\beta) \text{Tr}(\hat{P}_0 d_\gamma d_\alpha^\dagger) \\ &= \sum_\gamma U_{\gamma\beta}(\beta) (\delta_{\alpha\gamma} - n_{\alpha\gamma}) \\ &= U_{\alpha\beta}(\beta) - \sum_\gamma n_{\alpha\gamma} U_{\gamma\beta}(\beta), \end{aligned}$$

namely,

$$\hat{U}(\beta) = (1 - \hat{n})^{-1} \hat{n} = -1 + (1 - \hat{n})^{-1}, \quad (\text{A2})$$

which relates $\hat{U}(\beta)$ to \hat{n} . It also follows that

$$\hat{U}(-\beta) = (1 - \hat{n})\hat{n}^{-1} = \hat{n}^{-1}(1 - \hat{n}) = \hat{n}^{-1} - 1. \quad (\text{A3})$$

The renormalization coefficients R are obtained by solving for any α and γ ,

$$\text{Tr}\left(\sqrt{\hat{P}_0} \hat{\Phi}^\dagger \hat{c}_\alpha^\dagger \hat{\Phi} \frac{1}{\sqrt{\hat{P}_0}} \hat{d}_\gamma\right) = \sum_\beta \text{Tr}(\hat{\Phi}^\dagger \hat{\Phi} \hat{d}_\beta^\dagger \hat{d}_\gamma) R_{\alpha\beta}^*, \quad (\text{A4})$$

where

$$\begin{aligned} \frac{1}{\sqrt{\hat{P}_0}} \hat{d}_\gamma \sqrt{\hat{P}_0} &= e^{\beta\hat{H}/2} \hat{d}_\gamma e^{-\beta\hat{H}/2} \\ &= \hat{d}_\gamma(\beta/2) = \sum_\beta U_{\beta\gamma}(\beta/2) \hat{d}_\beta. \end{aligned}$$

Therefore, once we define

$$Q_{\alpha\beta}^* \equiv \text{Tr}(\hat{\Phi}^\dagger \hat{c}_\alpha^\dagger \hat{\Phi} \hat{d}_\beta),$$

then Eq. (A4) is equivalent to

$$\sum_\beta Q_{\alpha\beta}^* U_{\beta\gamma}(\beta/2) = \sum_\beta R_{\alpha\beta}^* n_{\beta\gamma},$$

or, in matrix form, and observing that $\hat{n} = \hat{U}(\beta) - \hat{n}\hat{U}(\beta)$,

$$\begin{aligned} \hat{Q}^* \hat{U}(\beta/2) &= \hat{R}^* \hat{n} = \hat{R}^* [\hat{U}(\beta) - \hat{n}\hat{U}(\beta)] \\ &= \hat{R}^* \hat{U}(\beta) - \hat{Q}^* \hat{U}(\beta/2), \end{aligned}$$

so that, multiplying both sides on the right by $\hat{U}(-\beta)$, we finally get

$$\begin{aligned} \hat{R}^* &= \hat{Q}^* [\hat{U}(\beta/2) + \hat{U}(-\beta/2)] \\ &= \hat{Q}^* (\sqrt{\hat{U}(\beta)} + \sqrt{\hat{U}(-\beta)}) \\ &= \hat{Q}^* \left(\sqrt{\frac{\hat{n}}{1-\hat{n}}} + \sqrt{\frac{1-\hat{n}}{\hat{n}}} \right) \\ &= \hat{Q}^* (\sqrt{\hat{n}(1-\hat{n})})^{-1}. \end{aligned}$$

We denote as

$$\hat{S}^* = (\sqrt{\hat{n}(1-\hat{n})})^{-1} = \hat{S}^T,$$

since $\hat{S} = \hat{S}^\dagger$, so that

$$\hat{R}^* = \hat{Q}^* \hat{S}^* \longrightarrow \hat{R}^\dagger = \hat{S}^\dagger \hat{Q}^\dagger = \hat{S} \hat{Q}^\dagger,$$

namely, the desired result,

$$\hat{R} = \hat{Q} \hat{S}. \quad (\text{A5})$$

One can rewrite

$$4\hat{S}^{-2} = 4\hat{n}^T(1-\hat{n}^T) = 1 - (1-2\hat{n}^T)^2 \equiv 1 - \hat{\Delta}^2,$$

where the matrix elements of $\hat{\Delta}$ are

$$\Delta_{\alpha\beta} = \delta_{\alpha\beta} - 2\text{Tr}(\hat{\Phi}^\dagger \hat{\Phi} \hat{d}_\beta^\dagger \hat{d}_\alpha) = \text{Tr}(\hat{\Phi}^\dagger \hat{\Phi} [\hat{d}_\alpha^\dagger, \hat{d}_\beta^\dagger]). \quad (\text{A6})$$

At equilibrium,

$$\Delta_{\alpha\beta}^{(0)} = \delta_{\alpha\beta} (1 - 2n_\alpha^{(0)}), \quad (\text{A7})$$

$$S_{\alpha\beta}^{(0)} = \delta_{\alpha\beta} / \sqrt{n_\alpha^{(0)}(1-n_\alpha^{(0)})} \equiv \delta_{\alpha\beta} S_\alpha^{(0)}, \quad (\text{A8})$$

are diagonal, which allows an explicit evaluation of the matrix derivatives. It follows that the equilibrium renormalization matrix has elements

$$R_{\alpha\beta}^{(0)} = \text{Tr}(\hat{\Phi}_0^\dagger \hat{c}_\alpha^\dagger \hat{\Phi}_0 \hat{d}_\beta^\dagger) S_\beta^{(0)} \equiv Q_{\alpha\beta}^{(0)} S_\beta^{(0)}. \quad (\text{A9})$$

Derivatives of \hat{R}

We write

$$\hat{\Phi} = \sum_n \phi_n \hat{\Phi}_n, \quad \hat{\Phi}^\dagger = \sum_n \phi_n^* \hat{\Phi}_n^\dagger,$$

where $\hat{\Phi}_n$ is a basis set,

$$\text{Tr}(\hat{\Phi}_n^\dagger \hat{\Phi}_m) = \delta_{nm},$$

with $\hat{\Phi}_0$ the equilibrium solution. By inspection, we realize that

$$\frac{\partial R_{\alpha\beta}}{\partial \hat{\Phi}^\dagger} = \hat{\Gamma}_{\alpha\beta}[\hat{\Phi}, \hat{\Phi}^\dagger] \hat{\Phi},$$

where the tensor $\hat{\Gamma}_{\alpha\beta}[\hat{\Phi}, \hat{\Phi}^\dagger]$ is still a functional of $\hat{\Phi}$ and $\hat{\Phi}^\dagger$. Therefore,

$$\frac{\partial R_{\alpha\beta}}{\partial \phi_n^*} = \text{Tr}(\hat{\Phi}_n^\dagger \hat{\Gamma}_{\alpha\beta}[\hat{\Phi}, \hat{\Phi}^\dagger] \hat{\Phi}).$$

The equilibrium value is obtained by setting $\phi_n = \delta_{n0}$.

In particular, exploiting the fact that \hat{S} is diagonal at equilibrium, the first-order derivatives evaluated at equilibrium explicitly read

$$\frac{\partial R_{\alpha\beta}}{\partial \phi_n^*} = \frac{\partial Q_{\alpha\beta}}{\partial \phi_n^*} S_\beta^{(0)} + \sum_\gamma Q_{\alpha\gamma}^{(0)} S_\gamma^{(0)} F_{\gamma\beta} \frac{\partial \Delta_{\gamma\beta}}{\partial \phi_n^*}, \quad (\text{A10})$$

$$\frac{\partial R_{\alpha\beta}}{\partial \phi_n} = \frac{\partial Q_{\alpha\beta}}{\partial \phi_n} S_\beta^{(0)} + \sum_\gamma Q_{\alpha\gamma}^{(0)} S_\gamma^{(0)} F_{\gamma\beta} \frac{\partial \Delta_{\gamma\beta}}{\partial \phi_n}, \quad (\text{A11})$$

while the second derivative, still calculated at equilibrium, is

$$\begin{aligned} \frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m} &= \text{Tr}(\hat{\Phi}_n^\dagger \hat{\Gamma}_{\alpha\beta}[\hat{\Phi}_0, \hat{\Phi}_0^\dagger] \hat{\Phi}_m) \\ &+ \text{Tr}\left(\hat{\Phi}_n^\dagger \frac{\partial \hat{\Gamma}_{\alpha\beta}[\hat{\Phi}, \hat{\Phi}^\dagger]}{\partial \phi_m} \Big|_0 \hat{\Phi}_0\right) \\ &\equiv \left(\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m}\right)_1 + \left(\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m}\right)_2, \end{aligned} \quad (\text{A12})$$

where

$$\left(\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m}\right)_1 = \sum_\gamma \left[\frac{\partial^2 Q_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m} S_\beta^{(0)} + Q_{\alpha\gamma}^{(0)} F_{\gamma\beta} \frac{\partial^2 \Delta_{\gamma\beta}}{\partial \phi_n^* \partial \phi_m} \right], \quad (\text{A13})$$

$$\begin{aligned} \left(\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m}\right)_2 &= \sum_\gamma \left[\frac{\partial Q_{\alpha\gamma}}{\partial \phi_n^*} F_{\gamma\beta} \frac{\partial \Delta_{\gamma\beta}}{\partial \phi_m} + \frac{\partial Q_{\alpha\gamma}}{\partial \phi_m} F_{\gamma\beta} \frac{\partial \Delta_{\gamma\beta}}{\partial \phi_n^*} \right. \\ &\left. + Q_{\alpha\gamma}^{(0)} \left(\frac{\partial^2 S_{\gamma\beta}}{\partial \phi_n^* \partial \phi_m}\right)_2 \right]. \end{aligned} \quad (\text{A14})$$

The terms that appear in the above equations are

$$\begin{aligned} \frac{\partial Q_{\alpha\beta}}{\partial \phi_n^*} &= \text{Tr}(\hat{\Phi}_n^\dagger \hat{c}_\alpha \hat{\Phi}_0 \hat{d}_\beta^\dagger), \\ \frac{\partial Q_{\alpha\beta}}{\partial \phi_n} &= \text{Tr}(\hat{\Phi}_0^\dagger \hat{c}_\alpha \hat{\Phi}_n \hat{d}_\beta^\dagger), \\ \frac{\partial \Delta_{\alpha\beta}}{\partial \phi_n^*} &= \text{Tr}(\hat{\Phi}_n^\dagger \hat{\Phi}_0 [\hat{d}_\alpha, \hat{d}_\beta^\dagger]), \end{aligned}$$

$$\frac{\partial \Delta_{\alpha\beta}}{\partial \phi_n} = \text{Tr}(\hat{\Phi}_0^\dagger \hat{\Phi}_n [\hat{d}_\alpha, \hat{d}_\beta^\dagger]),$$

$$F_{\alpha\beta} = \frac{1}{2} \frac{(S_\alpha^{(0)} S_\beta^{(0)})^2}{S_\alpha^{(0)} + S_\beta^{(0)}} (1 - n_\alpha^{(0)} - n_\beta^{(0)}),$$

$$\frac{\partial^2 Q_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m} = \text{Tr}(\hat{\Phi}_n^\dagger \hat{c}_\alpha \hat{\Phi}_m \hat{d}_\beta^\dagger),$$

$$\frac{\partial^2 \Delta_{\alpha\beta}(i)}{\partial \phi_n^* \partial \phi_m} = \text{Tr}(\hat{\Phi}_n^\dagger \hat{\Phi}_m [\hat{d}_\alpha, \hat{d}_\beta^\dagger]),$$

and, lastly,

$$\begin{aligned} &\left(\frac{\partial^2 S_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m}\right)_2 \\ &= \frac{(S_\alpha^{(0)} S_\beta^{(0)})^2}{S_\alpha^{(0)} + S_\beta^{(0)}} \sum_\gamma \left[\frac{\partial \Delta_{\alpha\gamma}}{\partial \phi_n^*} \frac{\partial \Delta_{\gamma\beta}}{\partial \phi_m} + \frac{\partial \Delta_{\alpha\gamma}}{\partial \phi_m} \frac{\partial \Delta_{\gamma\beta}}{\partial \phi_n^*} \right] \\ &\times \left[\frac{1}{4} + F_{\alpha\gamma} F_{\gamma\beta} \frac{S_\alpha^{(0)} S_\gamma^{(0)} + S_\gamma^{(0)} S_\beta^{(0)} + S_\beta^{(0)} S_\alpha^{(0)}}{(S_\alpha^{(0)} S_\gamma^{(0)} S_\beta^{(0)})^2} \right]. \end{aligned}$$

In addition,

$$\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m^*} = \left(\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n^* \partial \phi_m^*}\right)_2, \quad (\text{A15})$$

$$\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n \partial \phi_m} = \left(\frac{\partial^2 R_{\alpha\beta}}{\partial \phi_n \partial \phi_m}\right)_2, \quad (\text{A16})$$

where the right-hand sides are obtained straightforwardly through Eq. (A14). The above derivatives calculated at the equilibrium solution allow calculation of the Taylor expansion of \hat{R} . In particular, through Eqs. (A10) and (A11), the first-order expansion is

$$\hat{R}^{(1)} = \sum_n \left[\phi_n^* \frac{\partial R_{\alpha\beta}}{\partial \phi_n^*} + \phi_n \frac{\partial R_{\alpha\beta}}{\partial \phi_n} \right], \quad (\text{A17})$$

while the second-order expansion, mentioned in Eq. (50), is

$$\hat{R}^{(2)} = \hat{R}_1^{(2)} + \hat{R}_1^{(2)}, \quad (\text{A18})$$

where, explicitly,

$$\hat{R}_1^{(2)} = \sum_{nm} \phi_n^* \phi_m \left(\frac{\partial^2 \hat{R}}{\partial \phi_n^* \partial \phi_m}\right)_1, \quad (\text{A19})$$

and

$$\begin{aligned} \hat{R}_2^{(2)} &= \frac{1}{2} \sum_{nm} \left[2 \phi_n^* \phi_m \left(\frac{\partial^2 \hat{R}}{\partial \phi_n^* \partial \phi_m}\right)_2 + \phi_n^* \phi_m^* \left(\frac{\partial^2 \hat{R}}{\partial \phi_n^* \partial \phi_m^*}\right)_2 \right. \\ &\left. + \phi_n \phi_m \left(\frac{\partial^2 \hat{R}}{\partial \phi_n \partial \phi_m}\right)_2 \right]. \end{aligned} \quad (\text{A20})$$

- [1] M. C. Gutzwiller, *Phys. Rev. Lett.* **10**, 159 (1963).
- [2] M. C. Gutzwiller, *Phys. Rev.* **137**, A1726 (1965).
- [3] H. Yokoyama and H. Shiba, *J. Phys. Soc. Jpn.* **56**, 1490 (1987).
- [4] M. Capello, F. Becca, M. Fabrizio, S. Sorella, and E. Tosatti, *Phys. Rev. Lett.* **94**, 026406 (2005).
- [5] W. F. Brinkman and T. M. Rice, *Phys. Rev. B* **2**, 4302 (1970).
- [6] W. Metzner and D. Vollhardt, *Phys. Rev. Lett.* **62**, 324 (1989).
- [7] J. Bünemann, W. Weber, and F. Gebhard, *Phys. Rev. B* **57**, 6896 (1998).
- [8] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (1996).
- [9] X. Deng, X. Dai, and Z. Fang, *Europhys. Lett.* **83**, 37008 (2008).
- [10] K. M. Ho, J. Schmalian, and C. Z. Wang, *Phys. Rev. B* **77**, 073101 (2008).
- [11] X. Y. Deng, L. Wang, X. Dai, and Z. Fang, *Phys. Rev. B* **79**, 075114 (2009).
- [12] G. T. Wang, Y. Qian, G. Xu, X. Dai, and Z. Fang, *Phys. Rev. Lett.* **104**, 047002 (2010).
- [13] M.-F. Tian, X. Deng, Z. Fang, and X. Dai, *Phys. Rev. B* **84**, 205124 (2011).
- [14] Y. X. Yao, C. Z. Wang, and K. M. Ho, *Int. J. Quantum Chem.* **112**, 240 (2012).
- [15] T. Schickling, F. Gebhard, J. Bünemann, L. Boeri, O. K. Andersen, and W. Weber, *Phys. Rev. Lett.* **108**, 036406 (2012).
- [16] N. Lanatà, Y.-X. Yao, C.-Z. Wang, K.-M. Ho, J. Schmalian, K. Haule, and G. Kotliar, *Phys. Rev. Lett.* **111**, 196801 (2013).
- [17] N. Lanatà, Y.-X. Yao, C.-Z. Wang, K.-M. Ho, and G. Kotliar, *Phys. Rev. B* **90**, 161104 (2014).
- [18] G. Borghi, M. Fabrizio, and E. Tosatti, *Phys. Rev. B* **90**, 125102 (2014).
- [19] M.-F. Tian, H.-F. Song, H.-F. Liu, C. Wang, Z. Fang, and X. Dai, *Phys. Rev. B* **91**, 125148 (2015).
- [20] N. Lanatà, Y. Yao, C.-Z. Wang, K.-M. Ho, and G. Kotliar, *Phys. Rev. X* **5**, 011008 (2015).
- [21] E. Runge and E. K. U. Gross, *Phys. Rev. Lett.* **52**, 997 (1984).
- [22] E. K. U. Gross, J. F. Dobson, and M. Petersilka, in *Density Functional Theory II*, edited by R. F. Nalewajski, Topics in Current Chemistry, Vol. 181 (Springer, Berlin, 1996), pp. 81–172.
- [23] D. Vollhardt, *Rev. Mod. Phys.* **56**, 99 (1984).
- [24] G. Seibold and J. Lorenzana, *Phys. Rev. Lett.* **86**, 2605 (2001).
- [25] G. Seibold, F. Becca, and J. Lorenzana, *Phys. Rev. B* **67**, 085108 (2003).
- [26] G. Seibold, F. Becca, P. Rubin, and J. Lorenzana, *Phys. Rev. B* **69**, 155113 (2004).
- [27] M. Schiró and M. Fabrizio, *Phys. Rev. B* **83**, 165105 (2011).
- [28] J. Bnemann, M. Capone, J. Lorenzana, and G. Seibold, *New J. Phys.* **15**, 053050 (2013).
- [29] E. v. Oelsen, G. Seibold, and J. Bünemann, *Phys. Rev. Lett.* **107**, 076402 (2011).
- [30] E. von Oelsen, G. Seibold, and J. Bnemann, *New J. Phys.* **13**, 113031 (2011).
- [31] M. Schiró and M. Fabrizio, *Phys. Rev. Lett.* **105**, 076401 (2010).
- [32] G. Kotliar and A. E. Ruckenstein, *Phys. Rev. Lett.* **57**, 1362 (1986).
- [33] F. Lechermann, A. Georges, G. Kotliar, and O. Parcollet, *Phys. Rev. B* **76**, 155102 (2007).
- [34] M. Fabrizio, in *New Materials for Thermoelectric Applications: Theory and Experiment*, edited by V. Zlatic and A. Hewson, NATO Science for Peace and Security Series B: Physics and Biophysics (Springer, Netherlands, 2013), pp. 247–273.
- [35] M. Fabrizio, *Phys. Rev. B* **76**, 165110 (2007).
- [36] N. Lanatà, P. Barone, and M. Fabrizio, *Phys. Rev. B* **78**, 155127 (2008).
- [37] N. Lanatà, H. U. R. Strand, X. Dai, and B. Hellsing, *Phys. Rev. B* **85**, 035133 (2012).
- [38] N. Read, *J. Phys. C* **18**, 2651 (1985).
- [39] T. Jolicœur and J. C. Le Guillou, *Phys. Rev. B* **44**, 2403 (1991).
- [40] R. Raimondi and C. Castellani, *Phys. Rev. B* **48**, 11453 (1993).
- [41] M. Sandri, M. Schiró, and M. Fabrizio, *Phys. Rev. B* **86**, 075122 (2012).
- [42] T. Li, P. Wölfle, and P. J. Hirschfeld, *Phys. Rev. B* **40**, 6817 (1989).
- [43] S. D. Huber and A. Rüegg, *Phys. Rev. Lett.* **102**, 065301 (2009).
- [44] A. Rüegg, S. D. Huber, and M. Sigrist, *Phys. Rev. B* **81**, 155118 (2010).