

Symmetry breaking in occupation number based slave-particle methods

Alexandru B. Georgescu^{1,2} and Sohrab Ismail-Beigi^{1,2,3,4}

¹*Department of Physics, Yale University, New Haven, Connecticut 06520, USA*

²*Center of Research on Interface Structure and Phenomena, Yale University, New Haven, Connecticut 06520, USA*

³*Department of Applied Physics, Yale University, New Haven, Connecticut 06520, USA*

⁴*Department of Mechanical Engineering and Materials Science, Yale University, New Haven, Connecticut 06520, USA*

(Received 4 April 2017; revised manuscript received 2 August 2017; published 17 October 2017)

We describe a theoretical approach to finding spontaneously symmetry-broken electronic phases due to strong electronic interactions when using recently developed slave-particle (slave-boson) approaches based on occupation numbers. We describe why, to date, spontaneous symmetry breaking has proven difficult to achieve in such approaches. We then provide a total energy based approach for introducing auxiliary symmetry-breaking fields into the solution of the slave-particle problem that leads to lowered total energies for symmetry-broken phases. We point out that not all slave-particle approaches yield energy lowering: the slave-particle model being used must explicitly describe the degrees of freedom that break symmetry. Finally, our total energy approach permits us to greatly simplify the formalism used to achieve a self-consistent solution between spinon and slave modes while increasing the numerical stability and greatly speeding up the calculations.

DOI: [10.1103/PhysRevB.96.165135](https://doi.org/10.1103/PhysRevB.96.165135)

I. INTRODUCTION

The effects of strong electronic interactions and electronic correlations on materials properties is a subject with a considerable history. The most celebrated textbook example is the Mott transition, where with increasing strength of localized electronic repulsions, electrons in the material lose band mobility and instead localize at atomic sites (i.e., loss of wave behavior). However, electronic correlations also underlie many other ordered electronic phases such as various forms of magnetism as well as superconductivity. A canonical model Hamiltonian for correlated electrons is the (extended) Hubbard model, where electrons can hop between localized orbitals centered at atomic sites but multiple-electron occupancy of a given atomic site leads to a significant energy penalty U . By varying the ratio of U to the band hopping parameters, one can cover the range from weak to strong electronic interactions and correlations [1].

The workhorse in realistic first principles calculations in crystal and electronic structure calculations, density functional theory (DFT) [2,3], is fundamentally based on a description of noninteracting electrons, i.e., band theory. Due to its simple structure, band theory approaches cannot capture the effects of dynamical electronic fluctuations and localized correlations on electronic band spectra. Extensions of DFT to go beyond local exchange-correlation potentials and to include nonlocal Hartree-Fock-type electronic behavior, such as the DFT + U and hybrid functional approaches [4,5], can capture certain effects of electron-electron interactions, especially for strongly symmetry-broken situations. Nevertheless, these are still band theory descriptions incapable of leading, e.g., to electron localization without resorting to symmetry breaking.

More advanced computational many-body approaches for simulation of electronic correlations are based on Green's functions methods. One type of approach is the GW approximation to the electron self-energy [6–8], which is a fully *ab initio* approach that includes the physics of nonlocal and dynamical electronic screening and produces accurate results for electronic band energies of a wide variety of materials

[8,9]. However, the GW method is based on summation of a subset of many-body diagrams (RPA diagrams) and thus does not capture a number of physical effects; separately GW calculations are notoriously expensive in terms of computation time due to their fully *ab initio* nature and lack of a particular basis set. Another avenue of approach is represented by dynamical mean-field theory (DMFT) [10,11], which can include the effect of local interactions and dynamical fluctuations by solving a model Hamiltonian with local interactions exactly (i.e., all diagrams for the local interactions are included). However, DFT + DMFT calculations on realistic materials with large unit cells are still quite challenging, as they require large-scale parallel computations.

For all these reasons, approximate and efficient methods for solving correlated problems continue to be of interest to the computational many-body community. One set of methods of recent interest for solving Hubbard models is slave-particle (slave-boson) methods. These methods, which have a long background in condensed matter theory, have been used to study cases with infinitely strong repulsive interactions [12–18]. Dealing with finite interaction strengths was enabled by the Kotliar-Ruckenstein approach [17], whose variants and modifications have been applied to the study of high-temperature superconductors [19] as well as multiband models [20–22] to elucidate the effects of multiple orbitals, degeneracy, and Hund's coupling [20,21]. In these approaches, each bosonic slave degree of freedom tracks the occupancy of a particular electronic configuration of a correlated site: once multiple orbitals and multiple electron counts can exist at a site, the number of required bosons becomes large. These methods can and have been used to describe spontaneously broken electronic symmetry (e.g., magnetic) states [17,20,23]. In addition, fully rotationally invariant slave-boson formalisms have been designed that permit spontaneous breaking of particle-number conservation and superconducting solutions [24].

A recent set of more economical slave-particle methods has been developed and become of wider interest, such as the slave-rotor method [25,26] and its application to nickelate oxides [27] and the slave-spin method [28–30] and its application to

iron-based superconductors [31]. Recently, we have developed a generalized version of these methods that does not require the analogy with spin or angular momentum and introduces multiple intermediate slave-particle models [32]. These recent approaches use the slave degrees of freedom to track the electron occupation number at a site, and its distribution among orbital and spin channels, and thus require a much smaller number of bosons per site.

However, in all the previous literature in which these occupation number based methods have been used, spontaneous symmetry breaking has been achieved in multiorbital systems where both a Hubbard U and a nonzero Hund's J interaction have been operative [27,29,31]. For a system where only the repulsion U operates, spontaneous symmetry breaking has not been displayed, even when interaction-induced magnetism is a feature of the actual ground state of the model Hamiltonian [e.g., ground-state antiferromagnetic (AFM) order for a half-filled single-band Hubbard model]. Indeed, as we show, stabilizing a purely interaction-induced symmetry-broken phase is very difficult for slave-particle methods without the introduction of symmetry-breaking fields. Our work describes this issue in detail and provides a total energy approach that naturally produces symmetry breaking. We then show how one can make slave-particle self-consistency between spinon and slave modes much more efficient via a specific and exact decoupling of the two modes.

II. THE SLAVE-PARTICLE APPROACH

In this section we review the key aspects of the slave-particle formalism used in previous work to set up the notation and language used in subsequent sections. The general correlated-electron Hamiltonian we consider is an extended Hubbard model given by

$$\hat{H} = \sum_i \hat{H}_{\text{int}}^i + \sum_{im\sigma} \epsilon_{im\sigma} \hat{d}_{im\sigma}^\dagger \hat{d}_{im\sigma} - \sum_{ii'mm'\sigma} t_{ii'mm'\sigma} \hat{d}_{im\sigma}^\dagger \hat{d}_{i'm'\sigma}. \quad (1)$$

The \hat{d} are canonical fermion annihilation operators. The indices i and i' range over the localized sites in the system (usually atomic sites), mm' range over the localized spatial orbitals on each site, $\sigma = \pm 1$ denotes spin, \hat{H}_{int}^i is the local Coulombic interaction for site i , $\epsilon_{im\sigma}$ is the on-site energy of the state labeled $im\sigma$, and $t_{ii'mm'\sigma}$ is the spin-conserving hopping element term connecting orbital $im\sigma$ to orbital $i'm'\sigma$. A commonly used interaction term is given by the Slater-Kanamori form [33]

$$\begin{aligned} \hat{H}_{\text{int}}^i = & \frac{U_i}{2} (\hat{n}_i^2 - \hat{n}_i) + \frac{U_i - U'_i}{2} \sum_{m \neq m'} \hat{n}_{im} \hat{n}_{im'} \\ & - \frac{J_i}{2} \sum_{\sigma} \sum_{m \neq m'} \hat{n}_{im\sigma} \hat{n}_{im'\sigma} \\ & - \frac{J_i}{2} \sum_{\sigma} \sum_{m \neq m'} (\hat{d}_{im\sigma}^\dagger \hat{d}_{im\bar{\sigma}} \hat{d}_{im'\bar{\sigma}}^\dagger \hat{d}_{im'\sigma} \\ & + \hat{d}_{im\sigma}^\dagger \hat{d}_{im\bar{\sigma}}^\dagger \hat{d}_{im'\sigma} \hat{d}_{im'\bar{\sigma}}). \end{aligned} \quad (2)$$

While the Coulombic parameters U_i , U'_i , and J_i can in principle depend on the site index i , in practice in most models they are assumed to be the same for all correlated sites. Briefly, the U term describes repulsion between the same spatial orbitals on a site, U' describes repulsion between different orbitals, and J measures the strength of the Hund's interaction between different orbitals with the same spin state. The number operators are

$$\hat{n}_{im\sigma} = \hat{d}_{im\sigma}^\dagger \hat{d}_{im\sigma}, \quad \hat{n}_{im} = \sum_{\sigma} \hat{n}_{im\sigma}, \quad \hat{n}_i = \sum_{m\sigma} \hat{n}_{im\sigma}.$$

The interacting Hubbard problem is impossible to solve exactly and even difficult to solve approximately. Some of the complexity is due to the fact that the interacting fermions have both charge and spin degrees of freedom. In slave-boson approaches [12–18], one separates the spin from the charge degrees of freedom at each site by introducing a spinless charged bosonic “slave” degree of freedom at each site along with a spinful neutral fermion termed a spinon. The spinon and slave-boson annihilation operators are indicated by the \hat{f} and \hat{O} operators, respectively. Specifically, the electron-field operator is decomposed as

$$\hat{d}_{im\sigma} = \hat{f}_{im\sigma} \hat{O}_{i\alpha}, \quad \hat{d}_{im\sigma}^\dagger = \hat{f}_{im\sigma}^\dagger \hat{O}_{i\alpha}^\dagger. \quad (3)$$

The index α is part of our generalized notation [32] that permits us to unify different occupation number based slave-particle models. At each site, the α 's label disjoint sets of localized states $m\sigma$. The meaning of α depends on the type of slave-particle model chosen, and α refers to a subset of the $m\sigma$ indices that belong to a site i . For example, if we use a slave-rotor model for the correlated orbitals at a site [25,26], then α is nil: $\hat{O}_{i\alpha} = \hat{O}_i$. Namely, we have a single slave particle at each site i that only tracks the total number of electrons at that site. At the opposite limit, we can have a unique slave boson for each $m\sigma$ combination at a site (the “slave-spin” method [28,29]), so that in this case $\alpha = m\sigma$.

In the slave-particle approach used in this work, as well as prior literature on this class of methods [25,26,28,29,32], the \hat{O} operators are not standard bosonic-field annihilation operators and thus do not obey canonical bosonic commutation relations. Instead, they simply lower the number of particles, $\hat{O}|N\rangle = |N-1\rangle$ (there is no factor of \sqrt{N} present). This is required for the type of slave-boson approach we use to be faithful to the original description: i.e., the matrix elements of \hat{d} and $\hat{f}\hat{O}$ are identical between physical states and they both obey fermionic commutation relations. The equality of matrix elements is shown in the Appendix of Ref. [32] and a separate demonstration of faithfulness is found in the Appendix here.

The introduction of slave bosons by itself does not make solution of the Hubbard model any easier, as more degrees of freedom have been introduced to further enlarge the Hilbert space. To avoid sampling of unphysical states in the enlarged spinon + slave Hilbert space which have no correspondence to those in the original electronic Hilbert space, one must ensure that the number of slave particles and number of spinons track each other. More precisely, as Eq. (3) shows, spinon and slave particles are created or annihilated at the same time so that only state kets in the extended Hilbert space that obey this

condition are physical. Hence, one must ensure that

$$\hat{d}_{i\alpha}^\dagger \hat{d}_{i\alpha} = \hat{f}_{i\alpha}^\dagger \hat{f}_{i\alpha}$$

and also that the subset of physical states $|\Psi_{\text{phys}}\rangle$ must obey

$$\hat{n}_{i\alpha} |\Psi_{\text{phys}}\rangle = \hat{N}_{i\alpha} |\Psi_{\text{phys}}\rangle, \quad (4)$$

where $\hat{N}_{i\alpha}$ is the number counting operator for the slave particles and the corresponding particle count for spinons is

$$\hat{n}_{i\alpha} = \sum_{m\sigma \in \alpha} \hat{f}_{i\alpha}^\dagger \hat{f}_{i\alpha}. \quad (5)$$

The constraint of Eq. (4) ensures that the number of slave bosons and the number of spinons match exactly for each α index [25,26,28,32].

The key approximation that makes the slave-boson approach more tractable than the original problem is to assume a separable form for the overall wave function of the system, which takes a product form $|\Psi_f\rangle|\Phi_s\rangle$, where $|\Psi_f\rangle$ is a spinor-only state ket and $|\Phi_s\rangle$ is a slave-only state ket. This means that one can only enforce the above operator constraints on average

$$\langle \hat{n}_{i\alpha} \rangle_f = \langle \hat{N}_{i\alpha} \rangle_s, \quad (6)$$

where the spinon and slave averages for any operator \hat{A} are defined via

$$\langle \hat{A} \rangle_f = \langle \Psi_f | \hat{A} | \Psi_f \rangle, \quad \langle \hat{A} \rangle_s = \langle \Phi_s | \hat{A} | \Phi_s \rangle.$$

We point out that the matching condition of Eq. (6) can go beyond simply setting the total number of spinons and slaves equal at each site. When the index α is sufficiently fine-grained, the matching is a much stronger constraint: for example, if α distinguishes different spin directions, then the numbers of spinons and slaves must match for each spin direction separately. As we show below, this is crucial for correctly describing situations where symmetry is broken because it ensures that the broken symmetry appears in both spinon and slave sectors simultaneously.

This separability assumption means that one must solve two separate and easier eigenvalue problems,

$$\hat{H}_f |\Psi_f\rangle = E_f |\Psi_f\rangle, \quad \hat{H}_s |\Phi_s\rangle = E_s |\Phi_s\rangle,$$

in a self-consistent fashion. In the simplest case of the interaction Hamiltonian, where $U = U'$ and $J = 0$, the spinon Hamiltonian is given by

$$\begin{aligned} \hat{H}_f = & \sum_{i\alpha} \epsilon_{i\alpha} \hat{f}_{i\alpha}^\dagger \hat{f}_{i\alpha} - \sum_{i\alpha} h_{i\alpha} \hat{n}_{i\alpha} \\ & - \sum_{ii'\alpha'} \langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'} \rangle_s \sum_{\substack{m\sigma \in \alpha \\ m'\sigma \in \alpha'}} t_{imi'm'\sigma} \hat{f}_{i\alpha}^\dagger \hat{f}_{i'\alpha'}, \end{aligned} \quad (7)$$

while the slave-boson Hamiltonian takes the form

$$\begin{aligned} \hat{H}_s = & \sum_i \hat{H}_{\text{int}}^i + \sum_\alpha h_{i\alpha} \hat{N}_{i\alpha} \\ & - \sum_{ii'\alpha'} \left[\sum_{\substack{m\sigma \in \alpha \\ m'\sigma \in \alpha'}} t_{imi'm'\sigma} \langle \hat{f}_{i\alpha}^\dagger \hat{f}_{i'\alpha'} \rangle_f \right] \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'}, \end{aligned} \quad (8)$$

where the spinon averages $\langle \hat{f}_{i\alpha}^\dagger \hat{f}_{i'\alpha'} \rangle_f$ renormalize the slave boson hoppings (and vice versa). The slave-boson problem is one of interacting charged bosons without spin on a lattice. Self-consistency refers to the fact that the spinon Hamiltonian involves averaged quantities involving the slave wave function, and vice versa. In addition, the values of the Lagrange multipliers $h_{i\alpha}$ must be chosen to ensure average particle-number matching as per Eq. (6).

As stated above, the forms written in Eqs. (7) and (8) are for the simplest “ U -only” form of the interaction term \hat{H}_{int} . More generalized forms for \hat{H}_f and \hat{H}_s that describe the generic multiband case (where $U \neq U'$ and $J \neq 0$) can be found in Secs. II D 1 to II D 4 of Ref. [32]. We note that these slave-boson methods have successfully described multiband cases and the effects of Hund’s interactions on a number of physical properties (e.g., band narrowing, orbitally selective Mott transitions, magnetism, etc.) [27,28,30,32]. However, in the following, we focus on the simplest case ($U = U'$ and $J = 0$), which is already highly informative about the shortcomings of these slave-boson approaches regarding symmetry breaking: we use the simple forms in Eqs. (7) and (8) in the remainder of this work.

III. SINGLE-SITE MEAN-FIELD APPROXIMATION

In practice, the slave Hamiltonian, Eq. (8), represents a many-body interacting bosonic problem that has no exact solution. In what follows, when solving numerically for the ground state of a spinon + slave problem, we use a simple single-site mean-field approach: when dealing with site i in the slave problem, we replace the $\hat{O}_{i\alpha}$ slave operators at the other neighboring sites by their averages $\langle \hat{O}_{i\alpha} \rangle_s$. For the spinon Hamiltonian, this boils down to the simple replacement

$$\langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'} \rangle_s \rightarrow \langle \hat{O}_{i\alpha}^\dagger \rangle_s \langle \hat{O}_{i'\alpha'} \rangle_s$$

in the hopping term. The slave Hamiltonian turns into

$$\begin{aligned} \hat{H}_s = & \sum_i \hat{H}_{\text{int}}^i + \sum_\alpha h_{i\alpha} \hat{N}_{i\alpha} \\ & - \sum_{ii'\alpha'} \left[\sum_{\substack{m\sigma \in \alpha \\ m'\sigma \in \alpha'}} t_{imi'm'\sigma} \langle \hat{f}_{i\alpha}^\dagger \hat{f}_{i'\alpha'} \rangle_f \right] \\ & \times [\langle \hat{O}_{i\alpha}^\dagger \rangle_s \hat{O}_{i'\alpha'} + \text{H.c.}], \end{aligned} \quad (9)$$

which is a simple many-body system of isolated sites where the bosonic $\hat{O}_{i\alpha}$ and $\hat{O}_{i\alpha}^\dagger$ operators remove and add bosons to the site from an effective bosonic mean-field bath. We note that, for the simple model Hamiltonians we use below in this approach, the quasiparticle renormalization factor (or weight) Z is simply given by $Z_{i\alpha} = \langle \hat{O}_{i\alpha} \rangle_s^2$.

IV. DIFFICULTIES OBTAINING SYMMETRY-BROKEN PHASES

In this section, we explain why the current implementation of mean-field theory fails to obtain proper symmetry-broken

phases. We use the example of the well-understood one-dimensional (1D) Hubbard model at half-filling. Consider the Hamiltonian,

$$\hat{H} = \frac{U}{2} \sum_i (\hat{N}_i^2 - \hat{N}_i) - \sum_{i,\sigma} t (\hat{c}_{i,\sigma}^\dagger \hat{c}_{i+1,\sigma} + \hat{c}_{i+1,\sigma}^\dagger \hat{c}_{i,\sigma}), \quad (10)$$

where i is the site index, there is a single orbital per site, there are two spin channels per site, and we consider the case where we are at half-filling ($\langle \hat{N}_i \rangle = 1$). The ground state is well known. For $U = 0$, the ground state is nonmagnetic and metallic. For $U > 0$ but finite, the ground state is insulating and shows AFM correlations [34] but has finite quasiparticle weight $Z > 0$.

For $U = 0$ and $U \gg |t|$, the model's solutions are well described by existing slave-particle mean-field implementations. For the intermediate region $U \sim |t|$, we are aware of no published study using recent slave-spin, slave-rotor, or other formalisms from the same family that has obtained the correct AFM phase for this model. Namely, the AFM solution does not appear to be a self-consistent ground-state solution of the spinon + slave coupled Hamiltonians. In addition to being annoying, this is very worrisome since even a simple uncorrelated approach such as Hartree-Fock easily delivers an AFM ground state.

To understand where the problem lies, consider the spinon Hamiltonian of Eq. (7) and how one would achieve symmetry breaking, e.g., spin symmetry breaking and ordering, due to electron interaction effects. Since the electron interaction is handled by the slave sector, the only quantities that can be affected by the slave calculation that then feed into the spinon Hamiltonian are the Lagrange multipliers $h_{i\alpha}$ and the rescaling factors $\langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i\alpha'} \rangle_s$ of the spinon hopping.

In the simplest slave treatment, we have a single slave particle at the site: for example, the slave-number and slave-rotor treatments. In this case, the α label is nil so our Lagrange multipliers are only indexed by site h_i , and the rescaling factors as well, $\langle \hat{O}_i^\dagger \hat{O}_i \rangle_s$. Obviously, no spin symmetry breaking is possible in the spinon sector since these variables do not depend on spin in any way.

When we move to more elaborate slave-particle models where there are different slave modes for the different spin channels, one can imagine that symmetry breaking is possible. For example, in our single orbital per site, 1D Hubbard model, when we have one slave particle for each spin channel, then $\alpha = \sigma$. We can now imagine that the $h_{i\sigma}$ shift the on-site energies of the orbitals in such a way as to break spin symmetry or that the hopping rescaling factors are also spin dependent. In practice, however, we have not found this to be the case: starting from a strongly symmetry-broken initial guess, the self-consistency cycle between spinon and slave sectors drives the system towards a paramagnetic solution and the two spin channels become equivalent. Any initial magnetization disappears upon self-consistent iteration.

We have analyzed this failure and discovered the following situation. If at some point the spinon system has broken spin symmetry at a site i with net spin-up, then $h_{i\uparrow} > h_{i\downarrow}$ is what makes this true. However, although $h_{i\uparrow} > h_{i\downarrow}$ favors a higher spin \uparrow occupancy in the spinon sector [due to the negative




FIG. 1. $\Delta n = n_\uparrow - n_\downarrow$ as a function of $\Delta h = h_\uparrow - h_\downarrow$ at one site of the 1D, half-filled, single-band Hubbard model with $U = 2$ and $t = 1$. Top: FM phase. Bottom: AFM phase. The Δh dependences of the spinon and slave occupancies are shown separately. Self-consistency between the two requires zero occupancy difference.

sign in front of $h_{i\alpha}$ in Eq. (7)), it favors a higher occupancy of the spin \downarrow channel in the slave sector [positive sign of $h_{i\alpha}$ in Eq. (8)]. The two effects fight each other, and the final self-consistent solution has $h_{i\uparrow} = h_{i\downarrow}$. An explicit example is provided by the 1D single-band Hubbard model at half-filling, for which the dependences of slave and spinon occupancies on h are shown in Fig. 1. These plots are generated by providing $\Delta n_i = n_{i\uparrow} - n_{i\downarrow}$ at some fixed site i as input to the slave problem, which yields $\Delta h_i = h_{i\uparrow} - h_{i\downarrow}$ and $\langle \hat{O}_{i\sigma} \rangle$, which are then used to solve the spinon problem to get the spinon Δn_i . Figure 1 clearly shows that the only self-consistent solution where the slave and spinon particle numbers match is for $\Delta h_i = 0$, which is the symmetric paramagnetic state.

V. SYMMETRY-BREAKING FIELDS

In this section, we show how manually adding small external symmetry-breaking terms (“fields”) to the on-site energies can lead to electronic symmetry breaking and lower the energy of the self-consistent ground state. In the next section, we justify this apparently *ad hoc* approach.

Adding additional symmetry-breaking (“magnetic-field”) terms $b_{im\sigma}$ to the on-site energies of the orbitals in the spinon

Hamiltonian gives the simple modification

$$\begin{aligned} \hat{H}_f = & \sum_{im\sigma} \epsilon_{im\sigma} \hat{f}_{im\sigma}^\dagger \hat{f}_{im\sigma} - \sum_{i\alpha} h_{i\alpha} \sum_{m\sigma \in \alpha} \hat{f}_{im\sigma}^\dagger \hat{f}_{im\sigma} \\ & - \sum_{ii'\alpha\alpha'} \langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'} \rangle_s \sum_{\substack{m\sigma \in \alpha \\ m'\sigma' \in \alpha'}} t_{imim'\sigma} \hat{f}_{im\sigma}^\dagger \hat{f}_{i'm'\sigma} \\ & - \sum_{im\sigma} b_{im\sigma} \hat{f}_{im\sigma}^\dagger \hat{f}_{im\sigma}. \end{aligned} \quad (11)$$

We do not modify the slave Hamiltonian in any way in this *ad hoc* approach.

We note that a symmetry-breaking field indexed by $im\sigma$ can break symmetry between sites, between orbitals, and between spin directions (and any combination thereof). It is critical to note that the nature and type of symmetry-breaking fields predetermine the types of solutions one can describe: to generate more complex types of symmetry breaking (e.g., superconducting order with slave bosons [24]), one will first have to generalize the formalism to allow for the appropriate symmetry-breaking fields.

The addition of nonzero symmetry-breaking fields $b_{im\sigma}$ will modify the self-consistent solution to the spinon + slave problem. To gauge whether this improves the solution, we monitor the total electronic energy and see if it is lowered due to symmetry breaking. The total energy is the expectation value of the original Hubbard Hamiltonian, Eq. (1), with respect to the approximate spinon + slave wave function $|\Psi_f\rangle|\Phi_s\rangle$ and is equal to

$$\begin{aligned} E_{\text{total}} = \langle \hat{H} \rangle = & \sum_i \langle \hat{H}_{\text{int}}^i \rangle_s + \sum_{im\sigma} \epsilon_{im\sigma} \langle \hat{f}_{im\sigma}^\dagger \hat{f}_{im\sigma} \rangle_f \\ & - \sum_{ii'mm'\sigma} t_{imim'\sigma} \langle \hat{f}_{im\sigma}^\dagger \hat{f}_{i'm'\sigma} \rangle_f \langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'} \rangle_s. \end{aligned} \quad (12)$$

(Please note that in order for the expectation value of \hat{H} to give the associated energy, the state must be normalized, which is most convenient to achieve by setting $\langle \Psi_f | \Psi_f \rangle = \langle \Phi_s | \Phi_s \rangle = 1$.)

We now apply this approach to the 1D single-band Hubbard model at half-filling, Eq. (10). Without loss of generality, we choose $b_{i\uparrow} = -b_{i\downarrow}$ to break the spin symmetry at each site i . For ferromagnetic (FM) order, we choose aligned symmetry-breaking fields between neighboring sites $b_{i+1,\sigma} = b_{i\sigma}$, while AFM order requires staggered fields $b_{i+1,\sigma} = -b_{i\sigma}$. Hence, the field strength b for spin-up at one site is sufficient to specify the fields at all sites. We numerically solve the spinon + slave self-consistent equations using the single-site mean-field approximation described in Sec. III. In essence, we search broken-symmetry solutions parameterized by b to find the lowest energy state. As we have prespecified FM (or AFM) order with one (or two) site(s) per unit cell, we only find such solutions; more complex magnetic orderings will require more complex unit cells and symmetry-breaking fields with more degrees of freedom.

We begin our analysis with the most coarse-grained slave-boson representations, which only describe the total electron count at each site (i.e., give no information on the spin configuration). These are the slave-rotor and number-slave methods. The chief difference between them is that the




FIG. 2. Total energy per site and quasiparticle weight Z (renormalization factor) versus symmetry-breaking perturbation field strength b based on the slave-rotor method for the half-filled, single-band 1D Hubbard model with $U = 2$ and $t = 1$.

number count at a site can be any integer in the slave-rotor method, while the number-slave method corrects this by only permitting the electron count to be among the physically allowed values (e.g., 0, 1, or 2 for the single-band Hubbard model). Figure 2 shows the dependence of the total energy and quasiparticle weight Z (i.e., renormalization factor) on the field strength b within the slave-rotor approach. For the slave-rotor method, increasing b increases the total energy of both AFM and FM solutions: the nonmagnetic solution is the preferred ground state. The strength of electronic correlations, measured by how much Z deviates from its noninteracting value of unity, also increases with b . This b dependence is opposite to what one would expect for the actual model system: a more spin-polarized system should have smaller number fluctuations, as occupancies are driven towards 1 or 0 and the electron configuration becomes better described by a single Slater determinant. Finally, the slave rotor predicts an abrupt transition to a Mott insulator at finite b , which is peculiar (and wrong).

The number-slave results for total energy and Z versus b , displayed in Fig. 3, are somewhat improved over the slave-rotor results but are still fundamentally flawed. The energy is still minimized by the nonmagnetic solution at $b = 0$ (although the energy rises more gently with b) and Z drops with b (albeit more modestly). The failure of the slave-rotor and number-slave methods is tied to the fact that they do not consider the spin degree of freedom.




FIG. 3. Total energy per site and Z versus field strength b for the number-slave method for the single-band 1D Hubbard model at half-filling with $U = 2$ and $t = 1$.

Due to the simplicity of the single-band Hubbard model, the only remaining slave model is the spin + orbital-slave approach (called “spin-slave” in the literature [28,29,35]). At each site, each spin channel has its own dedicated slave particle. The energy versus b plot in Fig. 4 shows that we obtain an AFM ground state since a minimum appears at finite b . The figure also shows that the degree of electronic correlation is reduced with increasing b (and increasing strength of AFM order) as the occupancies get closer to 0 and 1: the system becomes less strongly interacting as b is strengthened. This is what we expect: with increasing AFM polarization, the electronic configuration of the system is driven to extremes of occupation (0 or 1 for each spin channel), meaning that one can describe the system more accurately with a single (noninteracting) Slater determinant. More details on the energetic behavior versus b are provided in Fig. 5, where the individual components of the total energy are shown versus b . The interaction energy (Hubbard U term) is reduced by the spin symmetry breaking since for both FM and AFM order the occupancies move away from half-filling, where occupancy fluctuation is largest. The band (hopping or kinetic) energy rises with b due to the splitting of bands upon symmetry reduction. Both behaviors are generic and as expected. However, the reason the AFM order shows a minimum total energy versus b is the fact that Z becomes larger with b in this case: a larger Z (i.e., larger $\langle O \rangle$) will enhance hopping, widen the bands, and thus offset the reduction in the total band energy due to the creation of spin polarization.




FIG. 4. Total energy per site and Z versus field b for the spin + orbital-slave approach for the single-band 1D Hubbard model at half-filling with $U = 2$ and $t = 1$. Unlike with the number-slave and slave-rotor approaches, correlations decrease with increasing b for the AFM phase and slowly increase with b for the FM phase.

The take-home message of this section is that the introduction of symmetry-breaking fields can succeed in stabilizing symmetry-broken ground states due to electronic correlations as long as the slave approach being used is able to describe the symmetry-breaking degree of freedom (spin in the 1D single-band Hubbard model). We are thus motivated to improve upon the *ad hoc* nature of the approach and put it on a firmer theoretical basis in the next section.

VI. SELF-CONSISTENT TOTAL ENERGY APPROACH

In this section, we justify the successful but *ad hoc* approach in the previous section. Namely, we describe a total energy functional that can be applied to any type of slave-particle problem and permits easy incorporation of the various types of desired constraints. Specifically, we show that the slave-particle approach is a variational approach to the interacting ground-state problem, and we provide an explicit form for the variational energy functional. We also show that this viewpoint provides significant practical benefits for efficient solution of the self-consistency problem between slave and spinon sectors.

The form of the energy functional F is given by

$$F = E_{\text{total}} + \text{constraints},$$

where E_{total} is from Eq. (12) and the constraint terms are enforced by Lagrange multipliers. Prior to the introduction of symmetry-breaking fields, the constraints we have enforced are




FIG. 5. Individual components of the total energy are shown versus b for the spin + orbital-slave approach for the single-band 1D Hubbard model at half-filling with $U = 2$ and $t = 1$.

that $\langle N_{i\alpha} \rangle_s = \langle \hat{n}_{i\alpha} \rangle_f$ as well as the normalization of the spinon and slave wave functions $\langle \Psi_f | \Psi_f \rangle = \langle \Phi_s | \Phi_s \rangle = 1$. To incorporate symmetry-breaking fields, we choose to parametrize the functional F by *target* spinon occupancies $v_{i\sigma}$: these numbers are the occupancies that we are constraining the spinons to obey, i.e., the constraints are $\langle n_{i\sigma} \rangle_f = v_{i\sigma}$. The associated Lagrange multipliers are $b_{i\sigma}$. Hence the energy functional has the form, where we write out E_{total} explicitly,

$$\begin{aligned}
 F(\{v_{i\sigma}\}) = & \sum_i \langle \hat{H}_{\text{int}}^i \rangle_s + \sum_{i\sigma} \epsilon_{i\sigma} \langle \hat{f}_{i\sigma}^\dagger \hat{f}_{i\sigma} \rangle_f \\
 & - \sum_{ii'mm'\sigma} t_{imim'\sigma} \langle \hat{f}_{i\sigma}^\dagger \hat{f}_{i'm'\sigma} \rangle_f \langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha} \rangle_s \\
 & - \lambda_f [\langle \Psi_f | \Psi_f \rangle - 1] \\
 & - \lambda_s [\langle \Phi_s | \Phi_s \rangle - 1] - \sum_{i\alpha} h_{i\alpha} [\langle \hat{n}_{i\alpha} \rangle_f - \langle \hat{N}_{i\alpha} \rangle_s] \\
 & - \sum_{i\sigma} b_{i\sigma} [\langle \hat{n}_{i\sigma} \rangle_f - v_{i\sigma}]. \quad (13)
 \end{aligned}$$

The Lagrange multipliers λ_f and λ_s enforce normalization of the spinon and slave wave functions, respectively (these are necessary so that the expectation values of the various Hamiltonian terms correspond to average energies). The $h_{i\alpha}$ enforce particle-number matching between slave and spinon sectors. The $b_{i\sigma}$ enforce spinon-particle matching to target values. As expected, when the constraints are obeyed, $F = E_{\text{total}}$.

The point of having an energy functional is that the minimizing variational conditions, which generate desired eigenvalue problems, are easily derived by differentiation. In addition, the value of F provides a variational estimate of the ground-state energy. Setting the derivative versus $\langle \Psi_f |$ to 0 gives the spinon eigenvalue equation

$$0 = \frac{\delta F}{\delta \langle \Psi_f |} = H_f | \Psi_f \rangle - \lambda_f | \Psi_f \rangle,$$

where the spinon Hamiltonian is Eq. (11), which includes the symmetry-breaking fields. Similarly, the minimum condition for $|\Phi_s\rangle$ gives a slave eigenvalue problem with the slave Hamiltonian, Eq. (8).

The above formalism shows that, once all the constraints are obeyed, $F(\{v_{i\sigma}\}) = E_{\text{total}}(\{v_{i\sigma}\})$. The remaining task is to search the target occupancies $v_{i\sigma}$ to find the minimum total energy. While theoretically straightforward, in practice such an approach is difficult and inefficient because for each specified $\{v_{i\sigma}\}$, one must find the fields $b_{i\sigma}$ that enforce those particular target occupancies: this requires solving the spinon + slave problem a great many times.

Practically, it is better to use the $b_{i\sigma}$ as the independent variables and to minimize the energy over them (formally, this corresponds to a Legendre transformation of F). Hence, we now view $v_{i\sigma}$ as whatever mean spinon occupancies are generated by the solution of the spinon + slave problem at fixed $\{b_{i\sigma}\}$ which makes that corresponding constraint form always vanish. Hence, in what follows, we use the symmetry-breaking fields as independent variables and consider the total energy functional $F(\{b_{i\sigma}\})$. Since we always obey the key constraints for a physical solution, $F(\{b_{i\sigma}\}) = E_{\text{total}}(\{b_{i\sigma}\})$ will be true. Hence, minimization of the total energy versus $\{b_{i\sigma}\}$ will coincide with minimization of F .

In summary, we have rewritten the slave-boson problem as a constrained variational minimization problem. When the constraints are obeyed, the minimization corresponds directly to minimizing the total energy expression, Eq. (12).

VII. SIMPLIFIED AND MORE EFFICIENT SLAVE-PARTICLE APPROACH

Up to this point, the slave-particle approaches we have developed require self-consistency between spinon and slave sectors in a specific manner: not only do the spinon expectations renormalize slave hopping terms (and conversely for slave expectations and spinon hoppings), but a shared set of Lagrange multipliers $h_{i\alpha}$ enforces particle-number matching $\langle \hat{n}_{i\alpha} \rangle_f = \langle \hat{N}_{i\alpha} \rangle_s$. The process of finding the $h_{i\alpha}$ is numerically challenging: the $h_{i\alpha}$ appear with opposite signs in the spinon H_f and slave H_s Hamiltonians, meaning that increasing $h_{i\alpha}$ decreases $\langle \hat{n}_{i\alpha} \rangle_f$ but increases $\langle \hat{N}_{i\alpha} \rangle_s$. Our general observation is that this “fighting” over $h_{i\alpha}$ between the slave and the spinon sectors leads to a time-consuming self-consistent process requiring many iterations to reach convergence.

Accelerating this process requires a simple change of variables that is motivated by three related observations: (i) in the total energy functional of Eq. (13), the spinon and slave number constraints are not treated symmetrically because the spinons have the added $b_{i\sigma}$ terms; (ii) in the spinon Hamiltonian, Eq. (11), we can add the $h_{i\alpha}$ and $b_{i\sigma}$ terms

together in a single term, whereas the slave Hamiltonian, Eq. (8), only has the $h_{i\alpha}$ terms; and (iii) in the end, these Lagrange multipliers $h_{i\alpha}$ and $b_{i\sigma}$ do not appear in the total energy so rearranging them in various ways does not change the total energy.

For the spinon Hamiltonian, we consider instead the new symmetry-breaking field given by the sum $B_{i\sigma} = h_{i\alpha} + b_{i\sigma}$. The spinon Hamiltonian is now

$$\begin{aligned} \hat{H}_f = & \sum_{i\sigma} \epsilon_{i\sigma} \hat{f}_{i\sigma}^\dagger \hat{f}_{i\sigma} - \sum_{i\sigma} B_{i\sigma} \hat{f}_{i\sigma}^\dagger \hat{f}_{i\sigma} \\ & - \sum_{ii'\alpha\alpha'} \langle \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'} \rangle_s \sum_{\substack{m\sigma \in \alpha \\ m'\sigma' \in \alpha'}} t_{imi'm'\sigma} \hat{f}_{i\sigma}^\dagger \hat{f}_{i'm'\sigma'}, \end{aligned} \quad (14)$$

while the slave Hamiltonian is unchanged:

$$\begin{aligned} \hat{H}_s = & \sum_i \hat{H}_{\text{int}}^i + \sum_\alpha h_{i\alpha} \hat{N}_{i\alpha} \\ & - \sum_{ii'\alpha\alpha'} \left[\sum_{\substack{m\sigma \in \alpha \\ m'\sigma' \in \alpha'}} t_{imi'm'\sigma} \langle \hat{f}_{i\sigma}^\dagger \hat{f}_{i'm'\sigma'} \rangle_f \right] \hat{O}_{i\alpha}^\dagger \hat{O}_{i'\alpha'}. \end{aligned}$$

The slave Hamiltonian H_s no longer shares a common Lagrange multiplier with the spinon Hamiltonian H_f .

Operationally, this means that when we solve the slave Hamiltonian problem, we are given specified $\langle \hat{n}_{i\alpha} \rangle_f$ as input, and we solve the slave problem while adjusting the $h_{i\alpha}$ so as to ensure that the slave-particle counts match the input: $\langle \hat{N}_{i\alpha} \rangle_s = \langle \hat{n}_{i\alpha} \rangle_f$. However, when solving the spinon problem in the presence of symmetry-breaking fields $B_{i\sigma}$, there is no need to do particle-number matching: the Lagrange multiplier $B_{i\sigma}$ simply make the spinon-particle counts match some free-floating values. In this way, particle-number matching between the slave and the spinon sectors is decoupled, which *greatly* simplifies the self-consistency process. Put another way, the symmetry-breaking fields $B_{i\sigma}$ specify a set of desired spinon-particle counts $\{v_{i\sigma}\}$, and the slave sector is required to match these particle numbers via the $h_{i\alpha}$ Lagrange multipliers.

We find that this simplified approach, which is equivalent to the standard approach of having $h_{i\alpha}$ appear in both Hamiltonians, is much more efficient in numerical calculations, as it greatly speeds up self-consistency. In this new approach, one achieves rapid self-consistency for a given set of $\{B_{i\sigma}\}$ which specify the spinon Hamiltonian and the target spinon occupancies $v_{i\sigma}$. One can then minimize $E_{\text{total}}(\{B_{i\sigma}\})$ over the $B_{i\sigma}$ to find the symmetry-broken ground state. In our experience, this new approach requires ~ 5 – 10 times fewer self-consistent steps to reach the same level convergence.

Using this method, we can rapidly scan over B in a stable, self-consistent way to obtain the ground-state energies. Figure 6 shows the dependence of the ground-state energy of the half-filled, single-band, 1D Hubbard model as a function of U/t : for each U/t , we easily scan over the new symmetry-breaking field strength B to find the AFM ground-state energy. The figure shows energy versus U/t for the AFM state as well as the $B = 0$ nonmagnetic solution compared to the exact Bethe ansatz solution for this problem [34]. Overall, the comparison between the AFM slave-spin solution (which




FIG. 6. Comparison of the total ground-state energies (in units of t) for the single-band 1D Hubbard model at half-filling based on the AFM Hartree-Fock solution, the PM slave-spin solution, the symmetry-broken (AFM) slave-spin ground-state solution, and the exact Bethe ansatz (AFM) solution as calculated using the method in Ref. [36].

is insulating in the spinon sector) and the exact Bethe ansatz is satisfactory given the simplicity of the single-site mean-field slave model used here. As expected, the AFM slave-spin method becomes very much like the AFM Hartree-Fock in the large- U/t limit of very strong spin polarization since both approaches essentially describe the system as a single Slater determinant. We note that the nonmagnetic ground state has an incorrect evolution from a metallic system at small U/t to a Mott-insulating phase at $U/t \geq 10$.

We provide a word of caution regarding the interpretation of the results. The results show that in this particular situation, the slave-boson approach provides an energy that is close to but below that of Hartree-Fock and above the true ground-state energy. However, to the best of our knowledge, there is no known principle guaranteeing that these slave-boson methods give an upper bound to the total energy (e.g., in the same way that the Hartree-Fock does). E_{total} in Eq. (12) does provide a variational upper bound of the energy, *but* in the extended spinon + slave Hilbert space. The state $|\Psi_f\rangle|\Phi_s\rangle$ is a valid and normalized state in the extended Hilbert space so that E_{total} , being the expectation of \hat{H} for this state, must be higher than the ground-state energy in the extended Hilbert space. However, the extended Hilbert space includes both physical and unphysical states, and there is no known guarantee that the ground-state energy in the extended space coincides with the ground state in the subspace of physical states. Nevertheless, we note that many useful electronic structure methods for solid-state systems (e.g., DFT with an approximate exchange-correlation functional) do not obey strict bounds on the total energy and yet can often outperform the Hartree-Fock (which does come with a bound). Hence, whether total energies from slave-boson methods lie above or below the true ground-state energy in more complex physical systems is an interesting and open question.

VIII. CONCLUSION

We have shown how slave-particle methods can be used to obtain spontaneously symmetry-broken electronic phases based on a total energy approach. We have described and

tested our ideas on the classic 1D Hubbard model Hamiltonian. Furthermore, we have shown how to enable symmetry breaking via the use of auxiliary symmetry-breaking fields in a self-consistent way that greatly lowers the computational burden and stability of the standard slave-particle calculation. Further, we have demonstrated that in order to obtain spontaneously symmetry-broken phases in the spinon sector, the slave sector must be allowed to break the corresponding symmetry explicitly by having different slave modes for the different degrees of freedom which may undergo symmetry breaking.

ACKNOWLEDGMENT

This work was supported by the National Science Foundation (USA) via Grant No. MRSEC DMR 1119826.

APPENDIX

We show, via a detailed derivation, that the slave-boson representation is faithful by showing that canonical commutation relations are obeyed. By definition, the commutators of the physical electron operators $\hat{d}_{im\sigma}$ satisfy fermionic anticommutation relations:

$$\{\hat{d}_{im\sigma}, \hat{d}_{jm'\sigma'}^\dagger\} = \delta_{ij}\delta_{mm'}\delta_{\sigma\sigma'}. \quad (\text{A1})$$

We show that this equality holds when we perform the slave-boson substitution $\hat{d}_{im\sigma} = \hat{f}_{im\sigma} \hat{O}_{i\alpha}$ if we deal only with physical states (i.e., states in the extended spinon + slave Hilbert space where the number of spinons and bosons match exactly). Namely, for a pair of physical states $|p_1\rangle$ and $|p_2\rangle$,

$$\langle p_1 | \{\hat{f}_{im\sigma} \hat{O}_{i\alpha}, \hat{f}_{jm'\sigma'}^\dagger \hat{O}_{j\beta}^\dagger\} | p_2 \rangle = \delta_{ij}\delta_{mm'}\delta_{\sigma\sigma'} \langle p_1 | p_2 \rangle. \quad (\text{A2})$$

To prove the above equality, we begin by remembering that α and β label disjoint sets of localized states each specified by the combination of labels $im\sigma$ describing site i , spatial orbital m , and σ . In addition, we have that $im\sigma \in \alpha$ and $jm'\sigma' \in \beta$. Also, by definition, the spinon operators \hat{f} are fermionic field operators obeying canonical commutation relations. As discussed in the text, the \hat{O} operators are bosonic but are not canonical field operators. The matrix representation of the lowering \hat{O} operator in the number basis is [32]

$$\hat{O}_{i\alpha} = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \\ C_{i\alpha} & 0 & 0 & \dots & 0 & 0 \end{pmatrix}. \quad (\text{A3})$$

The number of slave bosons $N_{i\alpha}$ is an integer obeying $0 \leq N_{i\alpha} \leq M_{i\alpha}$, where $M_{i\alpha}$ is the maximum slave boson count (i.e., maximum number of electrons) that can be placed into the set of orbitals α . For use below, we also have the matrix representations

$$\hat{O}_{i\alpha}^\dagger \hat{O}_{i\alpha} = \begin{pmatrix} |C_{i\alpha}|^2 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix} \quad (\text{A4})$$

and

$$[\hat{O}_{i\alpha}, \hat{O}_{i\alpha}^\dagger] = (1 - |C_{i\alpha}|^2) \cdot \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & -1 \end{pmatrix}. \quad (\text{A5})$$

To prove Eq. (A2), we use the anticommutation properties of the \hat{f} to arrive at

$$\{\hat{f}_{im\sigma} \hat{O}_{i\alpha}, \hat{f}_{jm'\sigma'}^\dagger \hat{O}_{j\beta}^\dagger\} = \hat{f}_{im\sigma} \hat{f}_{jm'\sigma'}^\dagger [\hat{O}_{i\alpha}, \hat{O}_{j\beta}^\dagger] + \hat{O}_{j\beta}^\dagger \hat{O}_{i\alpha} \delta_{ij} \delta_{mm'} \delta_{\sigma\sigma'}. \quad (\text{A6})$$

We examine the four cases of index combinations that can occur and show that in each case Eq. (A2) holds.

(a) When $i \neq j$, the bosonic operators at different sites commute by definition so the first term on the right-hand side of Eq. (A6) is 0. And the second term is 0 since $i \neq j$, so Eq. (A2) holds.

(b) When $i = j$ but $\alpha \neq \beta$, the bosonic operators at the same site refer to a disjoint set of states at the site and commute by definition. So the first term on the right-hand side of Eq. (A6) is 0. And the second term is also 0 since α and β are disjoint and do not share any states at the same site. So Eq. (A2) holds.

(c) When $i = j$ and $\alpha = \beta$ but $(m\sigma) \neq (m'\sigma')$, only the first term on the right-hand side of Eq. (A6) survives and equals

$$\{\hat{f}_{im\sigma} \hat{O}_{i\alpha}, \hat{f}_{im'\sigma'}^\dagger \hat{O}_{i\alpha}^\dagger\} = \hat{f}_{im\sigma} \hat{f}_{im'\sigma'}^\dagger \cdot (1 - |C_{i\alpha}|^2) \cdot \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & -1 \end{pmatrix}. \quad (\text{A7})$$

The action of the right-hand side on any physical state which has between 1 and $M_{i\alpha} - 1$ particles is 0. When acting on a state with 0 particles, the action of $\hat{f}_{im\sigma}$ will kill the state and yield 0. When acting on a state with the maximum number $M_{i\alpha}$ of particles, the action of $\hat{f}_{im'\sigma'}^\dagger$ gives 0. So Eq. (A2) holds.

(d) Finally, when all indices match ($i = j, \alpha = \beta, m = m', \sigma = \sigma'$), we have

$$\{\hat{f}_{im\sigma} \hat{O}_{i\alpha}, \hat{f}_{im\sigma}^\dagger \hat{O}_{i\alpha}^\dagger\} = \hat{f}_{im\sigma} \hat{f}_{im\sigma}^\dagger \cdot (1 - |C_{i\alpha}|^2) \cdot \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & -1 \end{pmatrix}$$

$$+ \begin{pmatrix} |C_{i\alpha}|^2 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}. \quad (\text{A8})$$

If the state $im\sigma$ is empty, the action of $\hat{f}_{im\sigma}\hat{f}_{im\sigma}^\dagger$ is the identity operation on this state, and adding the matrices we find the identity matrix except for the bottom right corner element, which is $|C_{i\alpha}|^2$ instead of 1; but this is irrelevant since if $im\sigma$ is empty, then the number of particles is less than the maximum $M_{i\alpha}$ so this matrix element is never accessed. If the

state $im\sigma$ is full, $\hat{f}_{im\sigma}\hat{f}_{im\sigma}^\dagger$ kills the state and we are again left with the identity matrix except for the top left corner element being $|C_{i\alpha}|^2$; this is irrelevant since if $im\sigma$ is full, the number of particles is greater than 0 so this element is never accessed. So Eq. (A2) holds.

-
- [1] J. Hubbard, *Proc. R. Soc. A* **276**, 238 (1963).
 [2] P. Hohenberg and W. Kohn, *Phys. Rev.* **136**, B864 (1964).
 [3] W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
 [4] A. D. Becke, *J. Chem. Phys.* **98**, 1372 (1993).
 [5] J. P. Perdew, M. Ernzerhof, and K. Burke, *J. Chem. Phys.* **105**, 9982 (1996).
 [6] L. Hedin, *Phys. Rev.* **139**, A796 (1965).
 [7] M. S. Hybertsen and S. G. Louie, *Phys. Rev. B* **34**, 5390 (1986).
 [8] F. Aryasetiawan and O. Gunnarsson, *Rep. Prog. Phys.* **61**, 237 (1998).
 [9] M. van Schilfgaarde, T. Kotani, and S. Faleev, *Phys. Rev. Lett.* **96**, 226402 (2006).
 [10] A. Georges, G. Kotliar, W. Krauth, and M. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (1996).
 [11] G. Kotliar, S. Y. Savrasov, K. Haule, V. S. Oudovenko, O. Parcollet, and C. A. Marianetti, *Rev. Mod. Phys.* **78**, 865 (2006).
 [12] S. E. Barnes, *J. Phys. F* **6**, 1375 (1976).
 [13] S. Barnes, *J. Phys. F* **7**, 2637 (1977).
 [14] P. Coleman, *Phys. Rev. B* **29**, 3035 (1984).
 [15] N. Read and D. M. Newns, *J. Phys. C* **16**, L1055 (1983).
 [16] N. Read, *J. Phys. C* **18**, 2651 (1985).
 [17] G. Kotliar and A. E. Ruckenstein, *Phys. Rev. Lett.* **57**, 1362 (1986).
 [18] P. A. Lee, N. Nagaosa, and X. G. Wen, *Rev. Mod. Phys.* **78**, 17 (2006).
 [19] M. Raczkowski, R. Fresard, and A. M. Oles, *Europhys. Lett.* **76**, 128 (2006).
 [20] R. Fresard and G. Kotliar, *Phys. Rev. B* **56**, 12909 (1997).
 [21] F. Lechermann, A. Georges, G. Kotliar, and O. Parcollet, *Phys. Rev. B* **76**, 155102 (2007).
 [22] J. Bünenmann, *Phys. Status Solidi A* **248**, 203 (2011).
 [23] R. Frésard, M. Dzierzawa, and P. Wölfle, *EPL (Europhys. Lett.)* **15**, 325 (1991).
 [24] A. Isidori and M. Capone, *Phys. Rev. B* **80**, 115120 (2009).
 [25] S. Florens and A. Georges, *Phys. Rev. B* **66**, 165111 (2002).
 [26] S. Florens and A. Georges, *Phys. Rev. B* **70**, 035114 (2004).
 [27] B. Lau and A. J. Millis, *Phys. Rev. Lett.* **110**, 126404 (2013).
 [28] L. de’Medici, A. Georges, and S. Biermann, *Phys. Rev. B* **72**, 205124 (2005).
 [29] S. R. Hassan and L. de Medici, *Phys. Rev. B* **81**, 035106 (2010).
 [30] Y. Komijani and G. Kotliar, *Phys. Rev. B* **96**, 125111 (2017).
 [31] L. de Medici, G. Giovannetti, and M. Capone, *Phys. Rev. Lett.* **112**, 177001 (2014).
 [32] A. B. Georgescu and S. Ismail-Beigi, *Phys. Rev. B* **92**, 235117 (2015).
 [33] J. Kanamori, *Prog. Theor. Phys.* **30**, 275 (1963).
 [34] E. H. Lieb and F. Y. Wu, *Exactly Solved Models: A Journey in Statistical Mechanics. Selected Papers with Commentaries (1963–2008)*, Vol. 321 (World Scientific, Singapore, 2009), p. 535.
 [35] L. De’Medici, *Phys. Rev. B* **83**, 205112 (2011).
 [36] M. M. Sánchez, A. Avella, and F. Mancini, *Europhys. Lett. (EPL)* **44**, 328 (1998).