Multichannel charge Kondo effect and non-Fermi-liquid fixed points in conventional and topological superconductor islands

Michał Papaj, Zheng Zhu, and Liang Fu

Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

(Received 23 July 2018; revised manuscript received 13 December 2018; published 14 January 2019)

We study multiterminal Majorana and conventional superconducting islands in the vicinity of the charge degeneracy point using bosonization and the numerical renormalization group. Both models map to the multichannel charge Kondo problem, but for noninteracting normal leads they flow to different non-Fermi-liquid fixed points at low temperatures. We compare and contrast both cases by numerically obtaining the full crossover to the low-temperature regime, and we predict distinctive transport signatures. The differences between the two types of islands result from a crucial distinction between charge-2*e* and charge-*e* transfer in the conventional and topological case, respectively. In the conventional case, our results establish *s*-wave islands as a platform to study the intermediate multichannel Kondo fixed point. In the topological setup, the crossover temperature to non-Fermi-liquid behavior is relatively high as it is proportional to the level broadening, and the transport results are not sensitive to channel coupling anisotropy, moving away from the charge degeneracy point or including a small Majorana hybridization, which makes our proposal experimentally feasible.

DOI: 10.1103/PhysRevB.99.014512

I. INTRODUCTION

The prospect of robust quantum computation using Majorana zero modes [1-5] sparked enormous experimental interest in the material platforms that enable direct observation and study of these topological quasiparticles. Among the leading platforms are proximitized semiconductor nanowires with spin-orbit coupling, which are predicted to become topological superconductors and host Majorana zero modes under external magnetic fields [6,7]. The immense experimental effort over the past several years has resulted in a significant improvement in material and device fabrication quality, and also helped to rekindle interest in mesoscopic superconductor tivity in semiconductor devices [8–24].

When considering the physics of mesoscopic conductors and superconductors, it is crucial to take into account the Coulomb blockade effect that arises due to electron-electron interaction. Since the charging energy of the island depends quadratically on the number of electrons it contains, it is possible to use an external gate to tune two charge states of the island to be equal in energy. In conventional superconductors, where putting an odd number of electrons on the island requires an extra energy cost of the superconducting gap, the ground state consists of an even number of electrons. This effect has been directly observed as an even-odd asymmetry in aluminum islands [25-28]. On the other hand, in topological superconductors, zero-energy Majorana bound states exist that can accommodate an unpaired electron without any additional energy. In this case, the degeneracy can occur between states with an even and odd number of electrons. This even-odd degeneracy underlies the phenomenon of electron teleportation [29], which involves phase-coherent transport of a single electron via the spatially separated Majorana modes. A recent experiment [13] on a proximitized InAs island connected to two normal leads via tunnel junctions

observed a transition from resonant Cooper pair transport to single-electron transport above a critical magnetic field, which is broadly consistent with the scenario of transition from a conventional to a topological superconducting island [30,31].

The degeneracy between the two charge states of the superconducting islands—2N and 2N + 2(2N + 1) in the conventional (topological) case-can be a source for Kondotype phenomena. These degenerate levels can be represented as a pseudospin-1/2 object, which enables observation of phenomena related to the multichannel Kondo effect [32,33]. In the topological case, when the superconductor is tuned into charge degeneracy, it has been shown [34,35] that the system exhibits quantized dc conductance $G_{ijdc} = \frac{2e^2}{Nh}$ in the T = 0 limit for N Majorana modes coupled to N normal leads by mapping the model onto quantum Brownian motion (QBM) on a honeycomb lattice [36,37]. Moreover, it has recently been shown that if the topological superconductor is time-reversalinvariant, the two-terminal island will realize the two-channel Kondo effect without fine-tuning [38]. In the s-wave case, the setup based on a two-terminal island at charge degeneracy has recently been shown [39] to map to the two-channel Kondo problem. These parallel developments not only enable a direct comparison of the properties of both a conventional and a topological setup, but also provide an attractive platform for studies of quantum criticality.

While initially the Kondo effect has been considered in the context of dilute magnetic impurities, the interest in this phenomenon was revived after theoretical proposals for its realization in normal state quantum dots [40-42] and subsequent experimental confirmation of the predictions [43,44]. The next step was then the extension to the elusive multichannel Kondo effect, which was again guided by theory [45-49]and culminated with detailed studies of various properties of this setup [50-54]. These theoretical and experimental results established highly tunable normal-state nanostructures as a perfect window into the world of strongly correlated electron systems, and so they are intensively studied in order to extract the essence of the physical phenomena without the picture being blurred by the complexity of the real materials. Now we want to extend the realm of device platforms to superconducting nanoislands.

Motivated by the above results on Majorana and conventional superconducting islands, we expand on these studies by comparing and contrasting the charge Kondo effects due to even-odd and even-even degeneracies in both types of mesoscopic islands using bosonization and numerical renormalization-group (NRG) methods. We provide a mapping of the N-terminal conventional superconductor island model to the N-channel charge Kondo problem in the bosonization language, and then we examine the differences in the treatment of the Majorana island. For noninteracting leads in the topological model, the system flows to a strongcoupling fixed point, as opposed to the flow toward intermediate coupling in the conventional case (see Fig. 2). The non-Fermi-liquid fixed point of the Majorana island is robust to channel coupling asymmetry (in contrast to anisotropy being a relevant perturbation at the intermediate fixed point in the conventional system). These differences between both types of islands in transport properties are due to the crucial distinction of charge 2e transfer in the Andreev processes in the conventional case versus charge e transfer by single electron tunneling in the topological island [29]. In the topological case, while each tunneling process transfers a single electron charge e, due to the statistical transmutation [34] the system behaves as if a charge-e boson was transferred, which enables a nontrivial mapping to a Kondo model [34,35].

Using the numerical renormalization group, we first support our bosonization results at T = 0 by calculating the residual entropy and conductance matrix elements. For the conventional island, we confirm that the dc conductance in T = 0 approaches the predicted value of 2 and $\frac{8}{2} \sin^2(\frac{\pi}{5}) e^2/h$ for two and three terminals, respectively. In the Majorana setup for three terminals, we obtain the anticipated dc con-ductance of $\frac{2e^2}{Nh} = 2/3 e^2/h$, which is robust against the tunnel coupling anisotropy (even if all three couplings are different) and moving away from the charge degeneracy point. More importantly, we go beyond the zero-temperature limit and obtain the full crossover to non-Fermi-liquid fixed points in both cases. In the conventional setup, our results establish the s-wave island as a platform for studying the physics of the intermediate multichannel Kondo fixed point. For Majorana islands, we demonstrate that the transition at the charge degeneracy point happens at a much higher temperature than in the Coulomb valley regime of the topological Kondo effect studied previously [55-67]. Our results will facilitate the future experimental observation of quantized conductance. For the three-terminal case, we predict a nontrivial crossover between the regimes dominated by two and three leads with an intermediate dc conductance plateau at $2/3 e^2/h$, which emerges at sufficiently low temperature while tuning the tunnel coupling of the third lead. This, together with the aforementioned robustness to variation in setup parameters, provides an experimental signature that can be used to verify our claims for the Majorana island.



FIG. 1. Multiterminal (a) conventional and (b) Majorana islands at charge degeneracy (charging energy is controlled by a gate). Both islands are connected to N normal leads (blue) via either Andreev reflection or tunneling into Majorana zero modes.

II. MODELS

In this work, we consider two types of setups with multiterminal superconducting islands (Fig. 1). We begin by describing the full Hamiltonian of the systems analyzed in the following sections, which consists of three parts:

$$H = H_C + H_L + H_T. \tag{1}$$

The central point of both setups considered in this paper is a mesoscopic superconducting island, either of an ordinary *s*-wave or topological nature with a gap Δ that is the largest energy scale of the problem. In the *s*-wave case, there are no quasiparticle excitations inside of the superconducting gap, and so in the usual BCS formalism, introducing an odd number of electrons into the island requires an energy cost of Δ . On the other hand, the topological superconductor hosts an even number of zero-energy Majorana bound states, and this allows us to put an additional electron into the island without paying the extra energy. Since we are studying a mesoscopic superconducting grain that is not grounded, we also have to consider the charging effects that arise due to Coulomb interactions. The electrostatics can be taken into account by including into the Hamiltonian a term

$$H_C = E_C (\hat{N} - N_g)^2,$$
 (2)

where E_C is the charging energy related to the capacitance of the island, \hat{N} is the number of charges in the superconductor, and N_g is the potential determined by the external gate. This tunability gives rise to a possible degeneracy between the two charge states of the island. However, the number of charges in the degenerate states differs in both of the considered cases. For the conventional superconductor, since we are working in the regime where $E_C \ll \Delta$, we can consider the states with an odd number of electrons to be unfavorable energetically, and so when we set N_g to be an odd integer $2N_0 + 1$, according to (2), the states with $2N_0$ and $2N_0 + 2$ electrons will be degenerate and lowest in energy. The situation is different in the topological superconductor, where there is no additional energy cost for the states with an odd number of electrons. There we can set N_g to $2N_0 + \frac{1}{2}$, and then states with $2N_0$ and $2N_0 + 1$ are degenerate. At very low temperatures, we can then restrict our Hilbert space to just those pairs of charge states of the island. The subspace of charge states can then be described by a pseudospin-1/2 object, with s_z eigenstates corresponding to $2N_0/2N_0 + 1$ or $2N_0/2N_0 + 2$ states. Then

the slight deviation from the charge degeneracy point can be taken into account by introducing a Zeeman-like term δs_z into the Hamiltonian, where δ can be tuned by the external gate.

A common part of both setups is a set of N normal leads, which are tunnel-coupled to the superconductor. In the conventional superconductor setup, they are described by the Hamiltonian of spinful fermions with dispersion linearized close to the Fermi energy:

$$H_L = -iv_F \sum_{a,\sigma=\uparrow/\downarrow}^N \int_0^\infty dx \,\psi_{a,R,\sigma}^\dagger \partial_x \psi_{a,R,\sigma} - \psi_{a,L,\sigma}^\dagger \partial_x \psi_{a,L,\sigma},$$

where $\psi_{a,r=L/R,\sigma=\uparrow/\downarrow}(x)$ are operators annihilating left/right -moving modes with spin σ at the point x of the lead a, combining into $\psi_{a,\sigma}(x) = \psi_{a,R,\sigma}(x)e^{ik_Fx} + \psi_{a,L,\sigma}e^{-ik_Fx}$. However, a difference arises in the topological case, because Majorana states couple only to one of two spin components [68,69]. This allows us to drop the spin index in this case and consider spinless fermions.

The leads are semi-infinite, ending at x = 0, where they are in contact with the superconductor. The exact form of the tunneling Hamiltonian depends then on the type of superconductor. In the case of the *s*-wave superconductor, the charge transfer into the island will occur due to the Andreev processes in which the incident electron in the lead is reflected as a hole and at the same time a single Cooper pair is added to the superconductor. Using the pseudospin-1/2 representation of the charge state of the island, we can write the tunneling Hamiltonian following Ref. [39]:

$$H_T = \sum_{a=1}^{N} t_a [\psi_{a,\uparrow}^{\dagger}(0)\psi_{a,\downarrow}^{\dagger}(0)s^- + \psi_{a,\downarrow}(0)\psi_{a,\uparrow}(0)s^+], \quad (4)$$

where we are either adding or removing two electrons of opposite spin at the x = 0 point of the lead a and at the same time changing the charge state of the island between $2N_0$ and $2N_0 + 2$. In writing this Hamiltonian, we assumed that the superconducting island is large enough so that the crossed Andreev reflection is suppressed. On the other hand, in the case of a topological superconductor, the tunneling will occur into the Majorana zero modes. We also use a pseudospin-1/2 representation of the charge state, with transitions between the states with $2N_0$ and $2N_0 + 1$ electrons. Then the tunneling Hamiltonian has the form [34,35]

$$H_T = \sum_{a=1}^{N} [t_a \psi_a^{\dagger}(0) s^{-} \gamma_a + \text{H.c.}], \qquad (5)$$

where t_a are tunnel couplings to the leads, $\psi_a^{\dagger}(0)$ are creation operators at the end of the leads, and $\gamma_a = \gamma_a^{\dagger}$ are Majorana operators.

Before turning to a bosonization analysis of these models, we would like to comment on the similarities and differences between superconducting islands and normal dots. In normal dots with small energy level spacing, transport at the charge degeneracy point is usually dominated by inelastic cotunneling events [47] and therefore is incoherent. In the case of an ordinary superconductor, transport occurs via resonant tunneling of Cooper pairs and is also incoherent. However, in topological islands, the resonant tunneling processes through Majorana states are phase-coherent [29] and therefore allow for interference effects to be used as probes for topological states.

III. BOSONIZATION ANALYSIS

Both setups can now be studied using bosonization by transforming the normal leads into Luttinger liquids, spinful in the case of an *s*-wave island and spinless when leads are coupled to Majorana zero modes. We derive the results for the ordinary superconductor and then highlight the differences that arise in the Majorana setup [34,35].

A. s-wave island

After spinful bosonization, the Hamiltonian of the leads now has the form [70]

$$H_{L} = \sum_{a=1}^{N} \frac{v}{2\pi} \sum_{j=\rho,\sigma} \int_{0}^{\infty} dx \, K_{j} (\nabla \theta_{a,j})^{2} + \frac{1}{K_{j}} (\nabla \phi_{a,j})^{2},$$
(6)

where we have used the following convention:

$$\psi_{a,r,\sigma}(x) = \frac{U_{a,r,\sigma}}{\sqrt{2\pi\alpha}} e^{-\frac{i}{\sqrt{2}} \{r\phi_{a,\rho}(x) - \theta_{a,\rho}(x) + \sigma[r\phi_{a,\sigma}(x) - \theta_{a,\sigma}(x)]\}},$$
(7)

with α being the short-distance cutoff, and $U_{a,r,\sigma}$ are the Klein factors. Using (7) we can now express the tunneling Hamiltonian using bosonic fields. Since the lead ends at x = 0, we can impose the open boundary condition $\psi_{a,L,\sigma}(0) = \psi_{a,R,\sigma}(0)$. This in turn means that $\phi_{\rho/\sigma}(0) = 0$ and that Klein factors for right and left movers of each spin are equal: $U_{a,R,\sigma} =$ $U_{a,L,\sigma} = U_{a,\sigma}$. An alternative approach would be to consider a single chiral bosonic field obtained by unfolding right- and left-moving modes onto a single axis extending from $-\infty$ to ∞ . Combining all of this together, we express the tunneling Hamiltonian (4) as

$$H_T = \sum_{a=1}^{N} \frac{2t_a}{\pi \alpha} (U_{a,\uparrow} U_{a,\downarrow} e^{-i\sqrt{2}\theta_{a,\rho}(0)} s^- + \text{H.c.}).$$
(8)

We can form a parity operator from the Klein factors $p_a = iU_{a,\uparrow}U_{a,\downarrow}$, and since $p_a^2 = 1$ we can use the identity $e^{i\gamma p_a} = \cos(\gamma) + ip_a \sin(\gamma)$. For $\gamma = \frac{\pi}{2}$ this translates to $ip_a = e^{i\frac{\pi}{2}p_a}$, so we have $U_{a,\uparrow}U_{a,\downarrow} = -ip_a = e^{-i\frac{\pi}{2}p_a}$. Thus the Klein factors translate to a phase shift, which can be absorbed into the bosonic field, because the parity in each lead is fixed as the only allowed tunneling process transfers pairs of electrons. The final form of the tunneling Hamiltonian is then

$$H_T = \sum_{a=1}^{N} \frac{2t_a}{\pi \alpha} (e^{-i\sqrt{2}\theta_{a,\rho}(0)}s^- + e^{i\sqrt{2}\theta_{a,\rho}(0)}s^+).$$
(9)

Therefore, both bosonic fields from the spin sector (θ_{σ} and ϕ_{σ}) and ϕ_{ρ} are not present in the tunneling Hamiltonian, and they are present only in the quadratic part of the action. This means that we can integrate them out from the imaginary-time action. Moreover, the field θ_{ρ} is taken only at x = 0 in H_T , so we can also integrate it out away from x = 0. After this procedure, we obtain the imaginary-time action:

2

$$S^{s-\text{wave}} = S_0^{s-\text{wave}} + S_T^{s-\text{wave}},\tag{10}$$

$$S_0^{s\text{-wave}} = \sum_{a=1}^N \frac{K_\rho}{2\pi} \int \frac{d\omega}{2\pi} |\omega| |\theta_{a,\rho}(\omega)|^2, \qquad (11)$$

$$S_T^{s\text{-wave}} = \sum_{a=1}^N \int_0^\beta d\tau \, \frac{J_{\perp,a}}{2} (e^{-i\sqrt{2}\theta_{a,\rho}(0)} s^- + \text{H.c.}).$$
(12)

In anticipation of the connection of this action to the multichannel Kondo problem, we have introduced the notation for the coupling $J_{\perp,a} = \frac{4t_a}{\pi a}$.

B. Majorana island

The procedure of obtaining the effective boundary action in the case of a Majorana island is essentially the same, with the important distinction that the leads now contain effectively spinless electrons and so the bosonization identity now takes the form

$$\psi_{a,r}(x) = \frac{U_{a,r}}{\sqrt{2\pi\alpha}} e^{-i[r\phi_a(x) - \theta_a(x)]}$$
(13)

with α again being the short-distance cutoff and $U_{a,r}$ are the Klein factors. The bosonized Hamiltonian of the leads is

$$H_{L} = \sum_{a=1}^{N} \frac{v}{2\pi} \int_{0}^{\infty} dx \ K(\nabla \theta_{a})^{2} + \frac{1}{K} (\nabla \phi_{a})^{2}$$
(14)

and the tunneling Hamiltonian is

$$H_T = \sum_{a=1}^{N} \frac{2t_a}{\sqrt{2\pi\alpha}} (e^{-i\theta_a(0)}s^- + e^{i\theta_a(0)}s^+), \qquad (15)$$

where the Klein factors hybridized with Majorana operators in a process of statistical transmutation [34,71].

When the bosonic field is integrated out away from x = 0, we obtain the imaginary-time action:

$$S^{\rm M} = S_0^{\rm M} + S_T^{\rm M}, (16)$$

$$S_0^{\rm M} = \sum_{a=1}^N \frac{K}{2\pi} \int \frac{d\omega}{2\pi} |\omega| |\theta_a(\omega)|^2, \qquad (17)$$

$$S_T^{\rm M} = \sum_{a=1}^N \int_0^\beta d\tau \, \frac{J_{\perp,a}}{2} (e^{-i\theta_a(0)} s^- + e^{i\theta_a(0)} s^+). \tag{18}$$

This time we made the identification $J_{\perp,a} = \frac{4t_a}{\sqrt{2\pi\alpha}}$. It is interesting to make a comparison between tunneling parts of the action for both cases. Equations (12) and (18) have virtually the same form, apart from the factor of $\sqrt{2}$ in the exponent for the *s*-wave superconducting island model. One can then perform the following transformation of the action (18): $\theta_a \rightarrow \sqrt{2\tilde{\theta}_a}$. To keep the quadratic part of the action the same under this transformation, we also have to rescale the Luttinger parameter $K: K \rightarrow \tilde{K}/2$. This means that the topological system will behave exactly the same as the conventional one, but with the interaction parameter rescaled by a factor of 2. This bears important consequences for the flow diagram of the perturbative RG close to the noninteracting value of K = 1.

C. Perturbative renormalization-group analysis

Because we have shown that there exists a direct correspondence between the actions for both *s*-wave and topological islands, it is sufficient to perform perturbative renormalization-group analysis of the action of the *s*-wave setup and then recover the behavior of the Majorana island by substituting \tilde{K} for K_{ρ} . During the RG procedure, an additional term is generated that is proportional to $\partial_x \phi_{a,\rho}$, even if its coupling is initially zero. Therefore, we add it into the action right from the beginning with J_z coupling:

$$S_z = -\sum_{a=1}^N \frac{v}{\sqrt{2}} J_z s_z \partial_x \phi_{a,\rho}.$$
 (19)

With this additional term and the relabeling of the couplings done in the previous section, the complete action for our problem has exactly the same form as the anisotropic multichannel Kondo problem action [36,37]. Therefore, the analysis steps follow directly from the standard procedure applied previously to the Kondo problem.

We begin by considering the isotropic limit when all $J_{\perp,a} = J_{\perp}$. In such a case, the RG equations for the couplings are

$$\frac{dJ_z}{dl} = J_\perp^2 \left(\frac{1}{K_\rho} - \frac{N}{2}J_z\right),\tag{20}$$

$$\frac{dJ_{\perp}}{dl} = \left(1 - \frac{1}{K_{\rho}}\right)J_{\perp} + J_z J_{\perp} \left(1 - \frac{NK_{\rho}}{4}J_z\right).$$
(21)

Those equations are exact in J_z and perturbative in J_{\perp} . We notice that in the isotropic case, the couplings flow to the Toulouse fixed point, where J_z becomes $\frac{2}{NK_{\rho}}$ and its flow stops. This means that we can perform a unitary transformation and eliminate the $\partial_x \phi_{\rho}$ term from the Hamiltonian

$$U = e^{iK_{\rho}J_{z}\sqrt{\frac{N}{2}}\Theta(0)s_{z}},$$
(22)

$$U^{\dagger}HU = H_L - \sum_{a=1}^{n} \frac{J_{\perp}}{2} (e^{-i\sqrt{2}[\theta_{a,\rho}(0) - \frac{1}{\sqrt{N}}\Theta(0)]} s^- + \text{H.c.}),$$
(23)

where $\Theta(0) = \frac{1}{\sqrt{N}} \sum_{j} \theta_{j,\rho}(0)$ is the global mode.

We can now determine the fixed points of the problem and understand them using the quantum Brownian motion (QBM) correspondence. In the QBM picture, the state of the system is described as a position of a fictitious particle placed in a periodic potential with a dissipative environment. This enables us to approach the problem from two dual perspectives: tunneling between the minima of a strong periodic potential, and free motion with weak potential as a perturbation. To make the mapping clearer, we can write the tunneling operators in the action as $e^{-i\sqrt{2}\vec{\theta}_{\rho}\vec{K}_{0}^{(a)}}s^{-}$, where $\vec{\theta}_{\rho} = (\theta_{1,\rho}, \dots, \theta_{N,\rho})$, and $\vec{R}_0^{(a)}$ is a vector with 1 on the *a*th component and 0 on the rest. In this notation, one can think of $\vec{\theta}_{\rho}$ as the momentum of the particle, and the number of charges in the leads (which is a variable conjugate to θ_{ρ}) describes the position of the particle. When the periodic potential is strong, the particle is mostly localized in the minima of the potential, which are connected by the lattice vectors $\vec{R}_0^{(a)}$ and only occasionally



FIG. 2. (a) Visualization of the QBM lattice for N = 3. Each axis corresponds to the number of Cooper pairs (or electrons in the topological island case) inside of a particular lead. Blue and red points form planes to which the motion of the particle is restricted for given \mathcal{N} . (b) Renormalization-group flow for *s*-wave and topological superconducting (TSC) islands connected to N < 5 leads as a function of the Luttinger parameter *K*. Stable and unstable fixed points are depicted as solid and dashed lines, respectively. The bottom line corresponds to the limit of weak tunneling $(t \rightarrow 0)$ and the top line corresponds to weak periodic potential $(v_{\vec{G}} \rightarrow 0)$. Due to a rescaling of *K*, the Majorana island case for noninteracting leads (K = 1) is of a strong-coupling nature, compared to intermediate coupling for the *s*-wave island.

tunnels between them. Since we have charging energy in our setup and the island can only accommodate a single additional Cooper pair, the total number of charges in the leads N_{tot} can only change between $\mathcal{N}/\mathcal{N} + 2$, and the particle's motion is restricted to two planes in the *N*-dimensional space. The corresponding lattices are a 1D zigzag chain for N = 2 channels and a (N - 1)-dimensional hyperhexagonal lattice for N > 2. Both lattice types are nonsymmorphic with a two-atom basis, which corresponds to the presence or absence of the additional Cooper pair in the superconducting island. An example of such a lattice for N = 3 leads is shown in Fig. 2(a). Each axis corresponds to N_i of Cooper pairs present in the leads, so $2N_1 + 2N_2 + 2N_3 = \mathcal{N}$. Blue points form a plane for one particular value of \mathcal{N} , while red points form a neighboring plane corresponding to $\mathcal{N} + 2$.

In this language, the global mode introduced after the unitary transformation at the Toulouse fixed point corresponds to the product of $\vec{\theta}_{\rho}$ and the vector $\vec{R}_{\perp} = \frac{1}{\sqrt{N}}(1, \ldots, 1)$ perpendicular to the planes to which the particle motion is confined. The tunneling operators after the transformation are $e^{-i\sqrt{2}\vec{\theta}_{\rho}\vec{R}_{\parallel}^{(a)}}s^{-}$, with $\vec{R}_{\parallel}^{(a)} = \vec{R}_{0}^{(a)} - \frac{1}{\sqrt{N}}\vec{R}_{\perp}$. The scaling dimension of the tunneling operator is then

$$\Delta[e^{-i\sqrt{2}\vec{\theta}_{\rho}\vec{R}_{\parallel}^{(a)}}s^{-}] = \frac{|\vec{R}_{\parallel}^{(a)}|^{2}}{K_{\rho}} = \frac{1}{K_{\rho}}\left(1 - \frac{1}{N}\right).$$
(24)

Therefore, determining whether the tunneling operator is relevant depends on the Luttinger parameter K_{ρ} —the condition for relevancy is $K_{\rho} > \frac{N-1}{N}$. Importantly, this means that for noninteracting electrons ($K_{\rho} = 1$), for all N the tunneling operator is relevant and the system will be flowing in the direction of decreasing periodic potential strength, away from the localized fixed point. Since the coupling increases substantially, the perturbation theory breaks down and we need to find the stable fixed-point properties in another way. To do this, we can use the dual perspective of looking at the QBM as a free motion with weak potential perturbation. In this case, the periodic potential can be expressed using its Fourier components $V(\vec{r}) = \sum_{\vec{G}} v_{\vec{G}} e^{i\vec{G}\cdot\vec{r}}$, where \vec{G} are reciprocal-lattice vectors of the honeycomb lattice. Then the scaling dimension of the most relevant $v_{\vec{G}}$ (corresponding to the shortest \vec{G}) is given by [34]

$$\Delta[e^{i\vec{G}\cdot\vec{r}}] = K_{\rho}|\vec{G}|^2 = K_{\rho}\left(1 - \frac{1}{N}\right).$$
 (25)

Again, the relevancy of the periodic potential perturbation depends on the value of K_{ρ} . The criterion in this case is $K_{\rho} < \frac{N}{N-1}$, which for noninteracting leads is always satisfied: the periodic potential is a relevant perturbation to the free motion fixed point. Therefore, there have to be additional fixed points between the localized and free motion, including at least one stable fixed point. This analysis is summarized for N < 5 in Fig. 2, which indicates stable and unstable fixed points as solid and dashed lines, respectively. For $N \ge 5$ there exists another unstable intermediate-coupling fixed point that has been analyzed in more detail by Yi [37]. The stable intermediate-coupling fixed point for noninteracting leads has been studied using conformal field theory in the context of the multichannel Kondo problem [72,73]. Applying those results to our model, we can immediately find the zero-temperature residual entropy:

$$S_{\rm imp}(T=0) = \ln\left[2\cos\left(\frac{\pi}{N+2}\right)\right].$$
 (26)

Moreover, we can also deduce the zero-temperature conductance matrix elements to be

$$G_{ij}(T=0) = 8\sin^2\left(\frac{\pi}{N+2}\right)\left(\frac{1}{N} - \delta_{ij}\right)\frac{e^2}{h}.$$
 (27)

The important distinguishing feature is that compared to the Kondo problem, the conductance matrix element here is quadrupled. Each Andreev reflection process transfers the charge of 2e between the leads and superconducting island, corresponding to doubling of the current compared to the conventional charge Kondo effect. This current operator is then used in the Kubo formula to obtain conductance as a currentcurrent correlation function, and the doubling translates in this way to quadrupling of $G_{ij}(T = 0)$. The conformal field theory also gives the scaling dimension of the leading irrelevant operator at the intermediate fixed point that translates to the leading temperature correction to $G_{ij}(T = 0)$:

$$G_{ij}(T) = G_{ij}(T=0) \left[1 - c \left(\frac{T}{T_K} \right)^{\lambda} \right], \qquad (28)$$

where λ is 1 for N = 2 [47,74,75] and $\frac{2}{5}$ for N = 3 [33,51,73], c is a constant on the order of unity, and T_K is the Kondo temperature. One important characteristic of the intermediate fixed point is that it is unstable to channel coupling asymmetry [76,77]: when one of the couplings is increased, the system will flow to a Fermi liquid fixed point that describes the single channel Kondo model, and when one of the couplings is decreased, the system will behave as an N - 1 channel setup in low temperatures. In general, the asymmetric system will behave as an N_{max} channel setup at low-energy scales,

where N_{max} is the number of leads with the largest coupling value. This constitutes a significant difficulty in performing experiments that verify the theoretical claims listed above.

Now we can turn to the case of a Majorana island in which we have to substitute $K_{\rho} \rightarrow \tilde{K} = 2K$. This change essentially shifts the flow diagram and redefines the condition for the relevancy of tunneling and weak periodic potential operators, which are now $K > \frac{N-1}{2N}$ and $K < \frac{N}{2(N-1)}$, respectively. This is also indicated in Fig. 2(b) (which again is valid for N < 5with a new unstable fixed point appearing for $N \ge 5$). The redefinition of the relevancy condition brings about a crucial change for the noninteracting leads: while the tunneling operator is still relevant for K = 1, the weak periodic potential becomes irrelevant for all N. This means that the free motion fixed point becomes stable and that conductance will assume the maximum value allowed by charge conservation. Remembering that in the Majorana island the tunneling processes carry a charge of 1*e*, we find that the conductance is

$$G_{ij}(T=0) = 2\left(\frac{1}{N} - \delta_{ij}\right)\frac{e^2}{h}.$$
(29)

The weak periodic potential becomes now the leading irrelevant operator, and its scaling dimension will now determine the exponent of the temperature correction of the conductance:

$$\Delta_{\rm irr} = 2\left(1 - \frac{1}{N}\right).\tag{30}$$

The form of the correction is still described by (28). The change of the nature of the low-temperature fixed point comes with another major difference: the channel coupling anisotropy, which corresponds to deformation of the periodic potential, becomes an irrelevant perturbation and does not cause the system to flow to the Fermi liquid fixed point. This will be explored in more detail in Sec. IV.

IV. NUMERICAL RESULTS

To verify the analytical results and obtain a fuller understanding of the crossover regime between the fixed points of the studied models, we employ the numerical renormalization-group (NRG) method, a powerful nonperturbative method for obtaining thermodynamics and correlation functions of quantum impurity systems, connected to non-interacting leads [78]. As we want to capture the universal physics of this setup, we simplify the problem by assuming that all the leads are identical with bandwidth 2D and a flat density of states $\rho = 1/2D$. Then we express the Hamiltonians in the form suitable for numerical calculations. More details of the numerical calculation are presented in Appendix A. The NRG simulations have been performed using the NRG Ljubljana code, which internally makes use of the SNEG library [79].

To directly relate our results to the experiment, we focus on the dc conductance in our calculations. We work in the framework of the linear-response theory and compute ac conductance using the Kubo formula as the correlation function of the number of electrons in one lead and current in the other lead. This allows us to avoid computation of the delicate limit present in the usual current-current correlation approach (see Appendix A). Finally, we obtain dc conductance as the limit $G_{jkdc}(T) = \lim_{\omega \to 0} G_{jkdc}(\omega, T)$ of the ac conductance.

A. Superconducting island

We begin by analyzing the numerical results obtained in the case of an *s*-wave superconductor island. Since this setup maps exactly to the multichannel Kondo problem (as shown above), which has been studied extensively using NRG, we only highlight that the Andreev reflection Hamiltonian indeed reproduces the key results of the Kondo effect. The Hamiltonian used in NRG simulations is

$$H_{\rm SC}^{\rm NRG} = H_{\rm leads} + \sum_{a=1}^{N} t_a c_{a0\uparrow}^{\dagger} c_{a0\downarrow}^{\dagger} f_{\downarrow} f_{\uparrow} + \text{H.c.}, \qquad (31)$$

where $c_{a0\sigma}$ are the fermionic operators at the end of the Wilson chain that are connected to the superconducting island, and f_{σ} describes the pair of fermionic states in the island that are only both occupied or both empty at the same time, simulating the two possible charge states of the island. First, we look at the entropy of the island at low temperatures [Fig. 3(a)]. For both the two- and three-channel cases tuned to the critical point, we observe residual entropy as in the usual Kondo effect. In the two-channel case, the entropy flows to $S_{2ch}(T = 0) =$ $\ln(2)/2$, which is explained by the observation of Emery and Kivelson [80] that the two-channel Kondo model maps to a resonant level system with only half of the impurity degrees of freedom coupled to the conduction electrons. For the threechannel case, the entropy flows to $S_{3ch}(T=0) = \ln(\frac{1+\sqrt{5}}{2})$, which is consistent with the conformal field theory result and previous numerical studies of the regular Kondo effect [81]. The inset of Fig. 3(a) shows scaling of the Kondo temperature for the two-channel model as the tunnel couplings are varied, and this dependence also exactly follows the behavior of the charge Kondo problem [74]:

$$T_K/D \sim \rho J \exp\left(-\frac{\pi}{4\rho J}\right).$$
 (32)

Next we move on to linear conductance between the normal leads. In Fig. 3(b) we show the ac conductance matrix element $G_{12}(\omega)$ for several temperatures for the case of two channels. All the curves follow the same universal behavior before saturating at their respective dc limit, which in the limit of T = 0 is equal to $2e^2/h$ as predicted by the lowenergy fixed point in the perturbative renormalization-group scheme and obtained previously by Pustilnik et al. [39]. The values of $G_{12}(\omega \to 0)$ are then determined for all the remaining temperatures and plotted in Fig. 3(c), together with corresponding values for the three-channel setup. For the three-channel setup, the predicted value of $\frac{8}{3}\sin^2(\frac{\pi}{5}) \approx 0.92$ is also observed. This calculated temperature dependence is then fitted with the low-temperature correction determined by the scaling dimension of the leading irrelevant operator at the intermediate fixed point. For $T \ll T_K$ we observe excellent agreement of the calculated curve with the predicted exponent $\Delta G \sim T$ in the case of two leads and $\Delta G \sim T^{2/5}$ in the case of three leads.

All of the results described above are unstable with respect to the tunnel coupling anisotropy, so if the values of t_a are



FIG. 3. (a) The impurity entropy $S_{imp}(T)$ curves showing the crossover between local moment $[S_{imp}(T) = \ln(2)]$ and non-Fermi-liquid fixed points $[S_{imp}(T) = \ln(\sqrt{2})$ for two channels and $\ln(\frac{1+\sqrt{5}}{2})$ for three channels]. The inset shows that T_K obtained at the center of the crossover (blue points) is given by $T_K \approx \rho J_{\perp} \exp(\frac{\pi}{4\rho J_{\perp}})$ (orange line). (b) $G_{12}(\omega)$ conductance matrix element for two channels in the isotropic case $t_1 = t_2 = 0.15D$ for several temperatures. (c) dc conductance $G_{12dc}(T)$ with a power-law correction given by the leading irrelevant operator for two ($\Delta G \sim T$) and three ($\Delta G \sim T^{2/5}$) channels.

detuned from a common value, the system in general flows to the Fermi-liquid fixed point of the single-channel Kondo model, as expected [76].

B. Majorana island

In the numerical analysis of the Majorana island model, we limit our considerations to the first nontrivial case with N = 3 leads. We consider a model that includes four Majorana modes in the island, one of which is not coupled to any lead. In such a system, the dimension of the Hilbert space is 4. It is then divided into two two-dimensional subspaces labeled by the fermion parity of the island. To transform the Hamiltonian to a form suitable for NRG calculations, we introduce a spinless fermion f^{\dagger} on the island to distinguish the two subspaces. Each of the two-dimensional parity subspaces is then described by a pseudospin-1/2 impurity $\vec{\sigma}$. We note that this pseudospin-1/2 object is different from \vec{s} used in the bosonization treatment, which was related to different charge states of the island. Then the Hamiltonian has the following form:

$$H = H_{\text{leads}} + \sum_{j=1}^{3} (t_j \psi_j^{\dagger} \sigma_j f + \text{H.c.}) + \delta \left(f^{\dagger} f - \frac{1}{2} \right). \quad (33)$$

We then define the level broadening $\Gamma = \rho t_{avg}^2$, where t_{avg} is the average tunnel coupling between the island and the leads. Even though Majorana hybridization is a relevant perturbation in our model, in most of our calculations we neglect it, motivated by experimental results [13] that suggest that minimizing the hybridization by using sufficiently long nanowires is possible and allows for performing satisfactory measurements. However, in order to test this assumption, we performed some calculations with an additional term $H_{hyb} = b_{jk}i\gamma_j\gamma_k$. In our mapping of the Majoranas to a pseudospin-1/2 object, this translates to $H_{hyb} = \vec{K} \cdot \vec{\sigma}$, an effective magnetic field for this spin. The Hamiltonian of Eq. (33) is now suitable for NRG treatment.

In our numerical analysis, we will be comparing the model of a topological superconductor island at charge degeneracy

(presented above) with the previously studied model [57] that describes the island in the Coulomb valley regime (topological Kondo regime). We begin our investigation with the first property that distinguishes the charge degeneracy point model from the topological Kondo regime, namely the temperature of the transition from the local moment fixed point to the non-Fermi-liquid fixed point. The dependence of the transition temperature T^* on the lead coupling parameter can be established in more detail by analyzing the flow of the entropy of the island to the non-Fermi-liquid fixed point, which is shown in Fig. 4(a). The entropy values flow from the local moment fixed point with $S_{imp}(T) = \ln(4)$ to the non-Fermi-liquid fixed point with $S_{imp}(T) = \ln(\sqrt{3})$. When the temperatures are expressed in units of the level broadening Γ , all the entropy curves collapse into one universal dependence. Now we define the transition temperature T^* as the temperature for which the impurity entropy attains the value $\left[\ln(4) + \ln(\sqrt{3})\right]/2$ that is in the middle between values at the two fixed points. We obtain it by numerically solving the equation $S_{imp}(T^*) = [\ln(4) + \ln(\sqrt{3})]/2$ and plot it as a function of the level broadening [inset of Fig. 4(a)]. The line on which the T^* points lie is defined as $T^* = c\Gamma$, where $c \approx 3.60$ is a constant coefficient determined from the fitting procedure. Since there is a direct relation between T^* and Γ , one can assess the transition temperature by estimating the value of level broadening as $g\Delta$ [30], with g being the sum of dimensionless conductances and Δ the superconducting gap in the Majorana island. In such a case, level broadening values are of the order of 10 μ eV. Such values translate to a temperature of about a few hundred mK. To contrast this with the previous proposals, in Fig. 4(b) we show the comparison between the crossover temperatures T^* for our model and the model in the topological Kondo regime (details of the model are provided in Appendix B) in the fully isotropic case $(t = t_1 = t_2 = t_3)$. Even for large tunnel couplings, T^* at the charge degeneracy point is at least three orders of magnitude higher than in the topological Kondo regime. Moreover, the Kondo temperature drops sharply with decreasing couplings $(T_K \sim (\rho t)^2 \exp[-1/(2\rho t)])$, while at the charge degeneracy point $T^* \sim t^2$, which can lead to a much easier



FIG. 4. (a) The collapsed (for *T* in units of Γ) island's entropy $S_{imp}(T)$ curves showing the crossover between local moment [$S_{imp}(T) = \ln(4)$] and non-Fermi-liquid fixed points [$S_{imp}(T) = \ln(\sqrt{3})$]. The inset shows the linear relation $T^* = c\Gamma$ with $c \approx 3.60$ obtained from fitting. (b) Crossover temperature comparison between the charge degeneracy point and the topological Kondo (Coulomb valley) regime for several values of $t = t_1 = t_2 = t_3$. The temperature at the charge degeneracy point is at least three orders of magnitude higher than in the topological Kondo regime.

experimental observation of the multiterminal teleportation. Furthermore, it would be possible to directly measure the dependence of T^* on the tunnel couplings by tuning them using external gates.

Next, we move to computing the transport properties of the three-terminal Majorana island. We start by analyzing the results exactly at the charge degeneracy point (when $\delta = 0$). In Fig. 5(a), we show the $G_{12}(\omega)$ ac conductance matrix element in the isotropic case ($t_1 = t_2 = t_3 = 0.05D$) for varying temperatures. All the computed curves follow a universal dependence, and at low temperature the fractional quantized value of $2/3e^2/h$ is attained as predicted by the quantum Brownian motion mapping. In Fig. 5(b) the temperature dependence of the G_{12dc} dc conductance is shown. The whole crossover happens over the span of approximately two orders of magnitude in temperature, which means it is much steeper than the crossover studied previously in the topological Kondo regime. This is another factor that can make the experiment possible—the increase of conductance should start at several Kelvins and approach the fractional quantized value for several mK. The quantum Brownian motion mapping provides a prediction of a universal power-law temperature correction to conductance at the strong-coupling fixed point, which has the form

$$G_{12dc}(T) = G_{12dc}(T=0) [1 - c'(T/T^*)^{2/3}].$$
 (34)

The curve presented in the plot is a fit of the predicted dependence, and it correctly describes a significant part of the crossover. This fact, together with the high crossover temperature, should allow for experimental verification of this low-temperature conductance correction.

However, in a real experiment, reaching the exact isotropic case requires fine-tuning. Therefore, it is important to verify the prediction of robustness to channel coupling asymmetry. In Fig. 5(c) we show the results for a fully anisotropic set of coupling constants ($t_1 = 0.0475D$, $t_2 = 0.0525D$, and t_3)



FIG. 5. (a) $G_{12}(\omega)$ conductance matrix element in the isotropic case $t_1 = t_2 = t_3 = 0.1D$ for several different temperatures showing a universal behavior. (b) $G_{12dc}(T)$ dc conductance with a fit of the universal temperature power-law correction with an exponent of 2/3. (c) $G_{12}(\omega)$ conductance matrix element in a fully anisotropic case $t_1 = 0.0475D$, $t_2 = 0.0525D$, and t_3 from the interval [0.00625D, 0.2D] with each curve increasing t_3 by a factor of 2. The inset shows the temperature dependence of $G_{12dc}(\omega)$ dc conductance for the case when $t_1 = t_2 = 0.05D$, $t_3 = 0.0125D$ with a nonmonotonic behavior that is a signature of crossover between two- and three-terminal teleportation. The curve is a fit of $T^{2/3}$ dependence.



FIG. 6. $G_{12dc}(\delta)$ dc conductance away from the charge degeneracy point for several temperatures. For low temperatures, the top of the curve becomes flattened at the value of $2/3 e^2/h$, which implies robustness against charge detuning.

varying in the range $[0.006\,25D, 0.2D]$ in T = 0, with each step increasing t_3 by a factor of 2). In this case, the dc conductance also reaches the value of $2/3 e^2/h$ independently of the initial value of t_3 , which is in stark contrast to the s-wave island model. Moreover, in the case of decreasing t_3 , one can observe a nontrivial crossover between the cases with two and three leads. For ω just below Γ , the value of conductance goes beyond the value of 2/3 and comes close to $1 e^2/h$, which is the value corresponding to the electron teleportation between only two leads. However, going further to lower frequencies decreases conductance and it again attains the fractional quantized value. This behavior is mimicked in the temperature dependence of dc conductance, which is shown in the inset of Fig. 5(c). We observe a nonmonotonic dependence, which first rises above the fractional value for intermediate temperatures, but in the low-temperature limit goes back to $2/3 e^2/h$. The curve is a fit of a $T^{2/3}$ dependence, in this case with a positive coefficient in front of it. This nonmonotonic behavior can be used as one of the experimental signatures of crossing between two- and three-terminal teleportation regimes. However, due to the slow decay of conductance back to the fractional value, reaching the low-temperature limit may prove to be more difficult.

Another important factor for the experimental verification of our claims is the sensitivity to tuning the system exactly to the charge degeneracy point. In Fig. 6 we present dc conductance of our system as a function of the energy shift δ away from the charge degeneracy point for four different temperatures. For the lowest temperature, the curve becomes flattened at the top, which corresponds to the conductance value of $2/3 e^2/h$. This flat top means that even when one moves away from the resonance, the observed conductance would still be equal to the fractional quantized value. For increased temperatures, the curves become narrower, but still it is reasonable to expect to observe a nonzero value of conductance even when being away from the charge degeneracy point. Nevertheless, this proves that tuning the system into the vicinity of the charge degeneracy point is crucial to observe fractional conductance at the temperatures within experimental reach.



FIG. 7. $G_{12}(\omega)$ in the isotropic case $t_1 = t_2 = t_3 = 0.1D$ for several values of hybridization strength $K_x \neq K_y \neq K_z$ with a constant ratio 0.84:1:1.11 (K being the proportionality constant). The hybridization affects conductance only for very small frequencies (and temperatures), so even a sizable overlap of Majorana states would not preclude experimental observation.

 ω/Γ

10

10

Finally, we study how the conductance is impacted by introducing Majorana hybridization into our Hamiltonian. Since hybridization is a relevant perturbation, one expects that in low temperatures it will significantly change the behavior of conductance. In Fig. 7 we show $G_{12}(\omega)$ in the isotropic case $t_1 = t_2 = t_3 = 0.1D$ for several generic values of hybridization strength $K_x \neq K_y \neq K_z$ with constant ratio 0.84:1:1.11 between the components of \vec{K} (additional results for different values of K components are presented in Appendix B). The conductance rises from 0, reaches a value of $2/3 e^2/h$, and then at lower energy scales changes to some nonuniversal value. The scale at which the crossover happens depends on the hybridization strength. We define the transition energy scale ω_H as the value for which conductance is in the middle between $2/3 e^2/h$ and the nonuniversal value. We can then extract $\omega_H \sim a K^{3/2}$, with the value of *a* dependent on the tunnel couplings in a particular simulation and in the cases presented here a lies in the range between 0.1 and 0.7. The most important fact is that even with a sizable magnitude of hybridization, it affects conductance only in the very lowenergy scales, and the fractional quantized conductance still prevails in the range of temperatures available in the experiment. This justifies neglecting the Majorana hybridization in the rest of the calculations.

Having verified the claim of robustness of our results with respect to the tunnel coupling anisotropy, charge degeneracy detuning, and showing that hybridization affects the results only at very low temperatures, we propose an experiment that yields a direct signature of the multiterminal Majoranaassisted electron teleportation. In Fig. 8 we show the dc conductance as a function of the tunnel coupling of the third lead for several different temperatures slightly off the charge degeneracy point to simulate the experimental conditions. At high temperatures, the conductance increases straight to the values close to $1 e^2/h$ while decreasing the tunnel coupling, as is expected for the electron teleportation between two leads. However, as the temperature is lowered, a plateau at 2/3 e^2/h emerges and it becomes wider in the process of cooling



FIG. 8. $G_{12dc}(t_3)$ dc conductance slightly away from the charge degeneracy point ($\delta = 0.0035D$) as a function of tunnel coupling of the third lead for several temperatures. In high temperatures there is a simple transition from 0 to over 0.8 e^2/h . When the system is cooled down, a plateau emerges at $2/3 e^2/h$, signifying a transition from two- to three-terminal electron teleportation.

down the system. Remarkably, the whole shape of the curve changes, the increase of conductance starting for larger tunnel couplings in lower temperatures, which allows us to observe the change for a large range of tunnel coupling strengths. This change of conductance curve shape provides direct evidence of entering the multiterminal teleportation regime.

V. SUMMARY

We have shown that both the conventional and the topological superconducting islands at the charge degeneracy point are interesting in their own right. By applying bosonization techniques, we demonstrated that the multiterminal s-wave superconductor island Hamiltonian maps to the multichannel Kondo problem. For the case of noninteracting leads, this means that at low temperatures the system is described by an intermediate coupling fixed point that displays non-Fermiliquid behavior and for which many observables are known from conformal field theory. We supported the mapping by a numerical renormalization-group calculation, which gives the residual entropy and conductance consistent with the analytical prediction. The intermediate fixed point is in general unstable to channel coupling asymmetry, and so experimental verification would require fine-tuning. On the other hand, due to Luttinger parameter rescaling, the topological superconductor island flows to a strong-coupling fixed point, which grants robustness to the anisotropy. This conclusion is backed by numerical calculation in which the conductance for N = 3leads reaches the value of $2/3 e^2/h$ independently of the initial tunnel couplings. Moreover, the crossover to a non-Fermiliquid fixed point happens at experimentally plausible temperatures, compared to the previous studies of the topological Kondo effect. Thanks to this robustness, for the topological island we have predicted distinctive experimental signatures of a crossover between two- and three-terminal cases: one is a nonmonotonic temperature dependence of dc conductance when the coupling of one of the leads is decreased, and the other is the change of the shape of the tunnel coupling strength dependence of dc conductance with a plateau emerging at

 $2/3 \ e^2/h$ while decreasing the temperature. As the experimental control of the hybrid semiconductor-superconductor structures sees rapid progress, our predictions may soon be verified in the laboratory.

ACKNOWLEDGMENTS

We thank Karen Michaeli, Aviad Landau, and Eran Sela for a previous collaboration on multiterminal Majorana island and Rok Žitko for helpful remarks about NRG. This work was supported by the DOE Office of Basic Energy Sciences, Division of Materials Sciences and Engineering under award de-sc0010526.

APPENDIX A: NRG CALCULATION DETAILS

We perform the spectral function calculations in the framework of full density matrix NRG [82] using a complete basis set [83] in order to properly account for finite-temperature effects. We also perform sliding parameter averaging [84–86] over four values of sliding parameter z to remove spurious oscillations in the results. In the calculations of the s-wave island model, we use the discretization parameter $\Lambda = 5$ for a two-channel system and $\Lambda = 10$ for a three-channel system. To make the three-channel case numerically tractable, we use the interleaved Wilson chain scheme [81,87], which requires some fine-tuning of the individual tunnel couplings to obtain the critical behavior. The cutoff energy has been set to $E_{\rm cut} =$ 14 for two leads and $E_{\rm cut} = 8.5$ for the three-lead case, with the maximum number of kept states 15 000 and 40 000, respectively. In the Majorana island part, we used $\Lambda = 3$ and kept up to 5000 states in each iteration.

We calculate conductance in the framework of linearresponse theory using the Kubo formula. To obtain the final expression for conductance, we follow a similar procedure as in Appendix B of Galpin *et al.* [57]. Therefore, we study how a perturbation of the form $H'(t) = \hat{O}_m f(t)e^{\eta t}$ (with $\eta \to 0$ to account for turning on the perturbation adiabatically) changes the equilibrium expectation value of an operator \hat{O}_n . We use a standard result in the first order of perturbation theory to express the change by

$$\delta\langle \hat{O}_n(t)\rangle = -\frac{i}{\hbar} \operatorname{Tr} \int_{-\infty}^t e^{\eta t'} [\hat{O}_m, \rho_{\text{eq}}] \hat{O}_n(t-t') f(t') dt',$$
(A1)

where we define $\delta \langle \hat{O}_n(t) \rangle = \text{Tr}[\rho(t)\hat{O}_n - \rho_{eq}\hat{O}_n]$, and $\rho_{eq} = e^{-\beta H}/Z$ and $\rho(t)$ are the density matrices in equilibrium and in the presence of the perturbation, respectively, and $\hat{O}_n(t - t')$ is defined in the interaction picture:

$$\hat{O}_n(t-t') = e^{\frac{i}{\hbar}\hat{H}(t-t')}O_n e^{-\frac{i}{\hbar}\hat{H}(t-t')}.$$
(A2)

To obtain conductance using the formula (A1), we have to study how current through a lead I_j changes when ac voltage V_k is applied to another lead. Therefore, we make the following substitutions: $\hat{O}_n \rightarrow I_j = e\langle \dot{N}_j \rangle = e\langle \frac{i}{\hbar}[H, N_j] \rangle$, $\hat{O}_n \rightarrow$ N_k , and $f(t) \rightarrow eV_k \cos(\omega t)$. This leads to a formula for the current present in the perturbed system:

$$I_j(t) = -\frac{ie^2 V_k}{\hbar} \operatorname{Tr} \int_{-\infty}^t e^{\eta t'} [N_k, \rho_{eq}] \dot{N}_j(t-t') \cos(\omega t') dt'.$$
(A3)

$$\mathcal{G}_{jk}(t,\omega) = \frac{\partial I_j}{\partial V_k}$$

= $-\frac{ie^2}{\hbar} \operatorname{Tr} \int_0^\infty e^{\eta(t-t'')} [N_k, \rho_{eq}] \dot{N}_j(t'')$
 $\times \cos[\omega(t-t'')] dt''.$ (A4)

To simplify the considerations, we focus on the value of conductance at t = 0. Using the cyclic property of a trace, we arrive at

$$\mathcal{G}_{jk}(t=0,\omega)$$

$$= -\frac{ie^2}{\hbar} \operatorname{Tr} \int_0^\infty e^{-\eta t''} \rho_{\text{eq}}[N_k, \dot{N}_j(t'')] \cos(\omega t'') dt''$$

$$= -\frac{ie^2}{2\hbar} \int_0^\infty e^{-\eta t''} \langle [N_k, \dot{N}_j(t'')] \rangle (e^{i\omega t''} + e^{-i\omega t''}) dt''.$$
(A5)

Now we insert the complete basis of energy states and compute the conductance using the Lehmann spectral representation. We finally arrive at

$$\mathcal{G}_{jk}(\omega) = \frac{e^2}{2\hbar} [\sigma_{jk}(\omega) + \sigma_{jk}(-\omega)]$$
(A6)

with

$$\sigma_{jk}(\omega) = \frac{1}{Z} \sum_{m,n} \frac{E_n - E_m + \omega - i\eta}{(E_n - E_m + \omega)^2 + \eta^2} (e^{-\beta E_m} - e^{-\beta E_n}) \times \langle m | N_k | n \rangle \langle n | \dot{N}_j | m \rangle.$$
(A7)

During the NRG simulation, we compute the imaginary part of $\sigma_{jk}(\omega)$. The real part can be obtained afterward by performing a Kramers-Kronig transformation. The quantity $G_{jk}(\omega)$ we show in the figures is

$$G_{jk}(\omega) = \operatorname{Im} \mathcal{G}_{jk}(\omega) = \frac{e^2}{h} \pi [\operatorname{Im} \sigma_{jk}(\omega) + \operatorname{Im} \sigma_{jk}(-\omega)].$$
(A8)

PHYSICAL REVIEW B 99, 014512 (2019)

The advantage of the method presented above is apparent when one calculates the dc conductance as a limit $\omega \rightarrow 0$. The usual approach is to compute

$$G_{jkdc} = -2\pi \lim_{\omega \to 0} \frac{\operatorname{Im} K(\omega)}{\omega}, \qquad (A9)$$

where

$$K(\omega) = -\frac{i}{\hbar} \int_0^\infty e^{i(\omega+i\eta)t} \langle [\dot{N}_j, \dot{N}_k(t)] \rangle dt.$$
(A10)

This approach involves calculation of a limit of a ratio of two very small quantities, which may prove to be unreliable numerically. The tradeoff of the method we used is that it requires computation of global operators N_j , which depend not only on the impurity, but also on the sites of the Wilson chains [88].

APPENDIX B: ADDITIONAL MAJORANA ISLAND NRG RESULTS

In this Appendix, we present additional NRG simulation results for the Majorana island. We begin with the details of the model to which we are comparing our results. The model describes the topological superconductor island in the Coulomb valley regime. The Hamiltonian in this case is [57]

$$\hat{H} = \hat{H}_{\text{leads}} + \frac{t_1}{\sqrt{2}} (\sigma^+ \psi_0^{\dagger} \psi_1 + \sigma^- \psi_1^{\dagger} \psi_0) + \frac{t_2}{\sqrt{2}} (\sigma^+ \psi_{-1}^{\dagger} \psi_0 + \sigma^- \psi_0^{\dagger} \psi_{-1}) + t_3 \sigma_z (\psi_1^{\dagger} \psi_1 - \psi_{-1}^{\dagger} \psi_{-1}), \qquad (B1)$$

where ψ_j are annihilation operators at the ends of the three spinless leads, and σ are the spin operators of the impurity formed on the island. The Hamiltonian is obtained by considering virtual transitions between leads in second-order perturbation theory. This results in a much stronger crossover energy scale dependence on the tunnel couplings and is one of the reasons for the many orders of magnitude difference



FIG. 9. (a) Entropy of the island's impurity $S_{imp}(T)$ for temperatures expressed in units of lead bandwidth *D* for several tunneling coupling strengths. Entropy flows from $S_{imp}(T) = \ln(4)$ for high temperatures to the non-Fermi-liquid fixed point with $S_{imp}(T) = \ln(\sqrt{3})$. (b) $G_{13}(\omega)$ conductance matrix element in the fully anisotropic case $t_1 = 0.095D$, $t_2 = 0.105D$, and t_3 from the interval $[0.125t_1, 4t_1]$ with each curve increasing t_3 by a factor of 2. (c) $G_{13dc}(t_3)$ dc conductance slightly away from the charge degeneracy point ($\Delta_g = 0.0035$) as a function of tunnel coupling of the third lead for several temperatures. The conductance has a nonmonotonic dependence, peaked at $t_3 \approx 0.05D = t_1 = t_2$. As the system is cooled down, the value at the peak increases until it reaches $2/3 e^2/h$. Further lowering the temperature develops a plateau at this value.



FIG. 10. Conductance matrix element (a) $G_{12}(\omega)$ and (b) $G_{13}(\omega)$ when a Majorana hybridization term is added to the Hamiltonian with $K_x = K_y = 0$, $K_z \neq 0$. (c) $G_{12}(\omega)$ conductance matrix element when a Majorana hybridization term is added to the Hamiltonian with $K_x = K_y = K_z = K$. In this case, $G_{12}(\omega) = G_{13}(\omega)$.

between the transition temperature at the charge degeneracy point and in the topological Kondo regime.

In Fig. 9(a) we show the entropy $S_{imp}(T)$ curves with temperature expressed in units of the lead bandwidth D, before collapsing all of them onto one curve as shown in the main text. In Fig. 9(b) we show the conductance matrix element $G_{13}(\omega)$ for several values of the tunnel coupling of the third lead t_3 [complementary plot to Fig. 5(c) from the main text]. In this case, one can also observe the transition to two-terminal teleportation: for $t_3 < t_1, t_2$, the conductance reaches the fractional quantized value of $2/3 \ e^2/h$ only for very low frequencies and analogously very low temperatures. For higher temperatures, the conductance is essentially 0 [in the same regime $G_{12}(\omega)$ is close to $1 \ e^2/h$]. In Fig. 9(c) we show the dc conductance $G_{13dc}(t_3)$ for several different temperatures [complementary plot to Fig. 6(a) from the main text]. In this case, the conductance forms a peak with the

- [1] A. Y. Kitaev, Ann. Phys. 303, 2 (2003).
- [2] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. 80, 1083 (2008).
- [3] C. Beenakker, Annu. Rev. Condens. Matter Phys. 4, 113 (2013).
- [4] J. Alicea, Rep. Prog. Phys. 75, 076501 (2012).
- [5] S. D. Sarma, M. Freedman, and C. Nayak, npj Quant. Info. 1, 15001 (2015).
- [6] R. M. Lutchyn, J. D. Sau, and S. Das Sarma, Phys. Rev. Lett. 105, 077001 (2010).
- [7] Y. Oreg, G. Refael, and F. von Oppen, Phys. Rev. Lett. 105, 177002 (2010).
- [8] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, Science 336, 1003 (2012).
- [9] A. Das, Y. Ronen, Y. Most, Y. Oreg, M. Heiblum, and H. Shtrikman, Nat. Phys. 8, 887 (2012).
- [10] L. P. Rokhinson, X. Liu, and J. K. Furdyna, Nat. Phys. 8, 795 (2012).
- [11] H. O. H. Churchill, V. Fatemi, K. Grove-Rasmussen, M. T. Deng, P. Caroff, H. Q. Xu, and C. M. Marcus, Phys. Rev. B 87, 241401(R) (2013).
- [12] A. D. K. Finck, D. J. Van Harlingen, P. K. Mohseni, K. Jung, and X. Li, Phys. Rev. Lett. 110, 126406 (2013).

maximum for t_3 close to the isotropic case. When the temperature is decreased, at first the height of the peak increases, but when it reaches $2/3 e^2/h$ the increase stops and instead a plateau is developed. This can also serve as an experimental signature of multiterminal electron teleportation.

In Figs. 10(a) and 10(b) we present the results of calculations with a hybridization term that includes only the *z* component of \vec{K} . Since $K_z \sigma_z \sim i \gamma_1 \gamma_2$, this term connects Majorana states coupled to leads 1 and 2, effectively decoupling the third lead. This in turn gives $1 e^2/h$ conductance at very low temperatures, the same as in the case of the two-terminal electron teleportation. At the same time, conductance $G_{13}(\omega)$ drops to 0 as a result of this decoupling. When the components of \vec{K} are all equal, the conductance is the same in the case of both $G_{12}(\omega)$ and $G_{13}(\omega)$ and is similarly equal to $2/3 e^2/h$ before decreasing to some nonuniversal value between 0 and 2/3.

- [13] S. M. Albrecht, A. P. Higginbotham, M. Madsen, F. Kuemmeth, T. S. Jespersen, J. Nygård, P. Krogstrup, and C. M. Marcus, Nature (London) 531, 206 (2016).
- [14] P. Krogstrup, N. L. B. Ziino, W. Chang, S. M. Albrecht, M. H. Madsen, E. Johnson, J. Nygård, C. M. Marcus, and T. S. Jespersen, Nat. Mater. 14, 400 (2015).
- [15] S. R. Plissard, I. van Weperen, D. Car, M. A. Verheijen, G. W. G. Immink, J. Kammhuber, L. J. Cornelissen, D. B. Szombati, A. Geresdi, S. M. Frolov, L. P. Kouwenhoven, and E. P. A. M. Bakkers, Nat. Nanotech. 8, 859 (2013).
- [16] W. Chang, S. M. Albrecht, T. S. Jespersen, F. Kuemmeth, P. Krogstrup, J. Nygård, and C. M. Marcus, Nat. Nanotech. 10, 232 (2015).
- [17] A. P. Higginbotham, S. M. Albrecht, G. Kiršanskas, W. Chang, F. Kuemmeth, P. Krogstrup, T. S. Jespersen, J. Nygård, K. Flensberg, and C. M. Marcus, Nat. Phys. 11, 1017 (2015).
- [18] O. Gül, H. Zhang, J. D. S. Bommer, M. W. A. de Moor, D. Car, S. R. Plissard, E. P. A. M. Bakkers, A. Geresdi, K. Watanabe, T. Taniguchi, and L. P. Kouwenhoven, Nat. Nanotech. 13, 192 (2018).
- [19] M. T. Deng, S. Vaitiekėnas, E. B. Hansen, J. Danon, M. Leijnse, K. Flensberg, J. Nygård, P. Krogstrup, and C. M. Marcus, Science 354, 1557 (2016).

- [20] M. Taupin, E. Mannila, P. Krogstrup, V. F. Maisi, H. Nguyen, S. M. Albrecht, J. Nygård, C. M. Marcus, and J. P. Pekola, Phys. Rev. Appl. 6, 054017 (2016).
- [21] D. Sherman, J. S. Yodh, S. M. Albrecht, J. Nygård, P. Krogstrup, and C. M. Marcus, Nat. Nanotech. 12, 212 (2017).
- [22] J. Chen, P. Yu, J. Stenger, M. Hocevar, D. Car, S. R. Plissard, E. P. A. M. Bakkers, T. D. Stanescu, and S. M. Frolov, Sci. Adv. 3, e1701476 (2017).
- [23] O. Gül, H. Zhang, F. K. de Vries, J. van Veen, K. Zuo, V. Mourik, S. Conesa-Boj, M. P. Nowak, D. J. van Woerkom, M. Quintero-Pérez, M. C. Cassidy, A. Geresdi, S. Koelling, D. Car, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, Nano Lett. 17, 2690 (2017).
- [24] H. Zhang, C.-X. Liu, S. Gazibegovic, D. Xu, J. A. Logan, G. Wang, N. van Loo, J. D. S. Bommer, M. W. A. de Moor, D. Car, R. L. M. O. het Veld, P. J. van Veldhoven, S. Koelling, M. A. Verheijen, M. Pendharkar, D. J. Pennachio, B. Shojaei, J. S. Lee, C. J. Palmstrøm, E. P. A. M. Bakkers, S. D. Sarma, and L. P. Kouwenhoven, Nature (London) 556, 74 (2018).
- [25] M. T. Tuominen, J. M. Hergenrother, T. S. Tighe, and M. Tinkham, Phys. Rev. Lett. 69, 1997 (1992).
- [26] P. Lafarge, P. Joyez, D. Esteve, C. Urbina, and M. H. Devoret, Phys. Rev. Lett. **70**, 994 (1993).
- [27] T. M. Eiles, J. M. Martinis, and M. H. Devoret, Phys. Rev. Lett. 70, 1862 (1993).
- [28] D. V. Averin and Y. V. Nazarov, Phys. Rev. Lett. 69, 1993 (1992).
- [29] L. Fu, Phys. Rev. Lett. 104, 056402 (2010).
- [30] B. van Heck, R. M. Lutchyn, and L. I. Glazman, Phys. Rev. B 93, 235431 (2016).
- [31] R. M. Lutchyn and L. I. Glazman, Phys. Rev. Lett. **119**, 057002 (2017).
- [32] P. Nozières and A. Blandin, J. Phys. (France) 41, 193 (1980).
- [33] D. L. Cox and A. Zawadowski, Adv. Phys. 47, 599 (1998).
- [34] K. Michaeli, L. A. Landau, E. Sela, and L. Fu, Phys. Rev. B 96, 205403 (2017).
- [35] L. Herviou, K. Le Hur, and C. Mora, Phys. Rev. B 94, 235102 (2016).
- [36] H. Yi and C. L. Kane, Phys. Rev. B 57, R5579(R) (1998).
- [37] H. Yi, Phys. Rev. B 65, 195101 (2002).
- [38] Z.-q. Bao and F. Zhang, Phys. Rev. Lett. 119, 187701 (2017).
- [39] M. Pustilnik, B. van Heck, R. M. Lutchyn, and L. I. Glazman, Phys. Rev. Lett. **119**, 116802 (2017).
- [40] L. I. Glazman and M. E. Raikh, JETP Lett. 47, 452 (1988).
- [41] T. K. Ng and P. A. Lee, Phys. Rev. Lett. 61, 1768 (1988).
- [42] A. Kawabata, J. Phys. Soc. Jpn. 60, 3222 (1991).
- [43] D. Goldhaber-Gordon, H. Shtrikman, D. Mahalu, D. Abusch-Magder, U. Meirav, and M. A. Kastner, Nature (London) 391, 156 (1998).
- [44] S. M. Cronenwett, T. H. Oosterkamp, and L. P. Kouwenhoven, Science 281, 540 (1998).
- [45] K. A. Matveev, JETP 72, 892 (1991).
- [46] K. A. Matveev, Phys. Rev. B 51, 1743 (1995).
- [47] A. Furusaki and K. A. Matveev, Phys. Rev. B 52, 16676 (1995).
- [48] Y. Oreg and D. Goldhaber-Gordon, Phys. Rev. Lett. 90, 136602 (2003).

- [49] M. Pustilnik, L. Borda, L. I. Glazman, and J. von Delft, Phys. Rev. B 69, 115316 (2004).
- [50] Z. Iftikhar, S. Jezouin, A. Anthore, U. Gennser, F. D. Parmentier, A. Cavanna, and F. Pierre, Nature (London) 526, 233 (2015).
- [51] Z. Iftikhar, A. Anthore, A. K. Mitchell, F. D. Parmentier, U. Gennser, A. Ouerghi, A. Cavanna, C. Mora, P. Simon, and F. Pierre, Science 360, 1315 (2018).
- [52] R. M. Potok, I. G. Rau, H. Shtrikman, Y. Oreg, and D. Goldhaber-Gordon, Nature (London) 446, 167 (2007).
- [53] H. T. Mebrahtu, I. V. Borzenets, D. E. Liu, H. Zheng, Y. V. Bomze, A. I. Smirnov, H. U. Baranger, and G. Finkelstein, Nature (London) 488, 61 (2012).
- [54] A. J. Keller, L. Peeters, C. P. Moca, I. Weymann, D. Mahalu, V. Umansky, G. Zaránd, and D. Goldhaber-Gordon, Nature (London) 526, 237 (2015).
- [55] B. Béri and N. R. Cooper, Phys. Rev. Lett. 109, 156803 (2012).
- [56] A. Altland and R. Egger, Phys. Rev. Lett. 110, 196401 (2013).
- [57] M. R. Galpin, A. K. Mitchell, J. Temaismithi, D. E. Logan, B. Béri, and N. R. Cooper, Phys. Rev. B 89, 045143 (2014).
- [58] I. Affleck and D. Giuliano, J. Stat. Mech. (2013) P06011.
- [59] A. Altland, B. Béri, R. Egger, and A. M. Tsvelik, J. Phys. A 47, 265001 (2014).
- [60] A. Altland, B. Béri, R. Egger, and A. M. Tsvelik, Phys. Rev. Lett. 113, 076401 (2014).
- [61] A. Zazunov, A. Altland, and R. Egger, New J. Phys. 16, 015010 (2014).
- [62] E. Eriksson, A. Nava, C. Mora, and R. Egger, Phys. Rev. B 90, 245417 (2014).
- [63] E. Eriksson, C. Mora, A. Zazunov, and R. Egger, Phys. Rev. Lett. 113, 076404 (2014).
- [64] O. Kashuba and C. Timm, Phys. Rev. Lett. 114, 116801 (2015).
- [65] D. I. Pikulin, Y. Komijani, and I. Affleck, Phys. Rev. B 93, 205430 (2016).
- [66] D. Meidan, A. Romito, and P. W. Brouwer, Phys. Rev. B 93, 125433 (2016).
- [67] S. Plugge, A. Zazunov, E. Eriksson, A. M. Tsvelik, and R. Egger, Phys. Rev. B 93, 104524 (2016).
- [68] K. T. Law, P. A. Lee, and T. K. Ng, Phys. Rev. Lett. 103, 237001 (2009).
- [69] A. Haim, E. Berg, F. von Oppen, and Y. Oreg, Phys. Rev. Lett. 114, 166406 (2015).
- [70] T. Giamarchi, *Quantum Physics in One Dimension*, International Series of Monographs on Physics (Oxford University Press, Oxford, 2004).
- [71] B. Béri, Phys. Rev. Lett. **110**, 216803 (2013).
- [72] I. Affleck and A. W. W. Ludwig, Nucl. Phys. B 360, 641 (1991).
- [73] I. Affleck and A. W. W. Ludwig, Phys. Rev. B 48, 7297 (1993).
- [74] A. K. Mitchell, L. A. Landau, L. Fritz, and E. Sela, Phys. Rev. Lett. 116, 157202 (2016).
- [75] L. A. Landau, E. Cornfeld, and E. Sela, Phys. Rev. Lett. 120, 186801 (2018).
- [76] I. Affleck, A. W. W. Ludwig, H.-B. Pang, and D. L. Cox, Phys. Rev. B 45, 7918 (1992).
- [77] M. Fabrizio, A. O. Gogolin, and P. Nozières, Phys. Rev. Lett. 74, 4503 (1995).
- [78] R. Bulla, T. A. Costi, and T. Pruschke, Rev. Mod. Phys. 80, 395 (2008).
- [79] R. Žitko, Comput. Phys. Commun. 182, 2259 (2011).

- [80] V. J. Emery and S. Kivelson, Phys. Rev. B 46, 10812 (1992).
- [81] A. K. Mitchell, M. R. Galpin, S. Wilson-Fletcher, D. E. Logan, and R. Bulla, Phys. Rev. B 89, 121105(R) (2014).
- [82] A. Weichselbaum and J. von Delft, Phys. Rev. Lett. 99, 076402 (2007).
- [83] F. B. Anders and A. Schiller, Phys. Rev. Lett. 95, 196801 (2005).
- [84] W. C. Oliveira and L. N. Oliveira, Phys. Rev. B 49, 11986 (1994).
- [85] R. Žitko and T. Pruschke, Phys. Rev. B 79, 085106 (2009).
- [86] R. Žitko, Comput. Phys. Commun. 180, 1271 (2009).
- [87] K. M. Stadler, A. K. Mitchell, J. von Delft, and A. Weichselbaum, Phys. Rev. B 93, 235101 (2016).
- [88] R. Žitko and T. Pruschke, New J. Phys. 12, 063040 (2010).