

Interaction-induced long-time tail of a nonlinear ac absorption in a localized system: A relay-race mechanism

Rajesh K. Malla and M. E. Raikh

Department of Physics and Astronomy, University of Utah, Salt Lake City, Utah 84112, USA

(Received 20 March 2019; revised manuscript received 3 May 2019; published 15 May 2019)

In conventional solid-state electron systems with localized states the ac absorption is linear since the inelastic widths of the energy levels exceed the drive amplitude. The situation is different in the systems of cold atoms in which phonons are absent. Then even a weak drive leads to saturation of the ac absorption within resonant pairs, so that the population of levels oscillates with the Rabi frequency. We demonstrate that, in the presence of weak dipole-dipole interactions, the response of the system acquires a long-time component which oscillates with frequency much smaller than the Rabi frequency. The underlying mechanism of this long-time behavior is that the fields created in the course of the Rabi oscillations serve as resonant drive for the *second-generation* Rabi oscillations in pairs with level spacings close to the Rabi frequency. The frequency of the second-generation oscillations is of the order of interaction strength. As these oscillations develop, they can initiate the next-generation Rabi oscillations, and so on. Formation of the second-generation oscillations is facilitated by the nondiagonal component of the dipole-dipole interaction tensor.

DOI: [10.1103/PhysRevB.99.195427](https://doi.org/10.1103/PhysRevB.99.195427)

I. INTRODUCTION

A transparent physical picture of absorption of the ac electric field in a system with localized electron states was proposed by Mott [1]. According to Mott, absorption takes place within pairs of states with energy spacing $\hbar\omega$, where ω is the driving frequency. Frequency dependence of the ac conductivity within this picture is $\sigma(\omega) \propto \omega^2 \ln^2 \omega$ (in one dimension), where one power of ω comes from the photon energy, while the other comes from the restriction that the pair is singly occupied. Finally, $\ln \omega$ comes from the overlap integral between initial and final states. Later, Mott's formula was rigorously derived from the Kubo linear-response formalism by Berezinskii [2].

The condition of applicability of the linear response is that inelastic widths of the localized levels are much bigger than the absorption matrix element. This condition is satisfied in conventional solid-state systems where inelastic widths are due to the phonon emission.

Regime of strong ac drive, opposite to the linear response, can be realized in cold-atom systems, where phonons are absent. The ac drive in these systems is implemented by the synchronous modulation of the intensity of laser beams which create a quasirandom one-dimensional (1D) on-site energy profile [3].

Possibility to realize the regime of strong drive in a localized system without thermal bath raises a number of conceptual questions which, with rare exceptions [4–6], were not addressed in earlier studies. These questions can be conventionally divided into three groups:

(i) On the single-particle level [7–11], the fundamental question is: does the localization persist in the presence of strong drive, when electron states evolve into the Floquet eigenstates? Anderson localization is the result of interference of the backscattering amplitudes in the course of

multiple scattering [2]. Floquet states can be viewed as a combination of satellites with energies separated by $n\hbar\omega$. Development of satellites upon increasing drive leads to the new channels of interference, and thus suppresses the localization, like in multichannel wires.

(ii) Another physical mechanism relevant for nonlinear ac response of localized noninteracting systems is the adiabatic Landau-Zener transitions [9,11]. This mechanism comes into play when the drive is strong and slow. In this limit, the effect of drive can be viewed as periodic modulation of energies of the localized states [12–14]. It was proposed in Refs. [12–14] that the effective absorption in this limit can be captured within the random-matrix description. As the levels corresponding to neighboring states slowly pass by each other, an electron can adiabatically change the level. This, in turn, can lead to the long-time component of the ac absorption [15]. Spreading of the electron due to the level crossings illustrates the tendency of drive to suppress the localization.

(iii) The third group of papers is the most numerous, see, e.g., Refs. [15–22], and addresses the physics of ac driven localized interacting systems. They are focused on the dynamics of heating and on the long-times properties of nonequilibrium state. In particular, the question of interest is whether or not the long-time behavior of an interacting many-body system is sensitive to its initial state.

When the drive amplitude is much smaller than the drive frequency ω , resonant pairs get saturated after the time of the order of the inverse Rabi period. Higher harmonics in the pair dynamics are small in this regime [23]. Landau-Zener transitions also do not take place when the drive is fast. It is argued in Ref. [15] that long-time dynamics in this limit is due to interaction between the pairs. Namely, a group of n interacting pairs can be engaged into collective Rabi




FIG. 1. Schematic illustration of the relay-race mechanism: resonant drive with frequency ω engages spin a into the Rabi oscillations. As a result, the Zeeman levels of a get split by Ω_a , the Rabi frequency. If Ω_a is close to the Zeeman splitting of spin b , then the Rabi oscillations of a will serve as a resonant drive for b via the nondiagonal component of the dipole-dipole interaction D_{ab} . If, in turn, the Rabi frequency Ω_b of the second-generation Rabi oscillations is close to the Zeeman splitting of spin c , the third-generation Rabi oscillations are initiated as a result of dipole-dipole interaction D_{bc} , and so on.

oscillations, whose frequency is proportional to n th power of drive.

In the present paper we propose an alternative mechanism of long-time dynamics in a system of weakly interacting pairs under a weak drive. Namely, a saturated pair, executing the Rabi oscillations, creates a field which plays the role of drive for a distant pair, thus causing the *second-generation* Rabi oscillations (Fig. 1). At resonance, the level spacing of the second-generation pair is equal to the Rabi frequency of the first-generation pair. If this frequency is much smaller than the pair-pair interaction, then the second-generation Rabi oscillations are slow. We perform statistical averaging analytically and find the slow component of the absorption.

II. DYNAMICS OF TWO INTERACTING DRIVEN SPINS

To illustrate the proposed mechanism we employ the simplest model. Namely, as is common in the literature, we employ the language of spins to describe two-level systems and, correspondingly, the ac magnetic field to describe the drive. Consider two spins a and b subject to magnetic fields B_a and B_b , respectively. Since the drive amplitude B_1 is much smaller than ω , the rotating-wave approximation applies. Then the Hamiltonian of the pair reads

$$\hat{H} = \frac{B_a}{2} S_z^a + \frac{B_b}{2} S_z^b + 2B_1(S_x^a + S_x^b) \cos \omega t + 2B_1(S_y^a + S_y^b) \sin \omega t - D(r) \left[\frac{3(\mathbf{S}^a \cdot \mathbf{r})(\mathbf{S}^b \cdot \mathbf{r}) - (\mathbf{S}^a \cdot \mathbf{S}^b)r^2}{r^2} \right], \quad (1)$$

where we have incorporated the dipole-dipole interaction of a magnitude,

$$D(r) = D_0 \left(\frac{a}{r} \right)^3, \quad (2)$$

with a being the distance between two neighboring spins, while \mathbf{r} is the vector distance between spin a and spin b . In the following we adopt the convention that all magnetic fields and the dipole-dipole interactions are measured in the units of frequency.

In the absence of interaction, only spin a is in resonance with the drive $|\omega - B_a| \ll \omega$, while spin b is off-resonance $B_b \ll \omega$, and does not respond to the drive. Components S_x^a and S_y^a of the driven spin a oscillate with frequencies close

to ω . Thus, the fields produced by these components on spin b via dipole-dipole interaction do not induce the dynamics of b . On the other hand, the z component of spin a oscillates with much smaller frequency

$$\Omega_a = [B_1^2 + (\omega - B_a)^2]^{1/2}. \quad (3)$$

These slow oscillations of S_z^a translate into the field acting on b . The field generated by the z - z component of the interaction is also inefficient, since spin b is already directed along z . Spin b can be set in motion via the z - x and z - y components of the dipole-dipole interaction when Ω_a is close to B_b . This is why we will keep only the z - x component.

Summarizing, the relevant components of the field acting on a are $(B_1 \cos \omega t, B_1 \sin \omega t, B_a + DS_x^b)$, while the field acting on b has an x component, equal to DS_z^a , and z component B_b .

The equations of motion for the projections of a which follow from $\frac{d\mathbf{S}}{dt} = \mathbf{B} \times \mathbf{S}$, read

$$\frac{dS_x^a}{dt} = B_1 \sin \omega t S_z^a - (B_a + DS_x^b) S_y^a, \quad (4)$$

$$\frac{dS_y^a}{dt} = -B_1 \cos \omega t S_z^a + (B_a + DS_x^b) S_x^a, \quad (5)$$

$$\frac{dS_z^a}{dt} = B_1 \cos \omega t S_y^a - B_1 \sin \omega t S_x^a, \quad (6)$$

while the equations of motion for the components of spin b have the form

$$\frac{dS_x^b}{dt} = -B_b S_y^b, \quad \frac{dS_y^b}{dt} = -DS_z^a S_z^b + B_b S_x^b, \quad (7)$$

$$\frac{dS_z^b}{dt} = DS_z^a S_y^b. \quad (8)$$

To analyze the coupled equations of motion for a and b it is convenient to cast them into the integral form. First, we express S_x^a and S_y^a in terms of S_z^a and S_x^b . Substituting the result into the equation for S_z^a and taking into account the initial condition $S_z^a(0) = 1$, we get

$$\frac{dS_z^a}{dt} = -B_1^2 \int_0^t dt' S_z^a(t') \cos \left[(\omega - B_a)(t - t') - D \int_{t'}^t dt'' S_x^b(t'') \right]. \quad (9)$$




FIG. 2. Numerical example illustrating the formation of the long-time component of the ac response. First generation of the Rabi oscillations (blue) is plotted from (12) for the drive frequency $\omega = 2B_1$ and the Zeeman splitting $B_a = 1.8B_1$. Assuming that the Zeeman energies are homogeneously distributed between 0 and 1.5ω , the ensemble-averaged $\langle S_z(t) \rangle$ is calculated from (34) and is plotted with black. Second-generation Rabi oscillations are shown with red. They are calculated from (15) and (16) in which we chose the Zeeman energy of spin b to be $B_b = 0.5B_1$ and the magnitude of interaction to be $D_0 = 0.05B_1$.

Similarly, we express S_x^b and S_y^b in terms of S_z^a and S_z^b and, using $S_z^b(0) = 1$, substitute them into the equation for S_z^b . This yields

$$\frac{dS_z^b}{dt} = -D^2 \int_0^t dt' S_z^b(t') [S_z^a(t) S_z^a(t')] \cos B_b(t - t'). \quad (10)$$

To get the closed system, we also invoke the expression for S_x^b obtained in the course of solving the system (7),

$$S_x^b(t) = -\frac{D}{B_b} (1 - \cos B_b t) + D \int_0^t dt' S_z^a(t') S_z^b(t') \sin B_b(t - t'). \quad (11)$$

Three equations (9), (10), and (11) describe fully the dynamics of both spins.

For $D = 0$, spin b points along z , while spin a executes the Rabi nutations. In course of these nutations $S_z^a(t)$ follows the seminal Rabi formula

$$S_z^a(t) = \frac{(\omega - B_a)^2}{\Omega_a^2} + \frac{B_1^2}{\Omega_a^2} \cos \Omega_a t, \quad (12)$$

which also follows from Eq. (9).

For a finite D spin b is also set into motion. This motion causes a “feedback” on spin a , reflected by the term proportional to D in the argument of cosine. Most importantly, comparison of (9) and (10) quantifies our main message that the motion of spin a plays the role of drive for spin b ; the role of the driving field is played by $DS_z^a(t)$.

For spin a the resonant drive corresponds to the frequency $\omega = B_a$. Under this condition, S_z^a oscillates with frequency B_1 . Thus, for spin b , the resonant condition is $B_1 = B_b$. We also expect that, within a factor, the nutation frequency of spin b at resonance is equal to D , as illustrated in Fig. 2. Note that the nutation frequency D of the second-generation Rabi oscillations does not depend on B_1 . However, this is

valid only at exact resonance. We will see below that, for any small deviation from resonance, both the amplitude and the frequency of the second-generation Rabi oscillations acquire the B_1 dependence. In particular, the amplitude vanishes in the limit $B_1 \rightarrow 0$.

Upon substituting (12) into (10), the product $S_z^a(t)S_z^a(t')$ assumes the form

$$S_z^a(t)S_z^a(t') = \frac{(\omega - B_a)^4}{\Omega_a^4} + \frac{B_1^4}{\Omega_a^4} \cos \Omega_a t \cos \Omega_a t' + \frac{(\omega - B_a)^2 B_1^2}{\Omega_a^4} (\cos \Omega_a t + \cos \Omega_a t'). \quad (13)$$

The second term of (13) generates the sum $\cos \Omega_a(t + t') + \cos \Omega_a(t - t')$. It is the second cosine that acts as a resonant drive for spin b . Keeping only $\cos \Omega_a(t - t')$ term in (10), we get

$$\frac{dS_z^b}{dt} = -\frac{D^2 B_1^4}{4\Omega_a^4} \int_0^t dt' S_z^b(t') \cos[(\Omega_a - B_b)(t - t')]. \quad (14)$$

This equation has a solution

$$S_z^b = \frac{(\Omega_a - B_b)^2}{\Omega_b^2} + \frac{B_2^2}{\Omega_b^2} \cos \Omega_b t, \quad (15)$$

where the “second-generation” drive and the second-generation Rabi frequency are defined as

$$B_2 = \frac{DB_1^2}{2\Omega_a^2}, \quad \Omega_b = [B_2^2 + (\Omega_a - B_b)^2]^{1/2}. \quad (16)$$

Equations (15) and (16) constitute the main result of the present paper. We analyze this result below.

III. STATISTICAL AVERAGING, LONG-TIME TAIL OF THE AC ABSORPTION

The energy absorbed is proportional to averaged $S_z^b(t)$. We should average $S_z^b(t)$ over B_a , B_b and over D , which is equivalent to averaging over distances to the neighbors. We notice that B_a enters into the $S_z^b(t)$ only in combination $(B_a - \omega)^2 + B_1^2$, so that B_a does not affect the result of averaging.

We now average $S_z^b(t)$, from Eq. (15), assuming the density of spin states g to be constant. The averaging amounts to the twofold integral

$$1 - \langle S_z^b(t) \rangle = g \int_a^\infty dr 4\pi r^2 \times \int_0^\infty dB_b \frac{\left(\frac{D(r)B_1^2}{4\Omega_a^2}\right)^2}{\left(\frac{D(r)B_1^2}{4\Omega_a^2}\right)^2 + (\Omega_a - B_b)^2} \times 2 \sin^2 \left\{ \left[\left(\frac{D(r)B_1^2}{4\Omega_a^2}\right)^2 + (\Omega_a - B_b)^2 \right]^{1/2} \frac{t}{2} \right\}, \quad (17)$$

where a is the distance to the neighboring spin. Angular averaging is not important, so we choose $D(r) = D_0 \left(\frac{a}{r}\right)^3$, as prescribed by (2).

To proceed further, it is convenient to introduce, instead of variables r and B_b , new variables u and v defined as

$$r = \left(\frac{D_0 B_1^2 a^3}{8 \Omega_a^2 u} t \right)^{1/3}, \quad B_b = \Omega_a - \frac{2v}{t}. \quad (18)$$

Then the integral assumes the form

$$1 - \langle S_z^b(t) \rangle = \frac{2\pi g a^3 D_0 B_1^2}{3 \Omega_a^2} \int_0^{D_0 t/2} du \times \int_{-\Omega_a t}^{\infty} dv \frac{\sin^2(u^2 + v^2)^{1/2}}{u^2 + v^2}. \quad (19)$$

In the long-time limit $D_0 t \gg 1$, $\Omega_a t \gg D_0 t$, we can replace $\Omega_a t$ with infinity in the lower limit of the v integral. In order to see the asymptotic behavior of $\langle S_z^b(t) \rangle$, we substitute v as $v = \frac{u}{\tan \psi}$. With new variable ψ the integral (19) can be rewritten as

$$1 - \langle S_z^b(t) \rangle_{3D} = \frac{4\pi g a^3 D_0 B_1^2}{3 \Omega_a^2} \int_0^{D_0 t/2} \frac{du}{u} \Phi(u), \quad (20)$$

with $\Phi(u)$ defined as

$$\Phi(u) = \int_0^{\pi/2} d\psi \sin^2\left(\frac{u}{\sin \psi}\right). \quad (21)$$

In one and two dimensions, the corresponding expressions for $1 - \langle S_z^b(t) \rangle$ are similar to (20) and read

$$1 - \langle S_z^b(t) \rangle_{1D} = \frac{4gaD_0}{3\left(\frac{D_0 t}{2}\right)^{2/3}} \left(\frac{B_1^2}{4\Omega_a^2}\right)^{1/3} \int_0^{D_0 t/2} \frac{du}{u^{1/3}} \Phi(u), \quad (22)$$

$$1 - \langle S_z^b(t) \rangle_{2D} = \frac{8\pi g a^2 D_0}{3\left(\frac{D_0 t}{2}\right)^{1/3}} \left(\frac{B_1^2}{4\Omega_a^2}\right)^{2/3} \int_0^{D_0 t/2} \frac{du}{u^{2/3}} \Phi(u). \quad (23)$$

The function $\Phi(u)$ in (21) can be calculated analytically in two limits. For small $u \ll 1$, only small $\psi \sim u$ contribute to the integral. This allows us to replace $\sin \psi$ by the argument and extend the integration to infinity. The resulting integral can be calculated explicitly, and one gets

$$\Phi(u)|_{u \ll 1} = \frac{\pi u}{2}. \quad (24)$$

For large $u \gg 1$, the typical argument of \sin^2 is big, so that \sin^2 can be replaced by $1/2$, leading to $\Phi(u)|_{u \gg 1} \approx \frac{\pi}{4}$. The leading u -dependent correction comes from the vicinity of $\psi = \frac{\pi}{2}$, and thus oscillates with u . The asymptote has the form

$$\Phi(u)|_{u \gg 1} = \frac{\pi}{4} - \left(\frac{\pi}{8}\right)^{1/2} \frac{\cos\left(2u + \frac{\pi}{4}\right)}{u^{1/2}}. \quad (25)$$

Using (24) and (25) we find the behavior of $\langle S_z^b(t) \rangle$ in three dimensions:

$$1 - \langle S_z^b(t) \rangle = \frac{8\pi g a^3 D_0 B_1^2}{\Omega_a^2} \times \begin{cases} D_0 t, & \frac{D_0 t}{2} \ll 1, \\ \frac{1}{2} \ln(D_0 t), & D_0 t \gg 1. \end{cases} \quad (26)$$

Note that, at $D_0 t \ll 1$, the average $\langle S_z^b(t) \rangle$ decreases *linearly* with time. This is despite the fact that, for any *given* spin b , the time deviation of S_z from $S_z = 1$ is quadratic. The reason is that, if one expands (17) at small t , then the integral over




FIG. 3. The average $1 - \langle S_z^b(t) \rangle$ is plotted versus dimensionless time $D_0 t$ from Eqs. (22), (23), and (20) corresponding to one, (red) two, (black), and three (blue) dimensions. The values of the coefficient C_d are $\frac{4gaD_0}{3} \left(\frac{B_1^2}{4\Omega_a^2}\right)^{1/3}$, $\frac{8\pi g a^2 D_0}{3} \left(\frac{B_1^2}{4\Omega_a^2}\right)^{2/3}$, and $\frac{4\pi g a^3 D_0 B_1^2}{3\Omega_a^2}$ for one, two, and three dimensions, respectively.

B_b will diverge. Overall, the characteristic timescale for the change of $\langle S_z^b(t) \rangle$ is D_0^{-1} .

In derivation of the expression for $S_z^b(t)$, we have already assumed that $D_0 \ll B_1$ when we neglected the feedback of b on a . Now we see that the same assumption insures that the evolution of the ensemble-averaged $S_z^b(t)$ is slow.

In Fig. 3 we show $\langle S_z^b(t) \rangle$ calculated numerically from Eq. (20). Logarithmic behavior is evident. One can also distinguish weak oscillations on the background of a log profile. These oscillations become more pronounced in lower dimensions. For example, in one dimension, Eq. (22) can be written in the following form:

$$1 - \langle S_z^b(t) \rangle = \frac{4gaD_0}{3\left(\frac{D_0 t}{2}\right)^{2/3}} \left(\frac{B_1^2}{4\Omega_a^2}\right)^{1/3} F(D_0 t), \quad (27)$$

where

$$F(D_0 t) = \int_0^{D_0 t/2} \frac{du}{u^{1/3}} \Phi(u). \quad (28)$$

If we substitute the leading asymptote $\Phi(u) = \frac{\pi}{4}$, we will find that $\langle S_z^b(t) \rangle$ is time independent. In order to capture the time dependence, we add and subtract $\pi/4$ from $\Phi(u)$. Then (28) takes the form

$$F(D_0 t) = \frac{3\pi}{8} \left(\frac{D_0 t}{2}\right)^{2/3} + \int_0^{D_0 t/2} \frac{du}{u^{1/3}} \left(\Phi(u) - \frac{\pi}{4}\right). \quad (29)$$

At long times, the first term describes the leading contribution, while the second term saturates. This saturation is accompanied by the oscillations. To establish the form of this oscillating correction, it is convenient to present the integral $\int_0^{D_0 t/2} du$ as the difference of integrals $\int_0^{\infty} du$ and $\int_{D_0 t/2}^{\infty} du$. Then, in the integral $\int_{D_0 t/2}^{\infty} du$ we can use the oscillating term from the large- u asymptote in (25). This generates the

following correction to $F(D_0t)$:

$$-\left(\frac{\pi}{8}\right)^{1/2} \frac{\sin\left(D_0t + \frac{\pi}{4}\right)}{2\left(\frac{D_0t}{2}\right)^{5/6}}.$$

Substituting this correction into (27) indicates that the amplitude of oscillations in $\langle S_z^b(t) \rangle$ falls off as $t^{-3/2}$. Numerical plots for $d = 1$ and $d = 2$ in Fig. 3 confirm the saturation of $\langle S_z^b(t) \rangle$ at long times, which is accompanied by slow-decaying oscillations. As follows from analytics and numerics, the small time behavior of $1 - \langle S_z^b(t) \rangle$ is linear in one and two dimensions as well.

IV. THE ROLE OF FEEDBACK

Unconventionally, we find that the coupling of two spins via dipole-dipole interaction is *unidirectional*: spin a drives spin b , while the feedback effect of b on a is negligible under the condition $D \ll B_1$. On the other hand, it is the domain $D \ll B_1$, which is of interest, since it is in this domain where the long-time tail of the ac absorption develops. If we take into account that spin b is dipole-dipole coupled to spin c , see Fig. 1, then b will drive c under the resonant condition, but with negligible feedback. This is why we identify this spin dynamics with relay race.

To estimate the effect of feedback, we take the expression for $S_z^a(t)S_z^b(t)$ obtained in the lowest order and substitute it into (11). The expression for $S_z^a(t)S_z^b(t)$ has the form similar to the expression for $S_z^a(t)S_z^a(t')$ in (13) with $t' = t$,

$$\begin{aligned} S_z^a(t)S_z^b(t) &= \frac{(\omega - B_a)^2(\Omega_a - B_b)^2}{\Omega_a^2\Omega_b^2} + \frac{B_1^2B_2^2}{\Omega_a^2\Omega_b^2} \cos \Omega_a t \cos \Omega_b t \\ &+ \frac{(\Omega_a - B_b)^2B_1^2}{\Omega_a^2\Omega_b^2} \cos \Omega_a t \\ &+ \frac{(\omega - B_a)^2B_2^2}{\Omega_a^2\Omega_b^2} \cos \Omega_b t. \end{aligned} \quad (30)$$

As seen from (11), S_x^b contains a “free precession” term $\frac{D}{B_b}(1 - \cos B_b t)$, and the “drive-induced” term. The role of the free precession term is the shift of resonance $\omega = B_a$. Indeed, substituting this term into the argument of cosine in (9), and assuming that $B_b t$ is big (or, equivalently, that $\Omega_a t$ is big), results in replacement of $(\omega - B_b)$ by $(\omega - B_b - \frac{D^2}{B_b})$, i.e., the corrected resonance condition is

$$\omega = B_b \left(1 + \frac{D^2}{B_b^2}\right). \quad (31)$$

Since the relevant value of B_b is the Rabi frequency B_1 , we conclude that the shift is relatively small under the condition $B_1 \gg D$, which coincides with the condition that the second-generation Rabi oscillations are slow.

We neglected $\cos B_b t$ in free precession term because it leads to the oscillating contribution $\frac{D^2}{B_b^2}(\sin B_b t - \sin B_b t')$ in the argument of cosine in (9). This oscillating contribution results in effective renormalization of the drive amplitude [24] $B_1 \rightarrow B_1 J_0(\frac{D^2}{B_b^2})$, where J_0 is the Bessel function. This renormalization is small by virtue of the same condition $D \ll B_b \sim B_1$.

We now turn to the effect of feedback from the drive-induced term. As we have established in the course of statistical averaging, the second-generation Rabi oscillations essentially saturate at times $\sim \frac{1}{D}$. On the other hand, we do not expect significant feedback at times smaller than the period of the first-generation Rabi oscillations. This simplifies our task by restricting the time to the interval

$$\frac{1}{B_1} < t < \frac{1}{D}.$$

For further simplification, we consider the most “dangerous” situation $\Omega_a = B_b$. Under this condition, spin b is resonantly driven, so that the expected feedback is the strongest. Setting $\Omega_a = B_b$ in (30) we find that the first and the third terms vanish. In the two remaining terms it is sufficient to set $\cos \Omega_b t = 1$, since Ω_b is of the order of D . Also, with $\Omega_a = B_b$, we have $\Omega_b = B_2$. After that, (30) simplifies to

$$S_z^a(t)S_z^b(t) = \frac{B_1^2}{\Omega_a^2} \cos \Omega_a t + \frac{(\omega - B_a)^2}{\Omega_a^2}, \quad (32)$$

which is nothing but simply $S_z^a(t)$. Still, the behavior of $S_x^b(t)$ emerging upon substitution of (32) into (11) is nontrivial due to the beating of $\cos \Omega_a t$ and $\sin B_b(t - t')$. This beating generates a contribution to S_x^b equal to

$$\frac{B_1^2}{2\Omega_a^2} D t \sin \Omega_a t.$$

We see that this contribution exceeds the free-precession contribution and at $t \sim 1/D$ becomes of the order of 1, which could be expected under the resonant condition $\Omega_a = B_b$. Our main point is that, even under this condition, the feedback of S_x^b on the first-generation Rabi oscillations remains small. Indeed, performing the integration $\int_{t'}^t dt''$ in the argument of cosine in (9) generates the correction to this argument equal to

$$\frac{B_1^2 D^2}{2\Omega_a^3} (t \cos \Omega_a t - t' \cos \Omega_a t').$$

While this correction grows with t' it does not exceed 1 as long as t' is smaller than $1/D$. Thus we conclude that when the drive exceeds the interaction magnitude, the feedback effect is negligible.

V. DISCUSSION

(i) In a driven system of noninteracting spins in a random magnetic field (random B_a), only resonant spins respond to the drive. With Rabi frequencies depending on B_a , Rabi oscillations of different spins average out, so that the average $\langle S_z^a(t) \rangle$ approaches a constant. We note that this approach is accompanied by slow-decaying oscillations. Indeed, for a given spin, the oscillating part $S_z^a(t)$ has the form

$$S_z^a(t) - \overline{S_z^a(t)} = \frac{B_1^2 \cos [B_1^2 + (\omega - B_a)^2]^{1/2} t}{B_1^2 + (\omega - B_a)^2}. \quad (33)$$

Assuming the homogeneous distribution of B_a in the interval $0 < B_a < \Delta$, the disorder-average of (33) has the form

$$\langle S_z^a(t) - \overline{S_z^a(t)} \rangle = \frac{1}{\Delta} \int_0^\Delta dB_a \frac{B_1^2 \cos [B_1^2 + (\omega - B_a)^2]^{1/2} t}{B_1^2 + (\omega - B_a)^2}. \quad (34)$$

At long times, $B_1 t \gg 1$, only the resonant spins contribute to the integral. This allows us to expand the argument of cosine as

$$[B_1^2 + (\omega - B_a)^2]^{1/2} t \approx B_1 t + \frac{(\omega - B_a)^2}{2B_1} t. \quad (35)$$

We see that the relevant domain of $(\omega - B_a)$ is $\sim (\frac{B_1}{t})^{1/2} \ll B_1$. This allows us to set $B_a = \omega$ in the denominator of (34) and to extend the integration domain over $(\omega - B_a)$ to $(-\infty, \infty)$. Performing the Gaussian integration, we get

$$\langle S_z^a(t) - \overline{S_z^a(t)} \rangle|_{B_1 t \gg 1} = \left(\frac{\pi B_1}{\Delta^2 t} \right)^{1/2} \cos \left(B_1 t + \frac{\pi}{4} \right). \quad (36)$$

In Fig. 2 the result of numerical calculation of $\langle S_z^a(t) \rangle$ for a certain parameter set is shown. Numerics confirms the presence of slow-decaying periodic oscillations in average $\langle S_z^a \rangle$. The amplitude of these oscillations of average $S_z(t)$ coming from the sparse resonant spins should be compared to the S_z^b coming from the typical second-generation Rabi oscillations (26). The reasonable choice of Δ is ω . Characteristic t in (26) is $\sim 1/D_0$. Then S_z^b is $\sim g a^3 D_0$, while the oscillating part of $\langle S_z^a(t) \rangle$ can be presented as $\sim \frac{B_1}{\omega} (\frac{D_0}{B_1})^{1/2}$, which is the product of two small parameters. On the other hand, the fact that we considered the interaction of spin a with only one spin b , requires that the product $g a^3 D_0$ is also small.

(ii) Spin b can induce even slower Rabi nutations in spin c , see Fig. 1. The corresponding Rabi frequency of these third-generation oscillations will be

$$\Omega_c = [B_3^2 + (\Omega_b - B_c)^2]^{1/2}, \quad (37)$$

where B_3 is given by

$$B_3 = \frac{D_{bc} B_2^2}{2\Omega_b^2} = \frac{D_{bc} D_{ab}^2 B_1^4}{8\Omega_a^4 \Omega_b^2}. \quad (38)$$

$$B_3 = \frac{D_{bc} (D_{ab})^2 B_1^4}{8(\omega - B_a)^4 (\omega - B_a - B_b)^2}. \quad (39)$$

If we are away from resonance at each step, then drive amplitude at n th step will be

$$B_n = \frac{D_{n,n-1} B_{n-1}^2}{2(\omega - \sum_{i=\{a,b,c,\dots,n-1 \text{ terms}\}} B_i)^2}. \quad (40)$$

By contrast, if we are at resonance in each step, then the drive amplitude for the n th step will depend only on the dipole-dipole interaction between n th spin and the $n-1$ th spin. For example, the drive amplitude for $n=3$ is $D_{bc}/2$, and the drive amplitude for $n=2$ is $D_{ab}/2$, see Fig. 1.

Suppose that spin a is not in resonance with the drive, $(\omega - B_a) > B_1$. Then the amplitude of the first-generation Rabi oscillations is small. Still, spin b can oscillate with big

amplitude ~ 1 , provided that B_b is equal to $\Omega_a \approx (\omega - B_a)$. At the same time, the frequency of the oscillations of spin b will be approximately $\frac{D_{ab} B_1^2}{2(\omega - B_a)^2}$, which is much smaller than D_{ab} .

(iii) It follows from Eq. (26) that, while the contribution of the second-generation Rabi oscillations to the absorption is a slow function of time, the magnitude of this slow component contains a small parameter $g a^3 D_0$. Thus, as the dipole-dipole interactions increases, the amplitude of the slow component grows linearly with D_0 . On the other hand, the characteristic time before they saturate drops as $1/D_0$.

(iv) For slow second-generation Rabi oscillations to develop the drive amplitude should be bigger than the interaction strength. On the other hand, the drive is assumed to be weak, $B_1 \ll \omega$, which, in classification of Ref. [11], corresponds to the linear absorption regime. Such a weak drive cannot affect the overall many-body localized regime [25–29].

(v) In spirit, the relay-race mechanism considered in the present paper bares some similarity to the mechanism of delocalization of eigenmodes of dipole-dipole coupled oscillators or of the ensemble of two-level systems [30–33]. In Refs. [30–33] two undriven oscillators or two spins get hybridized when the corresponding frequencies match each other within the interaction magnitude. This hybridization can be mediated by z - z component of the interaction. In our notations, the frequencies of two hybridized oscillators can be expressed as

$$\omega^2 = \frac{B_a^2 + B_b^2}{2} \pm \left[\frac{(B_a^2 - B_b^2)}{4} + D_{ab}^2 \right]^{1/2}. \quad (41)$$

We see that, even at resonance $B_a = B_b$, hybridization does not result in a slow motion. By contrast, in our situation, the resonance is dictated by drive and hybridization takes place when Ω_a is close to B_b . In other words, the motion of a in the *rotated* frame is in resonance with b in the laboratory frame.

(vi) We introduced the relay-race mechanism using the language of spins driven by ac magnetic field. In Refs. [9,11,15] the ac absorption of electric field by localized electrons has been studied. The main difference between the two scenarios is that we considered the fields B_a to be random, but parallel to z . In the case of the ac electric field $\mathcal{E} \cos \omega t$, the Hamiltonian describing the drive has the form $\mathbf{P} \cdot \mathcal{E} \cos \omega t$, where \mathbf{P} is the dipole matrix element between the ground and excited states. In spin language, randomness of the directions of \mathbf{P} translates into the randomness of the directions of B_a . In a general case when $\mathbf{B}_a = \mathbf{n}_a B_a$, $\mathbf{B}_1 = \mathbf{n}_1 B_1$, where \mathbf{n}_a and \mathbf{n}_1 are the unit vectors, the drive amplitude in the above expressions should be modified as

$$B_1^2 \rightarrow B_1^2 (\mathbf{n}_1 \times \mathbf{n}_a)^2. \quad (42)$$

It is important to note that when the directions of the fields \mathbf{B}_a are random, we do not need the nondiagonal component of dipole-dipole interaction to induce the second-generation Rabi oscillations.

(vii) There is a similarity between the relay-race mechanism considered above and the Rabi-vibronic resonance studied in Ref. [34]. In the latter case, the Rabi oscillations are resonantly coupled to a vibronic mode rather than to the neighboring spin.

(viii) In the system we considered the pairs of Zeeman levels with random splittings were well separated in space. This should be contrasted to the situation of the ac absorption in the system where energy levels strongly overlap in space and almost evenly separated in energy. Then driven electron can “climb” the staircase of energy levels. Remarkably, in this situation, the absorption will still eventually saturate [4–6].

The reason is “dynamic localization,” i.e., trapping of electron in energy space due to disorder in the level positions.

ACKNOWLEDGMENTS

The work was supported by the Department of Energy, Office of Basic Energy Sciences, Grant No. DE-FG02-06ER46313.

-
- [1] N. F. Mott, Conduction in non-crystalline systems, *Philos. Mag.* **17**, 1259 (1968).
- [2] V. L. Berezinskii, Kinetics of a quantum particle in a one-dimensional random potential, *Sov. Phys. JETP* **38**, 620 (1974).
- [3] P. Bordia, H. Lüschen, U. Schneider, M. Knap, and I. Bloch, Periodically driving a many-body localized quantum system, *Nat. Phys.* **13**, 460 (2017).
- [4] Y. Gefen and D. J. Thouless, Zener Transitions and Energy Dissipation in Small Driven Systems, *Phys. Rev. Lett.* **59**, 1752 (1987).
- [5] D. M. Basko and V. E. Kravtsov, Dynamic Localization and the Coulomb Blockade in Quantum Dots under ac Pumping, *Phys. Rev. Lett.* **93**, 056804 (2004).
- [6] D. Cohen and T. Kottos, Quantum-Mechanical Nonperturbative Response of Driven Chaotic Mesoscopic Systems, *Phys. Rev. Lett.* **85**, 4839 (2000).
- [7] V. Khemani, R. Nandkishore, and S. L. Sondhi, Nonlocal adiabatic response of a localized system to local manipulations, *Nat. Phys.* **11**, 560 (2015).
- [8] R. Ducatez and F. Huvneers, Anderson localization for periodically driven systems, *Annales Henri Poincaré* **18**, 2415 (2017).
- [9] K. Agarwal, S. Ganeshan, and R. N. Bhatt, Localization and transport in a strongly driven Anderson insulator, *Phys. Rev. B* **96**, 014201 (2017).
- [10] S. Ray, A. Ghosh, and S. Sinha, Drive-induced delocalization in the Aubry-André model, *Phys. Rev. E* **97**, 010101(R) (2018).
- [11] D. T. Liu, J. T. Chalker, V. Khemani, and S. L. Sondhi, Mott, Floquet, and the response of periodically driven Anderson insulators, *Phys. Rev. B* **98**, 214202 (2018).
- [12] M. Wilkinson, Statistical aspects of dissipation by Landau-Zener transitions, *J. Phys. A: Math. Gen.* **21**, 4021 (1988).
- [13] M. Wilkinson, Adiabatic transport of localised electrons, *J. Phys. A* **24**, 2615 (1991).
- [14] M. Wilkinson and E. J. Austin, Dynamics of a generic quantum system under a periodic perturbation, *Phys. Rev. A* **46**, 64 (1992).
- [15] S. Gopalakrishnan, M. Knap, and E. Demler, Regimes of heating and dynamical response in driven many-body localized systems, *Phys. Rev. B* **94**, 094201 (2016).
- [16] S. Gopalakrishnan, M. Müller, V. Khemani, M. Knap, E. Demler, and D. A. Huse, Low-frequency conductivity in many-body localized systems, *Phys. Rev. B* **92**, 104202 (2015).
- [17] P. Ponte, A. Chandran, Z. Papić, and D. A. Abanin, Periodically driven ergodic and many-body localized quantum systems, *Ann. Phys.* **353**, 196 (2015).
- [18] P. Ponte, Z. Papić, F. Huvneers, and D. A. Abanin, Many-Body Localization in Periodically Driven Systems, *Phys. Rev. Lett.* **114**, 140401 (2015).
- [19] A. Lazarides, A. Das, and R. Moessner, Fate of Many-Body Localization Under Periodic Driving, *Phys. Rev. Lett.* **115**, 030402 (2015).
- [20] M. Bukov, S. Gopalakrishnan, M. Knap, and E. Demler, Prethermal Floquet Steady States and Instabilities in the Periodically Driven, Weakly Interacting Bose-Hubbard Model, *Phys. Rev. Lett.* **115**, 205301 (2015).
- [21] V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, On the Phase Structure of Driven Quantum Systems, *Phys. Rev. Lett.* **116**, 250401 (2016).
- [22] E. Bairey, G. Refael, and N. H. Lindner, Driving induced many-body localization, *Phys. Rev. B* **96**, 020201(R) (2017).
- [23] J. H. Shirley, Solution of the Schrödinger equation with a Hamiltonian periodic in time, *Phys. Rev.* **138**, B979 (1965).
- [24] R. Glenn, M. E. Limes, B. Pankovich, B. Saam, and M. E. Raikh, Magnetic resonance in slowly modulated longitudinal field: Modified shape of the Rabi oscillations, *Phys. Rev. B* **87**, 155128 (2013).
- [25] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded Growth of Entanglement in Models of Many-Body Localization, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [26] M. Serbyn, Z. Papić, and D. A. Abanin, Local Conservation Laws and the Structure of the Many-Body Localized States, *Phys. Rev. Lett.* **111**, 127201 (2013).
- [27] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, Observation of many-body localization of interacting fermions in a quasirandom optical lattice, *Science* **349**, 842 (2015).
- [28] J.-Y. Choi, S. Hild, J. Zeiher, P. Schauß, A. Rubio-Abadal, T. Yefsah, V. Khemani, D. A. Huse, I. Bloch, and C. Gross, Exploring the many-body localization transition in two dimensions, *Science* **352**, 1547 (2016).
- [29] A. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated many-body system, *Science* **353**, 794 (2016).
- [30] L. S. Levitov, Absence of localization of vibrational modes due to dipole-dipole interaction, *Europhys. Lett.* **9**, 83 (1989).
- [31] L. S. Levitov, Delocalization of Vibrational Modes Caused by Electric Dipole Interaction, *Phys. Rev. Lett.* **64**, 547 (1990).
- [32] L. S. Levitov, Critical Hamiltonians with long range hopping, *Ann. Phys. (Leipzig)* **8**, 697 (1999).
- [33] A. L. Burin, Yu. Kagan, L. A. Maksimov, and I. Ya. Polishchuk, Dephasing Rate in Dielectric Glasses at Ultralow Temperatures, *Phys. Rev. Lett.* **80**, 2945 (1998).
- [34] R. Glenn and M. E. Raikh, Rabi-vibronic resonance with large number of vibrational quanta, *Phys. Rev. B* **84**, 195454 (2011).