

## Exchange-correlation thermal effects in shocked deuterium: Softening the principal Hugoniot and thermophysical properties

V. V. Karasiev,<sup>\*</sup> S. X. Hu, M. Zaghoo, and T. R. Boehly

Laboratory for Laser Energetics, University of Rochester, 250 East River Road, Rochester, New York 14623-1299, USA



(Received 27 February 2019; revised manuscript received 3 June 2019; published 28 June 2019)

Exchange-correlation (XC) thermal effects for transport and optical properties of deuterium along the principal Hugoniot are investigated. The study is performed using *ab initio* molecular dynamics simulations within the Mermin-Kohn-Sham density functional theory. XC thermal effects are taken into account via the temperature-dependent Karasiev-Dufty-Trickey generalized gradient approximation functional [V. V. Karasiev *et al.*, *Phys. Rev. Lett.* **120**, 076401 (2018)]. We find that XC thermal effects account for the softening of the Hugoniot at pressures  $P > 250$  GPa and improve agreement with recent experimental measurements. Also, XC thermal effects lead to the reflectivity increase by about 2% for shock speeds above 20 km/s. The calculated reflectivity for shock speeds up to 50 km/s is in excellent agreement with recent experimental measurements on the Omega Laser System. The dc conductivity is increased by about 4% due to XC thermal effects. The system evolution along the Hugoniot crosses the so-called warm-dense-matter regime, and XC thermal effects must be taken into account to accurately predict the thermophysical properties across warm-dense conditions.

DOI: [10.1103/PhysRevB.99.214110](https://doi.org/10.1103/PhysRevB.99.214110)

### I. INTRODUCTION

Reliably predicting the properties of hydrogen and its isotopes under extreme conditions remains a problem of great importance and broad scientific interest. Accurate knowledge of the equation of state and transport properties over a wide range of thermodynamic conditions of this simplest and most abundant element in the universe is used as input for planetary, brown dwarf, and stellar modeling [1] as well as the inertial confinement fusion simulations to design targets [2–4]. The most advanced theoretical and computational methods are used to interpret experimental results and to predict properties at thermodynamic conditions that are difficult to access experimentally.

On the other hand, new experimental measurements with improved accuracy [5,6] serve as an important benchmark to assess the accuracy of theoretical predictions. It was found [6] that recent shock-compression data for deuterium are well described by finite-temperature density functional theory (DFT) methods [7–9] and are in disagreement with recent quantum Monte Carlo (QMC) results [10]. The QMC method [10] fails to correctly describe the shock compression near the molecular-to-atomic (MA) transition for both the onset of dissociation and peak compression [6]. DFT calculations with exchange-correlation (XC) functionals that account for van der Waals (vdW) long-range interactions (vdW-DF1 [11] and vdW-DF2 [12]) provide an improved description of the onset of the dissociation process along the Hugoniot, while peak compression is underestimated by these functionals. Standard generalized gradient approximation (GGA) XC functionals such as the Perdew-Burke-Ernzerhof (PBE) functional [13] describe the peak compression reasonably well, but the

dissociation process begins at a slightly ( $\sim 3$  GPa) lower pressure compared to the experimental data (see details in Ref. [6]). At pressures above 250 GPa along the Hugoniot, the DFT calculations with the PBE functional predict a stiffer behavior than recent experimental data [5].

Furthermore, optical and transport properties such as reflectivity, thermal conductivity, and electrical conductivity are important material properties used to characterize the complex behavior of shocked deuterium, which includes, in particular, insulator-to-metal and degenerate-to-nondegenerate plasma transitions. *Ab initio* molecular dynamics (AIMD) simulations, when the classical molecular dynamics for the ionic degrees of freedom is combined with the DFT treatment for the electrons, take into account quantum effects and treat all electronic thermally occupied and empty states on the same basis. All of this is important to accurately predict transport coefficients via Kubo-Greenwood formalism [14,15], especially in the partially degenerate, strongly coupled regime when the chemical- and plasma-based models [16,17] become inaccurate [4].

All current DFT calculations of the Hugoniot data and transport coefficients are performed with temperature-independent XC functionals developed for the ground state [4,6,18–21]; therefore, XC thermal effects, which play an important role in warm-dense-matter (WDM) conditions [22], are not taken into account. Figure 1 shows the  $(r_s, t)$  domain where the  $T$  dependence of XC might be important for accurate predictions. The relative importance of XC thermal effects is shown as a function of the Wigner-Seitz radius  $r_s = (3/4\pi n)^{1/3}$ , and the reduced temperature  $t = T/T_F$  ( $T_F = [3\pi^2 n]^{2/3}/2k_B$  is the Fermi temperature, and  $n$  is the electron number density). At low  $r_s$  XC thermal effects might become important for  $t$  values between a few tenths and  $t \approx 1$ ; at large  $r_s$  the interval is expanded up to  $t \approx 10$  or even higher. At high density (low  $r_s$ ) and high temperatures

<sup>\*</sup>vkarasiev@lle.rochester.edu




FIG. 1. The relative importance of explicit temperature dependence in the XC free-energy functional for the homogeneous electron gas measured as  $\log_{10}\{|f_{xc}(r_s, t) - e_{xc}(r_s)|/|f_s(r_s, t) + e_{xc}(r_s)|\}$ , where  $f_{xc}$  is the XC free energy per particle given by the corrected Karasiev-Sjostrom-Dufty-Trickey parametrization [24,25],  $e_{xc}$  is the zero-temperature XC energy per particle [26], and  $f_s$  is the noninteracting free energy per particle [27].  $\Gamma = 2\lambda^2 r_s/t$  with  $\lambda = (4/9\pi)^{1/3}$  is the classical coupling parameter. The solid white line corresponds to the liquid-deuterium principal Hugoniot path; the end point corresponds to  $P = 1$  TPa.

the exchange correlation does not play any role (see Fig. 1 in Ref. [23]), and XC thermal effects become negligible as well. At low densities the relative magnitude of the XC term with respect to the noninteracting piece is increased; therefore, one may expect that XC thermal effects might be noticeable even at lower temperatures (light blue and dark green zone at  $r_s \geq 10$  and low  $t$  in Fig. 1). For deuterium, the reduced temperature reaches a value of 0.1 at pressures near 70 GPa along the principal Hugoniot; therefore, we expect that XC thermal effects might be important for pressures beyond this value.

Here we focus on the study of the optical and transport properties along the principal Hugoniot of deuterium with a temperature-dependent Karasiev-Dufty-Trickey (KDT16) generalized gradient approximation XC functional [24]. KDT16 in the zero-temperature limit by construction reduces to the standard ground-state PBE [13]. Therefore, a second set of calculations is performed with the ground-state PBE XC to assess the XC thermal effects calculated as the difference between the two sets of results for several thermophysical properties.

The next section provides details of the methodology used in this study, including computational details. Section III describes our main results, and Sec. IV provides a short summary and conclusions.

## II. METHOD

### A. Exchange-correlation free-energy functional

Treatment of electronic degrees of freedom via the finite-temperature DFT requires the use of an approximate XC

free-energy functional. The reliability of the DFT predictions under high-energy-density (HED) conditions depends crucially on having an XC free-energy functional that is accurate across temperature regimes. Reference [22] studied XC thermal effects at the local density approximation (LDA) level of refinement and found that the temperature-dependent Karasiev-Sjostrom-Dufty-Trickey (KSDT) XC functional [25] in particular yields improved agreement with experiment for the dc electrical conductivity of low-density aluminum.

For nonhomogeneous systems it is important to go beyond the LDA and take into account the XC nonhomogeneity and thermal effects simultaneously via the nonempirical KDT16 GGA XC free-energy functional [24]. The KDT16 functional is constrained by the correct temperature dependence in the slowly varying regime and the correct zero-temperature, high-temperature, and homogeneous electron gas limits. The correct zero-temperature limit means that the KDT16 at  $T = 0$  reduces by construction to the extremely widely used ground-state PBE functional [13]. The KDT16 uses the corrected KSDT parametrization as the LDA component. In this work we use the KDT16 XC free-energy functional in such a way that all the XC thermal and nonhomogeneity effects are taken into account at the generalized gradient level of approximation. The KDT16 XC free energy is given as an explicit temperature-dependent functional of density  $\mathcal{F}_{xc}[n, T]$ . Hugoniot calculations considered in Sec. III A require knowledge of XC internal-energy contribution, which is calculated via standard thermodynamic relations for the XC entropy,

$$\mathcal{S}_{xc}[n, T] = - \left. \frac{\partial \mathcal{F}_{xc}[n, T]}{\partial T} \right|_{N, V}, \quad (1)$$

and internal energy

$$\mathcal{E}_{xc}[n, T] = \mathcal{F}_{xc}[n, T] + T \mathcal{S}_{xc}[n, T]. \quad (2)$$

The derivative with respect to temperature at a constant number of particles and system volume in Eq. (1) is calculated analytically, and functionals (1) and (2) are evaluated together with  $\mathcal{F}_{xc}$  while solving the Mermin-Kohn-Sham DFT equations.

Calculations with the ground-state PBE, when the XC free energy is approximated by a ground-state functional without explicit  $T$  dependence,  $\mathcal{F}_{xc}[n, T] \approx \mathcal{E}_{xc}[n]$ , are also performed to assess the magnitude of XC thermal effects on calculated properties. It is important to note that only temperature-dependent XC functionals provide an explicit  $\partial \mathcal{E}_{xc}[n, T]/\partial T|_V$  contribution to the system's specific heats.

### B. Computational details

AIMD simulations are performed in the canonical ensemble by using the ground-state PBE and temperature-dependent KDT16 XC functionals as implemented in the PROFESS@QUANTUM-ESPRESSO interface [28–31]. The electron-ion interaction was treated by a standard projector augmented wave (PAW) pseudopotential data set [32] generated at  $T = 0$  K with the ground-state PBE XC. Such a PAW data set is compatible with the temperature-dependent KDT16 functional because the KDT16 functional, by construction, reduces to PBE in the zero-temperature limit at which the

TABLE I. Electronic pressure from AIMD simulations with the PBE and KDT16 XC functionals and with the use of the PAW and bare Coulomb external (electron-ion) potentials for deuterium at material density  $\rho_D = 0.7224 \text{ g/cm}^3$  (64 atoms in the simulation cell,  $(1/4, 1/4, 1/4)$  and only the  $\Gamma$  point for  $T = 50$  and  $100 \text{ kK}$ , respectively).

$v_{\text{ext}}$	$T$ (kK)	$P^{\text{PBE}}$ (GPa)	$P^{\text{KDT16}}$ (GPa)
PAW	50	62.9	56.2
Bare Coulomb	50	63.3	56.6
PAW	100	170.0	154.0
Bare Coulomb	100	170.4	154.1

PAW set was generated. The underlying assumption, common in WDM studies, that effective electron-ion interaction potentials generated at zero temperature are transferable to various thermodynamic conditions was confirmed for hydrogenic systems by comparison between the exact bare Coulomb and pseudopotential (given by the simple local form) calculations (see Fig. 5 in Ref. [33] and studies of the PAW transferability to high-density/high-temperature regimes in Ref. [34]). Additionally, we performed AIMD simulations at two representative temperatures (50 and 100 kK) near the maximum of the absolute and relative XC thermal effects on pressure (see Fig. 4 in Sec. III A) with the use of our PAW data set and the exact electron-ion interaction given by the bare Coulomb potential. The agreement between calculations with the PAW and bare Coulomb interactions, shown in Table I, is excellent: the largest pressure difference of 0.4 GPa is only 0.7% of the electronic pressure and is much smaller compared to the pressure differences of 7 and 16 GPa (relative difference of 12% and 10%, respectively) due to the XC thermal effects on pressure at  $T = 50$  and  $100 \text{ kK}$ , respectively. These calculations confirm transferability of the PAW data set generated at  $T = 0 \text{ K}$  to finite temperatures.

Calculations were performed for a range of temperatures between 6000 and 200 000 K. For each temperature, the AIMD simulations were performed for three densities to find a solution of the Rankine-Hugoniot equation (see Sec. III A). The ion temperature was controlled by an Andersen thermostat, and the electron temperature, set equal to that of the ions, was fixed according to the Fermi-Dirac statistics. At low temperatures ( $T \leq 20 \text{ 000 K}$ ), calculations included 256 atoms in the supercell and Baldereschi's mean value point (BMVP) [35]. The size of the supercell gradually decreased to 128 atoms (for  $25 \text{ 000 K} \leq T \leq 50 \text{ 000 K}$  using BMVP), 64 atoms (for  $50 \text{ 000 K} \leq T \leq 75 \text{ 000 K}$  using BMVP and for  $80 \text{ 000 K} \leq T \leq 150 \text{ 000 K}$  using only the  $\Gamma$  point), and 32 atoms for  $T = 200 \text{ 000 K}$  using only the  $\Gamma$  point. Additionally, at temperatures  $T = 75, 100,$  and  $150 \text{ kK}$  we performed simulations with 128 atoms and only the  $\Gamma$  point to assess the magnitude of finite-size effects compared to calculations with 64 atoms in the simulation cell.

A sample set of 5 to 20 statistically independent configurations was selected from the AIMD ionic trajectories (so-called snapshotting). For this set of snapshots, a single-point DFT calculation was performed with the ABINIT [36] and/or PROFESS@QUANTUM-ESPRESSO [28] to generate a set of electronic-state functions  $\{\psi_n\}$ . For selected configurations

we also performed calculations with PROFESS@QUANTUM-ESPRESSO [28] to ensure that the results obtained from the two different codes (the DFT and the subsequent Kubo-Greenwood ones are in agreement). The velocity dipole matrix elements  $\langle \psi_n | \nabla | \psi_m \rangle$  were used by postprocessing codes from Ref. [37] for ABINIT and from Ref. [38] for PROFESS@QUANTUM-ESPRESSO to calculate the frequency-dependent Onsager coefficients within the Kubo-Greenwood formalism [14,15] and, eventually, to calculate the real part of the dc and dynamic electrical conductivities  $[\sigma_{\text{dc}}$  and  $\sigma(\omega)$ , respectively], the electronic thermal conductivity  $\kappa(\omega)$ , the real and imaginary parts of the refractive index  $[n(\omega)$  and  $k(\omega)$ , respectively], the reflectivity, and, eventually, the Rosseland mean opacity and specific heat. Calculation of the thermophysical properties including opacity with the use of the Kubo-Greenwood formalism in combination with the AIMD snapshotting is a well-established procedure frequently used at extreme HED conditions [39–43].

The reflectivity is defined as

$$r(\omega) = \frac{[n(\omega) - n_0]^2 + k^2(\omega)}{[n(\omega) + n_0]^2 + k^2(\omega)}, \quad (3)$$

where  $n_0$  is the refractive index of the ambient environment (deuterium in the initial liquid state in the case of the reflectivity calculations along the principal Hugoniot). Near maximum compression in the temperature range of 6 to 15 kK, these calculations were performed for a fixed density of  $0.7568 \text{ g/cm}^3$  (4.4-fold compression); for  $T \geq 20 \text{ kK}$ , a density of  $0.7224 \text{ g/cm}^3$  (4.2-fold compression) was used (although small variations in material density have little effect on calculated properties). The  $\delta$  function in the real part of the electrical conductivity  $\sigma(\omega)$ , obtained using the Kubo-Greenwood formalism, is approximated by a Gaussian of width  $\Delta$  (see Ref. [38] for details). We find that results are stable for  $\Delta$  width in the range of a few tenths of eV; therefore, in our calculations  $\Delta = 0.5 \text{ eV}$  was used. Convergence with respect to the number of snapshots was tested for selected points by increasing this number by a factor of 2. Results for reflectivity and dc conductivity changed by 0.5% or less. Convergence of the reflectivity and dc conductivity with respect to the cell size was tested for two temperatures,  $T = 20$  and  $50 \text{ kK}$ , by increasing the number of atoms from 128 to 256. The results are shown in Figs. 6 and 8 below; changes do not exceed 0.5% compared to calculations with 128 atoms.

All simulations employed a plane-wave cutoff energy of 125 Ry to converge high-energy thermally occupied bands required for calculations at elevated temperature and Baldereschi's mean value point [35] or  $\Gamma$ -point sampling of the first Brillouin zone in the case of AIMD simulations and up to a  $3 \times 3 \times 3$  Monkhorst-Pack  $k$  grid [44] for Kubo-Greenwood calculations. To ensure convergence with the number of bands, we used a minimum Fermi-Dirac occupation number threshold of the order of  $10^{-6}$  to  $10^{-7}$ .

It is important to notice that the KDT16 functional, employed in this work, is implemented in locally modified versions of the two codes, such that XC thermal effects are taken into account in both the AIMD simulations providing “structural” properties and in the Kubo-Greenwood calculations providing transport properties.




FIG. 2. (a) AIMD snapshot of liquid deuterium at initial conditions and (b) radial D-D distribution function.

### III. RESULTS

#### A. The principal Hugoniot of deuterium up to 1 TPa

The Hugoniot curve is described by solutions of the Rankine-Hugoniot equation

$$\mathcal{E} - \mathcal{E}_0 - \frac{1}{2}(P + P_0)\left(\frac{1}{\rho} - \frac{1}{\rho_0}\right) = 0, \quad (4)$$

where  $\mathcal{E}$ ,  $P$ , and  $\rho$  are the specific *internal* energy, pressure, and bulk density, respectively, of a state derived by single-shock compression from an initial state at  $\rho_0$ ,  $\mathcal{E}_0$ , and  $P_0$ . We performed an AIMD simulation for liquid deuterium at initial conditions  $\rho_0 = 0.172 \text{ g/cm}^3$ ,  $T_0 = 20 \text{ K}$  with 32  $\text{D}_2$  molecules in a simulation cell and obtained an average pressure  $P_0 = 0.15 \text{ kbar}$ , which is negligible in comparison to the final pressure  $P$ . The difference between the average energy per molecule from the AIMD simulation and the energy of an isolated  $\text{D}_2$  molecule (a single molecule in a large simulation box) is only  $0.006 \text{ eV}$ . Therefore, an AIMD simulation for the initial state is not required unless one is interested in the corresponding optical properties. In this work the initial-state AIMD snapshots were used to calculate the real part of the refractive index  $n_0$  required for reflectivity calculations [Eq. (3)] (see Sec. III B). The AIMD snapshot and radial distribution function (RDF) shown in Fig. 2 confirm that the initial state represents a pure  $\text{D}_2$  molecular liquid. The RDF has a very sharp peak at the  $\text{D}_2$  molecule equilibrium distance  $r \approx 1.4 \text{ bohr}$  and a small tail starting at a relatively large  $r \approx 4.5 \text{ bohr}$  distance.

The deuterium principal Hugoniot was calculated from simulations with the KDT16 and PBE XC functionals. Calculations were performed over the temperature range of 6 kK ( $P = 34 \text{ GPa}$ , near the MA transition) to 200 kK ( $P = 1010 \text{ GPa}$ ). Figure 3 compares our theoretical predictions, experimental results across the MA transition (low-pressure range  $P < 150 \text{ GPa}$ ; [6] shown as blue diamonds), and recent experimental measurements over the high-pressure range between 87 and 550 GPa [5] (shown as orange circles). The KDT16 and PBE curves (obtained by combining results from simulations with 256, 128, 64, and 32 atoms; see Fig. 3 caption and Sec. II B) are indistinguishable up to a relatively high temperature  $T = 60 \text{ kK}$  ( $P \approx 250 \text{ GPa}$ ).

According to Fig. 1, XC thermal effects might become noticeable starting at a lower temperature of  $\sim 0.2T_F$ . Figures 4(a) and 4(b) show pressure and specific internal energy differences between KDT16 and PBE,  $\Delta P = P^{\text{KDT16}} - P^{\text{PBE}}$  and  $\Delta \mathcal{E} = \mathcal{E}^{\text{KDT16}} - \mathcal{E}^{\text{PBE}}$ , along the Hugoniot. These




FIG. 3. Deuterium principal Hugoniot derived from the initial state  $\rho_0 = 0.172 \text{ g/cm}^3$  and  $T_0 = 20 \text{ K}$ . The PBE (black dashed) and KDT16 (red dashed) curves are obtained by combining results from simulations with 256 atoms ( $6 \leq T \leq 20 \text{ kK}$ , pressure range between 34 and 86 GPa for both functionals), 128 atoms ( $25 \leq T \leq 50 \text{ kK}$ , pressure range between 104 and 208 GPa for KDT16), 64 atoms ( $60 \leq T \leq 150 \text{ kK}$ , pressure range between 253 and 736 GPa for KDT16), and 32 atoms for  $T = 200 \text{ kK}$ . The red solid curve corresponds to the KDT16 results from simulations with 256 atoms ( $6 \leq T \leq 20 \text{ kK}$ ) and 128 atoms ( $25 \leq T \leq 150 \text{ kK}$ ).

differences represent XC thermal effects on pressure and energy and, as functions of  $T$ , have the same sign: negative for  $T < 250 \text{ kK}$  and positive for  $T > 250 \text{ kK}$ . Consequently, the differences effectively cancel out on the left-hand side of Eq. (4), the solution of which remains unchanged for





FIG. 4. (a) Pressure and (b) specific internal energy differences and (c) relative pressure difference  $[(P^{\text{KDT16}} - P^{\text{PBE}})/P^{\text{KDT16}} \times 100\%]$  and (d) total pressure along the deuterium Hugoniot as functions of temperature.

$T < 60$  kK. The Hugoniot curve's similar insensitivity to the use of the LDA temperature-dependent XC functional rather than the ground-state LDA XC was previously observed in Ref. [22]. In spite of the Hugoniot curve (pressure versus compression) being weakly affected by XC thermal effects (due to effective cancellation between the energy and pressure terms as just discussed), the pressure along the Hugoniot is lowered because of XC thermal effects, and the relative pressure difference reaches about 4% at  $T$  near 50 kK [see Figs. 4(c) and 4(d)].

Returning to discussing Fig. 3, both functionals, KDT16 and PBE, are in good agreement with experimental measurements in the range of pressure up to 200 GPa [6]. At higher pressures, however, the PBE curve becomes noticeably stiffer compared to the recent experimental data [5]; the disagreement reaches about 4% at  $P = 550$  GPa. The KDT16 predicts a curve that is softer by slightly more than 1% beyond 250 GPa compared to PBE. Increasing the simulation cell size from 64 to 128 atoms in this range of pressure leads to further softening of the Hugoniot. Simulations with 128 atoms at  $T > 50$  kK computationally are very demanding, so only calculations with the KDT16 functional were performed, and XC thermal effects in this range of pressure are estimated from simulations with 64 atoms. Calculated KDT16 and experimental compression does not vary in this range of  $P$ . The KDT16 compressibility is within the experimental uncertainty in the entire pressure range (including high pressures  $P > 250$  GPa). Therefore, the inclusion of XC thermal effects in AIMD calculations makes the deuterium Hugoniot softer at  $P > 250$  GPa and improves agreement with the experimental data. Reference [5] found that good agreement with the experimental data for low pressure  $P < 200$  was observed for the PBE DFT predictions and for a range of pressure between 200 and 550 GPa good agreement was observed for two models: the chemical livermore equation of state (LEOS) 1014 model [45] and the first-principles EOS (FPEOS) model [2], which in this range of  $P$  is based on the path-integral Monte Carlo (PIMC) simulations; the authors concluded that no single model consistently describes the deuterium principal Hugoniot throughout the entire pressure range. The KDT16 Hugoniot curve at low pressure is identical to the PBE curve and at high pressure has the quality of the PIMC FPEOS model. Therefore, by including XC thermal effects in the consideration, the KDT16 XC functional is able to describe the principal Hugoniot of liquid deuterium consistently over the entire pressure range.

### B. Thermophysical properties

Deuterium along the principal Hugoniot experiences changes in material structure that can, in turn, change the thermophysical properties such as dc and dynamic electrical conductivity, reflectivity, and thermal conductivity. Reduced temperature (the degeneracy parameter) along the Hugoniot changes between  $3 \times 10^{-4}$  (degenerate limit) and 1 (non-degenerate classical system; see Fig. 1). The range of the classical coupling parameter  $\Gamma$  is between 5000 (strongly coupled) and near 1 (moderately coupled). Consequently, the system's evolution occurs across a wide range of degeneracy and coupling parameters from degenerate, strongly coupled



TC14633J1

FIG. 5. Radial distribution function for a few selected temperatures along the deuterium Hugoniot.

quantum to nondegenerate, moderately coupled classical regimes.

Figure 5 illustrates structural changes. A relatively broad peak near 1.5 bohrs presented at  $T = 6$  kK indicates that the system is still a molecular liquid with a  $D_2$  bond-length range of  $\sim 1.2$  to 1.8 bohrs, but the dissociation process is in progress. At  $T = 15$  kK, the molecular-to-atomic transition is already completed (see Ref. [6]), even though some long-range order remains. This structural change leads to a significant increase in reflectivity (from 0.34 to 0.43; details will be discussed below). Further evolution of the system leads to a completely structureless radial distribution function and an increase in reflectivity.

In this section we present our theoretical predictions for reflectivity, the dynamic and dc electrical conductivities, thermal conductivity, the Lorentz number, the Rosseland mean opacity, and specific heat along the deuterium principal Hugoniot performed within the DFT and Kubo-Greenwood formalism by using the standard ground-state PBE XC and temperature-dependent KDT16 XC to assess the role of XC thermal effects.

Figure 6(a) presents reflectivity as a function of the shock speed calculated with the two XC functionals. As a result of XC thermal effects, the reflectivity is increased by about 2% for shock speeds above 20 km/s ( $T > 12$  kK). Figure 6(b) presents the real part of the index of refraction  $n_0$  of liquid deuterium in the initial state ( $\rho_0 = 0.172$  g/cm<sup>3</sup> and  $T = 20$  K) as a function of photon frequency. The refractive index is flat at low  $\omega$  and takes a value of 1.17 at 532 and 808 nm, the probe wavelengths used in experiments. This result is in good agreement with the reference value of 1.14 [46]. To compare with the experimental data, the reflectivity was calculated at 532 and 808 nm with the KDT16 XC functional and our predicted value of the refractive index. Results of recent experiments [47] on Omega and previous measurements [48] are shown in Fig. 7. There is excellent agreement between the KDT16 values and experimental data at the 808-nm wavelength for the range of shock speeds considered in the calculations, even though the experimental data have relatively large error bars. The KDT16 results at 532 nm




FIG. 6. Reflectivity (for a wavelength of 532 nm) along the deuterium Hugoniot as a function of the shock speed calculated with the ground-state PBE and temperature-dependent KDT16 XC functionals by using Eq. (3) and (a)  $n_0 = 1$  and (b) the real part of the refractive index for liquid deuterium in the initial state. The reflectivity curves are obtained by combining results from simulations with 256 atoms ( $16 \leq U_s \leq 22$  km/s), 128 atoms ( $25 \leq U_s \leq 42$  km/s), and 64 atoms ( $44 \leq U_s \leq 75$  km/s).

are in very good agreement with recent Omega experimental data for shock speeds below 50 km/s. The reflectivity is underestimated by the DFT calculations at high shock speeds  $U_s > 50$  km/s compared to the experiment. Experimental reflectance as a function of shock speed changes the slope at  $U_s$  near 45 km/s ( $T \approx 0.4T_F = 60$  kK); this change of the slope is related to the lifting of the Fermi degeneracy. The system starts to behave as a classical one at a significantly lower temperature compared to  $T_F$  (see details in Ref. [47]). Calculated KDT16 reflectivity at the same 532-nm wavelength rises very quickly from 0.29 at 16 km/s ( $T \approx 6$  kK) to 0.39 at 20 km/s ( $T \approx 12$  kK), which roughly corresponds to maximum compression near molecular-to-atomic transition; it slowly continues to increase, and near 43 km/s the slope also increases. This qualitative behavior of DFT predictions is in agreement with the experimental data, but the experimental data slope increase is larger. Our PBE reflectivity data (not shown in Fig. 7) lie slightly lower than the KDT16 curve [see Fig. 6(a)], and the agreement with experiment is a bit worse.




FIG. 7. Reflectivity of shocked deuterium. Theoretical DFT predictions and experimental data are compared.




FIG. 8. (a) The dc electrical conductivity, (b) thermal conductivity, (c) Lorentz number, and (d) Rosseland mean opacity along the deuterium Hugoniot as functions of the shock speed (dc conductivity and Lorentz number) or as a function of temperature (thermal conductivity and opacity) calculated with the ground-state PBE and temperature-dependent KDT16 XC functionals. All curves are obtained by combining results from simulations with 256 atoms ( $16 \leq U_s \leq 22$  km/s,  $T$  between 6 and 15 kK), 128 atoms ( $25 \leq U_s \leq 42$  km/s,  $T$  between 20 and 55 kK), and 64 atoms ( $44 \leq U_s \leq 75$  km/s,  $T$  between 60 and 150 kK).

Figure 7 also shows PBE results of previous studies [4,21] that observed a similar change of the reflectivity slope at a shock speed near 45 km/s. These calculations used  $n_0 = 1$ ; as a consequence, the reflectivity at 532 nm is overestimated in comparison with experimental measurements.

To gain further insights into the properties of the shocked deuterium, the dc electrical conductivity, thermal conductivity, Lorentz number, Rosseland mean opacity, and specific heat were calculated. The four panels in Fig. 8 show the results. Using the temperature-dependent exchange correlation has an effect on the dc electrical conductivity:  $\sigma_{dc}$  is increased by 4% at intermediate shock speeds between 25 and 40 km/s. The dc conductivity increases very fast across the molecular-to-atomic transition (the  $U_s$  range is  $\approx 16$  to 20 km/s; the  $T$  range is between 6 and 12 kK), similar to the reflectivity increase shown in Fig. 7. After reaching the maximum at the end of the molecular-to-atomic transition, the system behaves as a liquid metal:  $\sigma_{dc}$  decreases with an increase in temperature for  $U_s$  between 20 and 30 km/s ( $T$  between 12 and 30 kK). For  $U_s > 30$  km/s ( $T > 30$  kK), the electrical dc conductivity increases with temperature as in the plasma regime, which is consistent with an observed degenerate-to-nondegenerate plasma transition [47]. The thermal conductivity  $\kappa$  is practically identical for calculations with ground-state and finite-temperature XC functionals;  $\kappa$  increases with a slight increase of the temperature dependence at  $T \approx 60$  kK, and  $\kappa$  behaves as  $\kappa \propto T^{1.6}$  at  $T \gtrsim 0.4T_F = 60$  kK. The Lorentz number, defined as the ratio  $L = \kappa / \sigma_{dc} T$ , is close to the degenerate limit value of  $2.44 \times 10^{-8}$  given by the Wiedemann-Franz law at  $U_s$  near 20 km/s and grows




FIG. 9. (a) The specific heat  $C_V^{\text{el}}$  (in units of  $k_B$  per atom) as a function of temperature for deuterium at material density  $\rho_D = 0.7224 \text{ g/cm}^3$  calculated from AIMD simulation data (circles and solid line) and for the homogeneous electron gas (dashed line) and (b) the radial distribution function for the three representative temperatures at  $\rho_D = 0.7224 \text{ g/cm}^3$ . All were calculated with the KDT16 XC functional.

with temperature. At  $U_s \approx 43 \text{ km/s}$ , this trend is reversed, and at high shock speeds (high temperatures), the Lorentz ratio approaches the nondegenerate classical limit.

The Rosseland mean opacity is shown in Fig. 8(d) as a function of temperature. At intermediate temperatures, it behaves according to the Kramers law (shown by the black dashed line) when bound-free and free-free contributions dominate. At  $T \approx 0.43T_F = 65 \text{ kK}$  ( $U_s \approx 45 \text{ km/s}$ ) it starts to deviate from  $\sim T^{-7/2}$  behavior toward saturation when electron scattering is the dominant opacity contribution. Saturation is expected to occur at very high temperatures when the system is completely ionized.

Figure 9(a) shows the heat capacity  $C_V^{\text{el}} = (\partial \mathcal{E}^{\text{el}} / \partial T)_V$ , where  $\mathcal{E}^{\text{el}}$  is the electronic structure internal energy

$$\mathcal{E}^{\text{el}} = \mathcal{T}_s + \mathcal{E}_H + \mathcal{E}_{xc} + \mathcal{E}_{\text{ion-el}} + \mathcal{E}_{\text{ion-ion}}, \quad (5)$$

calculated from the AIMD simulation data for deuterium at material density  $\rho_D = 0.7224 \text{ g/cm}^3$  ( $r_s = 2$ ) as a sum of the noninteracting kinetic energy of electrons, electronic Hartree, electronic XC, ion-electron interaction, and ion-ion classical electrostatic interaction terms. Energy  $\mathcal{E}^{\text{el}}$  extracted from AIMD simulations was represented in the form of Padé approximants [49] by fitting to two temperature intervals, below and above 20 kK, to reduce fitting errors that unfortunately affect the accuracy of the results for  $C_V^{\text{el}}$ . The total internal energy of the system is given by  $\mathcal{E}_{\text{tot}} = \mathcal{T}_{\text{kin-ion}} + \mathcal{E}^{\text{el}}$ , where the classical kinetic energy of the ions depends on (ionic) temperature  $T$  as  $\mathcal{T}_{\text{kin-ion}} = (3/2)Nk_B T$ , where  $N$  is the number of atoms. Therefore, the ionic kinetic energy makes a temperature-independent contribution to the heat capacity  $C_V^{\text{ion-kin}} = (3/2)k_B$  per atom, and the total heat capacity is given by  $C_V^{\text{tot}} = (3/2)k_B + C_V^{\text{el}}$ .

Results for the homogeneous electron gas (HEG) of the same density (same  $r_s$ ) are shown for comparison (dashed line in Fig. 9). At  $T = 1 \text{ kK}$ , the system represents a pure diatomic molecular liquid,  $C_V^{\text{el}} = 0.7k_B$ , and increases with an increase in temperature due to the growing contribution of the molecular vibrational degrees of freedom.  $C_V^{\text{el}}$  reaches a maximum in the predissociation state at  $T = 4 \text{ kK}$  and drops down to a minimum after dissociation is over ( $T \approx 10 \text{ kK}$ ). Figure 9(b) shows RDF at these three representative

temperatures and illustrates that at  $T = 1 \text{ kK}$  the system is a pure diatomic liquid, at  $T = 4 \text{ kK}$  the dissociation process begins, and at  $T = 10 \text{ kK}$  the dissociation is complete. At  $T > 10 \text{ kK}$  the  $C_V^{\text{el}}$  behavior is very close to the HEG curve. This means that the pure electronic contribution to  $C_V^{\text{el}}$  dominates after the molecular-to-atomic liquid transition ends at  $T > 10 \text{ kK}$ . At high temperature ( $T \geq 0.65T_F = 100 \text{ kK}$ ), the electronic contribution becomes close to the ideal gas value of  $3/2k_B$  per particle.

#### IV. SUMMARY AND CONCLUSIONS

In this work we studied the thermophysical properties of shocked deuterium along the principal Hugoniot up to 1 TPa by using the temperature-dependent XC functional within the Mermin-Kohn-Sham DFT and Kubo-Greenwood approach. It was found that inclusion of XC thermal effects leads to softening of the principal Hugoniot at high pressure  $P > 250 \text{ GPa}$ , which improves agreement with recent experimental measurements. Although XC thermal effects on the Hugoniot curve (pressure versus compression) are reduced by an effective cancellation between the pressure and energy differences, the XC functional that included thermal effects provides a consistent description of the principal Hugoniot throughout the entire pressure range.

Changes in material structure and associated changes of reflectivity and the electrical conductivity along the deuterium Hugoniot qualitatively are consistent with previous studies [20,21]. Quantitatively, however, the results are different. The magnitude of XC thermal effects on various properties is different. The reflectivity is increased by about 2% at intermediate temperatures above 12 kK, leading to a slightly better agreement with experiment (the KDT16 curve in Fig. 6 passes right through the middle of the experimental points for  $U_s < 50 \text{ km/s}$ ). However, the experimental error bars are larger than 2%; therefore, the reflectivity without XC thermal effects taken into account will be within the error bars as well. The electrical dc conductivity is increased by 4% at intermediate temperatures, and based on a previous study of dc conductivity in warm-dense aluminum [22], we might expect that this increase goes in the right direction and improves the reliability of theoretical predictions.

The deuterium system along the Hugoniot experiences transformations from an insulating molecular liquid to atomic poor metallic liquid and finally to nondegenerate classical plasma. The signature of the molecular-to-atomic transition is found in a sharp increase of electrical dc conductivity and reflectivity at shock speeds in the range between 16 and 20 km/s (a range of temperature between 6 and 12 kK). An increase in the slope of calculated reflectivity at  $U_s \approx 43 \text{ km/s}$  ( $T \approx 0.4T_F = 60 \text{ kK}$ ), related to the breakdown of the electrons' degeneracy and emergence of classical statistics [47], is in agreement with experimental measurements. Thermal conductivity slightly changes behavior at the same temperature,  $T \approx 0.4T_F$ , although its temperature dependence  $\kappa \propto T^{1.6}$  remains slow compared to the ideal plasma limit  $\kappa \propto T^{2.5}$ . Rosseland mean opacity is another property that exhibits change in behavior toward the classical system at  $U_s \approx 45 \text{ km/s}$  ( $T \approx 0.43T_F = 65 \text{ kK}$ ). However, the electronic structure heat capacity  $C_V^{\text{el}}$ , which at  $T > 10 \text{ kK}$  is

dominated by the pure electronic contribution, approaches the classical  $3/2k_B$  value at a higher temperature  $T = 0.65T_F = 100$  kK. Electrical dc conductivity changes behavior and starts to increase with temperature as in the plasma regime at a shock speed slightly above 30 km/s (temperature above 30 kK). Our results confirm that the crossover between the quantum and classical statistics occurs below the  $T = T_F$  limit. This is apparent in the observed change in the transport and the thermodynamic properties of the deuterium fluid in the region of  $0.4T_F - 0.65T_F$ . Future work should investigate the dependence of the onset of this crossover on density.

The KDT16 functional takes into account XC thermal effects at the GGA level of sophistication. This functional inherited limitations of its zero-temperature PBE limit, in particular the limitation of the energy band-gap underestimation. Therefore, development of more advanced temperature-dependent GGAs and/or meta-GGAs based, for example, on a deorbitalized meta-GGA [50] and also inclusion of the long-range van der Waals interaction [51] might provide a better description of thermophysical properties across the entire temperature-pressure range.

## ACKNOWLEDGMENTS

This material is based upon work supported by the Department of Energy National Nuclear Security Administration under Award Number DE-NA0003856, US National Science Foundation PHY Grant No. 1802964, the University of Rochester, and the New York State Energy Research and Development Authority.

This report was prepared as an account of work sponsored by an agency of the U.S. Government. Neither the U.S. Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the U.S. Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the U.S. Government or any agency thereof.

- 
- [1] J. J. Fortney and N. Nettelmann, The interior structure, composition, and evolution of giant planets, *Space Sci. Rev.* **152**, 423 (2010).
- [2] S. X. Hu, B. Militzer, V. N. Goncharov, and S. Skupsky, First-principles equation-of-state table of deuterium for inertial confinement fusion applications, *Phys. Rev. B* **84**, 224109 (2011).
- [3] S. X. Hu, L. A. Collins, V. N. Goncharov, T. R. Boehly, R. Epstein R. L. McCrory, and S. Skupsky, First-principles opacity table of warm dense deuterium for inertial-confinement-fusion applications, *Phys. Rev. E* **90**, 033111 (2014).
- [4] S. X. Hu, V. N. Goncharov, T. R. Boehly, R. L. McCrory, S. Skupsky, L. A. Collins, J. D. Kress, and B. Militzer, Impact of first-principles properties of deuterium-tritium on inertial confinement fusion target designs, *Phys. Plasmas* **22**, 056304 (2015).
- [5] A. Fernandez-Pañella, M. Millot, D. E. Fratanduono, M. P. Desjarlais, S. Hamel, M. C. Marshall, D. J. Erskine, P. A. Sterne, S. Haan, T. R. Boehly, G. W. Collins, J. H. Eggert, and P. M. Celliers, Shock Compression of Liquid Deuterium up to 1 TPa, *Phys. Rev. Lett.* **122**, 255702 (2019).
- [6] M. D. Knudson and M. P. Desjarlais, High-Precision Shock Wave Measurements of Deuterium: Evaluation of Exchange-Correlation Functionals at the Molecular-To-Atomic Transition, *Phys. Rev. Lett.* **118**, 035501 (2017).
- [7] N. D. Mermin, Thermal properties of the inhomogeneous electron gas, *Phys. Rev.* **137**, A1441 (1965).
- [8] M. V. Stoitsov and I. Zh. Petkov, Density functional theory at finite temperatures, *Ann. Phys. (NY)* **185**, 121 (1988).
- [9] R. M. Dreizler, Density functional theory at finite temperatures, in *The Nuclear Equation of State, Part A: Discovery of Nuclear Shock Waves and the EOS*, edited by W. Greiner and H. Stöcker, NATO Advanced Studies Institute, Series B: Physics Vol. 216 (Plenum, New York, 1989), p. 521.
- [10] N. M. Tubman, E. Liberatore, C. Pierleoni, M. Holzmann, and D. M. Ceperley, Molecular-Atomic Transition Along the Deuterium Hugoniot Curve with Coupled Electron-Ion Monte Carlo Simulations, *Phys. Rev. Lett.* **115**, 045301 (2015).
- [11] M. Dion, H. Rydberg, E. Schröder, D. C. Langreth, and B. I. Lundqvist, Van der Waals Density Functional for General Geometries, *Phys. Rev. Lett.* **92**, 246401 (2004).
- [12] K. Lee, E. D. Murray, L. Kong, B. I. Lundqvist, and D. C. Langreth, Higher-accuracy van der Waals density functional, *Phys. Rev. B* **82**, 081101(R) (2010).
- [13] J. P. Perdew, K. Burke, and M. Ernzerhof, Generalized Gradient Approximation Made Simple, *Phys. Rev. Lett.* **77**, 3865 (1996); **78**, 1396(E) (1997).
- [14] R. Kubo, General theory and simple applications to magnetic and conduction problems, *J. Phys. Soc. Jpn.* **12**, 570 (1957).
- [15] D. A. Greenwood, The Boltzmann equation in the theory of electrical conduction in metals *Proc. Phys. Soc.* **71**, 585 (1958).
- [16] Y. T. Lee and R. M. More, An electron conductivity model for dense plasma, *Phys. Fluids* **27**, 1273 (1984).
- [17] B. Wilson, V. Sonnad, P. Sterne, and W. Isaacs, A new implementation of the inferno algorithm, *J. Quantum Spectrosc. Radiat. Transfer* **99**, 658 (2006).
- [18] B. Holst, R. Redmer, and M. P. Desjarlais, Thermophysical properties of warm dense hydrogen using quantum molecular dynamics simulations, *Phys. Rev. B* **77**, 184201 (2008).
- [19] L. Caillabet, S. Mazevet, and P. Loubeyre, Multiphase equation of state of hydrogen from *ab initio* calculations in the range 0.2 to 5 g/cc up to 10 eV, *Phys. Rev. B* **83**, 094101 (2011).
- [20] L. A. Collins, S. R. Bickham, J. D. Kress, and S. Mazevet, T. J. Lenosky, N. J. Troullier, and W. Windl, Dynamical and optical properties of warm dense hydrogen, *Phys. Rev. B* **63**, 184110 (2001).
- [21] L. A. Collins, J. D. Kress, and D. E. Hanson, Reflectivity of warm dense deuterium along the principal Hugoniot, *Phys. Rev. B* **85**, 233101 (2012).



- [22] V. V. Karasiev, L. Calderín, and S. B. Trickey, Importance of finite-temperature exchange correlation for warm dense matter calculations, *Phys. Rev. E* **93**, 063207 (2016).
- [23] V. V. Karasiev, S. B. Trickey, and J. W. Dufty, Status of free-energy representations for homogeneous electron gas, *Phys. Rev. B* **99**, 195134 (2019).
- [24] V. V. Karasiev, J. W. Dufty, and S. B. Trickey, Nonempirical Semilocal Free-Energy Density Functional for Matter under Extreme Conditions, *Phys. Rev. Lett.* **120**, 076401 (2018).
- [25] V. V. Karasiev, T. Sjöstrom, J. Dufty, and S. B. Trickey, Accurate Homogeneous Electron Gas Exchange-Correlation Free Energy for Local Spin-Density Calculations, *Phys. Rev. Lett.* **112**, 076403 (2014).
- [26] J. P. Perdew and A. Zunger, Self-interaction correction to density-functional approximations for many-electron systems, *Phys. Rev. B* **23**, 5048 (1981).
- [27] R. P. Feynman, N. Metropolis, and E. Teller, Equations of state of elements based on the generalized Fermi-Thomas theory, *Phys. Rev.* **75**, 1561 (1949).
- [28] V. V. Karasiev, T. Sjöstrom, and S. B. Trickey, Finite-temperature orbital-free DFT molecular dynamics: Coupling profess and quantum espresso, *Comput. Phys. Commun.* **185**, 3240 (2014).
- [29] G. S. Ho, V. L. Lignères, and E. A. Carter, Introducing PROFESS: A new program for orbital-free density functional theory calculations, *Comput. Phys. Commun.* **179**, 839 (2008).
- [30] L. Hung, C. Huang, I. Shin, G. S. Ho, V. L. Lignères, and E. A. Carter, Introducing PROFESS 2.0: A parallelized, fully linear scaling program for orbital-free density functional theory calculation, *Comput. Phys. Commun.* **181**, 2208 (2010).
- [31] P. Giannozzi, S. Baroni, N. Bonini, M. Calandra, R. Car, C. Cavazzoni, D. Ceresoli, G. L. Chiarotti, M. Cococcioni, I. Dabo, A. Dal Corso, S. de Gironcoli, S. Fabris, G. Fratesi, R. Gebauer, U. Gerstmann, C. Gougoussis, A. Kokalj, M. Lazzeri, L. Martin-Samos, N. Marzari, F. Mauri, R. Mazzarello, S. Paolini, A. Pasquarello, L. Paulatto, C. Sbraccia, S. Scandolo, G. Sclauzero, A. P. Seitsonen, A. Smogunov, P. Umari, and R. M. Wentzcovitch, QUANTUM ESPRESSO: A modular and open-source software project for quantum simulations of materials, *J. Phys.: Condens. Matter* **21**, 395502 (2009).
- [32] F. Jollet, M. Torrent, and N. Holzwarth, Generation of projector augmented-wave atomic data: A 71 element validated table in the XML format, *Comput. Phys. Commun.* **185**, 1246 (2014).
- [33] V. V. Karasiev, T. Sjöstrom, and S. B. Trickey, Generalized-gradient approximation noninteracting free-energy functionals for orbital-free density functional calculations, *Phys. Rev. B* **86**, 115101 (2012).
- [34] V. V. Karasiev, T. Sjöstrom, and S. B. Trickey, Comparison of density functional approximations and the finite-temperature Hartree-Fock approximation in warm dense lithium, *Phys. Rev. E* **86**, 056704 (2012).
- [35] A. Baldeschi, Mean-value point in the Brillouin zone, *Phys. Rev. B* **7**, 5212 (1973).
- [36] X. Gonze, B. Amador, P.-M. Anglade, J.-M. Beuken, F. Bottin, P. Boulanger, F. Bruneval, D. Caliste, R. Caracas, M. Côté, T. Deutsch, L. Genovese, Ph. Ghosez, M. Giantomassi, S. Goedecker, D. R. Hamann, P. Hermet, F. Jollet, G. Jomard, S. Leroux, M. Mancini, S. Mazevet, M. J. T. Oliveira, G. Onida, Y. Pouillon, T. Rangel, G.-M. Rignanese, D. Sangalli, R. Shaltaf, M. Torrent, M. J. Verstraete, G. Zerah, and J. W. Zwanziger, A brief introduction to the ABINIT software package, *Comput. Phys. Commun.* **180**, 2582 (2009); X. Gonze, G.-M. Rignanese, M. Verstraete, J.-M. Beuken, Y. Pouillon, R. Caracas, F. Jollet, M. Torrent, G. Zerah, M. Mikami, Ph. Ghosez, M. Veithen, J.-Y. Raty, V. Olevano, F. Bruneval, L. Reining, R. Godby, G. Onida, D. R. Hamann, and D. C. Allan, A brief introduction to the ABINIT software package, *Z. Kristallogr.* **220**, 558 (2005).
- [37] S. Mazevet, M. Torrent, V. Recoules, and F. Jollet, Calculations of the transport properties within the PAW formalism, *High Energy Density Phys.* **6**, 84 (2010).
- [38] L. Calderín, V. V. Karasiev, and S. B. Trickey, Kubo-Greenwood electrical conductivity formulation and implementation for projector augmented wave datasets, *Comput. Phys. Commun.* **221**, 118 (2017).
- [39] M. P. Desjarlais, J. D. Kress, and L. A. Collins, Electrical conductivity for warm, dense aluminum plasmas and liquids, *Phys. Rev. E* **66**, 025401(R) (2002).
- [40] S. Mazevet, L. A. Collins, N. H. Magee, J. D. Kress, and J. J. Keady, Quantum molecular dynamics calculations of radiative opacities, *Astron. Astrophys.* **405**, L5 (2003).
- [41] S. Mazevet, J. D. Kress, and L. A. Collins, and P. Blottiau, Quantum molecular-dynamics study of the electrical and optical properties of shocked liquid nitrogen, *Phys. Rev. B* **67**, 054201 (2003).
- [42] S. Mazevet, M. P. Desjarlais, L. A. Collins, J. D. Kress, and N. H. Magee, Simulations of the optical properties of warm dense aluminum, *Phys. Rev. E* **71**, 016409 (2005).
- [43] V. Recoules and J.-P. Crocombette, *Ab initio* determination of electrical and thermal conductivity of liquid aluminum, *Phys. Rev. B* **72**, 104202 (2005).
- [44] H. J. Monkhorst and J. D. Pack, Special points for Brillouin-zone integrations, *Phys. Rev. B* **13**, 5188 (1976).
- [45] G. I. Kerley, Equations of state for hydrogen and deuterium, Sandia National Laboratories, Technical Report No. SAND2003-3613, 2003.
- [46] P. C. Souers, *Hydrogen Properties for Fusion Energy* (University of California Press, Berkeley, 1986).
- [47] M. Zaghoo, T. R. Boehly, J. R. Rygg, P. M. Celliers, S. X. Hu, and G. W. Collins, Breakdown of Fermi Degeneracy in the Simplest Liquid Metal, *Phys. Rev. Lett.* **122**, 085001 (2019).
- [48] P. M. Celliers, G. W. Collins, L. B. Da Silva, D. M. Gold, R. Cauble, R. J. Wallace, M. E. Foord, and B. A. Hammel, Shock-Induced Transformation of Liquid Deuterium into a Metallic Fluid, *Phys. Rev. Lett.* **84**, 5564 (2000).
- [49] *The Pade Approximant in Theoretical Physics*, edited by G. A. Baker, Jr., and J. L. Gammel (Academic, New York, 1970).
- [50] D. Mejia-Rodriguez and S. B. Trickey, Deorbitalization strategies for meta-generalized-gradient-approximation exchange-correlation functionals, *Phys. Rev. A* **96**, 052512 (2017).
- [51] J. Hinz, V. V. Karasiev, S. X. Hu, M. Zaghoo, and D. Mejia-Rodriguez, Deorbitalized Meta-GGA with the Long Range van der Waals Exchange-Correlation Functional Calculations of the Insulator-Metal Transition of Hydrogen, presented at the 59th Sanibel Symposium, St. Simons Island, GA, 17-22 February 2019, (unpublished).