

Diffusion of heavy quarks in the early stage of high-energy nuclear collisions at energies available at the BNL Relativistic Heavy Ion Collider and at the CERN Large Hadron Collider

J. H. Liu,^{1,2} S. Plumari^{3,2}, S. K. Das,⁴ V. Greco^{3,2} and M. Ruggieri^{1,*}

¹*School of Nuclear Science and Technology, Lanzhou University, 222 South Tianshui Road, Lanzhou 730000, China*

²*INFN-Laboratori Nazionali del Sud, Via S. Sofia 62, I-95123 Catania, Italy*

³*Department of Physics and Astronomy, University of Catania, Via S. Sofia 64, I-95125 Catania*

⁴*School of Physical Sciences, Indian Institute of Technology Goa, Ponda-403401, Goa, India*



(Received 18 December 2019; revised 9 May 2020; accepted 29 July 2020; published 7 October 2020)

We study the diffusion of charm and beauty in the early stage of high-energy nuclear collisions at RHIC and LHC energies, considering the interaction of these heavy quarks with the evolving Glasma by means of the Wong equations, in the color- $SU(2)$ case. In comparison with previous works, we add the longitudinal expansion of the gluon medium as well as we estimate the diffusion coefficient in momentum space, and the effect of energy loss due to gluon radiation. We find that heavy quarks diffuse in the strong transverse color fields in the very early stage (0.2–0.3 fm/c) and this leads to a suppression at low p_T and enhancement at intermediate low p_T . The shape of the observed nuclear suppression factor obtained within our calculations is in qualitative agreement with the experimental results of the same quantity for D mesons in proton-nucleus collisions. We compute the nuclear suppression factor in nucleus-nucleus collisions as well, for both charm and beauty, finding a substantial impact of the evolving Glasma phase on these, suggesting that initialization of heavy quarks spectra in the quark-gluon plasma phase should not neglect the early evolution in the strong gluon fields.

DOI: [10.1103/PhysRevC.102.044902](https://doi.org/10.1103/PhysRevC.102.044902)

I. INTRODUCTION

The study of the initial condition in high-energy collisions is an interesting problem related to the physics of relativistic heavy ion collisions (RHICs), as well as to that of high-energy proton-nucleus (pA) and proton-proton (pp) collisions. At very high collision energy the two colliding nuclei can be described within the color-glass-condensate (CGC) effective theory [1–7], in which fast partons dynamic is frozen by time dilatation and they act as static sources for low momentum gluons: their large occupation number allows to treat them as classical fields. The collision of the two CGC sheets, each representing one of the colliding objects in high-energy collisions, leads to the formation of strong gluon fields in the forward light cone called the Glasma [8–18]. The Glasma consists of longitudinal color-electric and color-magnetic fields in the weak coupling regime and characterized by large gluon occupation number, $A_\mu^a \simeq 1/g$ with g the QCD coupling, so they are described by classical field theory namely the classical Yang-Mills (CYM) theory. Finite coupling bring up fluctuations on the top of the Glasma [19–33] that we do not consider in the present manuscript.

Heavy quarks are considered as noble probes of the system produced in high-energy nuclear collisions, both for the pre-equilibrium phase and for the thermalized quark-gluon plasma (QGP); see Refs. [34–54] and references therein. Their formation time is very small in comparison with the one of

light quarks: indeed, this can be estimated as $\tau_{\text{form}} \approx 1/(2m)$ with m the heavy quark mass which gives $\tau_{\text{form}} \leq 0.1$ fm/c for charm and beauty quarks. Since heavy quarks are produced immediately after the collision, they can propagate in the evolving Glasma fields and probe its evolution.

Diffusion of heavy quarks in the early stage of high-energy nuclear collisions has been studied previously in Ref. [50] within a simplified approach based on a Fokker-Planck equation. A similar study has been performed in Refs. [51,52] where emphasis has been put on the calculation of the nuclear modification factor in p-Pb collision, R_{pPb} . In comparison with Ref. [50], the work in Refs. [51,52] goes beyond the small momentum exchange approximation, solving the heavy quarks dynamics in the evolving Glasma fields via the Wong equations; this method is equivalent to solve the Boltzmann-Vlasov equations for the heavy quarks in a collisionless plasma. As a matter of fact, the Boltzmann-Vlasov equations can be solved by means of the test particles method which analogous to solve the classical equations of motion of the test particles, here represented by the heavy quarks, and these classical equations are the Wong equations. Then in Ref. [34] the effect of this dynamics on the elliptic flow in Pb-Pb collisions has been considered. However, in Refs. [34,51,52] the longitudinal expansion of the medium has been ignored.

The main purpose of the present article is to report on a more complete study of the problem of diffusion of charm and beauty in the early stage of high-energy nuclear collisions, improving the work already presented in Refs. [34,51] by including the longitudinal expansion of the gluon medium. For the sake of computational simplicity we consider the

*ruggieri@lzu.edu.cn

color- $SU(2)$ case here. Although the expansion dilutes the energy density and makes the effect on the modification factors smaller than the one found in Refs. [34,51], we still find a substantial contribution of the early stage to these quantities both in pA and in AA collisions.

In addition to this necessary improvement, we also estimate the diffusion coefficient in momentum space of the heavy quarks in the evolving Glasma, which was not done in Refs. [34,51]; finally, we present an estimate of the energy loss due to gluon radiation that acts as a drag force on the heavy quarks. We find that unless we artificially take a low average energy density in the full evolution, the effect of the drag force is not strong enough to cancel the tilt of the spectrum of charm and beauty. We remark that we do not include cold nuclear matter effects in our calculations [35,36,55–60]: inclusion of these will be important, in particular, in pA collisions in the forward rapidity region; we will consider these in future studies.

The plan of the paper is as follows: in Sec. II we review the Glasma and the Classical Yang-Mills equations; in Sec. III we review the Wong equations; in Sec. IV we discuss our results for R_{pPb} and R_{PbPb} ; in Sec. V we estimate the effect of energy loss due to gluon radiation; finally, in Sec. VI we draw our conclusions.

II. GLASMA AND CLASSICAL YANG-MILLS EQUATIONS

In this section, we briefly describe the McLerran-Venugopalan (MV) model [1–3,61] which is used to prepare the initial condition known as the Glasma. We mention that in our notation the gauge fields have been rescaled by the QCD coupling $A_\mu \rightarrow A_\mu/g$.

In the MV model, the static color charge densities ρ_a on the nucleus A , the colliding objects, are assumed to be random variables that are normally distributed with zero mean and variance described by the equation

$$\langle \rho_A^a(\mathbf{x}_T) \rho_A^b(\mathbf{y}_T) \rangle = (g^2 \mu_A)^2 \varphi_A(\mathbf{x}_T) \delta^{ab} \delta^{(2)}(\mathbf{x}_T - \mathbf{y}_T); \quad (1)$$

here A corresponds to either the Pb nucleus or the proton, a and b denote the adjoint color index. In this work we limit ourselves, for simplicity, to the case of the $SU(2)$ color group therefore $a, b = 1, 2, 3$. In Eq. (1) $g^2 \mu_A$ indicates the color charge density and it is of the order of the saturation momentum Q_s [62].

The function $\varphi_A(\mathbf{x}_T)$ in Eq. (1) allows for a nonuniform probability distribution of the color charges in the transverse plane. For the Pb nucleus we assume a uniform probability and take $\varphi(\mathbf{x}_T) = 1$. For the proton we use a gaussian $\varphi_A(\mathbf{x}_T)$ that mimics the distribution of color charges obtained after rapidity evolution from the constituent quark model [63–66], namely,

$$\varphi_p(\mathbf{x}_T) = e^{-x_T^2/2B}. \quad (2)$$

The parameter in Eq. (2) is $B = 3 \text{ GeV}^{-2}$.

For the proton $g^2 \mu_p \varphi_p(\mathbf{x}_T)^{1/2}$ can be understood as an \mathbf{x}_T -dependent $g^2 \mu$. Following the result of Ref. [62], we fix $g^2 \mu_p$ for each event assuming that $\langle g^2 \mu_p \varphi_p(\mathbf{x}_T)^{1/2} \rangle / Q_s = 0.57$ where the average is defined with $\varphi_p(\mathbf{x}_T)$ as a weight function, then estimating Q_s at the relevant energy by using

the standard GBW fit [67–69]

$$Q_s^2 = Q_{s,0}^2 \left(\frac{x_0}{x} \right)^\lambda, \quad (3)$$

where $\lambda = 0.277$, $Q_0 = 1 \text{ GeV}$, and $x_0 = 4.1 \times 10^{-5}$. The relevant value of x for the two colliding objects can be obtained at midrapidity as $\langle p_T \rangle / \sqrt{s}$ where $\langle p_T \rangle$ corresponds to the average p_T of the gluons produced in that collision. For example, at the RHIC colliding energy for $x = 0.01$ we find $Q_s = 0.47 \text{ GeV}$ in agreement with the estimate of Ref. [70]. At the LHC energy, $\sqrt{s} = 5.02 \text{ TeV}$, we obtain $Q_s = 0.80 \text{ GeV}$ which gives $\langle g^2 \mu_p \varphi_p(\mathbf{x}_T)^{1/2} \rangle = 1.41 \text{ GeV}$.

For the Pb nucleus the uncertainty on the Q_s and on $g^2 \mu$ comes from the different model used to estimate Q_s for a large nucleus. In this case the GBW fit is modified as

$$Q_s^2 = f(A) Q_{s,0}^2 \left(\frac{x_0}{x} \right)^\lambda, \quad (4)$$

where

$$f(A) = A^{1/3} \quad (5)$$

within a naive scaling hypothesis, and

$$f(A) = cA^{1/3} \log A \quad (6)$$

within the IP-Sat model [71]. While other forms of $f(A)$ are possible [72,73], the two given above indicate the higher and lower value of Q_s at the RHIC energy [62]; therefore, we take these two to set the upper and lower estimate of Q_s . Again using $Q_s/g^2 \mu = 0.57$ we obtain $g^2 \mu_{Pb} = 2 \text{ GeV}$ and $g^2 \mu_{Pb} = 3 \text{ GeV}$ at the RHIC energy taking, respectively, the IP-Sat and naive forms; at the LHC energy $\sqrt{s} = 5.02 \text{ TeV}$ the modified GBW fit then leads to $g^2 \mu_{Pb} = 3.4 \text{ GeV}$ and $g^2 \mu_{Pb} = 5 \text{ GeV}$ for the two cases.

Before going ahead, we remark that the parameters quoted above (that are widespread in the literature), correspond to the color- $SU(3)$ case; however, in this work we analyze for simplicity the evolution of Glasma for the color- $SU(2)$ case. We have checked that if we rescale the $g^2 \mu$ from $SU(3)$ to $SU(2)$, we get an effect on the nuclear modification factors of at most the 10%: given this tiny effect of the rescaling, and to avoid confusion with the existing literature, we use the $SU(3)$ parameters here; in the Appendix we show one result for the $SU(2)$ parameter set.

The static color sources $\{\rho\}$ generate pure gauge fields which in the forward light cone combine and give the initial boost-invariant Glasma fields. To compute these fields we first solve the Poisson equations for the gauge potentials generated by the color charge distribution of the nuclei A and B , namely,

$$-\partial_\perp^2 \Lambda^{(A)}(\mathbf{x}_T) = \rho^{(A)}(\mathbf{x}_T) \quad (7)$$

(a similar equation holds for the color charge distribution belonging to B). Wilson lines are computed as $V^\dagger(\mathbf{x}_T) = e^{i\Lambda^{(A)}(\mathbf{x}_T)}$, $W^\dagger(\mathbf{x}_T) = e^{i\Lambda^{(B)}(\mathbf{x}_T)}$ and the pure gauge fields of the two colliding objects are given by $\alpha_i^{(A)} = iV \partial_i V^\dagger$, $\alpha_i^{(B)} = iW \partial_i W^\dagger$. In terms of these given fields the solution of the CYM in the forward light cone at initial time, namely, the Glasma gauge potential, are given by $A_i = \alpha_i^{(A)} + \alpha_i^{(B)}$ for $i = x, y$ and $A_\eta = 0$, and the initial longitudinal Glasma fields

are [8,9]

$$E^\eta = i \sum_{i=x,y} [\alpha_i^{(B)}, \alpha_i^{(A)}], \quad (8)$$

$$B^\eta = i([\alpha_x^{(B)}, \alpha_y^{(A)}] + [\alpha_x^{(A)}, \alpha_y^{(B)}]), \quad (9)$$

while the transverse fields are vanishing at initial time.

The dynamical evolution of the fields that we study here is given by the Classical Yang-Mills (CYM) equations in the case of a box with longitudinal expansion, which are very well known; see, for example, Refs. [18,20]. The Hamiltonian density is given by

$$H = \text{Tr} \left[\frac{1}{\tau^2} E^i E^i + E^\eta E^\eta + \frac{1}{\tau^2} B^i B^i + B^\eta B^\eta \right], \quad (10)$$

where the trace is over the adjoint color indices and τ denotes the proper time. The equations of motion for the fields and conjugate momenta are

$$E^i = \tau \partial_\tau A_i, \quad (11)$$

$$E^\eta = \frac{1}{\tau} \partial_\tau A_\eta, \quad (12)$$

$$\partial_\tau E^i = \frac{1}{\tau} D_\eta F_{\eta i} + \tau D_j F_{ji}, \quad (13)$$

$$\partial_\tau E^\eta = \frac{1}{\tau} D_j F_{j\eta}. \quad (14)$$

These equations are solved in a 4×4 fm box with periodic boundary conditions in the transverse plane, while rapidity independence is assumed; the lattice spacing is $a = 0.04$ fm.

III. CHARM AND BEAUTY IN THE EVOLVING GLASMA

We initialize the momentum distribution of charm and beauty with the prompt distribution obtained within fixed order + next-to-leading log (FONLL) QCD which describes the D-mesons and B-meson spectra in pp collisions after fragmentation [74–76],

$$\left. \frac{dN}{d^2 p_T} \right|_{\text{prompt}} = \frac{x_0}{(1 + x_3 p_T^{x_1})^{x_2}}; \quad (15)$$

the parameters that we use in the calculations are $x_0 = 20.2837$, $x_1 = 1.95061$, $x_2 = 3.13695$ and $x_3 = 0.0751663$ for c quark; $x_0 = 0.467997$, $x_1 = 1.83805$, $x_2 = 3.07569$, and $x_3 = 0.0301554$ for b quark; the slope of the distribution has been calibrated to a collision at 5.02 TeV. Moreover, we assume that the initial longitudinal rapidity share the same value of space-time rapidity $y = \eta$. Initialization in coordinate space is done as follows: for the pA collisions we use the gaussian weight in Eq. (2) to distribute heavy quarks in the transverse plane, while these are distributed with uniform probability in the rapidity direction; for AA collisions we also distribute heavy quarks with uniform probability in the transverse plane. Finally, we assume that the formation proper time of heavy quark is $\tau_{\text{formation}} = 1/(2m_{HQ}) \approx 0.06$ fm/c for charm quarks having mass $m_c = 1.5$ GeV; for beauty we use $m_b = 4.5$ GeV which gives $\tau_{\text{formation}} \approx 0.02$ fm/c.

The momentum evolution of heavy quarks in the evolving Glasma is studied by the Wong equations [77,78], that for a single heavy quark can be written as

$$\frac{dx_i}{dt} = \frac{p_i}{E}, \quad (16)$$

$$E \frac{dp_i}{dt} = Q_a F_{iv}^a p^v, \quad (17)$$

$$E \frac{dQ_a}{dt} = Q_c \varepsilon^{cba} A_{b\mu} p^\mu, \quad (18)$$

where $i = x, y, z$ and $E = \sqrt{\mathbf{p}^2 + m^2}$ with m is the mass of the heavy quark. In the above equations, the first two equations are the familiar Hamilton equations of motion for the coordinate and its conjugate momentum, while the third equation corresponds to the gauge invariant color current conservation. In the third equation Q_a corresponds to the classical heavy quarks color charge: We initialize this by a uniform distribution with support in the range $(-1, +1)$. For each c, b quark we produce a \bar{c}, \bar{b} quark as well: for this we assume the same initial position of the companion quark, opposite momentum and a new random color charge (color singlet condition of the produced quark-antiquark pair is not a requirement because most heavy quarks are produced via gluon fusion). Solving the Wong equations is equivalent to solve the Boltzmann-Vlasov equations for the heavy quarks in a collisionless plasma, which propagate in the evolving Glasma. In fact, the Boltzmann-Vlasov equation can be solved by means of the test particle method which amounts to solve the classical equations of motion of the particles in the background of the evolving gluon field.

We neglect the effect of the color current carried by the heavy quarks on the gluon field: this approximation is usually used to study the propagation of heavy quarks in a thermal QGP bath and sounds quite reasonable due to the small number of heavy quarks produced by the collision, as well as due to their large mass, both of these factors eventually leading to a negligible color current density. The term on the right hand side of Eq. (17) is responsible of diffusion in momentum space [34,51,79]; besides, we should add a dissipative term, $-E \Gamma p_i$, that takes into account the energy loss due to gluon radiation. We will estimate roughly the effect of gluon radiation in Sec. V in the case of charm quark only, by first computing the diffusion coefficient then using the fluctuation-dissipation theorem to compute the drag which describes the energy loss. We anticipate here that, within our approximations, the effect of the drag is not substantial for the range of p_T that we consider in our study unless the average effective temperature of the gluon bath is very low.

Whether a quark-gluon plasma is formed or not in pA collisions is still an open question. For the pA collisions we assume that all the dynamics can be described by the interaction with the evolving Glasma fields, without the formation of a quark-gluon plasma; eventually heavy quarks hadronize by fragmentation. At the end of this evolution we adopt a standard fragmentation for the heavy quark to D/B-meson




FIG. 1. Color electric fields, measured in lattice units, as a function of proper time for the expanding box (solid lines) and the static box (dashed and dot-dashed lines).

[80], with

$$f(z) \propto \frac{1}{z \left(1 - \frac{1}{z} - \frac{\epsilon_c/\epsilon_b}{1-z}\right)^2}, \quad (19)$$

where $z = p_D/p_c(p_B/p_b)$ is the momentum fraction of the D(B)-meson fragmented from the heavy quark and $\epsilon_c(\epsilon_b)$ is a free parameter to fix the shape of the fragmentation function to reproduce the D/B-meson production in pp collisions [81]. However, for AA collisions we do not perform any fragmentation since we can describe only the evolution in the early stage of the collision: Our result for the spectrum should eventually be fed to a relativistic transport code as an initialization for heavy quarks in the quark-gluon plasma phase as done in Ref. [34].

IV. RESULTS

In this section we present our results. We first show the color fields, the energy density, and we define an effective average temperature of the evolving gluon fields. Then we show the R_{pPb} for D and B mesons and the R_{pPb} for charm and beauty quarks.

A. Evolving color fields

In Fig. 1 we plot the color electric field, measured in lattice units, as a function of proper time; the behavior of the color magnetic field is similar so we do not show it. In the figure, solid green line corresponds to the longitudinal field while the solid green thick line denotes the averaged transverse field, namely $E_T = \sqrt{E_x^2 + E_y^2}/2$. In all cases, the fields have been averaged over the box. For comparison, we also show the fields in the case of the static box. The calculations correspond to the setup for a Pb-Pb collision and $g^2\mu = 3.4$ GeV. We notice that regardless of the longitudinal expansion, E_T forms within 0.1 fm/c in proper time, and initially its magnitude is comparable with that of the static box. At larger proper times, the magnitude of both longitudinal and transverse fields decreases due to the expansion that dilutes the energy density; nevertheless, up to $\tau = 0.4$ fm/c the size of E_T is still




FIG. 2. Average energy density of the gluon field versus proper time, for a Pb-Pb collision with $g^2\mu = 3.4$ GeV (solid maroon line) and $g^2\mu = 5$ GeV (dashed green line). Blue thin line corresponds to τ^{-1} which we draw to emphasize the free streaming that develops for $\tau \gtrsim 0.05$ fm/c.

substantial, being $\approx 40\%$ of the analogous field obtained for the static box. This E_T causes the diffusion of the transverse momentum of heavy quarks.

In Fig. 2 we plot the average energy density of the evolving gluon field, ρ , as a function of the proper time, computed for a Pb-Pb collision with $g^2\mu_{pB} = 3.4$ GeV (solid maroon line) and $g^2\mu = 5$ GeV (dashed green line). In the figure we have also drawn the blue thin line that corresponds to τ^{-1} : it emphasizes the free streaming that develops for $\tau \gtrsim 0.05$ fm/c, corresponding to a vanishing longitudinal pressure [22,33].

For what we will discuss in the next sections, it is useful to define an effective temperature from the energy density, assuming the evolving gluon system can be represented as an ensemble of thermalized gluons with average temperature T and screening mass m_S , namely,

$$\rho = 2(N_c^2 - 1) \int \frac{d^3p}{(2\pi)^3} \frac{\sqrt{p^2 + m_S^2}}{e^{\sqrt{p^2 + m_S^2}/T} - 1}, \quad (20)$$

with N_c the number of colors. From the results shown in Fig. 2 we extract the effective temperature for three values of m_S , and we show the results in Fig. 3 for two values of $g^2\mu$: the green solid lines correspond to $m_S = 1$ GeV, the maroon dashed lines to $m_S = 3$ GeV and the blue dot-dashed lines to $m_S = 5$ GeV. The screening mass can be read from the gauge invariant correlators [33] and it has been found to be of the order of $g^2\mu$: we consider here three values of m_S to check how the predictions depend on m_S . The average temperature is pretty high at initial time, but it reaches $T \approx 1$ GeV within $\tau \approx 0.4$ fm/c for the lowest value of m_S , being a bit higher for larger values of m_S . Clearly choosing a smaller value of m_S would give an effective temperature which matches the initialization temperature of hydro and relativistic transport calculations, $T_{in} \approx 0.6$ GeV.




FIG. 3. Upper panel. Effective temperature versus proper time, computed for a Pb-Pb collision with $g^2\mu = 3.4$ GeV and for $m_S = 1$ GeV (green solid line), $m_S = 3$ GeV (maroon dashed line), $m_S = 5$ GeV (blue dot-dashed line). Lower panel. Same as upper panel, with $g^2\mu = 5$ GeV.

B. Nuclear modification factor for p-Pb collisions

In Fig. 4, we plot the D-meson nuclear suppression factor, R_{pPb} , at different time for D meson and for two different values of $g^2\mu$ in the Pb nucleus, namely, $g^2\mu = 3.4$ GeV (upper panel) and $g^2\mu = 5$ GeV (lower panel). The last is a large value of $g^2\mu$, but still within standard estimates. We notice that anyway the effect on R_{pPb} only mildly depends on the $g^2\mu$. The evolution of charm has been studied via the Wong equations, then at the time shown in the figure we have adopted the standard fragmentation procedure to get the D meson spectra. We observe that the R_{pPb} substantial deviation from one because of the interaction of the charm with the gluon fields. We notice that due to the heavy quark interaction with the gluon fields, low momentum charm quarks are shifted to high momentum. This effect was first observed in Ref. [51] and named as the cathode tube effect. Due to Cathode tube effect we observe an enhancement of the R_{pPb} at intermediate p_T associated to a depletion, a consequence of heavy quark number conservation, at low momenta. In comparison with Ref. [51], in this present work we have studied the effect in a more realistic scenario including the longitudinal expansion. In Fig. 4, we consider $g^2\mu = 3.4$ GeV for the Pb nucleus. Although there is uncertainty on the life time of Glasma




FIG. 4. Nuclear suppression factor ($|y| < 0.2$) for the D meson versus p_T at different times. Upper panel corresponds to $g^2\mu_{Pb} = 3.4$ GeV while in the lower panel $g^2\mu_{Pb} = 5$ GeV. Experimental data correspond to the backward rapidity side of the LHCb collaboration [82].

produced in the p-Pb collisions, we observe that the nuclear suppression factor stops evolving at $\tau \approx 0.3$ fm/c: this is mainly due to the expansion which was missing in the earlier calculations [51]. Such a property of self-quenching in a short time scale is likely to favor a smooth transition into the subsequent evolution of the system in the QGP. In the lower panel of Fig. 4, we plot the D-meson nuclear suppression factor in Glasma corresponding to $g^2\mu_{Pb} = 5$ GeV. Larger value of $g^2\mu_{Pb}$ corresponds to more color charges, hence, large energy density. This leads to the larger enhancement at intermediate momentum. The fact that the shape of our R_{pPb} can reproduce that measured in experiments, might suggest that at least part of the measured R_{pPb} comes from the propagation of the charm quarks in the Glasma, and might be considered as the signature of the Glasma itself.

The net effect that we find on charm spectra seems different from what is usually observed in a quark-gluon plasma medium, in which the nuclear modification factor is larger than one at low p_T then it decreases monotonically smaller at larger p_T . However, it should be considered that in the early stage the energy density of the gluon bath is very large,




FIG. 5. Nuclear suppression factor of charm versus p_T for a Pb-Pb collision, at different times. Upper panel corresponds to $g^2\mu_{Pb} = 3.4$ GeV while in the lower panel $g^2\mu_{Pb} = 5$ GeV.

therefore it is natural that low- p_T heavy quarks diffuse to higher p_T and energy loss plays a minor role here. We will show in the next section that including a term that describes energy loss by radiation, the main effect will be still the diffusion from low to high p_T unless we keep the average energy density of the gluon bath very low in the full evolution, comparable with that of the quark-gluon plasma at the initialization time. This is in agreement with Ref. [79] where it has been shown that assuming a fluctuation dissipation theorem, the diffusion of the heavy particle dominates over the energy loss if the initial momentum of the heavy particle is smaller than the average energy of the bulk.

C. Nuclear modification factor for Pb-Pb collisions

In nucleus-nucleus collisions the Glasma phase can act as the pre-equilibrium phase before the QGP phase. Recently the impact of initial static Glasma phase on heavy quark dynamics has been reported in Ref. [34]. It is observed that the dynamics in evolving Glasma stage leads to a larger final elliptic flow (v_2) inducing a relation between R_{AA} and v_2 that is quite close to the experimental measurements. Keeping its significance in mind, we extended our calculation to nucleus-nucleus collisions.




FIG. 6. Nuclear suppression factor of beauty versus p_T for a Pb-Pb collision, at different times. Upper panel corresponds to $g^2\mu_{Pb} = 3.4$ GeV while in the lower panel $g^2\mu_{Pb} = 5$ GeV.

In Fig. 5, we plot R_{PbPb} of charm in evolving Glasma for $g^2\mu_{Pb} = 3.4$ GeV (upper panel) and $g^2\mu_{Pb} = 5$ GeV (lower panel) at different proper times. The effect on the charm spectrum is much stronger in nucleus-nucleus collisions in comparison with p-nucleus collisions: this is due to the larger energy density produced in nucleus-nucleus collisions than p-nucleus collisions. After the evolution in the initial gluon fields, the quark-gluon plasma phase should be considered and our result should serve as the initial condition of the charm spectrum for the QGP phase evolution. Once again we observe a saturation of the R_{PbPb} at $\tau \approx 0.3$ fm/c. It is remarkable that although expansion is present, the impact of initial evolution in the gluon field is still quite substantial in comparison with the earlier results in which expansion was not considered [34].

We further extend our model to study the evolution of beauty. In Fig. 6, we present the R_{PbPb} of beauty for the aforementioned two values of $g^2\mu_{Pb}$. The trend we observe for beauty is the same we have found for charm: Quantitatively the effect on the spectrum of beauty is much milder, of course, because of the larger mass of this quark compared to that of charm. Overall, the effect on beauty is still significant and might impact the later dynamics in the QGP phase [34].




FIG. 7. Transverse momentum variance as a function of proper time, for a Pb-Pb collision. Indigo diamonds correspond to $g^2\mu = 3.4$ GeV while green squares to $g^2\mu = 5$ GeV. Dashed lines denote the linear fit Eq. (22).

V. ESTIMATE OF THE RADIATION REACTION

In the previous section we have shown the results obtained neglecting the energy loss of the heavy quarks: This can be modeled by modifying Eq. (17) as,

$$E \frac{dp_i}{dt} = Q_a F_{iv}^a p^v - E \Gamma p_i, \quad (21)$$

where Γ is the drag coefficient. In this section we give a rough estimate of the radiation reaction, then evaluate its effect on the nuclear modification factors. We limit ourselves to show the results obtained for the charm quark since those for the beauty quark are similar. Our strategy is to first evaluate an average diffusion coefficient for the transverse momentum, D , of charm, defined via the equation

$$\sigma^2 \equiv \langle (p_T - \langle p_T \rangle)^2 \rangle = 2D\tau + \text{constant}, \quad (22)$$

where the average is taken with the charm spectrum, then assume that the fluctuation-dissipation theorem (FDT),

$$D = \Gamma E T, \quad (23)$$

relates D to Γ with E corresponding to the kinetic energy of the heavy quark. A similar assumption between drag and diffusion coefficients, although via a more rigorous implementation, has been already adopted in Ref. [50]. One major drawback of this assumption is that the FDT is rigorously valid only for a bath in thermal equilibrium, which is a condition that is not satisfied by the evolving gluon field considered here. However, the calculation of Γ without a FD theorem would require the calculation of the gluon radiation from quarks propagating in a random gluon field, which is beyond the scope of this study. Therefore, we follow [50] and we chose to estimate the drag coefficient by Eq. (23) and replacing T with the average temperature extracted from the energy density, see Fig. 2 and Eq. (20).

In Fig. 7 we plot the variance of the transverse momentum for the setup of a Pb-Pb collision. The result has been obtained by solving the equations of motion of charm in the evolving gluonic background, Eq. (17), which takes into account the




FIG. 8. Comparison of the nuclear modification factor for the setup of Pb-Pb collisions, obtained neglecting and considering the radiation reaction of the charm quark. Upper panel corresponds to $g^2\mu = 3.4$ GeV and three screening masses: $m_S = 1$ GeV (solid orange line), $m_S = 3$ GeV (maroon dot-dashed line), and $m_S = 5$ GeV (dotted green line). For comparison we also show the result without drag, see the blue dashed line. Lower panel shows the results $g^2\mu = 5$ GeV and the same values of the screening masses.

diffusion only. In the figure, the green squares correspond to our calculation for $g^2\mu = 5$ GeV while the indigo diamonds denote the results for $g^2\mu = 3.4$ GeV; the dashed lines correspond to the linear fit Eq. (22). We use Eq. (22) to fit the data in Fig. 7 and we find $D = 0.21$ GeV²/fm for $g^2\mu = 3.4$ GeV, while $D = 0.35$ GeV²/fm for $g^2\mu = 5$ GeV. We notice that these numbers are in the same ballpark of the pQCD estimates for an average $p_T \approx 1.5$ –2 GeV in the temperature range 0.4–1 GeV [48]. Using these results in Eq. (23) we estimate Γ for each charm and use it in Eq. (21).

In Fig. 8 we plot the nuclear modification factor computed with the drag force estimated by the FDT, using three different screening masses of the bath. For comparison we also show R_{PbPb} computed without taking the drag into account, see the blue dashed line. Upper panel corresponds to $g^2\mu = 3.4$ GeV, lower panel to $g^2\mu = 5$ GeV. In both panels we show the results obtained for $m_S = 1$ GeV (solid orange line), $m_S = 3$ GeV (maroon dot-dashed line), and $m_S = 5$ GeV (dotted green line). As expected, the smaller the average T is, the larger the effect of the drag. On the one hand, these results confirm that as long as the energy density of the gluon medium remains large in the initial stage, the drag force can be ne-

glected for low- p_T heavy quarks; on the other hand, when the energy density becomes smaller the drag might be important so energy loss by gluon radiation might play a role even in the very early stage of the collision. This last circumstance pushes us to study the problem of gluon radiation in more detail, and we leave a detailed study to a future project.

VI. CONCLUSIONS AND OUTLOOK

We have studied the diffusion of charm and beauty quarks by means of the Wong equations in the evolving strong gluon fields produced in the early stage of high-energy nuclear collisions. For the sake of computational simplicity we have addressed the problem in the color- $SU(2)$ case. Initialization of the gluon field has been achieved via the Glasma, carrying longitudinal color-electric and color-magnetic fields thus neglecting fluctuations. For the heavy quark initialization we have considered the standard FONLL perturbative production tuned to reproduce the D and B meson spectrum in proton-proton collisions. We have setup the saturation scale for both the proton and the Pb nucleus to reproduce the expected one at $\sqrt{s} = 5.02$ TeV collisions. The novelties in comparison with previous works [34,51] are that have added the longitudinal expansion of the gluon medium (in Ref. [51] we studied the diffusion in a static box), and performed a systematic comparative study of pA and AA collisions that is lacking in the literature. In particular, we have focused on the nuclear modification factors, R_{pA} and R_{AA} , showing how this is affected by the interaction of the heavy quarks with the gluon medium. We have also roughly estimated the effect of the radiation reaction on R_{AA} , by using a fluctuation-dissipation-theorem-like relation between the drag and the diffusion coefficient, leaving a complete study to a future project. We have found that the spectrum of charm and beauty is tilted toward higher p_T because of the interaction with the gluon fields in the early stage: This effect was named the cathode tube in Ref. [51].

We have first computed R_{pPb} . The main effect of the interaction of the charm quarks with the gluon field is to shift the low p_T charm quarks to high p_T states. This ensured an enhancement of the charm quarks yields in the intermediate p_T . We have found that the shape of our R_{pPb} is able to reproduce that measured by the LHCb collaboration on the proton side where the effect of shadowing is marginal. Since in our calculation the shape comes directly from the propagation of the charm quarks in the evolving Glasma fields, we suggest that at least part of the measured R_{pPb} is the signature of the Glasma formed in high-energy collisions. We have repeated the calculation for the beauty quarks, finding qualitatively similar results to those of the charm quarks. We have then studied R_{PbPb} for both charm and beauty. We have found that even when the longitudinal expansion is included, the impact of the early stage is substantial both for charm and beauty. It is not possible to compare R_{PbPb} coming from the early stage with experimental data, since the late evolution in the quark-gluon plasma cannot be neglected. However, even if the evolution in the gluon fields happens at very early times, this can be related to the observables in Pb-Pb collisions [34].

In most of the present work we have neglected the effect of gluon radiation on the motion of the heavy quarks. The inclusion of this is a highly nontrivial step, due to the fact that any estimate of the radiation based on pQCD should take into account that charm and beauty evolve in a random gluon field and that gluons are screened with a screening mass $m_D \sim g^2 \mu$. Instead of attacking this problem from the pQCD point of view, we have used a relation inspired by the fluctuation-dissipation theorem to relate the drag force responsible of the energy loss, to an averaged diffusion coefficient. The numerical calculation of the latter presents no difficulty within our approach since it is possible to follow the evolution of $\langle (p_T - \langle p_T \rangle)^2 \rangle$ of heavy quarks with time, thus allowing us to define a diffusion coefficient for the transverse momentum. We have computed the drag coefficient assuming the evolving gluon bath is described by a gas of massive gluons thermalized at an average temperature, T , which can be estimated within our code from the average energy density. We have found that the effect of the drag coefficient, that was neglected in previous calculations [34,51], is certainly present but does not seem enough to cancel the cathode tube effect as long as the average temperature remains $T \approx 1$ GeV, as already anticipated in Ref. [79]. These results are encouraging since they show that neglecting energy loss, we overestimate the effect of diffusion on R_{AA} only when the system approaches the initialization time of hydro, while in the very early stage the purely diffusive motion is a fair approximation. Anyway, a more thorough study of the gluon radiation has to be pursued and this will be the subject of a forthcoming publication.

We suggest that the evolution of heavy quarks in the early stages of high-energy nuclear collisions is important as it can alter the FFNLO initialization commonly used to study the heavy quarks dynamics in these collisions. We remark that a simultaneous description of heavy quark R_{AA} and v_2 is a top challenge to almost all the models on heavy quark dynamics: the inclusion of a pre-equilibrium phase might improve the situation, offering a better understanding of this puzzle [34]. This present work will further boost the phenomenology as well as our understanding of the experimental results.

ACKNOWLEDGMENTS

M. R. acknowledges Navid Abbasi, Gabriele Coci, Gianluca Giuliani, Feng Li, David Mueller, John Petrucci, Yifeng Sun, Shen-Song Wan and Bo-Nan Zhang for inspiration, discussions and comments. The work of M.R. and S.K.D. is supported by the National Science Foundation of China (Grants No. 11805087 and No. 11875153) and by the Fundamental Research Funds for the Central Universities (Grant No. 862946). The work of J. H. Liu is supported by China Scholarship Council (Scholarship No. 201806180032).

APPENDIX: SCALING TO $SU(2)$ PARAMETERS

In this Appendix we present the result for R_{pPb} we obtain if we rescale the $g^2 \mu$ of the MV model from the $SU(3)$ value to the $SU(2)$ one: our goal is to show that this rescaling does not affect substantially the modification factors computed in the article.




FIG. 9. Comparison of the nuclear modification factor for the setup of p-Pb collisions, obtained for the $g^2\mu$ of $SU(3)$ used in the main text, and for the $g^2\mu$ rescaled to the $SU(2)$ case.

To rescale the $g^2\mu$, we take the saturation momentum, Q_s , as independent on N_c , since Q_s depends on the thickness function of a nucleus therefore it is sensitive to how nucleons are distributed, while the number of colors has to do with the substructure of the nucleons. However, the $g^2\mu$ measures the density of the color charges at the quark-gluon level therefore it is sensitive to N_c . In the literature, a relation between the Q_s and $g^2\mu$ can be found in Ref. [83], which shows an explicit dependence on N_c that neglecting logarithm is $Q_s^2/(g^2\mu)^2 \propto N_c$. Leaving the Q_s fixed, we change the ratio $g^2\mu/Q_s$ by using the aforementioned relation: in particular, this scaling from 3 to 2 colors brings to $g^2\mu_{SU(2)} \approx 1.22 \times g^2\mu_{SU(3)}$. Therefore, the $g^2\mu$ has to be increased of approximately the 20%.

In Fig. 9 we compare the R_{pPb} obtained with the two values of the parameter: the $N_c = 3$ case, which we use in the main text, and the $N_c = 2$ obtained rescaling the $g^2\mu$ as outlined above. We find that the net effect is less than 10%. We have checked that this modest shift occurs also in the other cases studied in the article.

- [1] L. D. McLerran and R. Venugopalan, *Phys. Rev. D* **49**, 2233 (1994).
- [2] L. D. McLerran and R. Venugopalan, *Phys. Rev. D* **49**, 3352 (1994).
- [3] L. D. McLerran and R. Venugopalan, *Phys. Rev. D* **50**, 2225 (1994).
- [4] F. Gelis, E. Iancu, J. Jalilian-Marian, and R. Venugopalan, *Ann. Rev. Nucl. Part. Sci.* **60**, 463 (2010).
- [5] E. Iancu and R. Venugopalan, in *Quark Gluon Plasma*, edited by R. C. Hwa (World Scientific, Singapore, 2004), pp. 249–363.
- [6] L. McLerran, [arXiv:0812.4989](https://arxiv.org/abs/0812.4989).
- [7] F. Gelis, *Int. J. Mod. Phys. A* **28**, 1330001 (2013).
- [8] A. Kovner, L. D. McLerran, and H. Weigert, *Phys. Rev. D* **52**, 6231 (1995).
- [9] A. Kovner, L. D. McLerran, and H. Weigert, *Phys. Rev. D* **52**, 3809 (1995).
- [10] M. Gyulassy and L. D. McLerran, *Phys. Rev. C* **56**, 2219 (1997).
- [11] T. Lappi and L. McLerran, *Nucl. Phys. A* **772**, 200 (2006).
- [12] R. J. Fries, J. I. Kapusta and Y. Li, [arXiv:nucl-th/0604054](https://arxiv.org/abs/nucl-th/0604054).
- [13] G. Chen, R. J. Fries, J. I. Kapusta and Y. Li, *Phys. Rev. C* **92**, 064912 (2015).
- [14] A. Krasnitz and R. Venugopalan, *Phys. Rev. Lett.* **86**, 1717 (2001).
- [15] A. Krasnitz, Y. Nara, and R. Venugopalan, *Phys. Rev. Lett.* **87**, 192302 (2001).
- [16] A. Krasnitz, Y. Nara, and R. Venugopalan, *Nucl. Phys. A* **727**, 427 (2003).
- [17] K. Fukushima, F. Gelis, and L. McLerran, *Nucl. Phys. A* **786**, 107 (2007).
- [18] H. Fujii, K. Fukushima, and Y. Hidaka, *Phys. Rev. C* **79**, 024909 (2009).
- [19] K. Fukushima, *Phys. Rev. C* **89**, 024907 (2014).
- [20] P. Romatschke and R. Venugopalan, *Phys. Rev. Lett.* **96**, 062302 (2006).
- [21] P. Romatschke and R. Venugopalan, *Phys. Rev. D* **74**, 045011 (2006).
- [22] K. Fukushima and F. Gelis, *Nucl. Phys. A* **874**, 108 (2012).
- [23] H. Iida, T. Kunihito, A. Ohnishi, and T. T. Takahashi, [arXiv:1410.7309](https://arxiv.org/abs/1410.7309).
- [24] T. Epelbaum and F. Gelis, *Phys. Rev. Lett.* **111**, 232301 (2013).
- [25] T. Epelbaum and F. Gelis, *Phys. Rev. D* **88**, 085015 (2013).
- [26] R. Ryblewski and W. Florkowski, *Phys. Rev. D* **88**, 034028 (2013).
- [27] M. Ruggieri, A. Puglisi, L. Oliva, S. Plumari, F. Scardina, and V. Greco, *Phys. Rev. C* **92**, 064904 (2015).
- [28] N. Tanji and K. Itakura, *Phys. Lett. B* **713**, 117 (2012).
- [29] J. Berges and S. Schlichting, *Phys. Rev. D* **87**, 014026 (2013).
- [30] J. Berges, K. Boguslavski, S. Schlichting, and R. Venugopalan, *Phys. Rev. D* **89**, 114007 (2014).
- [31] J. Berges, K. Boguslavski, S. Schlichting, and R. Venugopalan, *J. High Energy Phys.* **05** (2014) 054.
- [32] J. Berges, K. Boguslavski, S. Schlichting, and R. Venugopalan, *Phys. Rev. D* **89**, 074011 (2014).
- [33] M. Ruggieri, J. H. Liu, L. Oliva, G. X. Peng, and V. Greco, *Phys. Rev. D* **97**, 076004 (2018).
- [34] Y. Sun, G. Coci, S. K. Das, S. Plumari, M. Ruggieri, and V. Greco, *Phys. Lett. B* **798**, 134933 (2019).
- [35] F. Prino and R. Rapp, *J. Phys. G* **43**, 093002 (2016).
- [36] A. Andronic *et al.*, *Eur. Phys. J. C* **76**, 107 (2016).
- [37] R. Rapp *et al.*, *Nucl. Phys. A* **979**, 21 (2018).
- [38] G. Aarts *et al.*, *Eur. Phys. J. A* **53**, 93 (2017).
- [39] X. Dong and V. Greco, *Prog. Part. Nucl. Phys.* **104**, 97 (2019).
- [40] S. Cao *et al.*, *Phys. Rev. C* **99**, 054907 (2019).
- [41] S. K. Das, S. Plumari, S. Chatterjee, J. Alam, F. Scardina, and V. Greco, *Phys. Lett. B* **768**, 260 (2017).
- [42] S. K. Das, F. Scardina, S. Plumari, and V. Greco, *Phys. Lett. B* **747**, 260 (2015).
- [43] S. K. Das, M. Ruggieri, F. Scardina, S. Plumari, and V. Greco, *J. Phys. G* **44**, 095102 (2017).
- [44] S. K. Das, M. Ruggieri, S. Mazumder, V. Greco, and J. e. Alam, *J. Phys. G* **42**, 095108 (2015).
- [45] A. Beraudo, A. De Pace, M. Monteno, M. Nardi, and F. Prino, *J. High Energy Phys.* **03** (2016) 123.
- [46] Y. Xu, S. Cao, G. Y. Qin, W. Ke, M. Nahrgang, J. Auvinen, and S. A. Bass, *Nucl. Part. Phys. Proc.* **276-278**, 225 (2016).

- [47] V. Ozvenchuk, J. Aichelin, P. B. Gossiaux, B. Guiot, M. Nahrgang, and K. Werner, *J. Phys. Conf. Ser.* **779**, 012033 (2017).
- [48] S. K. Das, F. Scardina, S. Plumari, and V. Greco, *Phys. Rev. C* **90**, 044901 (2014).
- [49] V. Chandra and S. K. Das, *Phys. Rev. D* **93**, 094036 (2016).
- [50] S. Mrowczynski, *Eur. Phys. J. A* **54**, 43 (2018).
- [51] M. Ruggieri and S. K. Das, *Phys. Rev. D* **98**, 094024 (2018).
- [52] M. Ruggieri and S. K. Das, *EPJ Web Conf.* **192**, 00017 (2018).
- [53] S. Plumari, V. Minissale, S. K. Das, G. Coci, and V. Greco, *Eur. Phys. J. C* **78**, 348 (2018).
- [54] B. B. Abelev *et al.* (ALICE Collaboration), *Phys. Rev. Lett.* **113**, 232301 (2014).
- [55] K. J. Eskola, H. Paukkunen, and C. A. Salgado, *J. High Energy Phys.* 0904 (2009) 065.
- [56] A. H. Rezaeian, *Phys. Lett. B* **718**, 1058 (2013).
- [57] H. Fujii and K. Watanabe, *Nucl. Phys. A* **920**, 78 (2013).
- [58] B. Ducloué, T. Lappi, and H. Mäntysaari, *Phys. Rev. D* **91**, 114005 (2015).
- [59] J. L. Albacete *et al.*, *Int. J. Mod. Phys. E* **22**, 1330007 (2013).
- [60] J. L. Albacete *et al.*, *Int. J. Mod. Phys. E* **25**, 1630005 (2016).
- [61] Y. V. Kovchegov, *Phys. Rev. D* **54**, 5463 (1996).
- [62] T. Lappi, *Eur. Phys. J. C* **55**, 285 (2008).
- [63] B. Schenke and R. Venugopalan, *Phys. Rev. Lett.* **113**, 102301 (2014).
- [64] B. Schenke, S. Schlichting, and R. Venugopalan, *Phys. Lett. B* **747**, 76 (2015).
- [65] H. Mäntysaari, B. Schenke, C. Shen, and P. Tribedy, *Phys. Lett. B* **772**, 681 (2017).
- [66] H. Mäntysaari and B. Schenke, *Phys. Rev. D* **94**, 034042 (2016).
- [67] K. J. Golec-Biernat and M. Wusthoff, *Phys. Rev. D* **60**, 114023 (1999).
- [68] K. J. Golec-Biernat and M. Wusthoff, *Phys. Rev. D* **59**, 014017 (1998).
- [69] Y. V. Kovchegov and E. Levin, *Quantum Chromodynamics at High Energy*, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **33** (2012).
- [70] J. L. Albacete, A. Dumitru, H. Fujii, and Y. Nara, *Nucl. Phys. A* **897**, 1 (2013).
- [71] H. Kowalski, T. Lappi, and R. Venugopalan, *Phys. Rev. Lett.* **100**, 022303 (2008).
- [72] N. Armesto, C. A. Salgado, and U. A. Wiedemann, *Phys. Rev. Lett.* **94**, 022002 (2005).
- [73] A. Freund, K. Rummukainen, H. Weigert, and A. Schafer, *Phys. Rev. Lett.* **90**, 222002 (2003).
- [74] M. Cacciari, M. Greco, and P. Nason, *J. High Energy Phys.* 9805 (1998) 007; M. Cacciari, S. Frixione, and P. Nason, *ibid.* 0103 (2001) 006.
- [75] M. Cacciari, S. Frixione, N. Houdeau, M. L. Mangano, P. Nason, and G. Ridolfi, *J. High Energy Phys.* 10 (2012) 137.
- [76] M. Cacciari, M. L. Mangano, and P. Nason, *arXiv:1507.06197*.
- [77] S. K. Wong, *Nuovo Cim. A* **65**, 689 (1970).
- [78] A. D. Boozer, *Am. J. Phys.* **79**, 925 (2011).
- [79] M. Ruggieri, M. Frasca, and S. K. Das, *Chin. Phys. C* **43**, 094105 (2019).
- [80] C. Peterson, D. Schlatter, I. Schmitt, and P. M. Zerwas, *Phys. Rev. D* **27**, 105 (1983).
- [81] F. Scardina, S. K. Das, V. Minissale, S. Plumari, and V. Greco, *Phys. Rev. C* **96**, 044905 (2017).
- [82] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* 10 (2017) 090.
- [83] T. Lappi, *J. Phys. G* **35**, 104052 (2008).