

**Coriolis coupling effects in proton-pickup spectroscopic factors from  $^{12}\text{B}$** A. O. Macchiavelli , H. L. Crawford , R. M. Clark, P. Fallon, I. Y. Lee , C. Morse, C. M. Campbell, M. Cromaz, and C. Santamaria*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

J. Chen

*National Superconducting Cyclotron Laboratory, East Lansing, Michigan 48824, USA*

C. R. Hoffman and B. P. Kay

*Physics Division, Argonne National Laboratory, Argonne, Illinois 60438, USA*

(Received 10 December 2020; accepted 4 February 2021; published 8 March 2021)

Spectroscopic factors to low-lying negative-parity states in  $^{11}\text{Be}$  extracted from the  $^{12}\text{B}(d, ^3\text{He})^{11}\text{Be}$  proton-removal reaction are interpreted within the rotational model. Earlier predictions of the  $p$ -wave proton-removal strengths in the strong-coupling limit of the Nilsson model underestimated the spectroscopic factors to the  $3/2^-$  and  $5/2^-$  states and suggested that deviations in the  $1^+$  ground state of the odd-odd  $^{12}\text{B}$  due to Coriolis coupling should be further explored. In this work we use the particle rotor model to take into account these effects and obtain a good description of the level scheme in  $^{11}\text{B}$ , with a moderate  $K$  mixing of the proton Nilsson levels  $[110]1/2$  and  $[101]3/2$ . This mixing, present in the  $1^+$  bandhead of  $^{12}\text{B}$ , is key to explaining the proton-pickup data.

DOI: [10.1103/PhysRevC.103.034307](https://doi.org/10.1103/PhysRevC.103.034307)**I. INTRODUCTION**

Shortly after Bohr and Mottelson's development of the collective model [1], Morinaga showed that the spectroscopy of a number of nuclei in the  $p$  shell could be interpreted in terms of rotational bands [2]. Perhaps one of the most obvious examples is that of  $^8\text{Be}$ , as evident from the ground-state rotational band and its enhanced  $B(E2)$  transition probability [3]. The strong  $\alpha$  clustering in  $^8\text{Be}$  naturally suggests that deformation degrees of freedom play a role in the structure of the Be isotopes, a topic that has been extensively discussed in the literature (see Ref. [4] for a review).

Turning from a collective model to a shell model picture, very nearby to  $^8\text{Be}$ , the lightest example of a so-called island of inversion [5,6] is that at  $N = 8$ , where the removal of  $p_{3/2}$  protons from  $^{14}\text{C}$  results in a quenching of the  $N = 8$  shell gap [7–10]. This is evinced in many experimental observations, including the sudden drop of the  $E(2^+)$  energy in  $^{12}\text{Be}$  relative to the neighboring even-even isotopes and the change of the ground state of  $^{11}\text{Be}$  from the expected  $1/2^-$  to the observed positive-parity  $1/2^+$  state.

Connecting the collective and single-particle descriptions, Bohr and Mottelson [11] actually proposed that the shell inversion could be explained as a result of the convergence of the up-sloping  $[101]_{\frac{1}{2}}$  and down-sloping  $[220]_{\frac{1}{2}}$  Nilsson [12,13] levels with deformation as seen in Fig. 1. Building on these arguments, Hamamoto and Shimoura [14] explained energy levels and available electromagnetic data on  $^{11}\text{Be}$  and  $^{12}\text{Be}$  in terms of single-particle motion in a deformed poten-

tial. It is remarkable that the concept of a deformed rotating structure appears to be applicable, even when the total number of nucleons is small as in the case of light nuclei. In fact, level energies and electromagnetic properties that follow the characteristic rotational patterns emerge for  $p$ -shell nuclei in *ab initio* no-core configuration interaction calculations [15].

In a series of articles we have recently applied the collective model to understand the structure of nuclei in the  $N = 8$  island of inversion and spectroscopic factors obtained from direct nucleon addition and removal reactions [16–18]. The mean-field description seems to capture the main physics ingredients and provides a satisfactory explanation of spectroscopic data, in a simple and intuitive manner.

Here we extend this approach to discuss the results of a recent study of the  $^{12}\text{B}(d, ^3\text{He})^{11}\text{Be}$  reaction [19] in terms of the particle rotor model (PRM) [13,20,21]. Estimates of spectroscopic factors in the strong-coupling limit, given in Ref. [19], underestimated the experimental data and pointed out that Coriolis effects in the structure of the  $1^+$  ground state in  $^{12}\text{B}$  should be taken into account, for which the PRM framework is the framework of choice. We present this analysis here.

**II. THE GROUND STATE OF  $^{12}\text{B}$** 

In order to assess the structure of the ground state of  $^{12}\text{B}$ , we consider first the odd-neutron and odd-proton low-lying negative-parity states in  $^{11}\text{Be}$  and  $^{11}\text{B}$ , respectively. Considering  $^{10}\text{Be}$  as a core, an inspection of the Nilsson diagram [12]




FIG. 1. Nilsson levels relevant for the structure of negative-parity neutron states in  $^{11}\text{Be}$  and  $^{11}\text{B}$  (solid lines). Also shown (dashed lines) are the levels originating from the  $d_{5/2}$  spherical orbital. The shaded band indicates the anticipated range of  $\epsilon_2$  deformation for these nuclei and the horizontal lines indicate the approximate Fermi levels of the odd neutron (blue) and the odd proton (red). Energies are in units of the harmonic oscillator frequency,  $\hbar\omega_0$ .

in Fig. 1 suggests that for  $N = 7$  the last neutron is expected to occupy the  $[101]_{\frac{1}{2}}$  level, and for  $Z = 5$  the last proton will occupy the  $[101]_{\frac{3}{2}}$  level. We have used the standard parameters,  $\kappa = 0.12$  and  $\mu = 0.0$  [22], and adopted a deformation,  $\epsilon_2 \approx 0.45$  ( $\beta_2 \approx 0.6$ ), where the crossing of the  $[220]_{\frac{1}{2}}$  and  $[101]_{\frac{1}{2}}$  levels is expected to occur, explaining the inversion of the  $1/2^+$  and the  $1/2^-$  states in  $^{11}\text{Be}$ .

We have previously discussed the positive-parity states in  $^{11}\text{Be}$  as arising from the strongly coupled  $[220]_{\frac{1}{2}}$  state [17], an assignment supported by the calculated gyromagnetic factor  $g_K = -2.79$ , with decoupling and magnetic-decoupling parameters for the ground state of  $a = 1.93$  and  $b = -1.27$ , respectively. The low-lying negative-parity states, namely, the  $1/2^-$ ,  $3/2^-$ , and  $5/2^-$  states, can be assigned to a  $K = 1/2^-$  strongly coupled band built on the neutron  $[101]_{\frac{1}{2}}$  level, with a decoupling parameter,  $a = 0.5$ , in line with the Nilsson predictions. Further, the  $3/2^-$  state originates from a neutron hole in the  $[101]_{\frac{3}{2}}$  level.

The case of  $^{11}\text{B}$  is somewhat different and more complex, requiring an explicit consideration of Coriolis coupling. An attempt to fit the energies of the *yrast* negative-parity states,  $3/2^-$ ,  $5/2^-$ ,  $7/2^-$ ,  $\dots$ , to leading order,

$$E(I) = E_K + AI(I+1) + BI^2(I+1)^2 + \dots, \quad (1)$$

requires an additional term [11] arising from the Coriolis interaction that induces  $\Delta K = \pm 2K$  mixing:

$$\Delta E_{\text{rot}} = (-1)^{I+K} A_{2K} \frac{(I+K)!}{(I-K)!}, \quad (2)$$

giving  $A = 978$  keV,  $B = -17$  keV, and  $A_3 \approx 20$  keV. Therefore we carried out a PRM calculation [21]—the results, shown in Fig. 2, are in good agreement with the experimental




FIG. 2. Left: The experimental level scheme of  $^{11}\text{B}$  from Ref. [3]. Right: Results of the PRM calculations. Energies are in keV.

level scheme. Here we briefly discuss the physical inputs to the PRM calculation. We include the three orbits in Fig. 1 with the Fermi level  $\lambda$  and the pairing gap  $\Delta$  obtained from a BCS calculation using a coupling constant,  $G = 1.9$  MeV. The solution gives  $\lambda = 45.73$  MeV and  $\Delta = 3.3$  MeV. The adjusted rotational constant of the core<sup>1</sup> corresponds to a moment of inertia,  $\mathcal{J} = 0.57 \hbar^2/\text{MeV}$ , approximately 60% of the rigid body value and consistent with the Migdal estimate [23] for  $A = 11$  and the deformation and pairing parameters above. A fit of Eq. (2) to the PRM results gives  $A_3 \approx 25$  keV.

The Coriolis  $K$  mixing in  $^{11}\text{B}$  gives rise to wave functions of the form

$$\psi_I = \sum_K \mathcal{A}_{IK} |IK\rangle \quad (3)$$

in the strong-coupled basis spanned by the intrinsic proton states  $[110]_{\frac{1}{2}}$ ,  $[101]_{\frac{3}{2}}$ , and  $[101]_{\frac{1}{2}}$ . The percent contributions (squared amplitudes) of each intrinsic proton state for the states shown in Fig. 2 are given in Table I. The results of the PRM calculations indicate a moderate  $K$  mixing for the  $3/2^-$ ,  $5/2^-$ , and  $7/2^-$  states and the  $3/2^-$  states with dominant components of the  $[101]_{\frac{3}{2}}$  and  $[110]_{\frac{1}{2}}$  Nilsson levels in each case. The  $1/2^-$  state is essentially a pure  $[101]_{\frac{1}{2}}$  state. Note that the form of the Coriolis matrix elements favors the mixing of the Nilsson levels with  $p_{3/2}$  parentage. It is also worthwhile noting that, due to the fact that the Fermi level of

<sup>1</sup>In the PRM, we do not include a  $BI^2(I+1)^2$  term in the rotational energy but set the  $E(2^+)_{\text{core}} = A2(2+1) + B[2(2+1)]^2$  from Eq. (1).

TABLE I. Coriolis mixing amplitudes of the proton states in the PRM calculations for  $^{11}\text{B}$ .

$I^\pi$	Energy (MeV)	$\mathcal{A}_{IK}^2$ (%)		
		$[110]_{\frac{1}{2}}$	$[101]_{\frac{3}{2}}$	$[101]_{\frac{1}{2}}$
$\frac{3}{2}^-$	0.00	18	80	2
$\frac{1}{2}^-$	2.12	4	0	96
$\frac{5}{2}^-$	4.21	6	90	4
$\frac{3}{2}^-$	5.34	81	19	0
$\frac{7}{2}^-$	6.66	45	52	3

the odd proton in  $^{11}\text{B}$  is lower than that of the odd neutron in  $^{11}\text{Be}$ , there is no parity inversion in  $^{11}\text{B}$  and the lowest positive-parity state,  $1/2^+$ , lies at  $\approx 4.6$  MeV relative to the  $1/2^-$ .

In addition to the reproduction of the energy levels, the calculated magnetic moment of the ground state is  $\mu_{3/2^-} = 2.66 \mu_N$  to be compared to the experimental value of  $2.688\ 648\ 9(10) \mu_N$  [24].

There is, however, an intriguing discrepancy with the measured  $Q_{3/2^-} = 0.040\ 65(26) e\ b$  [24,25], which is consistent with the leading-order collective model estimate of  $0.04 e\ b$ , but the PRM result of  $0.028 e\ b$  is smaller due to the mixing of the two  $I^\pi = 3/2^-$  states with  $K^\pi = 1/2^-$  and  $K^\pi = 3/2^-$  that have  $Q$ 's of opposite signs. While one may be tempted to explain this with a larger deformation,  $\epsilon_2 \approx 0.60$  (as in Ref. [14]), it would be at the expense of losing the agreement in the energy levels. This is also the case for the mirror nucleus  $^{11}\text{C}$ . We do not have an explanation for this discrepancy except to speculate that, perhaps, the deformation is decreasing with spin and the PRM reflects an average. The tilted-axis cranking model [26] results for the Be isotopes [27] may support this kind of argument. It is also interesting to point out that a recent *ab initio* no-core shell model study [28] of  $^{10-14}\text{B}$  isotopes with realistic  $NN$  forces predicts  $Q_{3/2^-}$  in the range  $0.027-0.031 e\ b$  (depending on the interaction used), very close to our estimate.

In contrast to heavy nuclei, light nuclei are more vulnerable to changes in deformation just by the addition of one particle. Relevant to our discussions is the fact that  $^{12}\text{C}$  is oblate [29], and one question that comes to mind is: why not consider  $^{11}\text{B}$  as a hole coupled to an oblate core? In fact, in an early study [30], the energy level scheme of  $^{11}\text{B}$  was described in the collective model with the deformation  $\epsilon_2 \approx -0.4$  corresponding to the left side of the Nilsson levels shown in Fig. 1. We have carried out PRM calculations for oblate deformations and obtained an agreement similar to that in Fig. 2 for  $\epsilon_2 \approx -0.35$  and  $\mathcal{S} = 0.50 \hbar^2/\text{MeV}$ . However, this solution gives  $Q_{3/2^-} = -0.0063 e\ b$  in clear disagreement with experiment. In looking at the positive-parity states, we find that the lowest excitation corresponds to a  $5/2^+$  at  $\approx 10$  MeV from the  $3/2^-$  ground state, also inconsistent with the experimental level scheme.

Based on the discussion above, we adopt the prolate results to assess the structure of the ground state in  $^{12}\text{B}$ , as a neutron and a proton coupled to the  $^{10}\text{Be}$  core. The ground state will result from the coupling of the structures discussed

previously, namely, a neutron  $1/2^-$  based on the  $[101]_{\frac{1}{2}}$  intrinsic neutron level and a  $3/2^-$  proton state as described in Table I, which will give rise to  $2^+$  and  $1^+$  states with parallel or antiparallel coupling, respectively. Following the empirical Gallagher-Moszkowski rule [31], the lower member of the doublet corresponds to the  $1^+$  built from the  $K^\pi = 0^+$  and  $1^+$  components in agreement with the experimental observations. Further qualitative support comes from the lowest negative-parity states,  $2^-$  and  $1^-$ , expected from the coupling to the neutron  $[220]_{\frac{1}{2}}$ . On the oblate side, a  $1^-$  to  $4^-$  spin multiplet should result from the coupling to the  $[202]_{\frac{5}{2}}$  level. Additionally, the measured [24] magnetic moment  $\mu_{1^-} = 1.003\ 06(15) \mu_N$  and the quadrupole moment  $Q_{1^+} = 0.0134(14) e\ b$ , which compare well with the leading-order estimates  $\mu_{1^-} = 0.97 \mu_N$  and  $Q_{1^+} = 0.017 e\ b$ . With the ingredients above we proceed to calculate the  $p$ -wave proton-removal strengths, in terms of the Coriolis mixing amplitudes for the  $^{12}\text{B}$  ground state.

### III. SPECTROSCOPIC FACTORS

We apply the formalism reviewed in Ref. [32] to a proton-pickup reaction, such as  $(d, {}^3\text{He})$ . In the strong-coupling limit, the spectroscopic factors ( $S_{i,f}$ ) from an initial ground state  $|I_i K_i\rangle$  to a final state  $|I_f K_f\rangle$  can be written in terms of the Nilsson amplitudes [32]:

$$\begin{aligned} \theta_{i,f}(j\ell, K) &= \langle I_i j K \Omega_\pi | I_f 0 \rangle C_{j,\ell} \langle \phi_f | \phi_i \rangle, \\ S_{i,f} &= \theta_{i,f}^2(j\ell, K), \end{aligned} \quad (4)$$

where  $\langle I_i j K \Omega_\pi | I_f 0 \rangle$  is a Clebsch-Gordan coefficient,  $C_{j,\ell}$  is the Nilsson wave-function amplitude, and  $\langle \phi_f | \phi_i \rangle$  is the core overlap between the initial and final states, which we assume to be 1.

Due to the effects of Coriolis coupling discussed previously, Eq. (4) is generalized to [33]

$$S_{i,f}(j\ell) = \left( \sum_K \mathcal{A}_{IK} \theta_{i,f}(j\ell, K) \right)^2. \quad (5)$$

Following from the results in Table I, where the  $3/2^-$  ground state is dominated by two contributing Nilsson orbitals, we only consider the  $[110]_{\frac{1}{2}}$  and  $[101]_{\frac{3}{2}}$  proton levels which, in the spherical  $|j, \ell\rangle$  basis, have the following wave functions:

$$|[110]_{\frac{1}{2}}\rangle = -0.34|p_{1/2}\rangle + 0.94|p_{3/2}\rangle, \quad (6)$$

$$|[101]_{\frac{3}{2}}\rangle = |p_{3/2}\rangle. \quad (7)$$

When applied to the case of  $^{12}\text{B}$ , the PRM Hamiltonian for the  $1^+$  ground state is a  $3 \times 3$  matrix in the basis:

$$\begin{aligned} |1\rangle &= |v[101]_{\frac{1}{2}} \otimes \pi[110]_{\frac{1}{2}}\rangle_{K=0}, \\ |2\rangle &= |v[101]_{\frac{1}{2}} \otimes \pi[110]_{\frac{1}{2}}\rangle_{K=1}, \\ |3\rangle &= |v[101]_{\frac{1}{2}} \otimes \pi[101]_{\frac{3}{2}}\rangle_{K=1}, \end{aligned} \quad (8)$$

giving a wave function of the form

$$|^{12}\text{B}, 1^+\rangle_{\text{g.s.}} = \mathcal{A}_1|1\rangle + \mathcal{A}_2|2\rangle + \mathcal{A}_3|3\rangle. \quad (9)$$

TABLE II. Comparison among the experimental  $\ell = 1$  proton-removal spectroscopic factors, the Nilsson strong-coupling limit, and the PRM results. Note that these are relative to the transitions normalized to 1 and, as such, quenching effects largely cancel.

Initial state	Final state	Energy (MeV)	Expt	$S_{i,f}$	
				Strong coupling	Coriolis <sup>a,b</sup>
<sup>12</sup> B	<sup>11</sup> Be				
$1_1^-$	$1_1^-$	0.32	1	1	1
	$3/2_1^-$	2.35	2.6(10)	0.8	2.6
	$5/2_1^-$	3.89	1.7(6)	0.2	1.7
<sup>12</sup> C	<sup>11</sup> B				
$0_1^+$	$3_1^-$	0.00	1	1	1
	$1_1^-$	2.12	0.12(2)	0.9	0.1
	$2_1^-$	5.02	0.10(2)	0.9	0.3

<sup>a</sup>Note that for <sup>12</sup>B, having two data points and two unknowns, the minimization solution reproduces the data exactly.

<sup>b</sup>For <sup>12</sup>C these are based on the <sup>11</sup>B amplitudes in Table II.

The amplitudes  $\mathcal{A}_{1-3}$  were fit using a least-squares minimization to the experimental spectroscopic factor data, yielding  $\mathcal{A}_1 = -0.60(3)$ ,  $\mathcal{A}_2 = 0.70(3)$ , and  $\mathcal{A}_3 = 0.40(4)$  given by the normalization condition  $\mathcal{A}_1^2 + \mathcal{A}_2^2 + \mathcal{A}_3^2 = 1$ . The derived amplitudes confirm that Coriolis mixing is required to explain the experimental data, reflecting the PRM results for the  $3/2^-$  band in <sup>11</sup>B, since the  $[101]_{\frac{1}{2}}$  neutron can be seen to act as a spectator. The Coriolis effects appear to be somewhat larger in <sup>12</sup>B, which may suggest a core with smaller deformation and a reduced momenta of inertia, both favoring the increased mixing.

We note that within our framework, pickup to the  $3/2_2^-$  state is not possible since a neutron hole in the  $[101]_{\frac{3}{2}}$  level is not present in the ground state of <sup>12</sup>B. In any case, a contribution to the  $3/2_2^- + 5/2_1^-$  doublet due to  $K$  mixing in <sup>11</sup>Be is expected to be quite small.

The calculated relative  $S_{i,f}$  values in the strong-coupling limit and the PRM are compared to the experimental data in Table II, which also includes those for the <sup>12</sup>C( $p, 2p$ )<sup>11</sup>B reaction [34]. As already mentioned, the spectroscopic factors of the  $3/2_1^-$  and  $5/2_1^-$  states were underestimated in the strong-coupling limit and the inclusion of Coriolis coupling appears to solve the discrepancy, bringing the collective model predictions in line with those of the shell-model and the *ab initio* variational Monte Carlo results discussed in Ref. [19]. It would be of interest if the study of Ref. [28] could be extended to obtain spectroscopic factors for the reactions in Table II.

#### IV. CONCLUSION

We have analyzed spectroscopic factors extracted from the <sup>12</sup>B( $d, ^3\text{He}$ )<sup>11</sup>Be proton-removal reaction in the framework of the collective model. The PRM quantitatively explains the available structure data in <sup>11</sup>Be and <sup>11</sup>B and provides clear evidence of Coriolis coupling in the ground state of <sup>12</sup>B. The resulting  $K$  mixing in the wave function is key to understanding the experimental (relative) spectroscopic factors, which are underestimated in the strong-coupling limit. The amplitudes, empirically adjusted to reproduce the data, are in agreement with the PRM expectations. An application of our phenomenological description to the structure of <sup>12</sup>B [Eqs. (8) and (9)] with the two-particle plus rotor model would be an interesting extension to explore.

#### ACKNOWLEDGMENTS

We thank Prof. Ikuko Hamamoto and Prof. Stefan Frauendorf for enlightening discussions on the Nilsson and particle rotor models. This material is based upon work supported by the US Department of Energy, Office of Science, Office of Nuclear Physics under Contracts No. DE-AC02-05CH11231 (LBNL) and No. DE-AC02-06CH11357 (ANL).

- |   |  |
|---|--|
| <p>[1] A. Bohr and B. R. Mottelson, <i>Mat. Fys. Medd. Dan. Vid. Selsk.</i> <b>27</b>, 16 (1953).</p> <p>[2] H. Morinaga, <i>Phys. Rev.</i> <b>101</b>, 254 (1956).</p> <p>[3] <a href="https://www.nndc.bnl.gov/ensdf/">https://www.nndc.bnl.gov/ensdf/</a>.</p> <p>[4] W. Von Oertzen, M. Freer, and Y. Kanada-Enyo, <i>Phys. Rep.</i> <b>432</b>, 43 (2006).</p> <p>[5] A. Poves and J. Retamosa, <i>Phys. Lett. B</i> <b>184</b>, 311 (1987).</p> <p>[6] E. K. Warburton, J. A. Becker, and B. A. Brown, <i>Phys. Rev. C</i> <b>41</b>, 1147 (1990).</p> <p>[7] I. Talmi and I. Unna, <i>Phys. Rev. Lett.</i> <b>4</b>, 469 (1960).</p> <p>[8] O. Sorlin and M. Porquet, <i>Prog. Part. Nucl. Phys.</i> <b>61</b>, 602 (2008).</p> <p>[9] K. Heyde and J. L. Wood, <i>Rev. Mod. Phys.</i> <b>83</b>, 1467 (2011).</p> <p>[10] T. Otsuka, A. Gade, O. Sorlin, T. Suzuki, and Y. Utsuno, <i>Rev. Mod. Phys.</i> <b>92</b>, 015002 (2020).</p> <p>[11] A. Bohr and B. R. Mottelson, <i>Nuclear Structure Volume II</i> (Benjamin, Advanced Book Program, Reading, MA, 1975), p. 285.</p> | <p>[12] S. G. Nilsson, <i>Mat. Fys. Medd. Dan. Vid. Selsk.</i> <b>29</b>, 16 (1955).</p> <p>[13] S. G. Nilsson and I. Ragnarsson, <i>Shapes and Shells in Nuclear Structure</i> (Cambridge University, Cambridge, England, 1995).</p> <p>[14] I. Hamamoto and S. Shimoura, <i>J. Phys. G: Nucl. Part. Phys.</i> <b>34</b>, 2715 (2007).</p> <p>[15] M. A. Caprio, P. Maris, J. P. Vary, and R. Smith, <i>Int. J. Mod. Phys. E</i> <b>24</b>, 1541002 (2015), and references therein.</p> <p>[16] A. O. Macchiavelli, H. L. Crawford, C. M. Campbell, R. M. Clark, M. Cromaz, P. Fallon, M. D. Jones, I. Y. Lee, A. L. Richard, and M. Salathe, <i>Phys. Rev. C</i> <b>96</b>, 054302 (2017).</p> <p>[17] A. O. Macchiavelli, H. L. Crawford, C. M. Campbell, R. M. Clark, M. Cromaz, P. Fallon, M. D. Jones, I. Y. Lee, and M. Salathe, <i>Phys. Rev. C</i> <b>97</b>, 011302(R) (2018).</p> <p>[18] A. O. Macchiavelli, H. L. Crawford, P. Fallon, I. Y. Lee, R. M. Clark, C. M. Campbell, M. Cromaz, C. Morse, and C. Santamaria, <i>Phys. Rev. C</i> <b>101</b>, 044319 (2020).</p> |
|---|--|

- [19] J. Chen, K. Auranen, M. L. Avila, B. B. Back, M. A. Caprio, C. R. Hoffman, D. Gorelov, B. P. Kay, S. A. Kuvin, Q. Liu, J. L. Lou, A. O. Macchiavelli, D. G. McNeel, T. L. Tang, D. Santiago-Gonzalez, R. Talwar, J. Wu, G. Wilson, R. B. Wiringa, Y. L. Ye, C. X. Yuan, and H. L. Zang, *Phys. Rev. C* **100**, 064314 (2019).
- [20] S. E. Larsson, G. Leander, and I. Ragnarsson, *Nucl. Phys. A* **307**, 189 (1978).
- [21] I. Ragnarsson and P. B. Semmes, *Hyperfine Interact.* **43**, 425 (1988).
- [22] I. Ragnarsson and R. K. Sheline, *Phys. Scr.* **29**, 385 (1984).
- [23] A. B. Migdal, *Nucl. Phys.* **13**, 655 (1959).
- [24] P. Raghavan, *At. Data Nucl. Data Tables* **42**, 189 (1989).
- [25] R. K. Nesbet, *Phys. Rev. Lett.* **24**, 1155 (1970).
- [26] S. Frauendorf, *Nucl. Phys. A* **677**, 115 (2000).
- [27] B. Qi and S. Frauendorf (private communication) (unpublished).
- [28] P. Choudhary, P. C. Srivastava, and P. Navratil, *Phys. Rev. C* **102**, 044309 (2020).
- [29] W. J. Vermeer, M. T. Esat, J. A. Kuehner, H. Spear, A. M. Baxter, and S. Hinds, *Phys. Lett. B* **122**, 23 (1983).
- [30] A. B. Clegg, *Nucl. Phys.* **38**, 353 (1960).
- [31] C. G. Gallagher and S. A. Moszkowski, *Phys. Rev.* **111**, 1282 (1958).
- [32] B. Elbek and P. O. Tjøm, in *Advances in Nuclear Physics*, edited by M. Baranger and E. Vogt (Springer, Boston, MA, 1969), pp. 259–323.
- [33] R. F. Casten, P. Kleinheinz, P. J. Daly, and B. Elbek, *Phys. Rev. C* **3**, 1271 (1971).
- [34] V. Panin *et al.*, *Phys. Lett. B* **753**, 204 (2016).