

Microscopic self-consistent description of induced fission: Dynamical pairing degree of freedomJie Zhao (赵杰) ¹, Tamara Nikšić,² and Dario Vretenar ^{2,3}¹Center for Circuits and Systems, Peng Cheng Laboratory, Shenzhen 518055, China²Department of Physics, Faculty of Science, University of Zagreb, Bijenička Cesta 32, Zagreb 10000, Croatia³State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

(Received 30 June 2021; revised 30 August 2021; accepted 30 September 2021; published 12 October 2021)

The role of dynamical pairing in induced fission dynamics is investigated using the time-dependent generator coordinate method in the Gaussian overlap approximation, based on the microscopic framework of nuclear energy density functionals. A calculation of fragment charge yields for induced fission of ^{228}Th is performed in a three-dimensional space of collective coordinates that, in addition to the axial quadrupole and octupole intrinsic deformations of the nuclear density, also includes an isoscalar pairing degree of freedom. It is shown that the inclusion of dynamical pairing has a pronounced effect on the collective inertia, the collective flux through the scission hypersurface, and the resulting fission yields, reducing the asymmetric peaks and enhancing the contribution of symmetric fission, in better agreement with the empirical trend.

DOI: [10.1103/PhysRevC.104.044612](https://doi.org/10.1103/PhysRevC.104.044612)**I. INTRODUCTION**

Nuclear density functional theory is the only microscopic framework that can be used over the entire table of nuclides in a self-consistent description of phenomena ranging from ground-state properties and collective excitations to large-amplitude nucleonic motion, fission and low-energy collisions. In the case of nuclear fission [1], in particular, a fully quantum-mechanical many-body model can be constructed starting from the time-dependent generator coordinate method (TDGCM) [2]. In this approach the nuclear wave function is represented by a linear superposition of many-body generator states that are functions of collective coordinates. In most cases these coordinates parametrize the shape of the nuclear density. The Hill-Wheeler equation of motion determines the time evolution of the wave function in the restricted space of generator states [3]. By employing the Gaussian overlap approximation (GOA), the GCM Hill-Wheeler equation reduces to a local, time-dependent Schrödinger equation in the space of collective coordinates. The microscopic input for the collective Schrödinger equation, that is, the nuclear potential and collective inertia, are determined by self-consistent mean-field calculations for a choice of the energy density functional (EDF) or effective interaction. The TDGCM + GOA method can be applied to the dynamics of induced fission, starting from the ground state and following the time evolution of collective degrees of freedom all the way to scission and the emergence of fission fragments. This framework has been very successfully implemented in a number of fission studies based on nonrelativistic Skyrme and Gogny functionals [2–11]. Those studies have analyzed the dependence of the predicted fission dynamics on the choice of the EDF, initial conditions, form of the collective inertia, and the definition of scission configurations.

Relativistic energy density functionals [12–14] have also been employed in the description of both spontaneous [15,16] and induced nuclear fission [17–20]. The microscopic input for these studies is generated using either the multidimensionally constrained relativistic mean-field (MDC-RMF) [21] or the relativistic Hartree-Bogoliubov model [22]. By employing the TDGCM + GOA collective model, several interesting topics have been explored in this framework, such as the influence of static pairing correlations on fission yields, different approximations for the collective inertia tensor, and finite temperature effects.

Most applications of the TDGCM to fission dynamics have considered a two-dimensional (2D) space of collective coordinates, such as, for instance, quadrupole and octupole shape degrees of freedom. The recently developed computer code FELIX [5,6] offers the possibility of solving the TDGCM + GOA equation for an arbitrary number of collective variables. In Ref. [8] a preliminary calculation of induced fission dynamics of ^{240}Pu isotope in the three-dimensional (3D) space of shape variables (quadrupole, octupole, and hexadecupole intrinsic deformations) has been reported. In particular, this model can also be used for a quantitative analysis of the critical role of dynamical pairing correlations in the process of induced fission.

The importance of pairing correlations for both spontaneous and induced nuclear fission has been emphasized in a number of studies [16,23–31]. For spontaneous fission it has been shown that the coupling between shape and pairing degrees of freedom has a pronounced effect on the calculated fission lifetimes [16,25]. In particular, when the gap parameter is considered as a dynamical variable, pairing correlations are generally enhanced, thus, reducing the effective inertia and the action integral along the fission path. This effect can significantly reduce the estimated spontaneous fission life-

times, and it has also been noted that pairing fluctuations can restore axial symmetry in the fissioning system. A study of induced fission of ^{240}Pu , using the microscopic time-dependent superfluid local density approximation [30], has shown that both shape and pairing modes determine the dynamics of the final stage of the fission process, from configurations close to the outer fission barrier to full scission.

The influence of ground-state (static) pairing correlations on charge yields and total kinetic energy of fission fragments for the case of induced fission of ^{226}Th isotope was analyzed in Ref. [17] using the TDGCM + GOA framework. It has been shown that an increase in the strength of the pairing interaction, beyond the range determined by empirical pairing gaps obtained from the experimental masses of neighboring nuclei, reduces the asymmetric peaks and enhances the symmetric peak in charge yields distribution. This is a very interesting result, and, thus, it is important to explore dynamical pairing correlations in induced fission. In this paper we explicitly include the isoscalar pairing degree of freedom in the space of TDGCM + GOA collective coordinates and perform the first realistic three-dimensional calculation of induced fission of ^{228}Th . The theoretical framework and methods are reviewed in Sec. II. The details of the calculation and principal results are discussed in Sec. III. Section IV contains a short summary and outlook for future studies.

II. THE TDGCM + GOA METHOD

In the TDGCM + GOA framework-induced fission is described as a slow adiabatic process determined by a small number of collective degrees of freedom. The initial step in modeling the fission of a heavy nucleus is a self-consistent mean-field calculation of the corresponding deformation energy surface as a function of few selected collective coordinates. Such a calculation provides the microscopic input, that is, the single-quasiparticle states, energies, and occupation factors, that determine the parameters of a local equation of motion for the collective wave function.

The theoretical framework and specific model have been detailed in our previous studies [17–20]. For completeness, here we include a short outline and discuss in more detail the specific points that arise when considering pairing as a collective degree of freedom. The relativistic energy density functional DD-PC1 [32] is employed in the particle-hole channel, whereas pairing correlations are taken into account in the Bardeen-Cooper-Schrieffer (BCS) approximation by a separable pairing force of finite range [33]. The parameters of the pairing interaction have been adjusted to reproduce the empirical pairing gaps in the mass region considered in the present paper [19].

The self-consistent deformation energy surfaces are calculated using the multidimensionally constrained relativistic mean-field (MDC-RMF) model [16,21,34,35] with constraints on mass multipole moments $Q_{\lambda\mu} = r^\lambda Y_{\lambda\mu}$ and the particle-number dispersion operator $\Delta\hat{N}^2 = \hat{N}^2 - \langle\hat{N}\rangle^2$. The Routhian is, therefore, defined as

$$E' = E_{\text{RMF}} + \sum_{\lambda\mu} \frac{1}{2} C_{\lambda\mu} Q_{\lambda\mu} + \lambda_2 \Delta\hat{N}^2, \quad (1)$$

where E_{RMF} denotes the total RMF energy that includes static BCS pairing correlations. The amount of dynamic pairing correlations can be controlled by the Lagrange multipliers $\lambda_{2\tau}$ ($\tau = n, p$) [16,25,36,37]. To reduce the number of collective degrees of freedom and, therefore, the considerable computational task, here we only consider isoscalar dynamical pairing; $\lambda_{2n} = \lambda_{2p} \equiv \lambda_2$ is employed as the collective coordinate.

The dynamics of the fission process is governed by a local time-dependent Schrödinger-like equation in the space of collective coordinates \mathbf{q} . The collective Hamiltonian $\hat{H}_{\text{coll}}(\mathbf{q})$,

$$\hat{H}_{\text{coll}}(\mathbf{q}) = -\frac{\hbar^2}{2} \sum_{ij} \frac{\partial}{\partial q_i} B_{ij}(\mathbf{q}) \frac{\partial}{\partial q_j} + V(\mathbf{q}) \quad (2)$$

determines the time evolution of the nuclear wave function from an initial state at equilibrium deformation, up to scission and the formation of fission fragments. $B_{ij}(\mathbf{q})$ and $V(\mathbf{q})$ denote the inertia tensor and collective potential, respectively, that are computed using the self-consistent solutions for the RMF + BCS deformation energy surface. Here we assume axial symmetry with respect to the axis along which the two fragments eventually separate and consider the 3D collective space of quadrupole β_2 and octupole β_3 deformation parameters and the dynamical pairing coordinate λ_2 . The inertia tensor is the inverse of the mass tensor, that is, $B_{ij}(\mathbf{q}) = (\mathcal{M}^{-1})_{ij}(\mathbf{q})$. The mass tensor is calculated using the adiabatic time-dependent Hartree-Fock-Bogoliubov (ATDHFB) method in the cranking approximation [38],

$$\mathcal{M}_{ij}^C = \frac{\hbar^2}{2\dot{q}_i \dot{q}_j} \sum_{\mu\nu} \frac{F_{\mu\nu}^{i*} F_{\mu\nu}^j + F_{\mu\nu}^i F_{\mu\nu}^{j*}}{E_\mu + E_\nu}, \quad (3)$$

where

$$\frac{F^i}{\dot{q}_i} = U^\dagger \frac{\partial \rho}{\partial q_i} V^* + U^\dagger \frac{\partial \kappa}{\partial q_i} U^* - V^\dagger \frac{\partial \rho^*}{\partial q_i} U^* - V^\dagger \frac{\partial \kappa^*}{\partial q_i} V^*. \quad (4)$$

U and V are the self-consistent Bogoliubov matrices, and ρ and κ are the corresponding particle and pairing density matrices, respectively. The cranking expression Eq. (3) can be further simplified in the perturbative approach [39–43], and this leads to the perturbative cranking mass tensor,

$$\mathcal{M}^{Cp} = \hbar^2 \mathcal{M}_{(1)}^{-1} \mathcal{M}_{(3)} \mathcal{M}_{(1)}^{-1}, \quad (5)$$

where

$$[\mathcal{M}_{(k)}]_{ij} = \sum_{\mu\nu} \frac{\langle 0 | \hat{Q}_i | \mu\nu \rangle \langle \mu\nu | \hat{Q}_j | 0 \rangle}{(E_\mu + E_\nu)^k}. \quad (6)$$

$|\mu\nu\rangle$ are the two-quasiparticle states with the corresponding quasiparticle energies E_μ and E_ν .

At this point we introduce an approximation that is not entirely consistent but is necessary to reduce the computational task and stabilize the time evolution of the collective state. Because we consider the particle-number dispersion operator $\Delta\hat{N}^2 = \hat{N}^2 - \langle\hat{N}\rangle^2$, the pairing part of the mass tensor must be calculated using the nonperturbative cranking expression Eq. (3). In the recent study of the differences between the perturbative and the nonperturbative ATDHFB collective masses

in the TDGCM + GOA description of induced fission [20] using the axial quadrupole and octupole intrinsic deformation as dynamical variables, we have shown that the structure of non-perturbative collective masses is much more complex due to changes in the intrinsic shell structure across the deformation energy surface and is characterized by pronounced isolated peaks located at single-particle level crossings near the Fermi surface. It has been shown that the choice of nonperturbative cranking collective mass leads to a reduction of symmetric charge yields and, generally, to a better agreement with data. Even though in the 2D analysis of Ref. [20] both nonperturbative and perturbative mass tensors were used in modeling induced fission dynamics, in the present 3D study the number of mesh points required to accurately calculate all the nonperturbative collective masses becomes prohibitively large. The reason is the occurrence of pronounced peaks in the collective masses related to single-particle level crossings near the Fermi surface and the corresponding abrupt changes in the occupation factors of single-particle configurations. This would require a major refinement of the grid, leading to very large number of mesh points and possible instabilities. Therefore, to be able to quantitatively analyze the effect of dynamical pairing on induced fission, here we use the perturbative cranking expression Eq. (5) to calculate the elements of the mass tensor that correspond to the quadrupole and octupole deformations whereas, as noted above, the pairing element of the mass tensor must be computed using the nonperturbative cranking formula. With the indices 1–3 corresponding to the β_{20} , β_{30} , and λ_2 collective coordinates, the following elements of the mass tensor are used: \mathcal{M}_{11}^{Cp} , \mathcal{M}_{12}^{Cp} , \mathcal{M}_{22}^{Cp} , \mathcal{M}_{33}^C , and we neglect the coupling terms \mathcal{M}_{13} and \mathcal{M}_{23} .

To model the fission dynamics we follow the time evolution of an initial wave packet, built as a Gaussian superposition of quasibound states and with an average collective energy set 1 MeV above the fission barrier. The time propagation is modeled using the TDGCM + GOA computer code FELIX (version 2.0) [6]. The time step is $\delta t = 5 \times 10^{-4}$ zs (1 zs = 10^{-21} s), and the charge and mass distributions are calculated after 4×10^4 time steps, which correspond to 20 zs.

The collective space is divided into an inner region with a single nuclear density distribution and an external region that contains two separated fission fragments. The scission hypersurface that divides the inner and external regions is determined by the expectation values of the Gaussian neck operator $\hat{Q}_N = \exp[-(z - z_N)^2/a_N^2]$, where $a_N = 1$ fm and z_N is the position of the neck [44]. We define the prescission domain by $\langle \hat{Q}_N \rangle > 3$ and consider the frontier of this domain as the scission surface. The flux of the probability current through this hypersurface provides a measure of the probability of observing a given pair of fragments at time t [6],

$$F(\xi, t) = \int_{t_0}^t dt' \int_{q \in \xi} \mathbf{J}(\mathbf{q}, t') \cdot d\mathbf{S}, \quad (7)$$

where $\mathbf{J}(\mathbf{q}, t)$ is the current,

$$J_k(\mathbf{q}, t) = \hbar \sum_l B_{kl}(\mathbf{q}) \text{Im} \left(g^* \frac{\partial g}{\partial q_l} \right); \quad (8)$$




FIG. 1. Two-dimensional projections of the deformation-energy manifold of ^{228}Th on the quadrupole-octupole axially symmetric plane, calculated with the RMF + BCS model based on the functional DD-PC1 for selected values of the pairing coordinate λ_2 . Contours join points on the surface with the same energy, and the separation between neighboring contours is 1 MeV. The red curves denote static fission paths of minimum energy for each value of λ_2 .

and $g(\mathbf{q}, t)$ is the wave function of the collective variables \mathbf{q} and time t . The yield for the fission fragment with mass A is defined by

$$Y(A) \propto \sum_{\xi \in \mathcal{A}} \lim_{t \rightarrow \infty} F(\xi, t). \quad (9)$$

The set $\mathcal{A}(\xi)$ contains all elements belonging to the scission hypersurface such that one of the fragments has mass number A .

III. RESULTS AND DISCUSSION

As an illustrative example, the fission of ^{228}Th is considered. For this nucleus the charge distribution of fission fragments exhibits a coexistence of symmetric and asymmetric peaks [45]. In the first step a large-scale MDC-RMF calculation is performed to generate the potential-energy surface, single-nucleon wave functions, and occupation factors in the $(\beta_2, \beta_3, \lambda_2)$ collective space. The intervals for the values of the collective variables are as follows: $-1 \leq \beta_2 \leq 7$ with a step $\Delta\beta_2 = 0.04$; $0 \leq \beta_3 \leq 3.5$ with a step $\Delta\beta_3 = 0.05$; and $-0.1 \leq \lambda_2 \leq 2.0$ with a step $\Delta\lambda_2 = 0.1$.

Figure 1 displays the 2D projections of the 3D deformation energy surface of ^{228}Th on the collective plane (β_2, β_3) for several values of the collective coordinate λ_2 . Note that the value $\lambda_2 = 0$ corresponds to static BCS pairing, whereas positive λ_2 equates to enhanced pairing correlations. Only




FIG. 2. Pairing gaps for neutrons Δ_n (upper panel) and protons Δ_p (lower panel) along the static fission path as functions of the axial quadrupole deformation for selected values of the isoscalar pairing collective coordinate λ_2 .

configurations with $Q_N \geq 3$ are plotted, and the frontier of this domain determines the scission contour. The red curves correspond to static fission paths of minimum total energy. The deformation surfaces for $\lambda_2 = 0$ and $\lambda_2 = 0.3$ are very similar with a pronounced ridge separating the asymmetric and symmetric fission valleys. This ridge decreases with increasing values of the pairing coordinate λ_2 . The scission contour for $\lambda_2 = 0$ starts from an elongated symmetric point at $\beta_2 \approx 6$ and evolves to a minimal elongation $\beta_2 \approx 3$ as reflection asymmetry increases. For larger values of λ_2 the scission contour is not modified significantly, although the starting point of the scission contour on the quadrupole axis shifts to smaller elongations at $\beta_2 \approx 5$.

To illustrate the dynamical effect on the pairing correlations, in Fig. 2 we display the values of the neutron (upper panel) and proton (lower panel) pairing gaps along the static fission paths as functions of the quadrupole coordinate β_2 for several values of the isoscalar pairing collective coordinate λ_2 (cf. Fig. 1). In the interval of values of λ_2 considered here, the values of the pairing gaps increase by a factor of ≈ 2 to 3 and, characteristically, for $\lambda_2 \geq 0.6$ all traces of the underlying shell effects along the static fission path vanish.

In Fig. 3 we plot the values of the perturbative cranking collective masses M_{11}^{Cp} , M_{22}^{Cp} , and the nonperturbative cranking mass M_{33}^{C3} on a logarithmic scale along the static fission path as functions of the quadrupole deformation for different pairing collective coordinates λ_2 . As noted above, the indices 1–3 refer to the β_2 , β_3 , and λ_2 coordinates, respectively. The collective masses M_{11}^{Cp} and M_{22}^{Cp} exhibit a more complex structure for smaller values of deformations $\beta_2 < 1$ and decrease for larger deformations. One notes that M_{22}^{Cp} exhibits a sharper decrease for $\beta_2 > 1$, compared to M_{11}^{Cp} . The behavior of the M_{33}^{C3} collective mass is much more complex due to the fact that it has been calculated using the nonperturbative cranking formula Eq. (3). This leads to the appearance of prominent




FIG. 3. Perturbative cranking masses M_{11}^{Cp} , M_{22}^{Cp} , and the non-perturbative cranking mass M_{33}^{C3} (in $\hbar^2 \text{MeV}^{-1}$) (logarithmic scale) along the static fission path for several values of λ_2 .

peaks at the locations of single-particle level crossing near the Fermi surface. However, in general, also M_{33}^{C3} decreases for large deformations, and the peaks are less pronounced. As the value of the dynamical pairing coordinate λ_2 increases, all three collective masses decrease, and the underlying shell effects gradually vanish. This effect is consistent with the results shown in Fig. 6 of Ref. [17]. The scission contours in the (β_2, β_3) plane are shown in Fig. 4 for several collective pairing coordinates λ_2 . The contours are generally not very different especially for asymmetric fission. In particular, scission points that are close to the static fission path are not sensitive to dynamical pairing. For larger values of λ_2 , however, the scission contour is shifted towards smaller quadrupole deformations β_2 values for nearly symmetric fission.

In Fig. 5 we compare the theoretical predictions for the charge yields with the data for photoinduced fission of ^{228}Th . The 3D calculation employs the collective space built from the deformation β_2 , β_3 and pairing λ_2 coordinates, whereas the 2D calculation includes only the shape degrees of freedom β_2 and β_3 , and static pairing correlations. As in our study of Ref. [17], the calculation in the 2D collective space corresponds to normal and enhanced static pairings, that is, to 100% and 110% of the normal pairing strength, determined by the empirical pairing gaps. For completeness, in Fig. 6 we plot the corresponding 2D deformation energy surfaces on the β_2 - β_3 axially symmetric plane.

The theoretical predictions follow the general trend of the data, except that our model obviously cannot reproduce the odd-even staggering of the experimental charge yields. The calculation that includes only the 2D collective space with




FIG. 4. The scission contour of ^{228}Th on the (β_2, β_3) deformation plane for several values of the collective pairing coordinate λ_2 .

a static pairing strength adjusted to empirical ground-state pairing gaps in this mass region (100%) predicts yields that are entirely dominated by asymmetric fission with peaks at $Z = 35$ and $Z = 55$. By increasing static pairing (110%), the asymmetric peaks are reduced, and a contribution of symmetric fission develops but not strong enough to reproduce the




FIG. 5. Charge yields for induced fission of ^{228}Th , calculated in the 3D collective space built from the deformation β_2 , β_3 and dynamical pairing λ_2 coordinates (solid red curve). The yields are shown in comparison to the results obtained in the 2D space of shape degrees of freedom β_2 and β_3 with static pairing correlations adjusted to empirical ground-state pairing gaps (100% pairing strength) and enhanced by 10% (110% pairing strength). The data for photoinduced fission correspond to photon energies in the interval 8–14 MeV and peak value of $E_\gamma = 11$ MeV [45].




FIG. 6. Two-dimensional deformation-energy surface of ^{228}Th on the β_2 - β_3 axially symmetric plane, calculated with the RMF+BCS model based on the functional DD-PC1 and with normal (100%) and enhanced (110%) static pairing strengths as determined by the empirical pairing gaps. Contours join points on the surface with the same energy, and the separation between neighboring contours is 1 MeV.

data. It is interesting to note that a very similar distribution of charge yields is predicted by the 3D model calculation that includes dynamical pairing. On a quantitative level, even the 3D calculation does not completely reproduce the experimental yields. The model predicts tails of the asymmetric peaks that are not seen in experiment and, thus, fails to quantitatively match the symmetric contribution. It has to be noted, however, that in the present paper the collective potential and inertia tensor have been calculated at zero temperature. In our recent study of finite temperature effects in TDGCM + GOA [18], a calculation of induced fission of ^{226}Th has shown that, although the model can qualitatively reproduce the empirical triple-humped structure of the fission charge and mass distributions already at zero temperature, the position of the asymmetric peaks and the symmetric-fission yield can be described much better when the potential and collective mass are determined at a temperature that approximately corresponds to the internal excitation energy of the fissioning nucleus.

Finally, to illustrate the effect of dynamical pairing on the flux of the probability current through the scission hypersurface, in Fig. 7 we plot the time-integrated flux through the




FIG. 7. Time-integrated collective flux $B(\lambda_2)$ Eq. (10) through the scission contour as a function of the pairing collective coordinate λ_2 .

scission contour on the (β_2, β_3) plane for a given value of the pairing collective coordinate λ_2 ,

$$B(\lambda_2) \propto \sum_{\xi \in \mathcal{B}} \lim_{t \rightarrow \infty} F(\xi, \lambda_2, t). \quad (10)$$

The set $\mathcal{B}(\xi \equiv \beta_2, \beta_3)$ contains all elements of the scission contour with a given value λ_2 . Even though it appears that dynamical pairing does not significantly modify the scission contour (cf. Fig. 4), nevertheless, its effect on the collective flux and, therefore, on the occurrence of fission, is remarkable. For negative values of λ_2 , that is, for correlations weaker than static pairing at $\lambda_2 = 0$, the flux rapidly decreases to zero. For positive values of λ_2 the flux exhibits a steep increase and a prominent peak at $\lambda_2 \approx 0.3$. Note that this value corresponds to an increase in $\approx 20\%$ with respect to the static proton and neutron pairing gaps (cf. Fig. 2). The collective flux through the scission contour weakens with a further increase in pairing and eventually vanishes for $\lambda_2 > 1$.

IV. SUMMARY

The influence of dynamical pairing degrees of freedom on induced fission has been investigated in a unified theoretical framework based on the GCM with the GOA. In an illustrative calculation of fragment charge yields for induced fission of ^{228}Th , the collective potential and inertia tensor have been computed using the self-consistent multidimensionally constrained relativistic mean field model, based on the energy density functional DD-PC1 and with pairing correlations treated in the BCS approximation with a separable pairing force of finite range. The fission fragment charge distributions are obtained by propagating the initial collective state in time with the time-dependent GCM + GOA. The flux of the probability current through the scission hypersurface determines the probability of observing a given pair of fragments.

In this paper, for the first time, the dynamics of induced fission has been consistently described in a three-dimensional space of collective coordinates that in addition to the axial quadrupole and octupole intrinsic deformations of the nuclear density also includes an isoscalar pairing degree of freedom. A number of studies have already established the importance of dynamical pairing for the calculation of spontaneous fission lifetimes and static pairing correlations for modeling induced fission. As this paper has also shown a much more difficult problem is the inclusion of dynamical pairing degrees of freedom in a time-dependent description of induced fission. One expects, of course, that the model becomes more realistic as the Hilbert space of collective coordinates is expanded. However, as the present analysis has illustrated, it can be difficult to numerically stabilize the time evolution of the fissioning system when shape and pairing collective coordinates are considered simultaneously in a three-dimensional calculation. Here it was necessary to compute the collective inertia using different cranking approximations for the shape and pairing degrees of freedom. Nevertheless, this approach enables a qualitative study of the effect of dynamical pairing on induced fission.

This paper has clearly demonstrated the important effect that dynamical pairing correlations have on the induced-fission fragment distribution. In particular, the charge distribution of fission fragments of ^{228}Th is characterized by symmetric and asymmetric peaks, but this structure cannot be reproduced in a two-dimensional calculation that only includes shape collective variables. In that case, and with a static pairing strength adjusted to empirical ground-state pairing gaps, the calculated yields correspond to a completely asymmetric fission. Only by artificially increasing the static pairing correlations or, more naturally, by including the dynamical pairing degree of freedom in the three-dimensional calculation, the asymmetric peaks get reduced, and a contribution of symmetric fission develops in agreement with the empirical trend. It is also interesting to note that the time-integrated collective flux through the scission contour on the (β_2, β_3) plane, exhibits a characteristic functional dependence on the pairing collective coordinate with a prominent peak at a value that correspond to an increase in $\approx 20\%$ with respect to the static pairing gaps.

Future advances in computational capabilities will open the possibility of more quantitative applications of multidimensional TDGCM + GOA to fission dynamics. An immediate task will be to consider shape and pairing degrees of freedom on an equal footing and consistently compute the corresponding collective inertia tensor and metric. A more challenging problem is to include dynamical pairing degrees of freedom in recently developed fission models that attempt to incorporate restoration of symmetries broken by the intrinsic densities in constrained mean-field calculations (rotational, reflection, and particle number symmetry) [46–48].

ACKNOWLEDGMENTS

This work has been supported, in part, by the QuantiXLie Centre of Excellence, a Project cofinanced by the Croatian Government and European Union through the

European Regional Development Fund, the Competitiveness and Cohesion Operational Programme (Grant No. KK.01.1.1.01.0004), and the Croatian Science Foundation under the Project Uncertainty quantification within the nu-

clear energy density framework (Grant No. IP-2018-01-5987). It has also been supported by the National Natural Science Foundation of China under Grants No. 12005107 and No. 11790325.

-
- [1] M. Bender, R. Bernard, G. Bertsch, S. Chiba, J. Dobaczewski, N. Dubray, S. A. Giuliani, K. Hagino, D. Lacroix, Z. Li, P. Magierski, J. Maruhn, W. Nazarewicz, J. Pei, S. Péru, N. Pillet, J. Randrup, D. Regnier, P.-G. Reinhard, L. M. Robledo *et al.*, *J. Phys. G: Nucl. Part. Phys.* **47**, 113002 (2020).
- [2] J. Berger, M. Girod, and D. Gogny, *Comput. Phys. Commun.* **63**, 365 (1991).
- [3] M. Verriere and D. Regnier, *Front. Phys.* **8**, 233 (2020).
- [4] H. Goutte, J. F. Berger, P. Casoli, and D. Gogny, *Phys. Rev. C* **71**, 024316 (2005).
- [5] D. Regnier, M. Verriere, N. Dubray, and N. Schunck, *Comput. Phys. Commun.* **200**, 350 (2016).
- [6] D. Regnier, N. Dubray, M. Verriere, and N. Schunck, *Comput. Phys. Commun.* **225**, 180 (2018).
- [7] D. Regnier, N. Dubray, N. Schunck, and M. Verriere, *Phys. Rev. C* **93**, 054611 (2016).
- [8] D. Regnier, N. Dubray, N. Schunck, and M. Verriere, *EPJ Web Conf.* **146**, 04043 (2017).
- [9] M. Verriere, N. Dubray, N. Schunck, D. Regnier, and P. Dossantos-Uzarralde, *EPJ Web Conf.* **146**, 04034 (2017).
- [10] D. Regnier, N. Dubray, and N. Schunck, *Phys. Rev. C* **99**, 024611 (2019).
- [11] A. Zdeb, A. Dobrowolski, and M. Warda, *Phys. Rev. C* **95**, 054608 (2017).
- [12] D. Vretenar, A. Afanasjev, G. Lalazissis, and P. Ring, *Phys. Rep.* **409**, 101 (2005).
- [13] J. Meng, H. Toki, S. Zhou, S. Zhang, W. Long, and L. Geng, *Prog. Part. Nucl. Phys.* **57**, 470 (2006).
- [14] *Relativistic Density Functional for Nuclear Structure*, edited by J. Meng, International Review of Nuclear Physics Vol. 10 (World Scientific, Singapore, 2016).
- [15] J. Zhao, B.-N. Lu, T. Niksic, and D. Vretenar, *Phys. Rev. C* **92**, 064315 (2015).
- [16] J. Zhao, B.-N. Lu, T. Niksic, D. Vretenar, and S.-G. Zhou, *Phys. Rev. C* **93**, 044315 (2016).
- [17] H. Tao, J. Zhao, Z. P. Li, T. Niksic, and D. Vretenar, *Phys. Rev. C* **96**, 024319 (2017).
- [18] J. Zhao, T. Niksic, D. Vretenar, and S.-G. Zhou, *Phys. Rev. C* **99**, 014618 (2019).
- [19] J. Zhao, J. Xiang, Z.-P. Li, T. Niksic, D. Vretenar, and S.-G. Zhou, *Phys. Rev. C* **99**, 054613 (2019).
- [20] J. Zhao, T. Nikšić, D. Vretenar, and S.-G. Zhou, *Phys. Rev. C* **101**, 064605 (2020).
- [21] B.-N. Lu, J. Zhao, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **89**, 014323 (2014).
- [22] J. Zhao, B.-N. Lu, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **95**, 014320 (2017).
- [23] J. W. Negele, S. E. Koonin, P. Möller, J. R. Nix, and A. J. Sierk, *Phys. Rev. C* **17**, 1098 (1978).
- [24] Z. Łojewski and A. Staszczak, *Nucl. Phys. A* **657**, 134 (1999).
- [25] J. Sadhukhan, J. Dobaczewski, W. Nazarewicz, J. A. Sheikh, and A. Baran, *Phys. Rev. C* **90**, 061304 (2014).
- [26] A. Staszczak, A. Baran, K. Pomorski, and K. Böning, *Phys. Lett. B* **161**, 227 (1985).
- [27] R. Bernard, S. A. Giuliani, and L. M. Robledo, *Phys. Rev. C* **99**, 064301 (2019).
- [28] Y. A. Lazarev, *Phys. Scr.* **35**, 255 (1987).
- [29] J. Sadhukhan, W. Nazarewicz, and N. Schunck, *Phys. Rev. C* **93**, 011304 (2016).
- [30] A. Bulgac, P. Magierski, K. J. Roche, and I. Stetcu, *Phys. Rev. Lett.* **116**, 122504 (2016).
- [31] Y. Qiang, J. C. Pei, and P. D. Stevenson, *Phys. Rev. C* **103**, L031304 (2021).
- [32] T. Nikšić, D. Vretenar, and P. Ring, *Phys. Rev. C* **78**, 034318 (2008).
- [33] Y. Tian, Z. Y. Ma, and P. Ring, *Phys. Lett. B* **676**, 44 (2009).
- [34] B.-N. Lu, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **85**, 011301(R) (2012).
- [35] S.-G. Zhou, *Phys. Scr.* **91**, 063008 (2016).
- [36] N. L. Vaquero, T. R. Rodriguez, and J. L. Egidio, *Phys. Lett. B* **704**, 520 (2011).
- [37] N. L. Vaquero, J. L. Egidio, and T. R. Rodríguez, *Phys. Rev. C* **88**, 064311 (2013).
- [38] A. Baran, J. A. Sheikh, J. Dobaczewski, W. Nazarewicz, and A. Staszczak, *Phys. Rev. C* **84**, 054321 (2011).
- [39] M. Brack, J. Damgaard, A. S. Jensen, H. C. Pauli, V. M. Strutinsky, and C. Y. Wong, *Rev. Mod. Phys.* **44**, 320 (1972).
- [40] S. G. Nilsson, C. F. Tsang, A. Sobiczewski, Z. Szymanski, S. Wycech, C. Gustafson, I.-L. Lamm, P. Moller, and B. Nilsson, *Nucl. Phys. A* **131**, 1 (1969).
- [41] M. Girod and B. Grammaticos, *Nucl. Phys. A* **330**, 40 (1979).
- [42] D. Bés and Z. Szymański, *Nucl. Phys.* **28**, 42 (1961).
- [43] A. Sobiczewski, Z. Szymański, S. Wycech, S. G. Nilsson, J. R. Nix, C. F. Tsang, C. Gustafson, P. Möller, and B. Nilsson, *Nucl. Phys. A* **131**, 67 (1969).
- [44] W. Younes and D. Gogny, *Phys. Rev. C* **80**, 054313 (2009).
- [45] K.-H. Schmidt, S. Steinhauser, C. Bockstiegel, A. Grewe, A. Heinz, A. Junghans, J. Benlliure, H.-G. Clerc, M. de Jong, J. Muller, M. Pfitzner, and B. Voss, *Nucl. Phys. A* **665**, 221 (2000).
- [46] P. Marević and N. Schunck, *Phys. Rev. Lett.* **125**, 102504 (2020).
- [47] M. Verriere, N. Schunck, and T. Kawano, *Phys. Rev. C* **100**, 024612 (2019).
- [48] M. Verriere, N. Schunck, and D. Regnier, *Phys. Rev. C* **103**, 054602 (2021).