

Elliptic flow splittings in the Polyakov–Nambu–Jona-Lasinio transport modelWen-Hao Zhou ^{1,2}, He Liu,³ Feng Li,⁴ Yi-Feng Sun,⁵ Jun Xu ^{6,1,*} and Che Ming Ko ⁷¹*Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China*²*University of Chinese Academy of Sciences, Beijing 100049, China*³*Science School, Qingdao University of Technology, Qingdao 266000, China*⁴*School of Physical Science and Technology, Lanzhou University, Lanzhou, Gansu, 073000, China*⁵*Laboratori Nazionali del Sud, INFN-LNS, Via S. Sofia 62, I-95123 Catania, Italy*⁶*Shanghai Advanced Research Institute, Chinese Academy of Sciences, Shanghai 201210, China*⁷*Cyclotron Institute and Department of Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA*

(Received 20 May 2021; accepted 15 September 2021; published 4 October 2021)

To incorporate the effect of gluons on the evolution dynamics of the quark matter produced in relativistic heavy-ion collisions, we extend the three-flavor Nambu–Jona-Lasinio (NJL) transport model to include the contribution from the Polyakov loops. Imbedding the resulting pNJL partonic transport model in an extended multiphase transport (extended AMPT) model, we then study the elliptic flow splittings between particles and their antiparticles in relativistic heavy-ion collisions at beam energy scan energies. We find that a weak quark vector interaction in the partonic phase is able to describe the elliptic flow splitting between protons and antiprotons in heavy-ion collisions at $\sqrt{s_{NN}} = 7.7$ to 39 GeV. Knowledge of the quark vector interaction is useful for understanding the equation of state of quark matter at large baryon chemical potentials and thus the location of the critical point in the QCD phase diagram.

DOI: [10.1103/PhysRevC.104.044901](https://doi.org/10.1103/PhysRevC.104.044901)**I. INTRODUCTION**

Understanding the properties of the quark-gluon plasma (QGP) and the phase diagram of quantum chromodynamics (QCD) is among the main goals of the experiments with heavy ions at relativistic energies. For a QGP of low baryon chemical potentials produced in ultrarelativistic heavy-ion collisions, its transition to hadronic matter is a smooth crossover according to lattice QCD (LQCD) calculations [1–3]. Although LQCD has not been able to determine the properties of the QGP at high baryon chemical potentials, it was shown in various effective models for QCD, such as the Nambu–Jona-Lasinio (NJL) model [4–7], the Dyson-Schwinger approach [8,9], and the functional renormalization-group method [10,11], that its transition to the hadronic matter could be a first-order transition. Searching for the QCD critical point, at which the smooth crossover changes to a first-order transition, and determining its location in the QCD phase diagram are currently topics of great interest in high-energy nuclear physics, particularly in relativistic heavy-ion collisions that are carried out in the beam energy scan (BES) program at the Relativistic Heavy Ion Collider (RHIC) and will also be carried out in the compressed baryonic matter (CBM) experiment at the Facility for Antiproton and Ion Research (FAIR) as well as at the Nuclotron-Based Ion Collider Facility (NICA) and the High Intensity Heavy-Ion Accelerator Facility (HIAF). On the other hand, observables measured in heavy-ion collisions at lower

energies can also provide information on the properties of quark matter, especially its equation of state (EOS) at large baryon chemical potentials. Such information is helpful for understanding the properties of strange quark stars [12,13] and gravitational waves from compact-star mergers. We note that the QCD phase diagram and the quark matter EOS at large baryon chemical potentials are related through the underlying quark interactions.

To search for the QCD critical point from the many interesting phenomena observed in the experiments of the BES program at RHIC, there have been great theoretical efforts based on various transport [14–16] or hydrodynamic [17,18] approaches. Assuming that the trajectory traversed by the produced matter with a certain temperature and a baryon chemical potential in the QCD phase diagram could pass through the QCD critical region or the spinodal region, there will then be observational effects on conserved charge fluctuations [19–23] and light cluster productions [24–26] as well as other physical quantities. These studies also provide the possibility to study the quark-matter EOS and the underlying quark interactions, especially at finite baryon chemical potentials. For example, from the collective flow splittings between particles and their antiparticles, one can determine the different mean-field potentials acting on quarks and antiquarks in the baryon-rich quark matter and thus their interactions. Using a QCD effective model with these constrained quark interaction, one can then construct the QCD phase diagram in the temperature and chemical-potential plane, besides determining the location of the critical point.

* xujun@zjlab.org.cn

Using an extended multiphase transport (extended AMPT) model with its partonic stage replaced by the partonic transport model based on the three-flavor NJL model [27–29], we have previously studied the elliptic flow splittings between particles and their antiparticles in relativistic heavy-ion collisions at RHIC-BES energies. We have found that the data measured by the STAR Collaboration [30] can be explained by choosing a proper strength for the vector interaction in the NJL model. The vector interaction extracted from our study puts a significant constraint not only on the EOS of quark matter at finite baryon chemical potentials but also on the properties of the QCD phase diagram. It is, however, well known that the NJL model gives a relatively low temperature for the critical point due to the lack of the effects from gluons. This drawback can be resolved by modifying the NJL model to include the contribution from the Polyakov loops [31], which then allows for the existence of the deconfinement transition besides the chiral transition in the NJL model. In the present paper, we report on the results from an improved transport model study of relativistic heavy-ion collisions based on the Polyakov-loop extended NJL (pNJL) model. Details on the extended AMPT model as well as the NJL and the pNJL model are provided in Sec. II. Section III gives detailed results on the beam-energy dependence of heavy-ion collision dynamics and the v_2 splittings between particles and their antiparticles, as well as those on the QCD phase diagram. We conclude our study in Sec. IV and also include in the Appendixes the details on the treatment of the collision integral in the partonic transport model, particularly the one based on the pNJL model.

II. THEORETICAL FRAMEWORK

The present study is based on the framework of an extended AMPT model with its description of the partonic dynamics using the NJL transport model. The extended AMPT model has been used previously to study the splittings between the elliptic flows [28,29] as well as directed flows [32] of particles and their antiparticles in heavy-ion collisions at RHIC-BES energies. In the present study, we further extend the NJL transport model to include the effect of Polyakov loops and make additional improvements to the extended AMPT model.

A. An extended multiphase transport model

As in the original string-melting AMPT model [33], the momentum distribution of initial partons in the present study is obtained from the constituent quarks and antiquarks of hadrons that are produced from elastic and inelastic scatterings of participant nucleons in the heavy-ion jet interaction generator (HIJING) model [34]. The positions of these initial partons in the transverse plane are set to be the same as those of the colliding nucleons from which they are produced. For their longitudinal positions, they are taken to be uniformly distributed within $(-lm_N/\sqrt{s_{NN}}, lm_N/\sqrt{s_{NN}})$, where $m_N = 0.938$ MeV is nucleon mass, $\sqrt{s_{NN}}$ is the collision energy per nucleon pair in the center-of-mass frame, and $l = (D^2 - b^2)^{1/2}$ fm is the thickness parameter with D being the

diameter of the colliding nucleus ($D = 14$ fm for Au nucleus in the present study) and b being the impact parameter. With the initial partons gradually formed according to their formation time [33], the evolution of these partons is then described by the NJL transport model or its extension by the inclusion of Polyakov loops, where both the mean-field potentials on partons and their scatterings are properly taken into account. For the mean-field potentials, they are calculated by using the test-particle method with parallel events, while scatterings between partons are treated with the stochastic method [35] using an isotropic cross section determined by the specific shear viscosity of the quark matter and including the Pauli blocking factors for fermions, as described in Appendixes A and B. The value of the specific shear viscosity is adjusted so that the transport model reproduces the final charged-particle elliptic flow. Details of the NJL and pNJL model will be given in the next section.

The partonic phase ends when the parton energy density in the central region of the system drops to about 0.8 GeV/fm³. The hadronization of partons is achieved by coalescing partons close in phase space into hadrons [36], during which the baryon and electric charges as well as strangeness are conserved, although the numbers of mesons, baryons, and antibaryons can be different from the initial ones from HIJING. The evolution of the formed hadronic matter is described by a relativistic transport (ART) model [37], which includes various elastic and inelastic channels as well as decay channels for resonances, with the conserved electric charge [29] and mean-field potentials included for baryons, antibaryons, pions, and kaons [27]. Typically, nucleons are affected by a weakly attractive potential while antinucleons are affected by a strongly attractive potential in baryon-rich hadronic matter according to the study of subthreshold production of antiprotons [38]. For other baryons (antibaryons), their potentials are reduced with respect to nucleons (antinucleons) by their light quark (antiquark) fractions. For the kaon and antikaon potentials, they are weakly repulsive and strongly attractive in baryon-rich hadronic matter, respectively, according to the chiral effective-field theory [39].

B. Nambu–Jona-Lasinio model and its Polyakov-loop extension

We start from the Lagrangian of the three-flavor NJL model, which is written as

$$\begin{aligned} \mathcal{L}_{\text{NJL}} = & \bar{\psi}(i/\partial - \hat{m})\psi + \frac{G_S}{2} \sum_{a=0}^8 [(\bar{\psi}\lambda_a\psi)^2 + (\bar{\psi}i\gamma_5\lambda_a\psi)^2] \\ & - \frac{G_V}{2} \sum_{a=0}^8 [(\bar{\psi}\gamma_\mu\lambda_a\psi)^2 + (\bar{\psi}\gamma_5\gamma_\mu\lambda_a\psi)^2] \\ & - K[\det[\bar{\psi}(1 + \gamma_5)\psi] + \det[\bar{\psi}(1 - \gamma_5)\psi]]. \quad (1) \end{aligned}$$

In the above, $\psi = (u, d, s)^T$ and $\hat{m} = \text{diag}(m_u, m_d, m_s)$ are the quark fields and the current quark mass matrix for u , d , and s quarks, respectively; λ_a are the Gell-Mann matrices with $\lambda_0 = \sqrt{2/3}I$ in the three-flavor space; G_S and G_V are, respectively, the scalar and vector coupling constant; and the term with the coupling constant K represents the six-point Kobayashi-Maskawa- t' Hooft interaction, which breaks the

axial $U(1)_A$ symmetry. In the present study, we employ the parameters $m_u = m_d = 3.6$ MeV, $m_s = 87$ MeV, $G_S \Lambda^2 = 3.6$, $K \Lambda^5 = 8.9$, and the cutoff value in the momentum integral $\Lambda = 750$ MeV/ c given in Refs. [7,40]. The position of the critical point is sensitive to the value of G_V [4,5,7], which can thus be used to characterize the QCD phase diagram and the quark-matter EOS at large baryon chemical potentials, as detailed later. For the vector coupling G_V , we choose typical values of the reduced vector coupling constant $R_V = G_V/G_S = 0, 0.5$, and 1.1 , corresponding to the case without vector interaction, that from the Fierz transformation, and that from reproducing vector-meson masses [40], respectively.

From the mean-field approximation and assuming that the vector coupling is flavor independent, the thermodynamic potential Ω_{NJL} of a quark matter at temperature T and chemical potential μ_q for quark flavor q can be expressed in terms of its in-medium Dirac mass M_q and energy E_q as well as its effective chemical potential $\tilde{\mu}_q$ and condensate ϕ_q , i.e.,

$$\begin{aligned} \Omega_{\text{NJL}} = & -2N_c \sum_{q=u,d,s} \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} [E_q \\ & + T \ln(1 + e^{-\beta(E_q - \tilde{\mu}_q)}) + T \ln(1 + e^{-\beta(E_q + \tilde{\mu}_q)})] \\ & + G_S (\phi_u^2 + \phi_d^2 + \phi_s^2) - 4K \phi_u \phi_d \phi_s - \frac{1}{3} G_V \rho^2. \end{aligned} \quad (2)$$

In the above, the factor $2N_c = 6$ represents the spin and color degeneracy of the quark, and $\beta = 1/T$ is the inverse of the temperature.

The quark condensate ϕ_q is given by the integral,

$$\phi_q = -2N_c \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} \frac{M_q}{E_q} (1 - f_q - \bar{f}_q), \quad (3)$$

where f_q and \bar{f}_q denote, respectively, the phase-space distribution functions of quarks and antiquarks. For a static quark matter in thermal equilibrium, they are given by the Fermi-Dirac distributions,

$$\begin{aligned} f_q &= \frac{1}{\exp[(E_q - \tilde{\mu}_q)/T] + 1}, \\ \bar{f}_q &= \frac{1}{\exp[(E_q + \tilde{\mu}_q)/T] + 1}, \end{aligned} \quad (4)$$

where the effective chemical potential $\tilde{\mu}_q$ is related to the chemical potential μ_q through the relation

$$\tilde{\mu}_q = \mu_q - \frac{2}{3} G_V \rho. \quad (5)$$

In the above, $\rho = \rho_u + \rho_d + \rho_s$ is the total net quark number density, and for a single quark flavor q the net quark number density ρ_q can be calculated from

$$\rho_q = 2N_c \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} (f_q - \bar{f}_q). \quad (6)$$

The single-quark energy E_q in Eqs. (2)–(4) is given by $E_q = (M_q^2 + \vec{p}^2)^{1/2}$ with $\vec{p} = \vec{p}^* \mp \frac{2}{3} G_V \vec{j}$ being the momentum of the parton, where \vec{p}^* is the canonical momentum and $\vec{j} = \vec{j}_u + \vec{j}_d + \vec{j}_s$ is the net quark current with that for each

quark flavor q given by

$$\vec{j}_q = 2N_c \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} \frac{\vec{p}}{E_q} (f_q - \bar{f}_q). \quad (7)$$

The in-medium Dirac mass M_q in Eq. (3) is related to the quark condensate through the relations

$$\begin{aligned} M_u &= m_u - 2G_S \phi_u + 2K \phi_d \phi_s, \\ M_d &= m_d - 2G_S \phi_d + 2K \phi_s \phi_u, \\ M_s &= m_s - 2G_S \phi_s + 2K \phi_u \phi_d. \end{aligned} \quad (8)$$

Since the quark condensates and the Dirac masses are related to each other, Eqs. (3) and (8) need to be solved self-consistently through the iteration method. We note that the quark condensate or the Dirac mass is the order parameter for the chiral phase transition.

Equation (2) can be generalized to obtain the thermodynamic potential Ω_{pNJL} of the three-flavor pNJL model at finite temperature and quark chemical potential, and it is

$$\begin{aligned} \Omega_{\text{pNJL}} = & \mathcal{U}(\Phi, \bar{\Phi}, T) - 2N_c \sum_{q=u,d,s} \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} E_q \\ & - 2T \sum_{q=u,d,s} \int \frac{d^3 p}{(2\pi)^3} [\ln(1 + e^{-3\beta(E_q - \tilde{\mu}_q)}) \\ & + 3\Phi e^{-\beta(E_q - \tilde{\mu}_q)} + 3\bar{\Phi} e^{-2\beta(E_q - \tilde{\mu}_q)} \\ & + \ln(1 + e^{-3\beta(E_q + \tilde{\mu}_q)}) + 3\bar{\Phi} e^{-\beta(E_q + \tilde{\mu}_q)} \\ & + 3\Phi e^{-2\beta(E_q + \tilde{\mu}_q)}] + G_S (\phi_u^2 + \phi_d^2 + \phi_s^2) \\ & - 4K \phi_u \phi_d \phi_s - \frac{1}{3} G_V \rho^2. \end{aligned} \quad (9)$$

In the above, Φ ($\bar{\Phi}$) is the contribution from the Polyakov loop, which is related to the excess free energy due to a static quark (antiquark) in a hot gluon medium [41] and thus serves as an order parameter for the deconfinement phase transition. The form of the temperature-dependent effective potential $\mathcal{U}(\Phi, \bar{\Phi}, T)$ as a function of the Polyakov loops Φ and $\bar{\Phi}$ in Eq. (9) is taken from Ref. [5], i.e.,

$$\begin{aligned} \mathcal{U}(\Phi, \bar{\Phi}, T) = & -bT \{54e^{-a/T} \Phi \bar{\Phi} + \ln[1 - 6\Phi \bar{\Phi} \\ & - 3(\Phi \bar{\Phi})^2 + 4(\Phi^3 + \bar{\Phi}^3)]\}. \end{aligned} \quad (10)$$

The parameters $a = 664$ MeV and $b = 0.028 \Lambda^3$ are determined by the condition that the first-order phase transition in a pure gluon matter takes place at $T = 270$ MeV [5], and the simultaneous crossover of the chiral restoration and the deconfinement phase transition occurs around $T \approx 212$ MeV.

Minimizing the thermodynamic potential Ω_{pNJL} with respect to the quark condensates, i.e.,

$$\frac{\partial \Omega_{\text{pNJL}}}{\partial \phi_u} = \frac{\partial \Omega_{\text{pNJL}}}{\partial \phi_d} = \frac{\partial \Omega_{\text{pNJL}}}{\partial \phi_s} = 0, \quad (11)$$

leads to the similar expression for ϕ_u , ϕ_d , and ϕ_s as in Eq. (3) except the absence of the cutoff in the momentum integral and the replacement of the quark and antiquark phase-space distribution functions f_q and \bar{f}_q by F_q and \bar{F}_q , respectively,

given by

$$F_q = \frac{1 + 2\bar{\Phi}\xi_q + \Phi\xi_q^2}{1 + 3\bar{\Phi}\xi_q + 3\Phi\xi_q^2 + \xi_q^3},$$

$$\bar{F}_q = \frac{1 + 2\Phi\xi_q' + \bar{\Phi}\xi_q'^2}{1 + 3\Phi\xi_q' + 3\bar{\Phi}\xi_q'^2 + \xi_q'^3}, \quad (12)$$

with $\xi_q = e^{(E_q - \tilde{\mu}_q)/T}$ and $\xi_q' = e^{(E_q + \tilde{\mu}_q)/T}$. Similarly, the net quark density and current given by Eqs. (6) and (7), respectively, in the NJL model are modified by replacing f_q and \bar{f}_q with F_q and \bar{F}_q , respectively, and without the cutoff in the momentum integrals. For the values of the Polyakov loops Φ and $\bar{\Phi}$ in Eq. (12), they are determined by minimizing the thermodynamic potential Ω_{pNJL} with respect to Φ and $\bar{\Phi}$, i.e., $\frac{\partial \Omega_{\text{pNJL}}}{\partial \Phi} = \frac{\partial \Omega_{\text{pNJL}}}{\partial \bar{\Phi}} = 0$.

Starting from the thermodynamic potential, the energy density of the system can be obtained from the thermodynamical relation

$$\varepsilon = \Omega + \beta \frac{\partial}{\partial \beta} \Omega + \sum_{q=u,d,s} \mu_q \rho_q. \quad (13)$$

For the NJL model, the energy density is then

$$\varepsilon_{\text{NJL}} = -2N_c \sum_{q=u,d,s} \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} E_q (1 - f_q - \bar{f}_q) + G_S (\phi_u^2 + \phi_d^2 + \phi_s^2) - 4K \phi_u \phi_d \phi_s + \frac{1}{3} G_V \rho^2 - \varepsilon_0, \quad (14)$$

where the last term ε_0 is from the quark condensate contribution to the energy density in vacuum and is needed to ensure $\varepsilon_{\text{NJL}} = 0$ in vacuum.

Similarly, the energy density from the pNJL model can be expressed as

$$\varepsilon_{\text{pNJL}} = 54abe^{-a/T} \Phi \bar{\Phi} - 2N_c \sum_{q=u,d,s} \int_0^\Lambda \frac{d^3 p}{(2\pi)^3} E_q + 2N_c \sum_{q=u,d,s} \int \frac{d^3 p}{(2\pi)^3} E_q (F_q + \bar{F}_q) + G_S (\phi_u^2 + \phi_d^2 + \phi_s^2) - 4K \phi_u \phi_d \phi_s + \frac{1}{3} G_V \rho^2 - \varepsilon_0. \quad (15)$$

For both the NJL and the pNJL model, the pressure P can be calculated from the thermodynamic potential according to

$$P = -\Omega + \Omega_0, \quad (16)$$

where $\Omega_0 = \varepsilon_0$ is the thermodynamic potential in vacuum to ensure that the pressure is zero in vacuum.

The chiral phase transition boundary in the quark chemical potential versus temperature (μ - T) plane for different values of the reduced vector coupling constant R_V are displayed in the upper panels of Fig. 1 for both the NJL and the pNJL model. At small μ and low T , the system is in the chiral-symmetry-broken phase with large in-medium light




FIG. 1. (upper panel) Phase diagram in the μ - T plane from the NJL (left) and the pNJL (right) model with $\mu = \mu_u = \mu_d = \mu_s$ for different values of the reduced vector coupling constant $R_V = G_V/G_S$. (lower panel) Phase diagram as well as the $(\partial P / \partial \rho)_T < 0$ region in the ρ - T plane from the NJL (left) and the pNJL (right) model with $\mu = \mu_u = \mu_d = \mu_s$ without vector coupling. “CP” in the lower panels represents the critical point.

quark masses M_q . At large μ or high T , the system is in the chiral-symmetry-restored phase with small in-medium light quark masses $M_q \approx m_q$. The dashed and the solid lines in Fig. 1 denote, respectively, the boundary between the chiral-symmetry-broken and -restored phases that are connected by a crossover and a first-order transition. The point where the dashed and the solid line meets is the critical point in the QCD phase diagram. The corresponding critical temperature is seen to decrease with increasing value of R_V , as a result of the increasing stiffness of the EOS of baryon-rich quark matter. To better understand the relation between the EOS and the QCD critical point, we also plot in the lower panels of Fig. 1 the phase diagram in the ρ - T plane without vector coupling for both NJL and pNJL models. Also shown in these panels is the mechanical instability or the spinodal region with $(\partial P / \partial \rho)_T < 0$, where small density fluctuations are expected to grow exponentially in time. The spinodal region or the hadron-quark mixed phase shrinks with increasing temperature and disappears when the temperature is above the critical

temperature. With the increasing R_V , the spinodal region also shrinks with the critical point moving to a lower temperature. Due to the different phase-space distribution functions in the pNJL model [Eq. (12)] and the NJL model [Eq. (4)], the effective temperature of the quark matter is higher in the pNJL model than in the NJL model with the same vector coupling. This leads to a higher critical temperature in the pNJL model, which is more comparable with those predicted by the Dyson-Schwinger approach [8,9] and the functional renormalization-group method [10,11].

C. Nambu–Jona-Lasinio and Polyakov-Nambu-Jona-Lasinio parton transport models

For the NJL model, the single-particle Hamiltonian for a parton of flavor q ($q = u, d, s$) is given by

$$H_q = \sqrt{M_q^2 + \vec{p}^2} \pm \frac{2}{3} G_V \rho. \quad (17)$$

This leads to the following transport equation for the quark and antiquark phase-space distribution functions $f_q(\vec{r}, \vec{p})$ [42]:

$$\frac{\partial f_q}{\partial t} + \frac{\vec{p}_q}{E_q} \cdot \vec{\nabla} f_q - \vec{\nabla} H_q \cdot \vec{\nabla}_p f_q = I_{\text{coll}}, \quad (18)$$

where the collision integral I_{coll} describes the effect of quark scatterings on the quark phase-space distribution functions, and it depends on the scattering cross section and the Pauli-blocking factors in the final state of a scattering. Solving Eq. (18) by the test-particle method using an ensemble of parallel events [42,43], we obtain the following canonical equations of motion for partons between their scatterings:

$$\frac{d\vec{r}}{dt} = \frac{\vec{p}}{E_q}, \quad \frac{d\vec{p}}{dt} = -\frac{M_q}{E_q} \vec{\nabla} M_q \pm \left(\vec{E}_q + \frac{\vec{p}}{E_q} \times \vec{B}_q \right), \quad (19)$$

where

$$\vec{E}_q = -\frac{2}{3} G_V \left(\vec{\nabla} \rho + \frac{\partial \vec{j}}{\partial t} \right) \quad (20)$$

is the effective electric field, and

$$\vec{B}_q = \frac{2}{3} G_V \vec{\nabla} \times \vec{j} \quad (21)$$

is the effective magnetic field.

For the parton scattering cross section in the collision integral of the transport equation, it is obtained from the specific shear viscosity of the partonic matter as described in Appendix A. As to the treatment of parton scattering with the Pauli blocking factor, we use the stochastic method [35] based on a full ensemble of N_{TP} parallel events, as detailed in Appendix B.

The quark and antiquark phase-space distribution functions f_q and \bar{f}_q in the transport model simulations are calculated by summing the particles from N_{TP} parallel events in the local rest frame of the cells around (\vec{r}, \vec{p}) in the phase space, i.e.,

$$f_q(\vec{r}, \vec{p}) \sim \frac{1}{N_{TP}} \sum_{i \in q} \delta(\vec{r} - \vec{r}_i) \delta(\vec{p} - \vec{p}_i),$$

$$\bar{f}_q(\vec{r}, \vec{p}) \sim \frac{1}{N_{TP}} \sum_{i \in \bar{q}} \delta(\vec{r} - \vec{r}_i) \delta(\vec{p} - \vec{p}_i). \quad (22)$$




FIG. 2. Momentum distribution of u quarks at different times for quark matter in a box of volume V with periodic boundary conditions. The initial distribution at $t = 0$ is that of the thermally equilibrated distribution given by Eq. (4) for the NJL model and Eq. (12) for the pNJL model (dotted lines). At $t = 10$ fm/c, it reaches the Boltzmann distribution due to the neglect of Pauli blockings in the scatterings. After turning on Pauli blockings at this time, the distribution changes back to the initial distribution, as seen at $t = 30$ fm/c. See text for details.

The NJL-based transport equation given in Eq. (18) and the resulting equations of motion for quarks and antiquarks can be generalized to that based on the pNJL model by replacing the quark and antiquark phase-space distribution functions f_q and \bar{f}_q with F_q and \bar{F}_q , respectively. A modified treatment of the Pauli blocking factor in partonic scattering is, however, required to ensure the approach of F_q and \bar{F}_q to the thermally equilibrated values given in Eq. (12), as described in Appendix C. To illustrate the validity of this method, we consider the evolution of a partonic matter of uniform spatial distribution of density $\rho = 1.5 \text{ fm}^{-3}$ in a box with periodic boundary conditions. Starting from the Fermi-Dirac momentum distributions of Eq. (4) for the NJL model or the momentum distributions of Eq. (12) for the pNJL model at temperature $T = 200$ MeV, the system evolves in time by solving the NJL or pNJL transport model without the Pauli blocking factors in parton scatterings for the first 10 fm/c. As shown in Fig. 2 for u quarks as an example, the momentum distribution evolves into a Boltzmann distribution shown by the dashed line, which is obviously different from the initial momentum distribution shown by the dotted line. Turning on the Pauli blocking factors at $t = 10$ fm/c, it is seen that the momentum distribution gradually changes back to the initial ones, demonstrating the correctness of our treatment of the collision integral in the NJL and pNJL models.

III. RESULTS AND DISCUSSIONS

Based on the extended AMPT model with the pNJL transport model to describe the evolution of the partonic phase, we now study the elliptic flow v_2 splittings between particles and their antiparticles in relativistic heavy-ion collisions at RHIC-BES energies. Although there have been a number of possible explanations for the observed v_2 splittings in the STAR experiments [44–50], our studies using different mean-field potentials for particles and their antiparticles in the baryon-rich matter are the only one that is based on a comprehensive transport model [27–29]. Taking the case in the partonic phase as an example, antiquarks are generally affected by a more attractive mean-field potential than quarks in the almond-shaped baryon-rich partonic matter. Thus they tend to stay longer in the produced matter compared with quarks which move along the short axis of this matter. Because of the difference in their mean-field potentials, quarks then generally have a larger v_2 than antiquarks. Such v_2 splittings are expected to be preserved after hadronization with a proper coalescence algorithm [36] and after hadronic evolution with different hadronic mean-field potentials for baryons and antibaryons as well as for K^+ and K^- and other particle and antiparticle pairs [27].

A. Time evolution of matter produced in relativistic heavy-ion collisions

Besides scatterings, the evolution of a partonic matter is also affected by the mean-field potentials of quarks and antiquarks. In the nonrelativistic approximation, they can be expressed as

$$\begin{aligned} U_{u(\bar{u})} &= -2G_S\phi_u + 2K\phi_d\phi_s - U_u^0 \pm \frac{2}{3}G_V\rho, \\ U_{d(\bar{d})} &= -2G_S\phi_d + 2K\phi_u\phi_s - U_d^0 \pm \frac{2}{3}G_V\rho, \\ U_{s(\bar{s})} &= -2G_S\phi_s + 2K\phi_u\phi_d - U_s^0 \pm \frac{2}{3}G_V\rho, \end{aligned} \quad (23)$$

where the term U_q^0 ($q = u, d, s$) is introduced to ensure that the mean-field potential vanishes in vacuum, and the upper (lower) signs in these equations are for quarks (antiquarks). Their density dependence in a quark matter at temperature $T = 200$ MeV is shown in Fig. 3. It is seen that the scalar potentials shown by dash-dotted lines are attractive, which results from the smaller effective quark masses in the medium than in vacuum. Because the pNJL model has a different thermal distribution from that in the NJL model due to the contribution from Polyakov loops, quark condensates in the pNJL model are less negative than in the NJL model at same temperature and density, leading to a more attractive scalar potential, as seen in Fig. 3. The mean-field potentials of quarks and antiquarks become very different when the vector coupling constant has a finite value, and the difference increases with increasing net quark density. Since the mean-field potential affects the elliptic flow v_2 of particles, measuring the v_2 splitting in heavy-ion collisions allows us to study the mean-field potential difference between quarks and antiquarks, and thus their vector interactions, which can affect significantly the EOS of quark matter at large chemical potentials and the QCD phase diagram in the NJL or pNJL model. The different




FIG. 3. Mean-field potentials of quarks and antiquarks of different flavors from nonrelativistic reduction as a function of net quark density with $\mu_u = \mu_d = \mu_s$ and at the temperature $T = 200$ MeV for the NJL (left) model and the pNJL (right) model.

mean-field potentials between s quarks and anti s quarks are qualitatively similar to those between light quarks and their antiquarks.

The time evolution of average densities of quarks (ρ_Q) and antiquarks ($\rho_{\bar{Q}}$) in the central region of minibias Au + Au collisions at different collision energies is displayed in Fig. 4, with $\rho_Q - \rho_{\bar{Q}} = \rho$ being the net quark number density. These results are obtained with a parton scattering cross section that corresponds to the specific shear viscosity of 0.14 and 0.06 in the NJL and the pNJL transport model, respectively, in order to reproduce measured v_2 of charged hadrons, as




FIG. 4. Average central number densities of quarks (ρ_Q) and antiquarks ($\rho_{\bar{Q}}$) as a function of time in minibias (0%–80%) Au + Au collisions at different collision energies for both NJL (left) and pNJL (right) models.




FIG. 5. Time components of the vector potential $U_V^0 = \frac{2}{3}G_V\rho$ in the central region of heavy-ion collisions as a function of time at different collision energies for both NJL (left) and pNJL (right) models.

discussed later. It is seen that partons are generally formed after 0.2 fm/c. With the gradual formation of partons, the central density peaks at around 0.5 fm/c at higher collision energies, while at the collision energy of $\sqrt{s_{NN}} = 7.7$ GeV the duration of the peak density lasts for a longer time due to the finite thickness of initial collision zone mentioned in Sec. II A. The parton phase ends around 4–5 fm/c when the central energy density drops to about 0.8 GeV/fm³. In peripheral collisions, since the maximum energy density cannot reach 0.8 GeV/fm³, initial partons from the AMPT model are immediately converted back to hadrons without undergoing any scatterings. The average lifetime of the partonic phase is seen to generally decrease with increasing collision energy; as a result of the stronger expansion in high-energy collisions and the quick drop of the energy density. The densities are seen to be higher in the pNJL model than in the NJL model, where the parton densities are calculated by counting those with their momenta lower than Λ [see Eq. (6)], while the lifetime of the partonic phase is slightly longer in the pNJL model than in the NJL model due to the higher energy density from the contribution of Polyakov loops.

The time component of the vector potential $U_V^0 = \frac{2}{3}G_V\rho$ is the driving force for the v_2 splitting between quarks and antiquarks because the space component of the vector potential, which is proportional to the net quark current, takes time to develop and becomes important only at later stages when v_2 has mostly been developed. It is thus of importance to know how U_V^0 evolves with time, and this is displayed in Fig. 5 for the central region of the collision system. Although the U_V^0 has the longest duration at $\sqrt{s_{NN}} = 7.7$ GeV, it is interesting to see that its peak value depends nonmonotonically on the collision energy. The largest peak value of U_V^0 is reached at about $\sqrt{s_{NN}} = 11.5$ GeV, due to the balance between the smaller baryon chemical potential at higher collision energies and a thicker quark matter in the longitudinal direction formed at lower collision energies. This has observational consequences

on the collision energy dependence of v_2 splitting, as shown later. The net quark density ρ on which the vector potential depends is given by the difference between the quark and antiquark density shown in Fig. 4, where in the NJL model only partons with their momenta lower than Λ are counted. Since antiquarks have a much softer momentum spectrum than quarks due to their attractive mean-field potentials, the difference between the numbers of quarks and antiquarks is much larger for partons with high momenta than for those with low momenta. Thus, excluding partons with momenta higher than Λ in the NJL model generally results in a vector potential that is weaker than that in the pNJL model in which partons of all momenta are included in calculating the net quark density, as seen in Fig. 4.

It is of great interest to know the evolution trajectory of the partonic matter formed in heavy-ion collisions in the QCD phase diagram. Assuming that the central region of the produced matter is in thermal and chemical equilibrium, the temperature T and the quark chemical potential μ can be determined from the net quark density ρ and the energy density ε of quarks and antiquarks by using quark and antiquark momentum distributions given by Eq. (4) for the NJL model and Eq. (12) for the pNJL model. The resulting trajectories in the T - ρ plane and T - μ plane are displayed in Fig. 6 for collisions at various energies. It is seen that, from the AMPT initial conditions, all trajectories follow a counterclockwise path in the phase diagram. As expected, higher temperatures are generally reached at higher collision energies. Also, the trajectories from the NJL transport model go through a higher temperature compared with that from the pNJL transport model, as a result of the different forms for the momentum distribution [Eqs. (4) and (12)] and thus different “effective” temperatures. On the other hand, the partonic phase has a higher net quark density ρ from the pNJL transport model than from the NJL model, as already shown in Figs. 4 and 5. As seen from the phase diagram shown in Fig. 1, none of the trajectories passes through the spinodal region or the critical point for collision energies considered here.

As mentioned above, Pauli blockings are incorporated in the NJL transport model and also in the pNJL transport model using a modified treatment in order to ensure that correct thermal distributions are reached when the partonic matter is in thermal equilibrium. Due to the more attractive scalar potential in the pNJL model as shown in the Fig. 3, a smaller specific shear viscosity η/s and thus a larger partonic scattering cross section is used in this model than in the NJL model to obtain the same charged-particle v_2 in these two models. To better understand parton scatterings in the partonic phase of heavy-ion collisions, we show in Fig. 7 the rate of successful collisions that are not Pauli blocked as well as the ratio of successful to attempted collision rates. As expected, higher successful collision rates are observed at higher collision energies. Also, the successful collision rate generally increases with time due to the constant η/s used in the model calculation, which results in a larger parton scattering cross section when the expanding partonic matter is at lower temperatures and/or densities (see Fig. 15 in Appendix A). Although a smaller η/s is used in the pNJL transport model, the ratio of successful to attempted collision rates is much smaller than




FIG. 6. Evolution trajectories of the partonic matter in the plane of temperature T versus density ρ (upper) as well as those in the plane of temperature T versus chemical potential μ (lower) in minias (0%–80%) collisions at different collision energies for both NJL (left) and pNJL (right) models.

that in the NJL transport model using a larger η/s . This is again due to the different phase-space distributions and treatments of parton scatterings. Figure 7 further shows that while the Pauli blocking effect in the NJL model does not depend much on the collision energy, parton scatterings in the pNJL model are less blocked at higher collision energies.

For a complete description of the evolution of the matter produced in heavy-ion collisions, we show in Fig. 8 the time evolution of the average central baryon and antibaryon densities during the hadronic phase of a heavy-ion collision. Converting freeze-out partons via an improved hadronization scheme based on the phase-space coalescence model [36], hadrons are produced after about 5 fm/c after the production of initial partons in a heavy-ion collision. The maximum baryon density of about 0.1 fm^{-3} is reached soon after hadronization, and this value is approximately independent of the collision energy. For the maximum antibaryon density, it is reached at about the same time, but its magnitude increases with the increasing collision energy. The pNJL model




FIG. 7. Successful collision rate (upper) and ratios of successful to attempted collision rates (lower) as a function of time in the partonic phase in heavy-ion collisions at different collision energies in both NJL (left) and pNJL (right) models.

generally shows similar baryon and antibaryon density evolutions compared with the NJL model. In such baryon-rich hadronic matter, antiparticles are expected to have more attractive mean-field potentials than particles [27]. As a result, the v_2 splittings between quarks and antiquarks developed during the partonic phase can be further amplified in those between hadrons and their antiparticles.

B. Description of physical observables

Having the evolution of produced matter in heavy-ion collisions described in the above section using the NJL or pNJL transport model, we show in this section that both models can describe satisfactorily many measured observables in heavy-ion collisions, particularly those sensitive to the bulk properties of the matter produced in these collisions. Figure 9 shows the comparison of the transverse momentum (p_T) dependence of charged-particle v_2 at different collision energies with corresponding experimental data from the STAR Collaboration [51]. By using a constant specific shear viscosity of $\eta/s = 0.14$ for the NJL transport model and 0.06 for the




FIG. 8. Average central baryon (ρ_B) and antibaryon ($\rho_{\bar{B}}$) number densities in the hadronic phase as a function of time in minibus (0%–80%) Au + Au collisions at different collision energies from the extended AMPT model with the partonic phase described by the NJL (left) or the pNJL (right) model.

pNJL transport model, the p_T dependence of v_2 in midcentral (20%–30%) Au + Au collisions at RHIC-BES energies are reasonably well reproduced. The smaller η/s , which corresponds to a larger partonic scattering cross section, used in the pNJL transport model than that used in the NJL transport model is because partons in the pNJL model are affected by a more attractive scalar potential as shown in Fig. 3. Although the η/s used in the pNJL transport model is smaller than its




FIG. 9. Transverse momentum dependence of mid-pseudorapidity charged-particle elliptic flow in midcentral (20%–30%) Au + Au collisions at different collision energies from the extended AMPT model with the partonic phase described by the NJL or the pNJL model. Experimental data are from Ref. [51].




FIG. 10. Transverse momentum spectra of mid-pseudorapidity π^+ , K^+ , and protons in central (0%–5%) Au + Au collisions at different collision energies from the extended AMPT model with the partonic phase described by the NJL or the pNJL model. Experimental data are taken from Ref. [53].

theoretical lower bound [52], it is an effective one since we have neglected its dependence on temperature and chemical potential.

In Fig. 10, we compare the p_T spectra of representative particles, i.e., π^+ , K^+ , and protons, at different collision energies with the corresponding experimental data from the STAR Collaboration [53]. Since most hadrons are produced in their resonance states right after hadronization, these hadrons experience hadronic elastic and inelastic scatterings and decays during the hadronic evolution. Except for a slightly stiffer p_T spectra for π^+ at lower collision energies and a softer p_T spectra for protons at higher collision energies, the extended AMPT model describes reasonably well the experimental data from heavy-ion collisions at RHIC-BES energies.

Before discussing the v_2 splitting of final hadrons, we first show in Fig. 11 the p_T dependence of quark and antiquark v_2 at the end of the partonic evolution at $\sqrt{s_{NN}} = 11.5$ GeV. In the presence of a strong vector interaction with $R_V = 1.1$, a larger v_2 is observed for quarks than their antiquarks, consistent with the intuitive picture discussed at the beginning of this section. Although the vector potential is slightly stronger in the pNJL transport model than in the NJL model as shown in Fig. 5, the v_2 splittings between light quarks and their antiquarks are similar in the two models, and this is because there are more successful scatterings in the pNJL model than in the NJL model as shown in Fig. 7. In general, the v_2 splitting between high-momentum particles and antiparticles is larger in the pNJL transport model than that in the NJL transport model, since they are not affected by the vector potential due to the momentum cutoff in the latter case. This is especially so for strange quarks with a larger dynamical mass.

The splitting between p_T -integrated v_2 of quarks and their antiquarks, which are obtained by their respective average v_2 over all p_T , at the end of the partonic phase at different collision energies is displayed in Fig. 12. The v_2 splitting is seen to increase with increasing R_V . The negative v_2 splitting between s and \bar{s} in the case without the vector interaction shown in the




FIG. 11. Elliptic flows of mid-pseudorapidity quarks and antiquarks as a function of transverse momentum just before hadronization in heavy-ion collisions at $\sqrt{s_{NN}} = 11.5$ GeV from both NJL (left) and pNJL (right) transport models.

right panels of Fig. 12 is due to more successful scatterings for \bar{s} than s , as a result of a more diffusive momentum distribution for Pauli blockings for \bar{s} . For the case using a larger R_V , the v_2 splitting shows a nonmonotonic dependence on the collision energy, especially for that between light quarks and their antiquarks with a peak v_2 splitting at $\sqrt{s_{NN}} = 11.5$ GeV. This behavior is similar to the nonmonotonic energy dependence of the magnitude of the vector potential shown in Fig. 5,




FIG. 12. Difference between the p_T -integrated v_2 of quarks and antiquarks just before hadronization for collisions at different energies from both NJL (upper) and pNJL (lower) transport models.




FIG. 13. Difference between the p_T -integrated v_2 of nucleon resonances and their antiparticles, positively charged kaon-like mesons and negatively charged kaon-like mesons, and strange baryon resonances and their antiparticles, right after hadronization at different collision energies from the extended AMPT model with the partonic phase described by the NJL (upper) or the pNJL (lower) transport model with different values for the reduced vector coupling constant R_V .

and this is due to the competition between the collision energy dependence of various factors, e.g., the longitudinal Lorentz contraction, the net quark density, and the expansion speed of the produced matter. This behavior is different from that observed in Ref. [29], where the finite thickness of the quark matter in the longitudinal direction was not properly incorporated. Although the v_2 splitting between light quarks and their antiquarks is similar in the NJL and pNJL transport models, that between strange quarks and their antiquarks is larger in the pNJL transport model compared with that in the NJL transport model for the reason already given in the discussions on the splitting of differential v_2 shown in Fig. 11.

To see how the v_2 splittings between quarks and antiquarks are carried over to those between hadrons and their antiparticles, we display in Fig. 13 the v_2 splitting between nucleon resonances and their antiparticles, positively charged kaon-like mesons (K^+ and K^{*+}) and negatively charged kaon-like mesons (K^- and K^{*-}), and strange baryon resonances and their antiparticles, right after hadronization when most hadrons are in their resonance states. It is seen that the v_2 splitting between baryons and antibaryons first increases and then decreases with the increasing collision energy, consistent with the collision-energy dependence of the v_2 splitting between quarks and antiquarks shown in Fig. 12. The situation is more complicated for mesons, which are composed of a quark and an antiquark, due to the opposite effects from the vector potential on quark and antiquark v_2 . For the larger $R_V = 1.1$, although negatively charged kaon-like mesons have a slightly larger v_2 than that of positively charged kaon-like mesons at low collision energies due to the dominating effect of heavier strange quarks, their v_2 splitting is reversed at high collision energies. The v_2 splitting between nucleon resonances and




FIG. 14. Difference between the p_T -integrated v_2 of freeze-out nucleons and antinucleons, K^+ and K^- , and Λ and $\bar{\Lambda}$ at different collision energies from the extended AMPT model with the partonic phase described by the NJL (upper) or the pNJL (lower) model. Corresponding experimental data are taken from the STAR Collaboration [22].

their antiparticles as well as that between kaon-like mesons and their antiparticles are seen to be similar in both the NJL and pNJL transport models, while that between strange baryon resonances and their antiparticles is larger in the pNJL transport model than in the NJL transport model, consistent with the larger $v_2(s) - v_2(\bar{s})$ in the pNJL transport model shown in Fig. 12.

We finally compare in Fig. 14 the v_2 splittings between final nucleons and antinucleons,¹ K^+ and K^- , and Λ and $\bar{\Lambda}$ at different collision energies with corresponding experimental data from the STAR Collaboration [22]. It is seen that the v_2 splittings between nucleons and antinucleons as well as those between Λ and $\bar{\Lambda}$ after hadronic evolution are similar to those of baryons and antibaryons right after hadronization shown in Fig. 13, and this is because of the more attractive potentials for antibaryons than baryons [27]. This is very different for the v_2 splitting between K^+ and K^- , which has an opposite sign compared with that between positively charged and negatively charged kaon resonances shown in the middle panel of Fig. 13, as a result of the repulsive potential for K^+ and the attractive potential for K^- [27]. Even for $R_V = 0$, there are v_2 splittings between hadrons and their antiparticles from our model calculations, and this is mainly due to their different hadronic potentials. The collision energy dependence of the v_2 splittings between hadrons and their antiparticles displayed in Fig. 14 shows a similar nonmonotonic behavior as that of the parton v_2 splittings, especially for the v_2 splitting between Λ and $\bar{\Lambda}$. Compared with the experimental data from the STAR Collaboration, the v_2 splitting between nucleons and antinu-

cleons can be reproduced reasonably well within $R_V = 0-1.1$ but favors a smaller R_V at higher collision energies. Although our models reproduce the v_2 splitting between K^+ and K^- at $\sqrt{s_{NN}} = 7.7$ GeV, they overestimate their v_2 splitting at higher collision energies. Since the partonic and hadronic phase have different effects on the v_2 splitting between K^+ and K^- , a more accurate handling of the lifetimes for the two phases is called for. For the v_2 splitting between Λ and $\bar{\Lambda}$, our results underestimate the experimental data at $\sqrt{s_{NN}} = 7.7$ GeV, while the data at higher collision energies can be reproduced by the extended AMPT model with $R_V = 1.1$ in the NJL transport model and $R_V = 0.5$ in the pNJL transport model. A detailed fit of the value of R_V by taking into account its collision-energy dependence via the incorporation of a baryon chemical-potential-dependent R_V needs to be investigated in future studies.

IV. CONCLUSION

With the partonic phase of heavy-ion collisions described by a partonic transport model that is based on the three-flavor Nambu–Jona-Lasinio (NJL) model or its extension with the inclusion of Polyakov loops, we have revisited the elliptic flow (v_2) splittings between particles and their antiparticles in relativistic heavy-ion collisions at RHIC-BES energies. We have checked that for a partonic matter in a box with periodic boundary conditions, the NJL and pNJL transport models lead to their respective thermal momentum distributions if the Pauli-blocking factors are included in the collision integral of the NJL model and a properly modified treatment of parton scatterings is introduced in the pNJL transport model. In a partonic matter of a given temperature and baryon chemical potential, it is found that the pNJL transport model gives a higher net quark density and a stronger vector potential than those from the NJL transport model. As a result, a larger v_2 splitting is found between s and \bar{s} quarks in heavy-ion collisions, and this affects the v_2 splitting between final Λ and $\bar{\Lambda}$. On the other hand, due to the competition between effects from the longitudinal Lorentz contraction, the baryon chemical potential, and the expansion time at different collision energies, the magnitude of the vector potential and the v_2 splitting between quarks and antiquark changes non-monotonically with the increasing collision energy for a given vector coupling constant in both the NJL and the pNJL model. Consequently, the v_2 splitting between final hadrons and their antiparticles also shows a nonmonotonic dependence on the collision energy. The resulting v_2 splitting between K^+ and K^- reproduces the STAR data at $\sqrt{s_{NN}} = 7.7$ GeV but overestimates it at higher collision energies. For the v_2 splitting between protons and antiprotons, the STAR data at $\sqrt{s_{NN}} = 7.7$ GeV can be reproduced with the range of vector coupling in both NJL and pNJL transport models considered in the present study, while the STAR data at higher collision energies favor a weak vector interaction in these transport models. Although our calculations underestimate the STAR data on the v_2 splitting between Λ and $\bar{\Lambda}$ at $\sqrt{s_{NN}} = 7.7$ GeV, a strong (weak) vector interaction is favored in the NJL (pNJL) transport model for collisions at higher energies. Our study has therefore provided useful insight on the determination

¹Experimental data are for the v_2 splitting between protons and antiprotons, which are compared with that between nucleons and antinucleons from the model calculation for better statistics.

of the vector interaction in the partonic matter produced in heavy-ion collisions, which will help understand the equation of state of quark matter at large baryon chemical potentials and thus the QCD phase diagram.

ACKNOWLEDGMENTS

J.X. was supported by the National Natural Science Foundation of China under Grant No. 11922514. C.M.K. was supported by the US Department of Energy under Award No. DESC0015266 and the Welch Foundation under Grant No. A-1358.

APPENDIX A: RELATION BETWEEN THE PARTON SCATTERING CROSS SECTION AND THE SPECIFIC SHEAR VISCOSITY

The specific shear viscosity in the NJL model has been studied in Ref. [54] from the quark-meson interactions, and that in the pNJL model has been studied in Ref. [55] based on the relaxation-time approximation. In the present study, we determine the isotropic parton scattering cross section σ from the shear viscosity η through the empirical relation

$$\eta = \frac{4\langle p \rangle}{15\sigma_{tr}}, \quad (\text{A1})$$

where

$$\langle p \rangle = \frac{2N_c \sum_q \int \frac{d^3p}{(2\pi)^3} p(n_q + \bar{n}_q)}{2N_c \sum_q \int \frac{d^3p}{(2\pi)^3} (n_q + \bar{n}_q)} \quad (\text{A2})$$

is the average parton momentum with n_q and \bar{n}_q being the occupation probability for parton species $q = u, d, s$ quarks and their antiquarks, i.e., $n_q = f_q$ and $\bar{n}_q = \bar{f}_q$ for the NJL model and $n_q = F_q$ and $\bar{n}_q = \bar{F}_q$ for the pNJL model. The σ_{tr} in Eq. (A1) is the transport cross section defined as

$$\sigma_{tr} = \int d\Omega \frac{d\sigma}{d\Omega} (1 - \cos^2 \theta). \quad (\text{A3})$$

For an isotropic cross section σ , one has $\sigma_{tr} = \frac{2}{3}\sigma$. The entropy density s of a quark matter is given by

$$s = -2N_c \sum_q \int \frac{d^3p}{(2\pi)^3} [n_q \ln n_q + (1 - n_q) \ln (1 - n_q)] - 2N_c \sum_q \int \frac{d^3p}{(2\pi)^3} [\bar{n}_q \ln \bar{n}_q + (1 - \bar{n}_q) \ln (1 - \bar{n}_q)]. \quad (\text{A4})$$

By using a constant specific shear viscosity $\eta/s = 0.14$ for the NJL model and 0.06 for the pNJL model, the cross section σ at different net quark number densities ρ and temperatures T are displayed in Fig. 15. It is seen that the cross section is around 1 fm^2 in the high-temperature region, while it can be as large as 10 fm^2 at low temperatures and/or densities.




FIG. 15. The ρ and T dependence of parton scattering cross section determined from a constant specific shear viscosity $\eta/s = 0.14$ for the NJL (left) model and 0.06 for the pNJL (right) model.

APPENDIX B: STOCHASTIC TREATMENT OF PARTON SCATTERING WITH PAULI BLOCKINGS

We employ the stochastic method [35] to treat elastic scatterings between partons. The collision probability for a pair of partons in a volume $\Delta V = \Delta x \Delta y \Delta z$ and a time interval Δt is

$$P_c = v_{12} \sigma \frac{\Delta t}{\Delta V}, \quad (\text{B1})$$

with

$$v_{12} = \frac{\sqrt{[s - (M_1 + M_2)^2][s - (M_1 - M_2)^2]}}{2E_1 E_2} \quad (\text{B2})$$

being the Møller velocity. In the above, s is the square of the invariant mass of the parton pair, and $E_{1(2)} = (M_{1(2)}^2 + p_{1(2)}^2)^{1/2}$ is the energy of the parton, with $M_{1(2)}$ being the in-medium parton mass from the NJL or the pNJL model, and $p_{1(2)}$ being the momentum of the parton in the rest frame of the cell. The σ in Eq. (B1) is an isotropic cross section, determined by the specific shear viscosity as detailed in Appendix A. The volume of each cell is set as $\Delta x = \Delta y = 0.5 \text{ fm}$ and $\Delta z = 0.25 \text{ fm}$, and only parton pairs in the same cell can collide with each other. The time step $\Delta t = 0.02 \text{ fm}/c$ is used in the calculation. The case $P_c > 1$ is equal to the case of $P_c = 1$, and in this way the collision rate from a too large cross section is limited by the values of ΔV and Δt .

After an attempted collision, the momentum of each parton in the center-of-mass frame of the collision pair is sampled isotropically. After the momenta of the two partons are boosted to the rest frame of the local cell, they change from $\vec{p}_{1(2)}$ to $\vec{p}'_{1(2)}$. For a given flavor, the occupation probability in the rest frame of the local cell after the attempted collision is calculated from the test-particle method,

$$n_q(\vec{r}, \vec{p}) \sim \frac{1}{N_{TP}} \sum_{i \in q} \delta(\vec{r} - \vec{r}_i) \delta(\vec{p} - \vec{p}'_i), \quad (\text{B3})$$

where N_{TP} is the test particle number. The occupation probability is calculated separately for partons with different flavors and also separately for quarks and antiquarks. The

Pauli blocking probability is then $1 - [1 - n_q(\vec{r}, \vec{p}'_1)][1 - n_q(\vec{r}, \vec{p}'_2)]$. The collision is successful if a random number within $[0,1]$ is larger than the Pauli blocking probability, otherwise the momenta of the two partons $\vec{p}_{1(2)}$ are retained.

APPENDIX C: MODIFIED PAULI BLOCKING IN THE POLYAKOV-LOOP NAMBU-JONA-LASINIO TRANSPORT MODEL

The collision integral in the Boltzmann equation with quantum correction generally contains the factor $(f_3 f_4 g_1 g_2 - f_1 f_2 g_3 g_4)$ for the $1 + 2 \rightarrow 3 + 4$ process, where $g_{1,2,3,4}$ is equal to $1 - f_{1,2,3,4}$ if the Fermi-Dirac statistics is adopted. For the NJL transport model, as mentioned in Appendix B, the successful collision probability is $(1 - f_3)(1 - f_4)$ when the Pauli blocking is applied. For the Fermi-Dirac distribution given in Eq. (4), $g = \xi f$ with $\xi = e^{(E - \tilde{\mu})/T}$ (neglecting the subscript flavor index q) is satisfied. To obtain the thermal distribution F [Eq. (12)] in the pNJL transport model, a similar consideration can be applied by taking the successful collision probability as $g_3 g_4$ with $g = \xi F$. Although $F_{3,4}$ can be calculated using the test-particle method through Eq. (22),

ξ is a thermodynamic quantity, which needs to be calculated by assuming local thermal equilibrium as described below.

Taking the expression of $F = F_q$ as an example, Eq. (12) can be rewritten as

$$\xi^3 + A\xi^2 + B\xi + C = 0, \quad (C1)$$

with

$$A = \frac{\Phi(3F - 1)}{F}, \quad B = \frac{\bar{\Phi}(3F - 2)}{F}, \quad C = \frac{F - 1}{F}. \quad (C2)$$

For $0 < F < 1$, the only real solution to Eq. (C1) is

$$\xi = \frac{1}{3}[-A + (-D + \sqrt{D^2 + H^3})^{1/3} + (-D - \sqrt{D^2 + H^3})^{1/3}], \quad (C3)$$

with

$$D = \frac{27C - 9AB + 2A^3}{2}, \quad H = 3B - A^2. \quad (C4)$$

In this way, ξ can be expressed as a function of F , Φ , and $\bar{\Phi}$. Similarly, the ξ' in \bar{F}_q can be expressed as a function of \bar{F} , Φ , and $\bar{\Phi}$.

-
- [1] C. Bernard, C. Bernard, T. Burch, C. DeTar, J. Osborn, S. Gottlieb, E. B. Gregory, D. Toussaint, U. M. Heller, and R. Sugar, *Phys. Rev. D* **71**, 034504 (2005).
- [2] Y. Aoki *et al.*, *Nature (London)* **443**, 675 (2006).
- [3] A. Bazavov, T. Bhattacharya, M. Cheng, C. DeTar, H. T. Ding, S. Gottlieb, R. Gupta, P. Hegde, U. M. Heller, F. Karsch, E. Laermann, L. Levkova, S. Mukherjee, P. Petreczky, C. Schmidt, R. A. Soltz, W. Soeldner, R. Sugar, D. Toussaint, W. Unger, and P. Vranas, *Phys. Rev. D* **85**, 054503 (2012).
- [4] M. Asakawa and K. Yazaki, *Nucl. Phys. A* **504**, 668 (1989).
- [5] K. Fukushima, *Phys. Rev. D* **77**, 114028 (2008); **78**, 039902(E) (2008).
- [6] S. Carignano, D. Nickel, and M. Buballa, *Phys. Rev. D* **82**, 054009 (2010).
- [7] N. M. Bratovic, T. Hatsuda, and W. Weise, *Phys. Lett. B* **719**, 131 (2013).
- [8] X. Y. Xin, S. X. Qin, and Y. X. Liu, *Phys. Rev. D* **90**, 076006 (2014).
- [9] C. S. Fischer, J. Luecker, and C. A. Welzbacher, *Phys. Rev. D* **90**, 034022 (2014).
- [10] W. J. Fu, J. M. Pawłowski, and F. Rennecke, *Phys. Rev. D* **101**, 054032 (2020).
- [11] F. Gao and J. M. Pawłowski, *Phys. Rev. D* **102**, 034027 (2020).
- [12] P. C. Chu, X. Wang, L. W. Chen, and M. Huang, *Phys. Rev. D* **91**, 023003 (2015).
- [13] H. Liu, J. Xu, and C. M. Ko, *Phys. Lett. B* **803**, 135343 (2020).
- [14] E. L. Bratkovskaya, W. Cassing, P. Moreau, L. Oliva, O. E. Soloveva, and T. Song, [arXiv:1908.00451](https://arxiv.org/abs/1908.00451).
- [15] J. Aichelin, E. Bratkovskaya, A. Le Fèvre, V. Kireyeu, V. Kolesnikov, Y. Leifels, V. Voronyuk, and G. Coci, *Phys. Rev. C* **101**, 044905 (2020).
- [16] D. Oliinychenko, V. Steinberg, J. Weil, J. Staudenmaier, M. Kretz, A. Schäfer, H. E. Petersen, S. Ryu, J. Rothermel, J. Mohs, F. Li, A. Sorensen, D. Mitrovic, L. Pang, J. Hammelmann, A. Goldschmidt, M. Mayer, O. Garcia-Montero, N. Kübler, and Nikita, smash-transport/smash: Smash-2.0 (2020) [<https://doi.org/10.5281/zenodo.4336358>].
- [17] K. Rajagopal, G. W. Ridgway, R. Weller, and Y. Yin, *Phys. Rev. D* **102**, 094025 (2020).
- [18] C. Shen and L. Yan, *Nucl. Sci. Tech.* **31**, 122 (2020).
- [19] M. A. Stephanov, *Phys. Rev. Lett.* **102**, 032301 (2009).
- [20] M. Asakawa, S. Ejiri, and M. Kitazawa, *Phys. Rev. Lett.* **103**, 262301 (2009).
- [21] M. M. Aggarwal *et al.* (STAR Collaboration) *Phys. Rev. Lett.* **105**, 022302 (2010).
- [22] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **112**, 032302 (2014).
- [23] J. Adam *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **126**, 092301 (2020).
- [24] K. J. Sun, L. W. Chen, C. M. Ko, J. Pu, and Z. Xu, *Phys. Lett. B* **781**, 499 (2018).
- [25] K. J. Sun, C. M. Ko, F. Li, J. Xu, and L. W. Chen, [arXiv:2006.08929](https://arxiv.org/abs/2006.08929).
- [26] D. Zhang (STAR Collaboration), *Nucl. Phys. A* **1005**, 121825 (2021).
- [27] J. Xu, L. W. Chen, C. M. Ko, and Z. W. Lin, *Phys. Rev. C* **85**, 041901(R) (2012).
- [28] J. Xu, T. Song, C. M. Ko, and F. Li, *Phys. Rev. Lett.* **112**, 012301 (2014).
- [29] J. Xu and C. M. Ko, *Phys. Rev. C* **94**, 054909 (2016).
- [30] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **110**, 142301 (2013).
- [31] K. Fukushima, *Phys. Lett. B* **591**, 277 (2004).
- [32] C. Q. Guo, H. Liu, and J. Xu, *Phys. Rev. C* **98**, 024914 (2018).
- [33] Z. W. Lin, C. M. Ko, B. A. Li, B. Zhang, and S. Pal, *Phys. Rev. C* **72**, 064901 (2005).
- [34] X. N. Wang and M. Gyulassy, *Phys. Rev. D* **44**, 3501 (1991).
- [35] Z. Xu and C. Greiner, *Phys. Rev. C* **71**, 064901 (2005).

- [36] F. T. Wang and J. Xu, *Phys. Rev. C* **100**, 064909 (2019).
- [37] B. A. Li and C. M. Ko, *Phys. Rev. C* **52**, 2037 (1995).
- [38] G. Q. Li, C. M. Ko, X. S. Fang, and Y. M. Zheng, *Phys. Rev. C* **49**, 1139 (1994).
- [39] G. Q. Li, C. H. Lee, and G. E. Brown, *Phys. Rev. Lett.* **79**, 5214 (1997); *Nucl. Phys. A* **625**, 372 (1997).
- [40] M. Lutz, S. Klimt, and W. Weise, *Nucl. Phys. A* **542**, 521 (1992).
- [41] K. Fukushima and T. Hatsuda, *Rep. Prog. Phys.* **74**, 014001 (2011).
- [42] G. F. Bertsch and S. Das Gupta, *Phys. Rep.* **160**, 189 (1988).
- [43] C. Y. Wong, *Phys. Rev. C* **25**, 1460 (1982).
- [44] J. C. Dunlop, M. A. Lisa, and P. Sorensen, *Phys. Rev. C* **84**, 044914 (2011).
- [45] V. Greco, M. Mitrovski, and G. Torrieri, *Phys. Rev. C* **86**, 044905 (2012).
- [46] Yu. B. Ivanov, *Phys. Lett. B* **723**, 475 (2013).
- [47] J. Steinheimer, V. Koch, and M. Bleicher, *Phys. Rev. C* **86**, 044903 (2012).
- [48] Y. Hatta, A. Monnai, and B. W. Xiao, *Phys. Rev. D* **92**, 114010 (2015).
- [49] X. Sun, H. Masui, A. M. Poskanzer, and A. Schmah, *Phys. Rev. C* **91**, 024903 (2015).
- [50] Y. Burnier, D. E. Kharzeev, J. Liao, and H. U. Yee, *Phys. Rev. Lett.* **107**, 052303 (2011).
- [51] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. C* **86**, 054908 (2012).
- [52] P. K. Kovtun, D. T. Son, and A. O. Starinets, *Phys. Rev. Lett.* **94**, 111601 (2005).
- [53] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. C* **96**, 044904 (2017).
- [54] S. Ghosh, T. C. Peixoto, V. Roy, F. E. Serna, and G. Krein, *Phys. Rev. C* **93**, 045205 (2016).
- [55] O. Soloveva, D. Fuseau, J. Aichelin, and E. Bratkovskaya, *Phys. Rev. C* **103**, 054901 (2021).