

## Nucleonic shells and nuclear masses

Landon Buskirk <sup>1,2</sup>, Kyle Godbey,<sup>1</sup> Witold Nazarewicz <sup>1,2</sup> and Wojciech Satuła <sup>3</sup>

<sup>1</sup>Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA

<sup>2</sup>Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA

<sup>3</sup>Institute of Theoretical Physics, Faculty of Physics, University of Warsaw, PL-02093 Warsaw, Poland



(Received 28 September 2023; revised 10 February 2024; accepted 13 March 2024; published 5 April 2024)

The binding energy of an isotope is a sensitive indicator of the underlying shell structure as it reflects the net energy content of a nucleus. Since magic nuclei are significantly lighter, or more bound, compared to their neighbors, the presence of nucleonic shell structure makes an imprint on nuclear masses. In this work, using a carefully designed binding-energy indicator, we catalog the appearance of spherical and deformed shell and subshell closures throughout the nuclear landscape. After presenting experimental evidence for shell and subshell closures as seen through the lens of nuclear masses, we study the ability of global nuclear mass models to predict local binding-energy variations related to shell effects.

DOI: [10.1103/PhysRevC.109.044311](https://doi.org/10.1103/PhysRevC.109.044311)

### I. INTRODUCTION

Nuclei with 2, 8, 20, 28, 50, 82, and 126 nucleons have been found to be special by having an exceptionally high natural abundance or being locally lighter than their neighbors [1]. These *magic* nucleon numbers were explained by the nuclear shell model [2,3] in terms of completely filled nucleon shells. The nuclei with such numbers of nucleons are referred to as magic, like doubly magic  $^{48}_{20}\text{Ca}_{28}$  or semimagic  $^{120}_{50}\text{Sn}_{70}$ . Experimentally, there are numerous signatures of magic gaps of shell closures. They include: locally enhanced binding energies, rapid changes of separation energies, low-lying collective excitations, kinks in charge radii, and spectroscopic factors, among other things [4–6].

The quantal stability of the atomic nucleus is determined by the behavior of the single-particle level density  $\rho(e)$  of the mean-field (intrinsic) Hamiltonian. As the ground state for many-fermion systems should correspond to the lowest possible degeneracy, the nucleus is expected to be more bound if the nucleonic level density near the Fermi level is low. Exceptionally stable systems (doubly magic nuclei) are indeed those with the least degenerate single-particle level density around the Fermi level. Quantitatively, the extra stability due to the presence of shell gaps can be encapsulated in the microscopic shell energy  $E^{\text{shell}}$  [7–9] that fluctuates with particle number and reflects the nonuniformities of the single-particle level distribution. Formally, the shell energy can be approximated by

$$E^{\text{shell}} = \sum_{i=1}^A e_i - \int e \tilde{\rho}(e) de, \quad (1)$$

where  $e_i$ 's are single-particle (Hartree-Fock) energies and  $\tilde{\rho}(e)$  is the smoothed single-particle density that averages out single-particle energies within large energy interval of the order of the energy difference between major shells. The total

binding energy of a nucleus can be roughly given by [7,8]

$$B = B^{\text{macr}} + E^{\text{shell}}, \quad (2)$$

where  $B^{\text{macr}}$  is the “macroscopic” energy that gradually depends on the number of nucleons and thus associated with the smooth distribution of single-particle levels given by  $\tilde{\rho}(e)$ .

The behavior of  $E^{\text{shell}}$  changes periodically with particle number. The lowest shell energy is expected in the regions of low single-particle level density, e.g., for the spherical magic numbers 8, 20, 28, 50, 82, and 126. However, below and above these magic numbers, the level density becomes large [(2j + 1)-fold degeneracy of spherical orbitals] and a Jahn-Teller transition takes place towards deformed shapes [10,11]. The stabilization of deformed nuclei can be associated with energy gaps in deformed single particle (s.p.) levels, i.e., deformed subshell closures [8,9,12]. Examples of deformed s.p. diagrams, or Nilsson plots, can be found in, e.g., Appendix on Nuclear Structure of Ref. [13].

### II. BINDING-ENERGY INDICATORS

Empirical information on the magnitude of nucleonic correlations is often extracted from experimental data using binding-energy relations (filters, indicators) based on measured masses of neighboring nuclei [14,15].

Usually, the binding-energy indicators are the finite-difference expressions representing various derivatives of (positive) nuclear binding energy  $B(N, Z)$  with respect to  $N$  and  $Z$ . Their role is to isolate some specific parts of the correlation energy by filtering out that part of the binding energy which behaves as a polynomial of a required order in  $N$  and  $Z$ . The commonly used mass differences are one-nucleon separation energies  $S_\tau$  ( $\tau = n, p$ ). For neutrons,

$$S_n(N, Z) = B(N, Z) - B(N - 1, Z). \quad (3)$$

The two-neutron separation energy is

$$S_{2n}(N, Z) = B(N, Z) - B(N - 2, Z). \quad (4)$$

The difference  $\delta_{2n} = S_{2n}(N, Z) - S_{2n}(N + 2, Z)$  is the so-called two-neutron shell gap indicator that represents twice the gap in the corresponding single-particle spectrum [16]. The neutron chemical potential  $\lambda_n$  can be expressed through two-neutron separation energies [17–19]

$$\lambda_n(N - 1, Z) \approx -\frac{1}{2}S_{2n}(N = 2k, Z), \quad (5)$$

where  $2k$  indicates an even number. We note that  $\lambda_n$  is negative for bound systems. In addition,

$$S_n(N = 2k, Z) \approx -\lambda_n(N - 1, Z) - \frac{1}{2} \frac{\partial \lambda_n(N - 1, Z)}{\partial N} + \Delta_n(N - 1, Z), \quad (6)$$

where  $\Delta_n(N - 1, Z)$  is the average neutron pairing gap [17,18].

The single-particle (s.p.) neutron energy splitting at the Fermi level,  $\Delta e_n$ , can thus be defined in terms of one-nucleon separation energy differences [20,21]:

$$\Delta e_n(N = 2k, Z) = S_n(N, Z) - S_n(N + 2, Z). \quad (7)$$

As demonstrated in Refs. [20,21], if variations of the mean field and pairing are smooth along isotopic or isotonic chains, the filter  $\Delta e_\tau$  represents the energy difference between the lowest particle level and the highest hole (occupied) level. For instance,

$$\Delta e_n(N = 2k, Z) \approx e_{k+1}^n - e_k^n. \quad (8)$$

Similar relations to Eqs. (3)–(8) hold for protons. It directly follows from Eqs. (6) and (7) that  $\Delta e_\tau$  is proportional to the derivative of  $\lambda_\tau$  with respect to the particle number  $N_\tau$  ( $N_\tau = Z$  or  $N$  for  $\tau = p$  or  $n$ ), i.e., it is inversely proportional to the level density [22]. The indicator  $\Delta e_\tau$  is thus sensitive to small changes of the level density at the Fermi level. Indeed, the regions of the low level density are expected to correspond to increased values of  $\Delta e_\tau$ .

Since for the smoothly varying mean-field potentials the chemical potential gradually *increases* with particle number,  $\Delta e_\tau$  should be positive in general. The deviations from the monotonic behavior of  $\lambda_\tau(N_\tau)$  do occur, and are usually associated with the rapid change of nuclear mean fields due to configuration changes. In some cases, usually associated with shape transitions,  $\Delta e_\tau < 0$ ; this corresponds to a backbending in the gauge space of  $N_\tau(\lambda_\tau)$  [19,22,23].

As an illustrative example, Fig. 1 shows  $\Delta e_n$  for the Zr isotopic chain. The local maxima in  $\Delta e_n$  can be associated with spherical and deformed s.p. gaps discussed in Sec. V. The negative value of  $\Delta e_n$  at  $N = 58$  reflects the well-known spherical-to-deformed shape transition around  $^{98}\text{Zr}$  [24,25].

While the goal of our work is to demonstrate that  $\Delta e_\tau$  is a superb measure of spherical and deformed shell closures, this indicator can also be used to study mean level spacing, or mean level density, at the Fermi energy. Indeed, beyond the regions of low level density associated with gaps,  $\Delta e_\tau$  represents mean level splitting at the Fermi energy. In the simplest scenario assuming Kramers and isospin degeneracy, the




FIG. 1. Experimental values and model predictions of  $\Delta e_n$  across zirconium isotopes. Extrapolated values from experimental data are marked with stars. Strong peaks appear for the deformed gap at  $N = 40$ , the magic gap at  $N = 50$ , and the spherical gap at  $N = 56$ . See text for details.

mean level spacing equals  $\bar{\epsilon} = 4/\tilde{\rho}(\lambda)$ , where  $\tilde{\rho}(\lambda) = 6a/\pi^2$  and  $a$  stands for the level density parameter, the value of which is uncertain. In the simplest isoscalar scenario assuming dominant volume-like  $A$  dependence the estimates for  $a$  vary from  $A/10$  (which is the harmonic oscillator limit [26]) to  $A/8 \text{ MeV}^{-1}$  [27–29]. This, in turn, gives  $\bar{\epsilon} \approx (60 \pm 6)/A \text{ MeV}$ . Note, that for the Zr isotopic chain presented in Fig. 1 it varies from 0.75(8) MeV for  $A = 80$  to 0.60(6) MeV for  $A = 100$ . The estimates agree relatively well with the data shown in Fig. 1 outside the regions of low level density associated with deformed and spherical energy gaps.

### III. DATASETS AND MODELS

In our analysis we use the most recent measured values of nuclear binding energies from the AME2020 dataset [30]. In this analysis we do not consider experimental errors and theoretical uncertainties as their proper inclusion would require the knowledge of underlying covariances. While in many nuclei the experimental mass errors are well below theoretical uncertainties and can be ignored, this is no longer the case for very exotic nuclei far from stability. In the simplest case where one assumes completely uncorrelated errors, the total error of mass filters grows substantially as several masses are involved. A detailed error analysis of  $\Delta e_\tau$  and other mass filters will be a subject of forthcoming study.

As for prediction, we consider seven theoretical models based on the energy density functional theory (EDF) which are capable of describing the whole nuclear chart: SkM\* [31], SkP [32], SLy4 [33], SV-min [34], UNEDF0 [35], UNEDF1 [36], and UNEDF2 [37]. The above set of EDF models was augmented by a well-fitted mass model FRDM-2012 [38] that has significantly more parameters than the (less

phenomenological) DFT models, resulting in a better fit to measured masses.

For  $\Delta e_\tau$  extraction from the data, the Wigner energy has to be removed from experimental binding energies as it represents an irregularity (kink) in the mass surface around  $|N = Z|$  and hence impacts mass difference indicators aiming at extracting shell structure effects [39,40]. In Ref. [41], the Wigner term has been parametrized as

$$E_W(2) = a_W |N - Z|/A, \quad (9)$$

where  $a_W = 47$  MeV. However, this expression notably underestimates the Wigner energy for  $^{80}\text{Zr}$  and  $^{56}\text{Ni}$ , two locations of shell closures that are later discussed. For this reason, we supplement  $E_W(2)$  with the model of Ref. [42]:

$$E_W(1) = V_W e^{-\lambda_W (\frac{N-Z}{A})^2} + V'_W |N - Z| e^{-(\frac{A}{A_0})^2}, \quad (10)$$

where  $V_W = 1.8$  MeV,  $\lambda_W = 380$ ,  $V'_W = -0.84$  MeV, and  $A_0 = 26$ . In our analysis, the average of  $E_W(1)$  and  $E_W(2)$  has been subtracted from all experimental binding energies. The effect of such subtraction is illustrated in Fig. 1 for  $\Delta e_n$  along the Zr chain (see Ref. [40] for the discussion of the  $^{80}\text{Zr}$  case).

#### IV. BMEX WEB APPLICATION

The exploration of the experimental and theoretical data was performed using the Bayesian Mass Explorer (BMEX) [43] web application and the associated database. An evolution of the Mass Explorer project [44], BMEX contains a suite of online plotting and comparison tools that were used to produce the draft figures in the current work. The BMEX database and software are hosted in a cloud computing environment and do not require any downloads or installation by the end user to access the tool. To save the user's sessions, plot exporting and link sharing is also included without the need for any user accounts or logins. A screenshot of the application can be found in Fig. 8 in Appendix B.

#### V. SYSTEMATIC TRENDS

In order to remove the average mass and isospin dependence of shell gaps, we scale  $\Delta e_\tau$  by the average oscillator frequency [45]:

$$\hbar\omega_0 = 41A^{-1/3} \left( 1 \pm \frac{N - Z}{3A} \right) \text{MeV}, \quad (11)$$

where the plus sign holds for neutrons and the minus sign for protons. In the following, we discuss the dimensionless splittings

$$\Delta \tilde{e}_\tau \equiv \Delta e_\tau / \hbar\omega_0. \quad (12)$$

When interpreting the patterns of shell gaps in the  $(N, Z)$  plane, it is important to recall that nuclei close to the spherical magic gaps at  $Z = 20, 28, 50, 82,$  and  $126$  are nearly spherical and that the quadrupole collectivity primarily depends on the distance of  $Z$  and  $N$  to the closest magic proton and neutron number [46,47]. That is, the largest quadrupole deformations are expected in the regions between spherical magic gaps.

#### A. Experimental single-nucleon shell gaps

Figure 2 shows the proton shell gaps  $\Delta \tilde{e}_p$  extracted from experimental binding energies. The experimental neutron shell gaps  $\Delta \tilde{e}_n$  are displayed in Fig. 3. The spherical magic gaps are clearly seen for both protons and neutrons. In addition, isotopic and isotonic bands of locally enhanced values of  $\Delta \tilde{e}_\tau$  are present; they can be associated with local subshell closures, both spherical and deformed. They are discussed in the following.

##### 1. Spherical magic gaps

In the protons, the pronounced  $Z = 50$  gap extends across the nuclear landscape. The  $Z = 82$  gap is large for  $N \geq 126$  but it seems to gradually fade away in neutron deficient Pb isotopes. This is consistent with the presence of shape coexistence effects in these nuclei, in which spherical, prolate, and oblate structures coexist (and interact) at low energies [24,48]. While the  $Z = 28$  proton shell gap is generally pronounced, the  $Z = 20$  gap becomes fairly diluted below  $N = 24$ .

The neutron magic gaps  $N = 50, 82,$  and  $126$  are well pronounced. The  $N = 28$  gap deteriorates in the lightest isotones, and a similar situation is seen at  $N = 20$ . The disappearance of  $N = 20$  and  $28$  magic gaps in neutron-rich nuclei is supported by an appreciable experimental evidence for deformed structures below  $^{44}\text{S}$  and  $^{32}\text{Mg}$  [4,24].

##### 2. Spherical subshell closures

Several local spherical shell gaps can be identified in Figs. 2 and 3. They include:  $Z = 14$  subshell closure in the Si isotopes [49];  $Z = 64$  subshell closure in  $^{146}\text{Gd}$  [19];  $N = 16$  subshell closure in  $^{36}\text{Ca}$  [50] and  $^{24}\text{O}$  [51];  $N = 32$  subshell closure in  $^{52}\text{Ca}$  [52];  $N = 56$  subshell closure in  $^{96}\text{Zr}$  [53]; and  $N = 64$  subshell closure in Sn [54]. The single  $2p_{1/2}$  orbital separates the  $N = 126$  magic gap from the  $N = 124$  spherical subshell [55]. Consequently, these two shell closures overlap in Fig. 3.

##### 3. Deformed subshell closures

In the regions between spherical magic gaps, the indicator  $\Delta \tilde{e}_\tau$  provides important information about deformed shell gaps. The region of deformed nuclei around  $^{64}\text{Cr}$  [56–59] can be associated with the deformed subshell closures  $Z = 24$  and  $N = 40$  [59]. In Fig. 2, the proton shell gap  $\Delta \tilde{e}_p$  is well pronounced for neutron-rich Cr isotopes. Of particular interest are deformed shell closures at  $Z = 38, 40$  that are responsible for very large ground-state deformations around  $^{76}\text{Sr}$  [60],  $^{80}\text{Zr}$  [40], and  $^{102}\text{Zr}$  [24]. The  $Z = 80$  oblate gap is responsible for weakly deformed ground states of the Hg isotopes [55]. It is separated from the  $Z = 82$  magic gap by a single  $2s_{1/2}$  orbit so these two shell closures overlap in Fig. 2.

The deformed neutron gaps in the rare-earth nuclei seen in Fig. 3 include:  $N = 98$  gap in the Gd-Dy region [22,61];  $N = 104$  gap around  $^{174}\text{Yb}$  [19]; and  $N = 108$  gap known around  $^{182}\text{W}$  [22].

In the actinide and transfermium regions, the most pronounced deformed neutron closures are  $N = 152$  [62,63] and  $N = 162$  [19,65–67]. In the protons, the deformed shell gap




FIG. 2. Experimental values of the dimensionless splitting  $\Delta\tilde{e}_p$  Eq. (12) throughout the nuclear landscape. The nuclei for which the expression (8) involves binding energies extrapolated from systematic trends in [30] are marked by circles. Shell closures corresponding to the bands of locally elevated values of  $\Delta\tilde{e}_p$  are clearly seen.

at  $Z = 108$  is particularly pronounced [64,65,67–69]. These subshells are essential for the stabilization of nuclear binding in the transactinides.

In addition to the above list of shell and subshell closures that can be straightforwardly identified, there are other regions in Figs. 2 and 3 with moderately enhanced values of  $\Delta\tilde{e}_\tau$ . For




FIG. 3. Similar to Fig. 2 but for  $\Delta\tilde{e}_n$ . The nuclei with negative values of  $\Delta\tilde{e}_n$  are marked by an asterisk.




FIG. 4. Similar as in Fig. 2 but for the mass model UNEDF1. The range of experimental data is marked by a solid black line.

instance, the  $N = 92$  shell effect around  $^{152}\text{Nd}$  can probably be attributed to octupole correlations.

#### 4. Shape transitions

Negative values of  $\Delta e_\tau$  are associated with shape transition. Several regions of shape-transitional behavior are seen in Fig. 3. They include the region of shape coexistence around  $^{98}\text{Zr}$  and the transition regions to well deformed prolate shapes around  $N = 88$  [22,24].

It is interesting to notice that rapid shape transitions are clearly seen in  $\Delta \tilde{e}_n$  in Fig. 3 but not in  $\Delta \tilde{e}_p$ . Indeed, no regions of  $\Delta \tilde{e}_p < 0$  can be seen in Fig. 2, which indicates that the proton chemical potential  $\lambda_p$  increases monotonically with  $Z$  throughout the nuclear landscape.

#### 5. Two-nucleon shell gap indicator

The plots of experimental  $\delta_{2\tau}$  are shown in Figs. 6 and 7 in Appendix A. As seen, this indicator behaves in a similar way as  $\Delta e_\tau$ , though, in practice, the resolving power of the  $\delta_{2\tau}$  for identifying subshell closures is slightly below that of  $\Delta e_\tau$ . Indeed, as shown in Ref. [39],  $\delta_{2\tau} = \Delta e_\tau + \Delta P_\tau$  where  $\Delta P_\tau$  represents a pairing correction. Consequently,  $\delta_{2\tau}$  is more affected by correlations, which tend to smear out shell effects.

#### B. Model predictions

Figures 4 and 5 illustrate the performance of the representative UNEDF1 mass model with respect to  $\Delta \tilde{e}_p$  and  $\Delta \tilde{e}_n$ , respectively. The predictions of other models can be obtained by using the BMEX tool [43]. The predictions extend beyond the region of nuclei with experimentally known masses, and

hence provide useful guidance for the future experiments at radioactive ion beam facilities. For instance, it is seen that the magic gaps  $Z = 50$  and  $Z = 82$  are significantly weakened around  $N = 106$  and  $N = 150$ , respectively.

The overall performance of the mass models with respect to  $\Delta \tilde{e}_\tau$  is illustrated in Table I. As expected, FRDM-2012 performs fairly well overall. Several deformed subshell closures are robustly predicted in almost all models:  $Z = 70, 80, 92, 108$  and  $N = 92, 104, 162$ . The same holds for spherical subshell closure  $N = 56$ . While the peak at  $N = 56$  predicted by UNEDF1 is underestimated in Fig. 1 the subshell closure is clearly seen in Fig. 5.

Other shells are predicted by a subset of models. In some cases, the “theoretically fragile” gaps have been discussed in literature. See, e.g., Ref. [70] for the  $N = 152$  gap predictions. Interestingly, the models consistently predict deformed proton shell gaps at  $Z = 46$  around  $N = 70$  and  $Z = 56$  around  $N = 72$ , and the deformed neutron gap  $N = 72$  around  $Z = 62$ . These features are not clearly seen in the experimental data. In general, the predictive power of the mass models used in this study with respect to  $\Delta \tilde{e}_\tau$  is quite reasonable. Moreover, the experimental finding that  $\Delta \tilde{e}_p$  is usually positive is nicely confirmed by theory, see Fig. 4. The predicted regions of  $\Delta \tilde{e}_n < 0$  in Fig. 5 are broader than in experiment. This is to be expected as the shape transitions predicted by mean-field models are too abrupt due to the missing dynamical (zero-point) correlations. While the mean-field models are generally expected to reproduce shell and subshell closures at correct particle numbers, the actual size of the predicted  $\Delta \tilde{e}_\tau$  is expected to depend on zero-point correlations and small model differences (e.g., due to poorly known spin-isospin terms [25,70] of EDFs).




FIG. 5. Similar as in Fig. 3 but for the mass model UNEDF1. The range of experimental data is marked by a solid black line and asterisks denote negative values.

**VI. SUMMARY**

The s.p. energy splitting at the Fermi level  $\Delta e_\tau$  has been extracted from measured nuclear masses and compared with predictions of mean-field models. As demonstrated in this

work,  $\Delta e_\tau$  is indeed a superb indicator of shell closures in spherical and deformed nuclei. In particular, this quantity can be very useful when studying the appearance and disappearance of nucleonic shell gaps in exotic nuclei.

TABLE I. Performance of different mass models with respect to  $\Delta \tilde{e}_\tau$  corresponding different subshell closures seen in experimental data. The models are SM = SkM\*, SP=SkP, SL=SLy4, SV=SV-min, U0=UNEDF0, U1=UNEDF1, U2=UNEDF2, and FR=FRDM-2012.

$\Delta e_\tau$	SM	SP	SL	SV	U0	U1	U2	FR
Protons								
38	✓		✓	✓		✓	✓	✓
40	✓	✓	✓	✓		✓	✓	
44			✓					✓
70	✓	✓	✓	✓	✓	✓	✓	✓
74		✓						✓
80	✓		✓	✓	✓	✓	✓	✓
88		✓						
92	✓	✓	✓	✓	✓	✓	✓	✓
108	✓	✓	✓	✓	✓	✓	✓	✓
Neutrons								
56	✓	✓		✓	✓	✓	✓	✓
70	✓	✓	✓	✓	✓	✓	✓	✓
92	✓		✓	✓	✓	✓	✓	✓
98								✓
104	✓	✓	✓	✓	✓	✓	✓	
108								✓
124		✓		✓	✓	✓	✓	✓
152	✓		✓			✓	✓	✓
162	✓	✓	✓	✓	✓	✓	✓	✓




FIG. 6. Similar as in Fig. 2 but for experimental values of the dimensionless single-particle splitting  $\tilde{\delta}_{2p}$ .

After cataloging experimental shell and subshell closures obtained by means of  $\Delta\tilde{e}_\tau$ , we showed that EDF-based models yield the placements of s.p. energy splitting maxima consistently with experiment. Indeed, mean-field models are expected to perform well in this regard as the concept of intrinsic s.p. orbits and energies is naturally present there. In some cases, such as the deformed  $A \approx 80$  and  $A \approx 100$  regions, theory sometimes poorly predicts the spherical-to-deformed shape transition due to missing zero-point correlations [25]. This deficiency of current models will need to be addressed. In this context, we wish to emphasize that the intent of this work is not to exhaustively quantify the fidelity of theoretical models' predictions of  $\Delta e_\tau$ .

Additionally, this work highlights the potential for user-focused scientific software to aid discovery and provide guidance for future experimental campaigns. To this end, the BMEX tool used in this work will be continually updated to include new experimental data and extended to a broader set of nuclear models. A broader set of uncertainty estimates for both experimental and theoretical data will also be added to the tool. The new features will include estimates of experimental and theoretical errors on mass filters, and a Bayesian model mixing module that will combine the knowledge from multiple models [71–73].

#### ACKNOWLEDGMENTS

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Awards No. DE-SC0023688 and No. DOE-DE-SC0013365, the National Science Foundation under Award No. 2004601 (CSSI program, BAND collaboration), and by

the Polish National Science Centre (NCN) under Contract No. 2018/31/B/ST2/02220.

#### APPENDIX A: EXPERIMENTAL LANDSCAPES OF $\tilde{\delta}_{2\tau}$

The two-nucleon shell gap indicators are usually defined as  $\delta_{2n} = S_{2n}(N, Z) - S_{2n}(N + 2, Z)$  and  $\delta_{2p} = S_{2n}(N, Z) - S_{2p}(N, Z + 2)$ . Figures 6 and 7 show the dimensionless single-particle splitting  $\tilde{\delta}_{2\tau}$  extracted from experimental values of  $\delta_{2\tau}$  scaled as  $\tilde{\delta}_{2\tau} \equiv \delta_{2\tau}/2\hbar\omega_0$ , where  $\hbar\omega_0$  is given by Eq. (11).

#### APPENDIX B: THE BAYESIAN MASS EXPLORER

The Bayesian Mass Explorer (BMEX) project aims to provide a user-friendly interface to theoretical model predictions with quantified uncertainties. To enable this vision, BMEX utilizes a cloud-based infrastructure that allows for efficient data retrieval, plotting, and light computation to be performed server-side and then delivered to the user in their browser. The web application uses the Plotly Dash framework and a PYTHON back end that takes advantage of multiple server workers to better handle several users simultaneously. The application is continuously built as a Docker container which eases deployment to arbitrary server architectures and can also easily be self-hosted locally for development or for local deployments. Should the load on the server pass a certain threshold, independent instances of the container can also be deployed onto new servers and access to each one can be load balanced. This improves availability and stability of the application, at the cost of needing to have a separately hosted database instance to manage user-saved sessions. A separate mechanism for saving user sessions is also implemented via link encoding, though this is less scal-




FIG. 7. Similar as in Fig. 3 but for experimental values of the dimensionless single-particle splitting  $\tilde{\delta}_{2n}$ .

able and presents issues for backwards compatibility. Figure 8 presents a sample screenshot of the BMEX web interface in a

configuration similar to what was used for the present investigation of single-particle energy splittings.



FIG. 8. Sample screenshot of the Bayesian Mass Explorer web interface in a configuration similar to what was used for the present investigation. The configuration and quantity of interest for each plot is accessed through the tab functionality on the left and additional data series can be added to the isotopic chains in the sidebar. The current view can be saved and shared using the “Share View” button on the right sidebar. Vector graphics in the PDF format can also be exported through the web interface for easy integration into documents.



- [1] M. G. Mayer, On closed shells in nuclei, *Phys. Rev.* **74**, 235 (1948).
- [2] M. G. Mayer, On closed shells in nuclei. II, *Phys. Rev.* **75**, 1969 (1949).
- [3] O. Haxel, J. H. D. Jensen, and H. E. Suess, On the “magic numbers” in nuclear structure, *Phys. Rev.* **75**, 1766 (1949).
- [4] O. Sorlin and M.-G. Porquet, Nuclear magic numbers: New features far from stability, *Prog. Part. Nucl. Phys.* **61**, 602 (2008).
- [5] T. Otsuka, A. Gade, O. Sorlin, T. Suzuki, and Y. Utsuno, Evolution of shell structure in exotic nuclei, *Rev. Mod. Phys.* **92**, 015002 (2020).
- [6] F. Nowacki, A. Obertelli, and A. Poves, The neutron-rich edge of the nuclear landscape: Experiment and theory, *Prog. Part. Nucl. Phys.* **120**, 103866 (2021).
- [7] V. Strutinsky, Shells in deformed nuclei, *Nucl. Phys. A* **122**, 1 (1968).
- [8] M. Brack, J. Damgaard, A. S. Jensen, H. C. Pauli, V. M. Strutinsky, and C. Y. Wong, Funny Hills: The shell-correction approach to nuclear shell effects and its applications to the fission process, *Rev. Mod. Phys.* **44**, 320 (1972).
- [9] S. G. Nilsson and I. Ragnarsson, *Shapes and Shells in Nuclear Structure* (Cambridge University Press, Cambridge, 1995).
- [10] P.-G. Reinhard and E. W. Otten, Transition to deformed shapes as a nuclear Jahn-Teller effect, *Nucl. Phys. A* **420**, 173 (1984).
- [11] W. Nazarewicz, Microscopic origin of nuclear deformations, *Nucl. Phys. A* **574**, 27 (1994).
- [12] I. Ragnarsson and R. K. Sheline, Systematics of nuclear deformations, *Phys. Scr.* **29**, 385 (1984).
- [13] R. Firestone, C. Baglin, and S. Chu, *Table of Isotopes: 1999 Update*, A Wiley-Interscience publication No. Vol. 4 (Wiley, New York, 1999).
- [14] J. Jänecke and E. Comay, Properties of homogeneous and inhomogeneous mass relations, *Nucl. Phys. A* **436**, 108 (1985).
- [15] A. Jensen, P. Hansen, and B. Jonson, New mass relations and two- and four-nucleon correlations, *Nucl. Phys. A* **431**, 393 (1984).
- [16] M. Bender, T. Cornelius, G. Lalazissis, J. Maruhn, W. Nazarewicz, and P.-G. Reinhard, The  $Z = 82$  shell closure in neutron-deficient Pb isotopes, *Eur. Phys. J. A* **14**, 23 (2002).
- [17] M. Beiner and R. Lombard, The energy density formalism and the shell structure effects, *Ann. Phys.* **86**, 262 (1974).
- [18] M. Beiner, R. Lombard, and D. Mas, Self-consistent calculations of ground state properties for unstable nuclei, *Nucl. Phys. A* **249**, 1 (1975).
- [19] R. Bengtsson, P. Möller, J. R. Nix, and J. Zhang, Nuclear shapes and shape transitions, *Phys. Scr.* **29**, 402 (1984).
- [20] W. Satuła, J. Dobaczewski, and W. Nazarewicz, Odd-even staggering of nuclear masses: Pairing or shape effect? *Phys. Rev. Lett.* **81**, 3599 (1998).
- [21] J. Dobaczewski, P. Magierski, W. Nazarewicz, W. Satuła, and Z. Szymański, Odd-even staggering of binding energies as a consequence of pairing and mean-field effects, *Phys. Rev. C* **63**, 024308 (2001).
- [22] R. Bengtsson, J. Zhang, and S. Åberg, On the analogy between backbending in gauge space and in ordinary space, *Phys. Lett. B* **105**, 5 (1981).
- [23] J. Zhang, The analysis in gauge space, *Nucl. Phys. A* **421**, 353 (1984).
- [24] K. Heyde and J. L. Wood, Shape coexistence in atomic nuclei, *Rev. Mod. Phys.* **83**, 1467 (2011).
- [25] P.-G. Reinhard, D. J. Dean, W. Nazarewicz, J. Dobaczewski, J. A. Maruhn, and M. R. Strayer, Shape coexistence and the effective nucleon-nucleon interaction, *Phys. Rev. C* **60**, 014316 (1999).
- [26] A. Bohr and B. R. Mottelson, *Nuclear Structure*, Vol. I (W. A. Benjamin, New York, 1969).
- [27] A. Gilbert and A. Cameron, A composite nuclear-level density formula with shell corrections, *Can. J. Phys.* **43**, 1446 (1965).
- [28] S. K. Kataria and V. S. Ramamurthy, Macroscopic level density parameters of nuclei, *Phys. Rev. C* **22**, 2263(R) (1980).
- [29] S. Shlomo, Energy-level density of nuclei, *Nucl. Phys. A* **539**, 17 (1992).
- [30] M. Wang, W. Huang, F. Kondev, G. Audi, and S. Naimi, The AME 2020 atomic mass evaluation (II). Tables, graphs and references, *Chin. Phys. C* **45**, 030003 (2021).
- [31] J. Bartel, P. Quentin, M. Brack, C. Guet, and H.-B. Håkansson, Towards a better parametrisation of Skyrme-like effective forces: A critical study of the SkM force, *Nucl. Phys. A* **386**, 79 (1982).
- [32] J. Dobaczewski, H. Flocard, and J. Treiner, Hartree-Fock-Bogolyubov description of nuclei near the neutron-drip line, *Nucl. Phys. A* **422**, 103 (1984).
- [33] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer, New Skyrme effective forces for supernovae and neutron rich nuclei, *Phys. Scr.* **T56**, 231 (1995).
- [34] P. Klüpfel, P.-G. Reinhard, T. J. Bürvenich, and J. A. Maruhn, Variations on a theme by Skyrme: A systematic study of adjustments of model parameters, *Phys. Rev. C* **79**, 034310 (2009).
- [35] M. Kortelainen, T. Lesinski, J. Moré, W. Nazarewicz, J. Sarich, N. Schunck, M. V. Stoitsov, and S. Wild, Nuclear energy density optimization, *Phys. Rev. C* **82**, 024313 (2010).
- [36] M. Kortelainen, J. McDonnell, W. Nazarewicz, P.-G. Reinhard, J. Sarich, N. Schunck, M. V. Stoitsov, and S. M. Wild, Nuclear energy density optimization: Large deformations, *Phys. Rev. C* **85**, 024304 (2012).
- [37] M. Kortelainen, J. McDonnell, W. Nazarewicz, E. Olsen, P.-G. Reinhard, J. Sarich, N. Schunck, S. M. Wild, D. Davesne, J. Erler, and A. Pastore, Nuclear energy density optimization: Shell structure, *Phys. Rev. C* **89**, 054314 (2014).
- [38] P. Möller, A. Sierk, T. Ichikawa, and H. Sagawa, Nuclear ground-state masses and deformations: FRDM(2012), *At. Data Nucl. Data Tables* **109-110**, 1 (2016).
- [39] A. Gelberg, H. Sakurai, M. W. Kirson, and S. Heinze, Wigner energy and shell gaps in two-nucleon separation energies, *Phys. Rev. C* **80**, 024307 (2009).
- [40] A. Hamaker, E. Leistenschneider, R. Jain, G. Bollen, S. A. Giuliani, K. Lund, W. Nazarewicz, L. Neufcourt, C. R. Nicoloff, D. Puentes, R. Ringle, C. S. Sumithrarachchi, and I. T. Yandow, Precision mass measurement of lightweight self-conjugate nucleus  $^{80}\text{Zr}$ , *Nat. Phys.* **17**, 1408 (2021).
- [41] W. Satuła, D. Dean, J. Gary, S. Mizutori, and W. Nazarewicz, On the origin of the Wigner energy, *Phys. Lett. B* **407**, 103 (1997).
- [42] S. Goriely, N. Chamel, and J. M. Pearson, Further explorations of Skyrme-Hartree-Fock-Bogoliubov mass formulas. XIII. The 2012 atomic mass evaluation and the symmetry coefficient, *Phys. Rev. C* **88**, 024308 (2013).
- [43] K. Godbey, L. Buskirk, and P. Giuliani, BMEX - The Bayesian Mass Explorer (2023), <https://bmxex.dev>.
- [44] MassExplorer <http://massexplorer.frib.msu.edu>.

- [45] S. G. Nilsson, C. F. Tsang, A. Sobiczewski, Z. Szymański, S. Wycech, C. Gustafson, I.-L. Lamm, P. Möller, and B. Nilsson, On the nuclear structure and stability of heavy and superheavy elements, *Nucl. Phys. A* **131**, 1 (1969).
- [46] J. Jänecke, Simple parameterization of nuclear deformation parameters, *Phys. Lett. B* **103**, 1 (1981).
- [47] B. Pritychenko, M. Birch, B. Singh, and M. Horoi, Tables of E2 transition probabilities from the first  $2^+$  states in even-even nuclei, *At. Data Nucl. Data Tables* **107**, 1 (2016).
- [48] J. Wood, K. Heyde, W. Nazarewicz, M. Huyse, and P. van Duppen, Coexistence in even-mass nuclei, *Phys. Rep.* **215**, 101 (1992).
- [49] P. D. Cottle, Single proton energies in the Si isotopes and the  $Z = 14$  subshell closure, *Phys. Rev. C* **76**, 027301 (2007).
- [50] L. Lalanne, O. Sorlin, A. Poves, M. Assié, F. Hammache, S. Koyama, D. Suzuki, F. Flavigny, V. Girard-Alcindor, A. Lemasson, A. Matta, T. Roger, D. Beaumel, Y. Blumenfeld, B. A. Brown, F. D. O. Santos, F. Delaunay, N. de Séréville, S. Franchoo, J. Gibelin, J. Guillot, O. Kamalou, N. Kitamura, V. Lapoux, B. Mauss, P. Morfouace, J. Pancin, T. Y. Saito, C. Stodel, and J.-C. Thomas,  $N = 16$  magicity revealed at the proton drip line through the study of  $^{35}\text{Ca}$ , *Phys. Rev. Lett.* **131**, 092501 (2023).
- [51] K. Tshoo, Y. Satou, H. Bhang, S. Choi, T. Nakamura, Y. Kondo, S. Deguchi, Y. Kawada, N. Kobayashi, Y. Nakayama, K. N. Tanaka, N. Tanaka, N. Aoi, M. Ishihara, T. Motobayashi, H. Otsu, H. Sakurai, S. Takeuchi, Y. Togano, K. Yoneda, Z. H. Li, F. Delaunay, J. Gibelin, F. M. Marqués, N. A. Orr, T. Honda, M. Matsushita, T. Kobayashi, Y. Miyashita, T. Sumikama, K. Yoshinaga, S. Shimoura, D. Sohler, T. Zheng, and Z. X. Cao,  $N=16$  spherical shell closure in  $^{24}\text{O}$ , *Phys. Rev. Lett.* **109**, 022501 (2012).
- [52] F. Wienholtz, D. Beck, K. Blaum, C. Borgmann, M. Breitenfeldt, R. B. Cakirli, S. George, F. Herfurth, J. D. Holt, M. Kowalska, S. Kreim, D. Lunney, V. Manea, J. Menéndez, D. Neidherr, M. Rosenbusch, L. Schweikhard, A. Schwenk, J. Simonis, J. Stanja, R. N. Wolf, and K. Zuber, Masses of exotic calcium isotopes pin down nuclear forces, *Nature (London)* **498**, 346 (2013).
- [53] G. Molnár, The first submagic nucleus:  $^{96}\text{Zr}$ , *AIP Conf. Proc.* **238**, 227 (1991).
- [54] C. Piller, C. Gugler, R. Jacot-Guillarmod, L. A. Schaller, L. Schellenberg, H. Schnewly, G. Fricke, T. Hennemann, and J. Herberz, Nuclear charge radii of the tin isotopes from muonic atoms, *Phys. Rev. C* **42**, 182 (1990).
- [55] M. Vergnes, G. Berrier-Ronsin, G. Rotbard, J. Skalski, and W. Nazarewicz, Evidence for a change of structure in the heavy mercury isotopes around  $^{200}\text{Hg}$ , *Nucl. Phys. A* **514**, 381 (1990).
- [56] O. Sorlin, C. Donzaud, F. Nowacki, J. C. Angélique, F. Azaiez, C. Bourgeois, V. Chisté, Z. Dlouhy, S. Grévy, D. Guillemaud-Mueller, F. Ibrahim, K. L. Kratz, M. Lewitowicz, S. M. Lukyanov, J. Mrasek, Y. E. Penionzhkevich, F. de Oliveira Santos, B. Pfeiffer, F. Pougheon, A. Poves, M. G. Saint-Laurent, and M. Stanoiu, New region of deformation in the neutron-rich  $^{60}\text{Cr}$  and  $^{62}\text{Cr}$ , *Eur. Phys. J. A* **16**, 55 (2003).
- [57] C. Babcock, H. Heylen, M. Bissell, K. Blaum, P. Campbell, B. Cheal, D. Fedorov, R. Garcia Ruiz, W. Geithner, W. Gins, T. Day Goodacre, L. Grob, M. Kowalska, S. Lenzi, B. Maass, S. Malbrunot-Ettenauer, B. Marsh, R. Neugart, G. Neyens, Y. Nörtershäuser, T. Otsuka, R. Rossel, S. Rothe, R. Sánchez, Y. Tsunoda, C. Wraith, L. Xie, and X. Yang, Quadrupole moments of odd- $A$   $^{53-63}\text{Mn}$ : Onset of collectivity towards  $N = 40$ , *Phys. Lett. B* **760**, 387 (2016).
- [58] R. Silwal, C. Andreoiu, B. Ashrafkhani, J. Bergmann, T. Brunner, J. Cardona, K. Dietrich, E. Dunling, G. Gwinner, Z. Hockenbery, J. Holt, C. Izzo, A. Jacobs, A. Javaji, B. Kootte, Y. Lan, D. Lunney, E. Lykiardopoulou, T. Miyagi, M. Mougeot, I. Mukul, T. Murböck, W. Porter, M. Reiter, J. Ringuette, J. Dilling, and A. Kwiatkowski, Summit of the  $N = 40$  island of inversion: Precision mass measurements and ab initio calculations of neutron-rich chromium isotopes, *Phys. Lett. B* **833**, 137288 (2022).
- [59] H. Oba and M. Matsuo, Deformation around neutron-rich Cr isotopes in axially symmetric skyrme-Hatree-Fock-Bogoliubov method, *Prog. Theor. Phys.* **120**, 143 (2008).
- [60] W. Nazarewicz, J. Dudek, R. Bengtsson, T. Bengtsson, and I. Ragnarsson, Microscopic study of the high-spin behaviour in selected  $A \simeq 80$  nuclei, *Nucl. Phys. A* **435**, 397 (1985).
- [61] D. J. Hartley, F. G. Kondev, R. Orford, J. A. Clark, G. Savard, A. D. Ayangeakaa, S. Bottoni, F. Buchinger, M. T. Burkey, M. P. Carpenter, P. Copp, D. A. Gorelov, K. Hicks, C. R. Hoffman, C. Hu, R. V. F. Janssens, J. W. Klimes, T. Lauritsen, J. Sethi, D. Seweryniak, K. S. Sharma, H. Zhang, S. Zhu, and Y. Zhu, Masses and  $\beta$ -decay spectroscopy of neutron-rich odd-odd  $^{160,162}\text{Eu}$  nuclei: Evidence for a subshell gap with large deformation at  $N=98$ , *Phys. Rev. Lett.* **120**, 182502 (2018).
- [62] E. M. Ramirez, D. Ackermann, K. Blaum, M. Block, C. Droese, C. E. Düllmann, M. Dworschak, M. Eibach, S. Eliseev, E. Haettner, F. Herfurth, F. P. Heßberger, S. Hofmann, J. Ketelaer, G. Marx, M. Mazzocco, D. Nesterenko, Y. N. Novikov, W. R. Plaß, D. Rodríguez, C. Scheidenberger, L. Schweikhard, P. G. Thirolf, and C. Weber, Direct mapping of nuclear shell effects in the heaviest elements, *Science* **337**, 1207 (2012).
- [63] H. Makii, T. Ishii, M. Asai, K. Tsukada, A. Toyoshima, M. Matsuda, A. Makishima, J. Kaneko, H. Toume, S. Ichikawa, S. Shigematsu, T. Kohno, and M. Ogawa,  $Z$  dependence of the  $N = 152$  deformed shell gap: In-beam  $\gamma$ -ray spectroscopy of neutron-rich  $^{245,246}\text{Pu}$ , *Phys. Rev. C* **76**, 061301(R) (2007).
- [64] J. Dvorak, W. Brüche, M. Chelnokov, R. Dressler, C. E. Düllmann, K. Eberhardt, V. Gorshkov, E. Jäger, R. Krücken, A. Kuznetsov, Y. Nagame, F. Nebel, Z. Novackova, Z. Qin, M. Schädel, B. Schausten, E. Schimpf, A. Semchenkov, P. Thörle, A. Türler, M. Weggrzecki, B. Wierczinski, A. Yakushev, and A. Yeremin, Doubly magic nucleus  $^{270}_{108}\text{Hs}_{162}$ , *Phys. Rev. Lett.* **97**, 242501 (2006).
- [65] Y. T. Oganessian, V. K. Utyonkov, F. S. Abdullin, S. N. Dmitriev, R. Graeger, R. A. Henderson, M. G. Itkis, Y. V. Lobanov, A. N. Mezentsev, K. J. Moody, S. L. Nelson, A. N. Polyakov, M. A. Ryabinin, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. A. Stoyer, N. J. Stoyer, V. G. Subbotin, K. Subotic, A. M. Sukhov, Y. S. Tsyganov, A. Türler, A. A. Voinov, G. K. Vostokin, P. A. Wilk, and A. Yakushev, Synthesis and study of decay properties of the doubly magic nucleus  $^{270}\text{Hs}$  in the  $^{226}\text{Ra} + ^{48}\text{Ca}$  reaction, *Phys. Rev. C* **87**, 034605 (2013).
- [66] O. Kaleja *et al.*, Direct high-precision mass spectrometry of superheavy elements with SHIPTRAP, *Phys. Rev. C* **106**, 054325 (2022).
- [67] Z. Patyk and A. Sobiczewski, Main deformed shells of heavy nuclei studied in a multidimensional deformation space, *Phys. Lett. B* **256**, 307 (1991).

- [68] P. Möller and J. Nix, Stability and decay of nuclei at the end of the periodic system, *Nucl. Phys. A* **549**, 84 (1992).
- [69] S. Ćwiok, S. Hofmann, and W. Nazarewicz, Shell structure of the heaviest elements, *Nucl. Phys. A* **573**, 356 (1994).
- [70] J. Dobaczewski, A. Afanasjev, M. Bender, L. Robledo, and Y. Shi, Properties of nuclei in the nobelium region studied within the covariant, Skyrme, and Gogny energy density functionals, *Nucl. Phys. A* **944**, 388 (2015), Special Issue on Superheavy Elements.
- [71] L. Neufcourt, Y. Cao, W. Nazarewicz, E. Olsen, and F. Viens, Neutron drip line in the Ca region from Bayesian model averaging, *Phys. Rev. Lett.* **122**, 062502 (2019).
- [72] D. R. Phillips, R. Furnstahl, U. Heinz, T. Maiti, W. Nazarewicz, F. Nunes, M. Plumlee, S. Pratt, M. Pratola, F. Viens, and S. M. Wild, Get on the BAND wagon: A Bayesian framework for quantifying model uncertainties in nuclear dynamics, *J. Phys. G* **48**, 072001 (2021).
- [73] V. Kejzlar, L. Neufcourt, and W. Nazarewicz, Local Bayesian Dirichlet mixing of imperfect models, *Sci. Rep.* **13**, 19600 (2023).