

Contribution to differential π^0 and γ_{dir} modification in small systems from color fluctuation effectsDennis V. Perepelitsa **Department of Physics, University of Colorado Boulder, Boulder, Colorado 80309, USA*

(Received 6 May 2024; accepted 20 June 2024; published 1 July 2024)

A major complication in the search for jet quenching in proton- or deuteron-nucleus collision systems is the presence of physical effects which influence the experimental determination of collision centrality in the presence of a hard process. For example, in the proton color fluctuation picture, protons with a large Bjorken- x ($x \gtrsim 0.1$) parton interact more weakly with the nucleons in the nucleus, leading to a smaller (larger) than expected yield in large (small) activity events. A recent measurement by the PHENIX Collaboration compared the yield of neutral pion and direct photon production in $d + \text{Au}$ collisions, under the argument that the photon yields correct for such biases, and the difference between the two species is thus attributable to final-state effects (i.e., jet quenching). The main finding suggests a significant degree of jet quenching for hard processes in small systems. In this paper, I argue that the particular photon and pion events selected by PHENIX arise from proton configurations with significantly different Bjorken- x distributions, and thus are subject to different magnitudes of modification in the color fluctuation model. Using the results of a previous global analysis of data from the BNL Relativistic Heavy Ion Collider (RHIC) and the CERN Large Hadron Collider (LHC), I show that potentially all of the pion-to-photon difference in PHENIX data can be described by a proton color fluctuation picture at a quantitative level before any additional physics from final-state effects is required. This finding reconciles the interpretation of the PHENIX measurement with others at RHIC and LHC, which have found no observable evidence for jet quenching in small systems.

DOI: [10.1103/PhysRevC.110.L011901](https://doi.org/10.1103/PhysRevC.110.L011901)

Introduction. Measurements of hard processes in relativistic proton- or deuteron-nucleus ($p/d + A$) collisions at the BNL Relativistic Heavy Ion Collider (RHIC) and the CERN Large Hadron Collider (LHC) have a number of scientific purposes [1,2], including a precision determination of the parton densities in nuclei (see Ref. [3] for a recent example), constraints on dynamical processes in the initial state of the cold nucleus (such as those observed at lower energies [4]), and studies of the general interplay between soft and hard processes in collisions involving nuclei. Another key interest, given the robust experimental signatures of quark gluon plasma–like behavior in these “small” systems [5–9], is the search for evidence of final-state interactions between the hard-scattered parton and the dilute system. If the parton shower is modified in the final state as it propagates through the produced system, it may lead to a number of effects, the most direct of which is the decreased production of jets or hadrons at fixed transverse momentum (p_T), i.e., jet quenching [10–12].

By analogy to large, nucleus-nucleus collision systems, one wants to search in the most “central,” or highest-activity, collisions where the produced system is largest and longest lived. Unfortunately, the experimental selection on centrality in small systems is challenging due to strong autocorrelation biases and nontrivial physics effects, with magnitudes larger than that required for a precision constraint on jet quenching

effects. For events with a generic hard process, the selection is sensitive to upward multiplicity fluctuations (multiplicity vetoes), leading to an apparently enhanced yield in central (decreased yield in peripheral) collisions compared to that expected from the estimated number of nucleon-nucleon collisions, $\langle N_{\text{coll}} \rangle$, in the events [13–16].

A separate phenomenon observed in data is that, in extreme kinematic regions characterized by Bjorken- $x \gtrsim 0.1$, the pattern appears to reverse, with a significant depletion (enhancement) in the high-activity central (low-activity peripheral) events. This feature has been observed in measurements by ATLAS [17], CMS [18], STAR [19], and PHENIX [20] Collaborations in $p/d + A$ events. In a recent measurement by ATLAS [21], the produced dijet pair was used to estimate the parton-level kinematics in each event, strongly suggesting that the modifications follow a universal pattern in the Bjorken- x of the proton, with their magnitude systematically increasing with x . A particular quantitative interpretation of these observations in data is the QCD color fluctuation model implemented in Refs. [22,23], with references therein describing the theoretical development of this idea. The model proposes that proton configurations with a large- x parton interact more weakly with nucleons in the nucleus than configuration-averaged protons, thus leading to a decreased centrality signal and the particular pattern of x -dependent modifications described above.

To get around these challenges, experimental searches of jet quenching in $p/d + A$ collisions have therefore been performed in centrality-averaged (minimum-bias) collisions

*Contact author: dvp@colorado.edu

[24–26], or using techniques such as jet-hadron correlations [27,28] and dijet asymmetries [18,19] in central events, all giving null results for a detectable jet-quenching-like signature. A recent PHENIX measurement [29] in 200 GeV $d + Au$ collisions used a different technique and reported the relative yields of neutral pions (π^0 's) to those of direct photons (γ_{dir} 's), under the argument that both are subject to the same centrality bias effects, and thus any difference in the degree of modification should be attributed to centrality-dependent final-state effects (i.e., which would not be felt by the photons). The PHENIX measurement reports the quantity $R_{dAu, \text{EXP}}$, defined as the double ratio of the nuclear modification factors for π^0 's and γ_{dir} , $R_{dAu, \pi^0}/R_{dAu, \gamma_{\text{dir}}}$, i.e., conceptually defining $R_{dAu, \gamma_{\text{dir}}}$ to be unity for all centrality selections. The measurement reports $R_{dAu, \text{EXP}}$ in the most central 0–5% of $d + Au$ events as $0.77 \pm 0.03(\text{stat}) \pm 0.13(\text{syst})$, where the latter is dominated by a global uncertainty due to the normalization of the γ_{dir} measurement. If interpreted as a jet quenching effect, the data are challenging to understand given the extensive null results described above. For example, the PHENIX data would require a π^0 p_T spectrum shift of $\delta p_T \approx -0.21$ GeV in the most central events at RHIC [30], corresponding to a fractional energy loss of $\delta p_T/p_T \approx 1\text{--}2\%$ in $d + Au$ events, whereas the measurement by ATLAS in Ref. [28] sets an exclusion limit at 90% confidence level of $\delta p_T/p_T < 1.4\%$ for charged hadrons in $p + Pb$ collisions at the LHC. However, these two measurements probe partons with different initial energies and collisions with different energies and geometries, and thus may not be in explicit tension.

Notably, the π^0 and γ_{dir} were measured in the same p_T range of 7.5–18 GeV which, since the π^0 carries only a fraction of the tree-level parton p_T , actually corresponds to different Bjorken- x ranges. Thus, the measurement technique defined by PHENIX is directly sensitive to the x -dependent color fluctuation effects described above, in addition to any jet quenching effects. In particular, the strength of the effect changes quickly with x in the region of the measurement, and is thus only partially “calibrated out” with the γ_{dir} baseline. In this paper, I argue that the color fluctuation effect, as calculated using the results of Ref. [23] with no additional model modifications or re-tuning to new data, can potentially explain the full centrality dependence of the observable defined by PHENIX. When accounting for it, the possible magnitude of any jet quenching effect is significantly smaller, reconciling the interpretation of the PHENIX measurement with the established constraints on jet quenching by the other RHIC and LHC experiments.

Method. The PYTHIA Monte Carlo event generator [31] was used to simulate the nucleon–nucleon sub-collisions in $d + Au$ collisions which produce π^0 's and γ_{dir} 's in the PHENIX kinematic selections [29], and determine the distribution of Bjorken- x values for these events. The specific selections are $p_T = 7.5\text{--}18$ GeV and $|\eta| < 0.35$ for both species. PYTHIA version 8.307 was used to generate events at $\sqrt{s} = 200$ GeV, with a mixture of proton-proton, proton-neutron, and neutron–neutron collisions appropriate for $d + Au$. PYTHIA was configured with all hard QCD processes (“HardQCD”) and a $\hat{p}_{T, \text{min}}$ (minimum parton-parton transverse momentum exchange) of 5 GeV for the π^0 case. For the direct




FIG. 1. Distribution in PYTHIA8 of the values of Bjorken- x in the deuteron in 200 GeV $d + Au$ collisions, for events producing π^0 's (solid black) or γ_{dir} 's (dashed red) in the kinematic selection used by PHENIX in Ref. [29].

photon case, PYTHIA was configured with all prompt photon (“PromptPhoton”) processes, also with $\hat{p}_{T, \text{min}} = 5$ GeV, and photons which are the result of hadron decays were rejected (i.e., only those radiated by a quark were selected). For both samples, this $\hat{p}_{T, \text{min}}$ threshold was checked that it does not introduce a kinematic bias in the final-state p_T range of interest. One of the beam hadrons was chosen to represent the proton or neutron in the deuteron of a $d + Au$ collision and its Bjorken- x value, defined in PYTHIA as the x at which the PDF in the beam is defined [31], was recorded.

Figure 1 shows the distribution of Bjorken- x values in the deuteron for events with a π^0 or γ_{dir} within the PHENIX kinematic selection. For γ_{dir} , the distribution is peaked near $x \approx 7.5/100 = 0.075$, where 7.5 and 100 GeV are the minimum γ_{dir} p_T threshold and the beam energy, respectively, with a small tail of contributions from events with higher x . For π^0 , the distribution has a significantly harder tail to large- x values, reflecting the fact that they are produced by the fragmentation of hard scattered partons and thus carry only a fraction of their energy. Notably, the average x value is more than 50% larger for π^0 - than γ_{dir} -producing events, with 28% (9%) of the π^0 's (γ_{dir} 's) produced in events with $x > 0.2$. In principle, the magnitude of the color fluctuation effect is expected to be different depending on whether the high- x parton in the deuteron is a quark or a gluon. However, in these kinematics, the quark/gluon fraction is similar in the π^0 and γ_{dir} events and thus the difference in their x distributions is the main distinction.

The distributions of x values in Fig. 1 were then folded with the results of the color fluctuation model in Ref. [23] to produce a prediction for the results of the PHENIX measurement. First, the color fluctuation model paper primarily reports R_{CP} , the ratio of nuclear modification in central to peripheral events, which must be converted to the $R_{dAu, \text{EXP}}$ and other observables reported in the recent PHENIX paper. This is accomplished using the detailed information provided in an earlier PHENIX publication [13] on the geometric properties (e.g. the average number of colliding nucleon-




FIG. 2. Calculated nuclear modification factor R_{dAu} within the color fluctuation model as a function of the average number of colliding nucleon pairs, $\langle N_{\text{coll}} \rangle$. Calculations are shown for π^0 's (solid black) and γ_{dir} 's (dashed red) matching the PHENIX kinematic selection. Since this model does not include any overall modification, the minimum-bias R_{dAu} is unity by construction.

nucleon pairs ($\langle N_{\text{coll}} \rangle$) of centrality-selected $d + \text{Au}$ events. Importantly, while the results here are presented as a function of the average $\langle N_{\text{coll}} \rangle$, the calculations in Ref. [23] on which they are based were determined using the distribution of the N_{coll} values in each centrality bin. Second, the color fluctuation model presents results in exclusive Bjorken- x bins, which are then combined in a weighted average with the x distributions for π^0 and γ_{dir} events above. Since the model in Ref. [23] did not consider the region $x \lesssim 0.1$, I treat the color fluctuation effect as negligible for events with x in this range, i.e., for these all $R_{dAu} = 1$. I highlight that the parameters of this model, determined using previous RHIC and LHC data, have not otherwise been adjusted or updated before the comparison to the PHENIX data of interest.

Notably, the color fluctuation model in Ref. [23] does not itself include any centrality-independent modification, i.e., R_{dAu} for minimum-bias events is unity by construction. However, there are expected to be small modifications of the minimum-bias rate for both species due to the nuclear parton densities [3] and the ‘‘isospin effect’’ for direct photon production from the valence quark region [32,33]. Given the specific interest in the centrality dependence of the observable and the small magnitude of the minimum-bias-averaged effects compared to the large overall normalization uncertainties in PHENIX data, I do not model these effects explicitly in this work. If such effects were included, they would result in a global rescaling of all calculated R_{dAu} points in the same direction.

Figure 2 shows the calculated nuclear modification factors, R_{dAu} , within the color fluctuation model, applied to the π^0 and γ_{dir} events in the PHENIX kinematics selection, and within the PHENIX centrality intervals. Since the minimum-bias R_{dAu} is unity by construction (see above), the modifications appear as a relative enhancement (suppression) pattern around unity in peripheral (central) events. Importantly, the magnitude of the relative modifications is stronger for π^0 's than




FIG. 3. Comparison of the nuclear modification factor R_{dAu} for direct photons as a function of $\langle N_{\text{coll}} \rangle$, showing the measurement in PHENIX data (black points) and the calculation from the color fluctuation model (red line). The vertical bars and boxes around the data points indicate the statistical uncertainties and the overall 16.5% normalization uncertainty, respectively.

for γ_{dir} 's, reflecting the different x distributions in Fig. 1. Indeed, in Ref. [23], the quantity $\lambda(x)$, which describes the x -dependent weakening of the inelastic nucleon-nucleon interaction strength, falls rapidly over the Bjorken- x range relevant for γ_{dir} and π^0 production in the measurement, where it decreases from $\lambda \approx 0.8$ at $x \approx 0.1$ to $\lambda \approx 0.5$ at $x \approx 0.4$.

Comparison to data. Fig. 3 compares the calculated R_{dAu} for γ_{dir} integrated over the $p_T = 7.5\text{--}18$ GeV range within the color fluctuation model to that measured by PHENIX, where the quantity is styled as $(Y_{dAu}^{\gamma_{\text{dir}}}/Y_{pp}^{\gamma_{\text{dir}}})/N_{\text{coll}}^{\text{Glauber}}$. For this comparison, the measured minimum-bias R_{dAu} for γ_{dir} is essentially unity, and so no overall rescaling of the model calculation is applied. The model gives a good description of the modest centrality dependence of the data, with a relative enhancement and suppression pattern in going from peripheral to central events. As a reminder, the γ_{dir} selection typically results in events with $\langle x \rangle \approx 0.11$, and thus the impact of the color fluctuation physics on this observable is modest in the x range relevant for γ_{dir} production.

Figure 4 compares the measured and calculated ‘‘double ratio’’ of the R_{dAu} for π^0 's to that for γ_{dir} 's, as a function of $\langle N_{\text{coll}} \rangle$, which is the main result of the PHENIX analysis. The dashed red line shows the result of the color fluctuation model, under the assumption that the R_{dAu} in minimum-bias events is unity for both species. The calculation gives a good, quantitative description of the data within its significant normalization uncertainty, including the slow decrease of the double ratio with increasing $\langle N_{\text{coll}} \rangle$.

In the PHENIX analysis, the minimum bias $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{\text{dir}})$ is measured to be $0.92 \pm 0.02(\text{stat}) \pm 0.15(\text{syst})$. Reference [29] notes that this value is consistent with unity within the large overall scale uncertainty, which is dominated by uncertainties in the π^0 and γ_{dir} $p + p$ references. Thus, the data-model comparison here has a significant degree of freedom in the overall normalization of the data. The best match between the model and the central values of the data is achieved by adjusting the




FIG. 4. Comparison of the ratio of nuclear modification factors between that for neutral pions and direct photons, $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$, as a function of $\langle N_{coll} \rangle$, showing the measurement in PHENIX data (black points) and the calculation from the color fluctuation model (red lines). The vertical bars and boxes around the data points indicate the statistical uncertainties and the overall 16.5% normalization uncertainty, respectively. The dashed red line assumes that R_{dAu} in minimum-bias events for both π^0 's and γ_{dir} 's is unity. The solid red line adjusts the calculation by a factor of 0.9, to better match the actually measured $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$ in minimum-bias events by PHENIX.

model down by a factor 0.90, shown as a solid red line in Fig. 4. I note that this factor is close to the measured minimum-bias $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$ value of 0.92, and is well within the normalization uncertainty of the data. This may alternatively be thought of as scaling the data up by approximately two-thirds of its stated global uncertainty. With this normalization, the model now provides an excellent description of the central values of the PHENIX data. While not explored here, I note that taking into account the uncertainties in the model parameters themselves [23] may further improve the data-model agreement. Thus, no other physics effects, such as a significant centrality-dependent jet quenching, are needed to describe the PHENIX data.

In the future, several experimental avenues may help to separate the contribution of the color fluctuation effect from searches for the jet quenching physics of interest. First, one may design the kinematic selections with an aim to match

the Bjorken- x ranges accessed by the hadron/jet and photon processes, and thus cancel the impact of the color fluctuation effect. Second, as motivated in Ref. [34], the two effects are expected to have different sensitivities to changing the projectile, such as in $p + Au$ or ${}^3\text{He} + Au$ data previously recorded at RHIC. This has been performed for π^0 's in Ref. [35], but not yet for γ_{dir} . Third, one could attempt a centrality selection based on measuring spectator neutrons in a zero-degree calorimeter (e.g., as used in Refs. [27,28]), which may better isolate the underlying geometry and not be sensitive to physics effects which manifest as a bias on the produced multiplicity. Fourth, oxygen-oxygen (O + O) collisions at RHIC and the LHC [36] would allow for the study of jet quenching in a small system, in which a single weakly-interacting nucleon in the ${}^{16}\text{O}$ nucleus will have a much smaller impact on the overall centrality signal.

Conclusion. This paper examines a recent PHENIX measurement which found a different degree of centrality-dependent modification for π^0 's and γ_{dir} 's in $d + Au$ collisions at RHIC. The measurement strategy is ostensibly chosen to calibrate out any centrality-dependent biases with the γ_{dir} measurement. However, I show that the particular π^0 and γ_{dir} kinematic selections used in the measurement select events which arise from different distributions of Bjorken- x values. As such, the PHENIX measurement is directly sensitive to the physics effects described by the color fluctuation model, in which centrality-dependent modifications (which do not arise from jet quenching) systematically increase with x . Using the model parameters determined from RHIC and LHC measurements reported ten years ago, without any postdiction updating, I show that a straightforward application of the model to the PHENIX kinematics gives a good description of the experimental data. After accounting for the possible role of color fluctuation effects, the evidence for any remaining final-state effects, such as from jet quenching, is significantly more limited. This finding reconciles the interpretation of the PHENIX data with that from other measurements at RHIC and LHC, which have set stringent limits on the possible amount of parton energy loss in small collision systems.

Acknowledgments. D.V.P. acknowledges John Lajoie, Riccardo Longo, Jamie Nagle, Peter Steinberg, and Mark Strikman for useful discussions and input on an early draft of this paper. D.V.P.'s work is supported by the U.S. Department of Energy, Grant No. DE-FG02-03ER41244.

[1] C. A. Salgado *et al.*, Proton-nucleus collisions at the LHC: Scientific opportunities and requirements, *J. Phys. G* **39**, 015010 (2012).
 [2] C. A. Salgado and J. P. Wessels, Proton-lead collisions at the CERN LHC, *Annu. Rev. Nucl. Part. Sci.* **66**, 449 (2016).
 [3] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, EPPS21: A global QCD analysis of nuclear PDFs, *Eur. Phys. J. C* **82**, 413 (2022).
 [4] S. J. Paul (CLAS Collaboration) *et al.*, Observation of azimuth-dependent suppression of hadron pairs in electron scattering off nuclei, *Phys. Rev. Lett.* **129**, 182501 (2022).

[5] J. L. Nagle and W. A. Zajc, Small system collectivity in relativistic hadronic and nuclear collisions, *Annu. Rev. Nucl. Part. Sci.* **68**, 211 (2018).
 [6] R. D. Weller and P. Romatschke, One fluid to rule them all: Viscous hydrodynamic description of event-by-event central p+p, p+Pb and Pb+Pb collisions at $\sqrt{s} = 5.02$ TeV, *Phys. Lett. B* **774**, 351 (2017).
 [7] J. Adam (ALICE Collaboration) *et al.*, Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions, *Nat. Phys.* **13**, 535 (2017).
 [8] A. M. Sirunyan (CMS Collaboration) *et al.*, Elliptic flow of charm and strange hadrons in high-multiplicity $p + Pb$ col-

- lisions at $\sqrt{s_{NN}} = 8.16$ TeV, *Phys. Rev. Lett.* **121**, 082301 (2018).
- [9] G. Aad (ATLAS Collaboration) *et al.*, Transverse momentum and process dependent azimuthal anisotropies in $\sqrt{s_{NN}} = 8.16$ TeV $p + \text{Pb}$ collisions with the ATLAS detector, *Eur. Phys. J. C* **80**, 73 (2020).
- [10] K. Tywoniuk, Is there jet quenching in pPb? *Nucl. Phys. A* **926**, 85 (2014).
- [11] A. Huss, A. Kurkela, A. Mazeliauskas, R. Paatelainen, W. van der Schee, and U. A. Wiedemann, Discovering partonic rescattering in light nucleus collisions, *Phys. Rev. Lett.* **126**, 192301 (2021).
- [12] D. V. Perepelitsa, Connection between soft and hard probes of small collision systems at RHIC and LHC, *EPJ Web Conf.* **235**, 04003 (2020).
- [13] A. Adare (PHENIX Collaboration) *et al.*, Centrality categorization for $R_{p(d)+A}$ in high-energy collisions, *Phys. Rev. C* **90**, 034902 (2014).
- [14] J. Adam (ALICE Collaboration) *et al.*, Centrality dependence of particle production in $p\text{-Pb}$ collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. C* **91**, 064905 (2015).
- [15] D. V. Perepelitsa and P. A. Steinberg, Calculation of centrality bias factors in $p + A$ collisions based on a positive correlation of hard process yields with underlying event activity, [arXiv:1412.0976](https://arxiv.org/abs/1412.0976).
- [16] C. Loizides and A. Morsch, Absence of jet quenching in peripheral nucleus–nucleus collisions, *Phys. Lett. B* **773**, 408 (2017).
- [17] G. Aad (ATLAS Collaboration) *et al.*, Centrality and rapidity dependence of inclusive jet production in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector, *Phys. Lett. B* **748**, 392 (2015).
- [18] S. Chatrchyan, Jr. (CMS Collaboration) *et al.*, Studies of dijet transverse momentum balance and pseudorapidity distributions in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Eur. Phys. J. C* **74**, 2951 (2014).
- [19] STAR Collaboration, Event activity correlations and jet measurements in p+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, [arXiv:2404.08784](https://arxiv.org/abs/2404.08784).
- [20] A. Adare (PHENIX Collaboration) *et al.*, Centrality-dependent modification of jet-production rates in deuteron-gold collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* **116**, 122301 (2016).
- [21] G. Aad (ATLAS Collaboration) *et al.*, Measurement of the centrality dependence of the dijet yield in $p + \text{Pb}$ collisions at $\sqrt{s_{NN}} = 8.16$ TeV with the ATLAS detector, *Phys. Rev. Lett.* **132**, 102301 (2024).
- [22] M. Alvioli, B. A. Cole, L. Frankfurt, D. V. Perepelitsa, and M. Strikman, Evidence for x -dependent proton color fluctuations in pA collisions at the CERN Large Hadron Collider, *Phys. Rev. C* **93**, 011902(R) (2016).
- [23] M. Alvioli, L. Frankfurt, D. Perepelitsa, and M. Strikman, Global analysis of color fluctuation effects in proton- and deuteron-nucleus collisions at RHIC and the LHC, *Phys. Rev. D* **98**, 071502(R) (2018).
- [24] V. Khachatryan (CMS Collaboration) *et al.*, Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *J. High Energy Phys.* **04** (2017) 039.
- [25] S. Acharya (ALICE Collaboration) *et al.*, Nuclear modification factor of light neutral-meson spectra up to high transverse momentum in p–Pb collisions at $\sqrt{s_{NN}} = 8.16$ TeV, *Phys. Lett. B* **827**, 136943 (2022).
- [26] G. Aad (ATLAS Collaboration) *et al.*, Charged-hadron production in pp , $p+\text{Pb}$, $\text{Pb}+\text{Pb}$, and $\text{Xe}+\text{Xe}$ collisions at $\sqrt{s_{NN}} = 5$ TeV with the ATLAS detector at the LHC, *J. High Energy Phys.* **07** (2023) 074.
- [27] S. Acharya (ALICE Collaboration) *et al.*, Constraints on jet quenching in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured by the event-activity dependence of semi-inclusive hadron-jet distributions, *Phys. Lett. B* **783**, 95 (2018).
- [28] G. Aad (ATLAS Collaboration) *et al.*, Strong constraints on jet quenching in centrality-dependent $p + \text{Pb}$ collisions at 5.02 TeV from ATLAS, *Phys. Rev. Lett.* **131**, 072301 (2023).
- [29] N. J. Abdulameer (PHENIX Collaboration) *et al.*, Disentangling centrality bias and final-state effects in the production of high- $p_T \pi^0$ using direct γ in $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV, [arXiv:2303.12899](https://arxiv.org/abs/2303.12899).
- [30] A. Drees, PHENIX overview, Proceedings of the 7th International Conference on the Initial Stages in High-Energy Nuclear Collisions, June, 19–23, 2023, Copenhagen (unpublished).
- [31] C. Bierlich *et al.*, A comprehensive guide to the physics and usage of PYTHIA 8.3, *SciPost Phys. Codeb.* **8** (2022).
- [32] A. Adare (PHENIX Collaboration) *et al.*, Direct photon production in $d + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **87**, 054907 (2013).
- [33] M. Aaboud (ATLAS Collaboration) *et al.*, Measurement of prompt photon production in $\sqrt{s_{NN}} = 8.16$ TeV $p + \text{Pb}$ collisions with ATLAS, *Phys. Lett. B* **796**, 230 (2019).
- [34] D. McGlinchey, J. L. Nagle, and D. V. Perepelitsa, Consequences of high- x proton size fluctuations in small collision systems at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **94**, 024915 (2016).
- [35] U. A. Acharya (PHENIX Collaboration) *et al.*, Systematic study of nuclear effects in $p + \text{Al}$, $p + \text{Au}$, $d + \text{Au}$, and $^3\text{He} + \text{Au}$ collisions at $\sqrt{s_{NN}} = 200$ GeV using π^0 production, *Phys. Rev. C* **105**, 064902 (2022).
- [36] J. Brewer, A. Mazeliauskas, and W. van der Schee, Opportunities of OO and pO collisions at the LHC, [arXiv:2103.01939](https://arxiv.org/abs/2103.01939).