

## Proton-deuteron bremsstrahlung at 145 and 195 MeV

J. Clayton,\* W. Benenson, M. Cronqvist,<sup>†</sup> R. Fox, D. Krofcheck,<sup>‡</sup> R. Pfaff,  
T. Reposeur,<sup>§</sup> J. D. Stevenson,\* J. S. Winfield, and B. Young

*National Superconducting Cyclotron Laboratory and Department of Physics and Astronomy, Michigan State University,  
East Lansing, Michigan 48824*

M. F. Mohar

*National Superconducting Cyclotron Laboratory and Department of Chemistry, Michigan State University,  
East Lansing, Michigan 48824*

C. Bloch and D. E. Fields

*Indiana University Cyclotron Facility, Bloomington, Indiana 47405*

(Received 22 August 1991)

Energy spectra and angular distributions have been measured for high energy gamma rays ( $E_\gamma \geq 20$  MeV) from the  $p+d$  reaction at 145 and 195 MeV. Gamma rays were observed up to the maximum energy allowed by kinematics. A comparison is made with previous measurements for the  $p+d$  system at 140, 197, and 200 MeV. Below the free  $pn\gamma$  threshold the general shape of the energy spectra and angular distributions are in reasonable agreement with a recent calculation of the free  $pn\gamma$  elementary process. However, the magnitude of the predicted cross section is not in good agreement with the present data.

PACS number(s): 25.40. - h

### I. INTRODUCTION

Reports on hard photon production by several research groups [1–6] prompted investigations which covered the periodic table from light systems, such as  $d+C$  [7], to heavy systems, such as  $Xe+Sn$  [8]. Incident beam energies have ranged from  $E/A=10$  to 124 MeV. The characteristics of photon emission for the majority of these studies strongly suggest that the main source of the radiation is first-chance proton-neutron bremsstrahlung. Proton-proton bremsstrahlung is expected to play a minor role in this energy regime. With all the interest in the heavy-ion data, it is quite surprising that there is little information on proton-nucleus bremsstrahlung [9–11]. Unfortunately, there is even less data on free  $pn\gamma$ , and the published data that do exist is with poor statistics [12–14]. Lacking incontrovertible data on the free  $pn\gamma$  cross section, information on the cross section can still be gleaned by studying proton-nucleus bremsstrahlung. As was described in an accompanying paper on high energy gamma ray production in proton-nucleus collisions [15], the phase space problem is more tractable than in the

nucleus-nucleus case.

In this paper we report on  $p+d$  bremsstrahlung ( $pd\gamma$ ) at incident proton energies of 145 and 195 MeV. The  $pd\gamma$  reaction is important because the reaction makes the transition between the elementary  $pn\gamma$  reaction and proton-nucleus bremsstrahlung. For low and intermediate energy photons, the  $pd\gamma$  result should be a good approximation to the free  $pn\gamma$  cross section, while at energies above the kinematic limit for free  $pn\gamma$  the gamma ray production cross section should reflect the influence of the internal momentum distribution in the deuteron. Comparisons between the deuteron results and heavier targets is the best way to obtain information on the importance of multistep collision processes in proton-nucleus bremsstrahlung reactions and phase space considerations. These ideas can then be extended to explain the nucleus-nucleus bremsstrahlung data.

Previously, there have been three  $pd\gamma$  measurements reported in the literature [9,16,17] Edgington and Rose [9] used a 140 MeV proton beam at Harwell to study inclusive photon production. In this measurement they employed a ( $D_2O-H_2O$ ) subtraction and obtained a poor resolution gamma ray energy spectrum. They found a total  $pd\gamma$  gamma ray cross section of  $4.3 \pm 0.3 \mu b$  for gamma rays with energies  $E_\gamma \geq 40$  MeV. Koehler, Rothe, and Thorndike [16] used a 197 MeV proton beam in conjunction with a liquid-deuterium target. Photons above 40 MeV were detected in coincidence with one or two charged particles. Koehler, Rothe, and Thorndike measured a total  $pd\gamma$  cross section of  $\approx 23 \mu b$  for gamma rays above 40 MeV, which was clearly incompatible with the result of Edgington and Rose. It should be noted that there is a value of  $26 \mu b$  for the total cross section at 148 MeV [18] for gamma rays above 25 MeV. Using this

\*Present address: Science Applications International Corporation, 2950 Patrick Henry Drive, Santa Clara, CA 95054.

<sup>†</sup>On leave from Dept. of Physics, Chalmers Univ. of Tech., S-412 96 Göteborg, Sweden.

<sup>‡</sup>Present address: Lawrence Livermore National Laboratory, Livermore, CA 94550.

<sup>§</sup>Present address: Laboratoire de Physique Nucléaire, Univ. de Nantes, 2 rue de la Houssinière, 44072 Nantes CEDEX 03, France.

value to make a crude estimate of the cross section for gamma rays above 40 MeV at a beam energy of 148 MeV, we obtain a value of at least  $12 \mu\text{b}$ . The most recent measurement of  $pd\gamma$  bremsstrahlung was completed by Pinston *et al.* [17] with 200 MeV protons incident on targets of C and  $\text{CD}_2$ . The  $pd\gamma$  result was obtained by a ( $\text{CD}_2$ -C) subtraction. Inclusive gamma ray energy spectra above 20 MeV were measured in a large  $\text{NaI(Tl)} + \text{BaF}_2$  telescope [19]. They found the  $pd\gamma$  cross section for gamma rays with  $E_\gamma \geq 40$  MeV to be  $\approx 34 \mu\text{b}$ , which supports the value obtained by Koehler, Rothe, and Thorndike and casts serious doubt on the value obtained by Edgington and Rose. Earlier work by Kwato Njock *et al.* [10] and Pinston *et al.* [11] was also in disagreement with the values obtained by Edgington and Rose for heavier target data.

The objective of the experiment discussed in this paper was to study photon production for the  $pd\gamma$  reaction at 145 and 195 MeV and measure the angular distribution and total cross section for these bremsstrahlung photons. We also measured the angular distribution for this reaction and will compare our value to the earlier measurements mentioned above. We will also compare our data to the most recent calculations by Herrmann, Speth, and Nakayama on free  $pn\gamma$  [21].

## II. EXPERIMENT

The energy spectra and angular distribution of gamma rays were measured in the energy range between 20 and 170 MeV. Details of the experimental setup can be found in a separate paper on high energy gamma ray production in proton-induced reactions [15]. Proton beams of 145 and 195 MeV bombarded self-supporting foils of C and  $\text{CD}_2$ . The thicknesses of the targets ranged from 31 to  $51.3 \text{ mg/cm}^2$  for the  $\text{CD}_2$  target. The  $\text{CD}_2$  target was monitored by comparing the rates in four BGO detectors arranged around the target. The  $\text{BaF}_2$  detectors covered the angles between  $45^\circ$  and  $135^\circ$ . We also made several runs in which both detectors were operated simultaneously at the same angle and found that to within statistical uncertainties there was no discernible loss of deuterium in the  $\text{CD}_2$  targets.

The extraction of the bremsstrahlung cross section for deuterium from the  $\text{CD}_2$  target was made quite simple by the fact that there is an excited state gamma ray transition at 15.1 MeV in  $^{12}\text{C}$ . This discrete gamma ray state is clearly evident in Fig. 1 where the gamma ray energy spectra for both the  $\text{CD}_2$  and C targets are displayed. The procedure was then to find the 15.1 MeV yield for the two runs at each of the angles and subtract the properly normalized C spectra from the  $\text{CD}_2$  spectra. The yield of 15.1 MeV gamma rays from the C target is given by

$$Y = \sigma_{15.1} N_p \left( \frac{N_A \rho t \Delta\Omega}{A} \right), \quad (1)$$

where  $\sigma$  is the cross section for the 15.1 MeV state,  $N_p$  is the number of protons incident on the target, and  $N_A$  is Avogadro's number.  $A$  is the atomic number of either




FIG. 1. Gamma ray energy spectra at  $90^\circ$  in the laboratory for both the C (lower frame) and  $\text{CD}_2$  (upper frame) targets at 145 MeV. The 15.1 MeV gamma ray transition in  $^{12}\text{C}$  is clearly observed.

the C or  $\text{CD}_2$ ,  $\rho t$  is the target thickness in units of  $\text{mg/cm}^2$ , and  $\Delta\Omega$  is the solid angle covered by the detector. The normalization factor is therefore given by the ratio of the yield of the 15.1 MeV state with the  $\text{CD}_2$  target to that of the C target. Empirically, this ratio is given by

$$R = \frac{N_{p1}\rho_1 t_1 A_2}{N_{p2}\rho_2 t_2 A_1}. \quad (2)$$

This was used as a cross-check against the experimentally derived value which was determined by fits to the 15.1 MeV peak for both targets. A smooth polynomial background was assumed for both spectra. In general, the results for both methods showed small variations overall (less than 10%) with largest difference at 30% for the




FIG. 2. Bremsstrahlung gamma ray spectrum for deuterium at 145 MeV; also shown is a comparison of the radiative capture peak  $p + d \rightarrow {}^3\text{He} + \gamma$ . The capture peak is generated by using the value for the radiative capture process measured by Pikar *et al.* [20] and a procedure described in the text.

value at  $60^\circ$  for the 145 MeV data set. After the properly normalized C spectrum is subtracted from the  $\text{CD}_2$  spectrum, what remains is the bremsstrahlung spectrum for deuterons. The results of the subtraction are displayed in Fig. 2. The peak near the end point arises from the radiative capture process  $p + d \rightarrow {}^3\text{He} + \gamma$ . The magnitude of this cross section and its energy have been verified by comparing with previous measurements by Pikar *et al.* at 150 and 200 MeV [20]. The measurement by Pikar *et al.* was a coincidence experiment detecting both the  ${}^3\text{He}$  as well as the emitted gamma ray. Also shown in Fig. 2 is an energy spectrum constructed from the capture cross section at 150 MeV reported by Pikar *et al.* at  $90^\circ$  in the laboratory. The cross section was adjusted according to the observed systematic decrease in the yield as the incident energy is increased. This adjusted value of the cross section was assumed to be Gaussian and then transformed from the nucleon-nucleon center-of-mass to the laboratory frame. The transformed Gaussian function for the capture cross section was then folded with the measured response function for the  $\text{BaF}_2$  detector. The agreement demonstrates the validity of our subtraction method. All of the energy spectra are then corrected for this capture contribution using the data from Pikar *et al.* and the procedures outlined above.

### III. RESULTS AND DISCUSSION

A comparison of the deuteron energy spectra at  $90^\circ$  using the reduced variable  $E_\gamma/E_p$  at 145 and 195 MeV shows reasonable agreement between the two incident energies (Fig. 3), reflecting the consistency of the methods to derive the deuteron bremsstrahlung energy spectra. Figure 4 shows a direct comparison between the energy spectrum for our data at  $90^\circ$  in the laboratory for protons at 195 MeV to the data of Pinston *et al.* [11] at 200 MeV, and there is a reasonable agreement between the present measurements and data of Pinston *et al.* At the most backward angle,  $\approx 150^\circ$  all the data sets agree to within statistical uncertainty. However, the present data and




FIG. 3. Comparison of the energy spectra at  $90^\circ$  in the laboratory for deuterium at 145 and 195 MeV vs a reduced variable  $E_\gamma/E_p$ , which is the measured photon energy divided by the incident proton energy.




FIG. 4. Comparison of the gamma ray energy spectra for deuterium at 195 MeV to the data of Pinston *et al.* at 200 MeV [17]. Both energy spectra have been corrected for the radiative capture cross section.

data of Pinston *et al.* exhibit a discrepancy with the measurements of Koehler, Rothe, and Thorndike [16] at the forward angles. The magnitude of this discrepancy increases for smaller angles. It should be noted that the data from Pinston *et al.* required proper treatment for not only the capture process, but also for the contribution from  $\pi^0$  decay. The threshold for the reaction  $p + d \rightarrow {}^3\text{He} + \pi^0$  is 198.7 MeV [22], and Pinston *et al.* found the contribution to the total cross section for gamma rays with  $E_\gamma \geq 40$  MeV to be roughly 10%. The measurement by Koehler, Rothe, and Thorndike was performed at  $197 \pm 5$  MeV, and so this data may also contain some background contribution due to  $\pi^0$  decay. A comparison of the angular distributions measured by Koehler, Rothe, and Thorndike and Pinston *et al.* to the present data at 195 MeV is displayed in Fig. 5. The energy threshold is 40 MeV in the laboratory frame, but the cross section is in the nucleon-nucleon center-of-mass frame, and again it can be clearly seen that there is reasonable agreement between the measurements. It should be noted that the angular distribution in the nucleon-nucleon center-of-mass frame is not the expected




FIG. 5. Comparison of the measured angular distribution for  $pd\gamma$  at 195, 197 [16], and 200 MeV [17].

TABLE I. Comparison of the total cross section for gamma rays above 40 MeV from proton-induced reactions on deuterium at incident energies of 140 [9], 145, 148 [18], 195, 197 [16], and 200 MeV [11].

$E_p$	Target	$\sigma_{\text{tot}}$ ( $\mu\text{b}$ )	Reference
140	$^2\text{H}$	$4.3 \pm 0.3$	[9]
145	$^2\text{H}$	$21 \pm 2$	present data
148	$^2\text{H}$	12	[18]
195	$^2\text{H}$	$34 \pm 4$	present data
197	$^2\text{H}$	$23 \pm 12$	[16]
200	$^2\text{H}$	34	[11]

isotropic plus dipole component, but exhibits an isotropic plus  $\cos(\theta)$  dependence. The ratio of  $\sigma(45^\circ)/\sigma(135^\circ)$  in the nucleon-nucleon center-of-mass frame is  $2.5 \pm 0.3$  at 195 MeV and  $3.1 \pm 0.3$  at 145 MeV, which contrasts with value of  $1.1 \pm 0.1$  we found for our earlier measurements on a Pb target [15]. This forward peaking in the emitting frame may be due to the asymmetry of the collision, since in heavy-ion collisions it is generally believed that the reaction mechanism is incoherent proton-neutron bremsstrahlung and in these reactions there is an average over protons from the target colliding with projectile neutrons and vice versa. As a result, we are only left with isotropic and dipole terms, but in the proton-nucleus case that symmetry is lost. Herrmann, Speth, and Nakayama [21] predict a forward peaking in the nucleon-nucleon frame for the  $pn\gamma$  process. However, the effect is much smaller than that observed in the present data. One other explanation may be the influence of multiple collisions in the target nucleus, which could explain the difference between deuterium and lead. The value of the total cross section for  $E_\gamma \geq 40$  MeV that we obtain at 195 MeV is  $34 \pm 4 \mu\text{b}$ , which is in agreement with the value of  $23 \pm 12 \mu\text{b}$  reported by Koehler, Rothe, and Thorndike [16] and the  $\approx 34 \mu\text{b}$  value of Pinston *et al.* [11].

The value for the total cross section at 145 MeV is  $21 \pm 2 \mu\text{b}$ , which is 4.7 times greater than the value reported by Edgington and Rose at 140 MeV [9]. It is also higher than the estimated value of  $12 \mu\text{b}$ , which is based on the cross section of  $26 \mu\text{b}$  reported by Rothe, Koehler, and Thorndike for photons with  $E_\gamma \geq 25$  MeV. In that measurement Rothe, Koehler, and Thorndike [18] used a beam degrader to lower the incident beam energy from 197 to 148 MeV. The results from all the measurements are compiled in Table I.

Recent calculations on free  $pn\gamma$  by Herrmann, Speth, and Nakayama [21] use a meson-exchange potential model, and they also give a careful treatment of one-body rescattering. It is shown that the one-body term enhances the cross section near the photon maximum. Figure 6 shows a comparison of our data in the nucleon-nucleon center-of-mass frame for both incident proton energies. The calculation reproduces the shape of the low energy portion of spectrum and does well up to the kinematic limit for the free  $pn\gamma$  cross section. The calculation is in




FIG. 6. Comparison of the measured  $pd\gamma$  energy spectra at 145 and 195 MeV in the nucleon-nucleon center-of-mass frame with the free  $pn\gamma$  calculation of Herrmann, Speth, and Nakayama at 150 and 200 MeV [21].

much better agreement with the data at 200 MeV than the data at 145 MeV. In fact, the calculation underpredicts the magnitude of the data at 145 MeV by roughly 30%. At 195 MeV the agreement is quite good up to 80 MeV; after this energy it is not possible to compare  $pd\gamma$  with free  $pn\gamma$  if the internal momentum distributions is not taken into account. We see a decrease in the cross section for  $pd\gamma$  at high photon energies, which arises from Pauli blocking effects in the deuterium nucleus. The calculated values for the total cross section reported by Herrmann, Speth, and Nakayama [21] for photons above  $E_\gamma \geq 40$  MeV were  $14.5 \mu\text{b}$  at 140 MeV and  $22 \mu\text{b}$  at 197 MeV.

#### IV. CONCLUSIONS

In conclusion, the present measurement confirms the earlier findings of Pinston *et al.* [11] and is in strong disagreement with the cross sections reported by Edgington and Rose [9]. In contrast, we are also in agreement with the earlier measurement of Koehler, Rothe, and Thorndike [16]. Recent calculations by Nakayama [23] which take into account the deuteron wave function reproduce the data reasonably well.

#### ACKNOWLEDGMENTS

The authors would like to thank the operations group at IUCF for their assistance. We would also like to thank Professor G. F. Bertsch and Professor W. Bauer for their very helpful discussions. This work was supported in part by Grant No. PHY-89-13815 from the National Science Foundation.

- [1] K. B. Beard, W. Benenson, C. Bloch, E. Kashy, J. D. Stevenson, D. J. Morrissey, J. van der Plicht, B. Sherrill, and J. S. Winfield, *Phys. Rev. C* **32**, 1111 (1985).
- [2] E. Grosse, P. Grimm, H. Heckwolf, W. F. J. Mueller, H. Noll, A. Oskarsson, H. Stelzer, and W. Rosch, *Europhys. Lett.* **2**, 9 (1986).
- [3] N. Alamanos, P. Braun-Munzinger, R. F. Freifelder, P. Paul, J. Stachel, T. C. Awes, R. L. Ferguson, F. E. Obenshain, F. Plasil, and G. R. Young, *Phys. Lett. B* **173**, 392 (1986).
- [4] M. Kwato Njock, M. Maurel, E. Monnard, H. Nifenecker, J. Pinston, F. Schussler, and D. Barneoud, *Phys. Lett. B* **175**, 125 (1986).
- [5] J. D. Stevenson, K. B. Beard, W. Benenson, J. Clayton, E. Kashy, A. Lampis, D. J. Morrissey, M. Samuel, R. J. Smith, C. L. Tam, and J. S. Winfield, *Phys. Rev. Lett.* **57**, 555 (1986).
- [6] R. Hingmann, W. Kuhn, V. Metag, R. Muhlans, R. Novotny, A. Ruckelshausen, W. Cassing, B. Haas, H. Em-ling, R. Kulesa, H. J. Wallersheim, J. P. Vivien, A. Boullay, H. Delagrange, H. Doubre, C. Gregoire, and Y. Schutz, *Phys. Rev. Lett.* **58**, 759 (1987).
- [7] C. L. Tam, J. D. Stevenson, W. Benenson, Y. Chen, J. Clayton, E. Kashy, A. R. Lampis, D. J. Morrissey, T. K. Murakami, M. Samuel, and J. S. Winfield, *Phys. Rev. C* **38**, 2526 (1989).
- [8] J. Clayton, J. D. Stevenson, W. Benenson, Y. Chen, E. Kashy, A. R. Lampis, M. F. Mohar, D. J. Morrissey, T. K. Murakami, M. Samuel, B. Sherrill, C. L. Tam, and J. S. Winfield, *Phys. Rev. C* **40**, 1207 (1989).
- [9] J. A. Edgington and B. Rose, *Nucl. Phys.* **89**, 523 (1966).
- [10] M. Kwato Njock, M. Maurel, H. Nifenecker, J. Pinston, F. Schussler, D. Barneoud, S. Drissi, J. Kern, and J. P. Vorlet, *Phys. Lett. B* **207**, 269 (1988).
- [11] J. A. Pinston, D. Barneoud, V. Bellini, S. Drissi, J. Guil- lot, J. Julien, M. Kwato Njock, H. Nifenecker, M. Maurel, F. Schussler, and J. P. Vorlet, *Phys. Lett. B* **218**, 128 (1989).
- [12] F. P. Brady and J. C. Young, *Phys. Rev. C* **4**, 1579 (1970).
- [13] J. A. Edgington, V. J. Howard, I. M. Blair, B. E. Bonner, F. P. Brady, and M. W. McNaughton, *Nucl. Phys. A* **218**, 151 (1974).
- [14] C. Dupont, C. Deom, P. Leleux, P. Lipnik, P. Macq, A. Ninane, J. Pestieau, S. W. Kitwanga, and P. Wauters, *Nucl. Phys. A* **481**, 424 (1988).
- [15] J. Clayton, W. Benenson, M. Cronqvist, R. Fox, D. Krofcheck, R. Pfaff, T. Reposeur, J. D. Stevenson, J. S. Winfield, B. Young, M. F. Mohar, C. Bloch, and D. E. Fields, *Phys. Rev. C* **45**, 1815 (1992) the following paper.
- [16] P. F. M. Koehler, K. W. Rothe, and E. H. Thorndike, *Phys. Rev. Lett.* **18**, 933 (1967).
- [17] J. A. Pinston, D. Barneoud, V. Bellini, S. Drissi, J. Guil- lot, J. Julien, H. Nifenecker, and F. Schussler, *Phys. Lett. B* **249**, 402 (1990).
- [18] K. W. Rothe, P. F. M. Koehler, and E. H. Thorndike, contribution to the Williamsburg Conference on Inter- mediate Energy Physics (College of William and Mary, Williamsburg, VA, 1966), p. 677.
- [19] R. Bertholet, M. Kwato Njock, M. Maurel, E. Monnard, H. Nifenecker, P. Perrin, J. A. Pinston, F. Schussler, D. Barneoud, C. Guet, and Y. Schutz, *Nucl. Phys. A* **474**, 541 (1987).
- [20] M. A. Pikar, H. J. Karwowski, J. D. Brown, J. R. Hall, M. Hugi, R. E. Pollock, V. R. Cupps, M. Fatyga, and A. D. Bacher, *Phys. Rev. C* **35**, 37 (1987).
- [21] V. Herrmann, J. Speth, and K. Nakayama, *Phys. Rev. C* **43**, 394 (1991).
- [22] M. A. Pikar, in *Pion Production and Absorption in Nuclei—1981 (Indiana University Cyclotron Facility)*, Proceedings of the Conference on Pion Production and Absorption in Nuclei, edited by Robert D. Bent, AIP Conf. Proc. No. 79 (AIP, New York, 1982), p. 541.
- [23] K. Nakayama, *Phys. Rev. C* (to be published).