

$E2$ properties of nuclei far from stability and the proton-halo problem of ${}^8\text{B}$

H. Nakada^{1,3} and T. Otsuka^{2,3}

¹*Department of Physics, Juntendo University, Hiraga-gakueidai 1-1, Inba-mura, Inba-gun, Chiba 270-16, Japan*

²*Department of Physics, Faculty of Science, University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113, Japan*

³*Institute for Nuclear Theory, University of Washington, HN-12, Seattle, Washington 98195*

(Received 22 October 1993)

$E2$ properties of $A = 6-10$ nuclei, including those of nuclei far from stability, are studied by a $(0+2)\hbar\omega$ shell-model calculation which includes $E2$ core-polarization effects explicitly. The quadrupole moments and the $E2$ transition strengths in $A = 6-10$ nuclei are described quite well by the present calculation. This result indicates that the relatively large value of the quadrupole moment of ${}^8\text{B}$ can be understood without introducing the proton halo in ${}^8\text{B}$. An interesting effect of the $2\hbar\omega$ core-polarization is found for effective charges used in the $0\hbar\omega$ shell model; although isoscalar effective charges are almost constant as a function of nucleus, appreciable variations are needed for isovector effective charges which play important roles in nuclei with high isospin values.

PACS number(s): 21.10.Ky, 23.20.-g, 21.60.Cs, 27.20.+n

I. INTRODUCTION

The structure of light neutron-rich nuclei has recently attracted much interest. A good example is the ${}^{11}\text{Li}$ nucleus [1], which shows the exotic feature of the neutron halo, owing to loosely bound neutrons. Research on light proton-rich nuclei is also in progress. One then comes up with a question of whether or not the proton halo exists despite the Coulomb barrier. Recently Minamisono *et al.* succeeded in a precise measurement of the quadrupole moment of ${}^8\text{B}$ [2]. They pointed out [2,3] that the quadrupole moment of ${}^8\text{B}$ is considerably larger than the shell-model prediction. They claimed the existence of a proton halo [2] based on the analysis of Ref. [3]. This analysis, however, appears to be model dependent, as will be discussed later.

On the other hand, the observed interaction radius of ${}^8\text{B}$ shows no enhancement, compared with those of surrounding nuclei [4]. In contrast to the case of ${}^{11}\text{Li}$, this datum undoubtedly contradicts with the proton-halo hypothesis [2,3] of ${}^8\text{B}$. Thus, a serious conflict arises between Refs. [2] and [4].

In this paper, we shall consider $E2$ properties of light nuclei, from a more general perspective including nuclei far from stability. While the isoscalar degrees of freedom dominate the $E2$ properties of low-lying states of light stable nuclei, the isovector ones become important in unstable nuclei with higher isospin. This work will shed light on a new aspect in the structure of light unstable nuclei, namely, the $E2$ properties.

As will be shown later, it is difficult to avoid in the $0\hbar\omega$ shell-model calculation ambiguities arising from effective charges as well as from single-particle wave functions (i.e., single-particle matrix elements). It is of much importance to remove such ambiguities in effective charges. For this purpose, a shell-model calculation with explicit inclusion of the $2\hbar\omega$ excitation is very useful, because this

excitation is the major origin of the effective charges and therefore the effective charges are almost equal to the bare charges in the $(0+2)\hbar\omega$ shell model. We investigate the $E2$ properties in terms of the $(0+2)\hbar\omega$ shell-model calculation by using the interaction introduced by Wolters *et al.* [5]. Although several problems have been pointed out with respect to this interaction [6], it has a certain advantage in investigating $E2$ properties. We shall also discuss some effects of the $2\hbar\omega$ configuration on the isoscalar and isovector effective charges.

II. $(0+2)\hbar\omega$ SHELL-MODEL CALCULATION

The interaction of Wolters *et al.* [5] has been determined for the $(0+2)\hbar\omega$ shell-model calculation, so as to fit the experimental energies, including the binding energies, of $A = 4-16$ nuclei. In this paper, we shall primarily discuss results obtained by a $(0+2)\hbar\omega$ shell-model calculation with this interaction. The harmonic-oscillator basis is employed with a constant $\hbar\omega$ for all of those nuclei. Since the effective interaction is given in terms of the relative coordinates as the values of the Talmi integrals, it is free from the spurious center-of-mass motion. The energy levels of ${}^8\text{Li}$ and ${}^8\text{B}$, which are mirror nuclei, are shown in Fig. 1, in comparison with the data. The Coulomb energies are subtracted in the same manner as in Ref. [5]. Note that the isospin symmetry is maintained in the present calculation. The result of the $0\hbar\omega$ shell-model calculation with the Cohen-Kurath (8-16)TBME interaction [7] is also displayed in Fig. 1. The ground-state energy in this $0\hbar\omega$ result is adjusted to the experimental one. It is clear that the present $(0+2)\hbar\omega$ calculation reproduces the data quite well: The agreement is better than the $0\hbar\omega$ result with the Cohen-Kurath interaction.

Although in the $(0+2)\hbar\omega$ calculation of Ref. [5] Q mo-




FIG. 1. Energy levels of ^8Li and ^8B . Coulomb energies are subtracted in the same way as Ref. [5]. The levels labeled “Utr.” are obtained by the $(0+2)\hbar\omega$ calculation with the interaction of Wolters *et al.*, while those labeled “CK” are by the $0\hbar\omega$ calculation with the Cohen-Kurath interaction.

ments were adjusted to the data of Ref. [8], we shall not adopt those parameters because of the following reasons. While we investigate both Q moments and $B(E2)$ values, no transition was discussed in Ref. [5]. The experimental value of $Q(^8\text{Li})$, which is particularly important in discussing $Q(^8\text{B})$, was confirmed recently [2,9]. This value is considerably larger than that shown in Refs. [8,10]. The

parameters in Ref. [5] do not reproduce the presently confirmed value of $Q(^8\text{Li})$.

We restrict ourselves to $A = 6-10$ nuclei, so as to keep the parameters almost constant. Because the $E2$ core-polarization effect is included significantly in the $(0+2)\hbar\omega$ space, it is expected that one can reproduce the $E2$ properties to a great extent with the bare charges. The calculated Q moments and $B(E2)$'s with the bare charges are shown in Table I, in comparison with the data. The $0\hbar\omega$ results displayed in Table I will be discussed later. It is found that, by this $(0+2)\hbar\omega$ calculation, the Q moments are already reproduced to an appreciable extent with the bare charges. If we introduce a small value of the isoscalar effective-charge correction, $\delta e_{\text{IS}} = 0.05e$ (i.e., $e_{\pi}^{\text{eff}} = 1.05e$ and $e_{\nu}^{\text{eff}} = 0.05e$), the agreement with the data in $A = 6-8$ is improved further as shown in Table I. With this parameter we come to a very good agreement in $Q(^8\text{Li})$. It is found that $Q(^8\text{B})$ is reproduced by the same parameter within reasonable accuracy ($\sim 8\%$). Thus, one cannot claim that the Q moment of ^8B is anomalously enhanced. The structure of ^8B can be understood within the shell model, as the mirror nucleus of ^8Li . Because there is no evidence for halo in ^8Li , the proton-halo hypothesis for ^8B appears to be very unlikely. Although the 8% difference in $Q(^8\text{B})$ may originate from some halolike structure (i.e., slowly damping tail), it should occur with a much reduced amplitude compared with ^{11}Li or ^{11}Be [1,4].

As is discussed in the Appendix, the tail behavior of

TABLE I. Q moments ($e\text{fm}^2$) and $B(E2)$ values ($e^2\text{fm}^4$) in $A = 6-10$ nuclei. Cal.(A): $e_{\pi}^{\text{eff}} = e$, $e_{\nu}^{\text{eff}} = 0$; Cal.(B): $e_{\pi}^{\text{eff}} = 1.05e$, $e_{\nu}^{\text{eff}} = 0.05e$; Cal.(C): $e_{\pi}^{\text{eff}} = 1.5e$, $e_{\nu}^{\text{eff}} = 0.5e$ ($b = 1.77\text{ fm}$).

Nucleus	Quantity	$(0+2)\hbar\omega$		Exp.	$0\hbar\omega$ Cal.(C)
		Cal.(A)	Cal.(B)		
^6Li	$Q(1_1^+)$	0.09	0.10	-0.08 ± 0.01^a	-1.83
	$B(E2; 3_1^+ \rightarrow 1_1^+)$	5.95	7.21	10.7 ± 0.8^b	8.13
	$B(E2; 2_1^+ \rightarrow 1_1^+)$	9.17	11.10	4.4 ± 2.3^b	3.93
^7Li	$Q([\frac{3}{2}]_1^-)$	-3.79	-4.29	-4.06^a	-4.86
	$B(E2; [\frac{1}{2}]_1^- \rightarrow [\frac{3}{2}]_1^-)$	13.21	17.21	15.7 ± 1.0^b	22.74
	$B(E2; [\frac{7}{2}]_1^- \rightarrow [\frac{3}{2}]_1^-)$	5.96	7.83	3.42^b	8.53
^8Li	$Q(2_1^+)$	2.78	3.21	3.15 ± 0.05^a	3.24
	$B(E2; 1_1^+ \rightarrow 2_1^+)$	3.94	5.30	75 ± 17^d	6.72
^8B	$Q(2_1^+)$	5.84	6.27	$6.83 \pm 0.21^{c,e}$	5.17
^9Li	$Q([\frac{3}{2}]_1^-)$	-3.89	-4.36	$-3.6 \pm 0.7^{a,e}$	-5.05
^9Be	$Q([\frac{3}{2}]_1^-)$	5.46	5.98	5.3 ± 0.3^a	5.36
	$B(E2; [\frac{5}{2}]_1^- \rightarrow [\frac{3}{2}]_1^-)$	26.39	32.02	27.1 ± 2.0^b	31.94
	$B(E2; [\frac{7}{2}]_1^- \rightarrow [\frac{3}{2}]_1^-)$	9.75	11.63	7.0 ± 3.0^b	10.74
^{10}Be	$B(E2; 2_1^+ \rightarrow 0_1^+)$	13.48	16.26	10.2 ± 1.0^b	17.38
	$B(E2; 0_2^+ \rightarrow 2_1^+)$	5.87	7.20	3.2 ± 1.9^b	0.01
^{10}B	$Q(3_1^+)$	9.62	10.58	8.47 ± 0.06^a	10.70
	$B(E2; 1_1^+ \rightarrow 3_1^+)$	1.12	1.35	4.13 ± 0.06^b	9.76
	$B(E2; 1_2^+ \rightarrow 3_1^+)$	7.11	8.61	1.71 ± 0.26^b	1.35
	$B(E2; 1_2^+ \rightarrow 1_1^+)$	3.21	3.88	0.83 ± 0.40^b	2.03
	$B(E2; 3_2^+ \rightarrow 1_1^+)$	13.86	16.77	20.5 ± 2.6^b	9.68
^{10}C	$B(E2; 2_1^+ \rightarrow 0_1^+)$	12.54	15.22	12.3 ± 2.1^b	15.01

^aReference [9].

^bReference [10].

^cReference [2].

^dReference [11].

^eThe sign is speculated from the $0\hbar\omega$ and $(0+2)\hbar\omega$ shell-model calculations.

wave functions is connected with separation energies, although this connection is not taken into account in most shell-model calculations including the present one. It is clear that the proton distribution of ${}^8\text{B}$ is damped more slowly in the radial direction than that of ${}^8\text{Li}$, corresponding to the smaller separation energy. The slower damping of the proton density could lead to a stronger influence on the Q moment. Such an effect, on the other hand, is not contained in the present calculation. As discussed above, this might be a reason why we still have a slight underestimation of $Q({}^8\text{B})$ with $\delta e_{\text{IS}} = 0.05e$, by which we can reproduce $Q({}^8\text{Li})$ very well. It should be noticed, however, that the amplitude of the tail part of the present wave function is not connected with the separation energy, as pointed out in the Appendix. The results of Ref. [4] and the present work suggest consistently that the amplitude of the slowly damping part in the proton distribution of ${}^8\text{B}$ is too small to be regarded as a proton halo. Namely, its influence on radius and Q moment does not appear to be so significant. It is emphasized that, although we have introduced a single parameter δe_{IS} , the value of this parameter is quite small and therefore the resultant ambiguity becomes much less than that arising from the effective charges needed for the $0\hbar\omega$ space. The present small value of δe_{IS} may come from the configurations which are still outside the present shell-model space and/or from a mesonic effect.

A recent experiment [11] indicates that the $B(E2; 2_1^+ \rightarrow 1_1^+)$ value of ${}^8\text{Li}$ is quite large. This datum appears to be far beyond what can be explained by the present calculation or by the cluster-model calculation of Ref. [12].

It is interesting that the Q moments and $B(E2)$'s of $A = 9$ and 10 nuclei are reproduced quite well with the bare charges, while $\delta e = 0.05e$ is preferable in $A = 6-8$. There could be a slight change in the core-polarization effect which originates from the excitations to higher $\hbar\omega$ space.

The oscillator length b is derived from the $\hbar\omega$ value fixed in the effective Hamiltonian, in the present calculation of $E2$ properties. On the other hand, for computing charge radii and Q moments Wolters *et al.* adopted a smaller b parameter in Ref. [5], which is inconsistent with the $\hbar\omega$ value for calculating energy levels. This happened because Wolters *et al.* tried to adjust the b parameter to the measured radii [13] of the whole region of $A = 4-16$ nuclei. However, as long as we restrict ourselves to $A = 6-10$ nuclei, the b parameter adopted in Ref. [5] appears to be too small to reproduce the charge radii of these nuclei.

The b parameter used in the present work reproduces the charge radius of ${}^6\text{Li}$. The approximate constancy of the radii over the p -shell nuclei [13], however, cannot be reproduced. This is a common problem with the usual shell model, especially when the harmonic-oscillator single-particle wave functions are used. If we consider the $0\hbar\omega$ space, the radius must increase with the mass number, even with a constant b value, owing to the center-of-mass correction [14]. This tendency does not change essentially in the present $(0+2)\hbar\omega$ calculation, whereas the experimental values increase more slowly. There is an approach in which the b parameter is determined by the variation with a Skyrme-type interaction for each nucleus [15]. A nearly constant value of b was then obtained for $A = 6-12$, implying that the above problem remains. We shall not discuss the radii in further detail. It is basically beyond the scope of the usual shell-model calculations, including the present one, to reproduce the radii.

III. EFFECT OF $2\hbar\omega$ CONFIGURATION

We shall consider the following ratio of the matrix elements:

$$R_T(i \rightarrow f) \equiv \frac{\langle f | \mathcal{O}^{(T)} | i \rangle}{\langle f | P_{0\hbar\omega} \mathcal{O}^{(T)} P_{0\hbar\omega} | i \rangle} \bigg/ \sqrt{\langle f | P_{0\hbar\omega} | f \rangle \langle i | P_{0\hbar\omega} | i \rangle} \quad (T = 0, 1), \quad (1)$$

where the numerator is obtained from the present $(0+2)\hbar\omega$ shell-model calculation, $P_{0\hbar\omega}$ in the denominator represents the projection operator onto the $0\hbar\omega$ space (i.e., the space consisting only of the p -shell configuration), and $T = 0$ ($T = 1$) denotes the ratio with respect to the isoscalar (isovector) operator. Since $E2$ properties are under discussion, we consider the case of $\mathcal{O} = r^2 Y^{(2)}$. Apart from the possible changes in the single-particle wave functions between the $0\hbar\omega$ and the $(0+2)\hbar\omega$ calculations mentioned below, R_T expresses, to a certain extent, the relative ratio of the effective charge for the $0\hbar\omega$ calculation over the bare charge, because these effective charges incorporate the isoscalar and isovector $2\hbar\omega$ core-polarization effects. The present single-particle wave functions are determined so as to reproduce the energy levels in the $(0+2)\hbar\omega$ space. As a result, the p -shell

single-particle wave functions do not necessarily correspond to the ones suitable for the usual $0\hbar\omega$ shell-model calculation. The overlap between these two sorts of p -shell wave functions, however, is expected to be fairly large, and therefore R_T will serve as a good measure of the effective charges for the usual $0\hbar\omega$ shell-model calculation.

Figure 2 shows the R_T values for the $E2$ properties of $A = 6-10$ nuclei. For isovector matrix elements, $R_{T=1}$ is not shown for the matrix elements between $T = 0$ states, because it is indefinite. We do not show the $R_{T=1}$ values also when the isovector matrix element is less than 15% of the isoscalar one. Note that the isovector matrix element reaches 40% of the isoscalar one at maximum. The R_T values for $B(E2; 0_2^+ \rightarrow 2_1^+)$ of ${}^{10}\text{Be}$ are omitted in Fig. 2, since the 0_2^+ state is highly dominated by the




FIG. 2. R_T values defined in Eq. (1) for Q moments and $E2$ transitions. They are displayed according to the sequence of moment or transition probability in Table I, except for $B(E2; 0_2^+ \rightarrow 2_1^+)$ of ^{10}Be . Each $R_{T=1}$ value concerns the same transition (or Q moment) as the $R_{T=0}$ value shown right above. See the text for details.

$2\hbar\omega$ configuration (97%) in the present calculation.

It is remarkable that $R_{T=0}$ is fairly constant for $A = 6$ – 10 . Most of the $R_{T=0}$ values are in the 0.9–1.2 range, which leads to $e_{1S}^{\text{eff}} \equiv \frac{1}{2}(e_{\pi}^{\text{eff}} + e_{\nu}^{\text{eff}}) = 0.45e$ – $0.6e$ for the $0\hbar\omega$ space. We point out that $R_{T=0}$ for ^8Li (or ^8B) is not extraordinary, compared with their surrounding nuclei. Since there is no enhancement in the $R_{T=0}$ value for $Q(^8\text{Li})$ and $Q(^8\text{B})$, a strong quadrupole deformation is not likely to occur in ^8B . We here state that the probability of the $2\hbar\omega$ configuration shows no increase in the ground-state wave functions of ^8Li and ^8B . This result, showing no notable increase of the $2\hbar\omega$ mixing in ^8Li , is consistent with the result of a Hartree-Fock shell-model calculation in Ref. [16].

The $R_{T=1}$ value shown in Fig. 2 fluctuates considerably. The $R_{T=1}$ value changes from 0.9–1.5 for $A = 7$ and 8 to 0.0–0.5 for $A = 9$ and 10. The $R_{T=1}$ values for the Q moments of ^8Li and ^8B are larger than those for the other nuclei, which accounts for the relatively large difference between $Q(^8\text{B})$ and $Q(^8\text{Li})$. However, the R_T 's for ^8Li and ^8B still stay within the range of its fluctuation.

The present systematic study suggests that the isovector core-polarization effect changes from nucleus to nu-

cleus in lighter mass region, probably because the nuclear system is not large enough and therefore the mean field is less developed. The R_T values shown here indicate that the $2\hbar\omega$ core-polarization effect on isovector effective charges seems to have a significant nucleus dependence. Such variation of the core-polarization effect, however, cannot be taken into consideration within the $0\hbar\omega$ configuration space, as far as constant (or almost constant) effective charges are used. The present extension of the model space has a remarkable advantage to take into account this variation of the core-polarization effect.

IV. DISCUSSION

A. Assessments of proton-halo hypothesis

It has been shown that $Q(^8\text{B})$ can be understood without introducing a rather exotic feature such as a proton halo, contrary to the analysis in Refs. [2,3]. Being consistent with the interaction-radius measurement [4], the proton halo in ^8B is not likely to exist. At this stage we should reconsider the soundness of the proton-halo hypothesis in Refs. [2,3].

The proton separation energy (S_p) of ^8B is very small (~ 0.14 MeV). This was probably one of the basic motivations for the proton-halo interpretation in Refs. [2,3]. It is known that the tail form of the ^8B wave function is connected with S_p , as is discussed in the Appendix for a more general case. We should keep in mind, however, that the separation energies do not fix the amplitude of the tail part, which is denoted by ξ in Eq. (A9). In usual cases, this amplitude is considered to be small enough to be neglected. Only the halo nuclei, in which the nucleon occupation number in the tail region is of order of magnitude one, have appreciable contributions of the tail part to various physical quantities.

In Ref. [3], Kitagawa and Sagawa compared two results with different sets of single-particle wave functions. One is obtained from the harmonic-oscillator potential, while a Woods-Saxon potential is assumed in the other. They applied the Cohen-Kurath shell-model density matrix [7] to both cases. Their procedure to determine the single-particle wave functions from the Woods-Saxon potential is explained in the following. The ground state of ^8B was expanded in the products of the ^7Be (^7B) “core” states and last proton (neutron). For the $^7\text{Be}+p$ channel, this expansion is expressed as

$$\psi(^8\text{B}; \text{g.s.}) = \sum_{i,j} c_{i,j}^p [\psi(^7\text{Be}; i) \varphi_p(j)], \quad (2)$$

where $\psi(^7\text{Be}; i)$ denotes the i th eigenstate of ^7Be , $\varphi_p(j)$ the single-particle orbit j of the last proton, and $c_{i,j}^p$ stands for the spectroscopic amplitude. The single-particle wave functions of the last proton were fixed by the observed S_p and the Woods-Saxon potential as shown below. Note that the wave function $\varphi_p(j)$ is determined by this method not only for the tail part but also for

the inner part, and is used for calculating physical observables. The proton single-particle energies $\epsilon_p(j)$ are determined by the condition

$$E(^8\text{B}; \text{g.s.}) = E(^7\text{Be}; i) + \epsilon_p(j), \quad (3)$$

for each set of i and j in Eq. (2). Note that $E(^8\text{B}; \text{g.s.})$ and $E(^7\text{Be}; i)$ are obtained from observed energies of the relevant states of ^8B and ^7Be . The depth of the Woods-Saxon potential was varied for each combination of i and j separately, so that the proton single-particle energy should be adjusted to this value. Since the proton separation energy is defined by $S_p(^8\text{B}) = E(^7\text{Be}; \text{g.s.}) - E(^8\text{B}; \text{g.s.})$, the value of $\epsilon_p(j)$ was fitted to $-S_p(^8\text{B})$, when the state i refers to the ground state in Eq. (3). Associated with the small S_p , this configuration led to the proton halo. It is noticed that the wave function of the last proton, as well as the Woods-Saxon potential depth, depend on the ^7Be -core states in Eq. (2). It was claimed in Refs. [2,3] that this adjusted Woods-Saxon approach can reproduce $Q(^8\text{B})$ with a set of effective charges similar to those of heavier nuclei.

The “adjusted Woods-Saxon” prescription described above was originally developed by Millener *et al.* [17] in order to reproduce the tail form of the total wave function. We point out here that, through the condition of Eq. (3), the following two approximations are made in determining $\varphi_p(j)$ by the “adjusted Woods-Saxon” approach: (a) The residual interaction between the ^7Be core and the last proton is ignored (likewise the interaction between ^7B and the last neutron). (b) The coupling among different configurations was ignored. In other words, off-diagonal matrix elements of the nuclear force are neglected among different configurations in Eq. (3). The separation like Eq. (3) could be valid when the quantum number of the state is owed by the last proton. This requires for the core state to be 0^+ . Thus the “adjusted Woods-Saxon” manipulation may work better for a loosely bound particle decoupled from its 0^+ inert core. The states discussed in Ref. [17] seem to be the cases of this kind. On the other hand, several problems occur in ^8B .

We list the problems on the proton-halo hypothesis in Refs. [2,3] below.

(i) As is stated in Sec. I, no enhancement of the interaction radius is observed in ^8B [4].

(ii) The lowest 2^+ , 1^+ , and 3^+ levels of ^8B correspond quite well to those of ^8Li [10], thus indicating good isospin symmetry. Note that those isobaric analog states are also observed as excited states of ^8Be [10]. Since the neutron halo is not expected in ^8Li , the proton-halo hypothesis for the ^8B ground state would destroy these isospin multiplets to a certain extent.

(iii) In ^8B , we have three protons in the p shell, as far as we work within the $0\hbar\omega$ shell-model space. In the theoretical analysis of Refs. [2,3], Kitagawa and Sagawa seem to have distinguished the orbits for the last proton from those for the other two valence protons. Hence, when the ^7Be core stays in the ground state in the expansion of Eq. (2), they assumed a loosely bound orbit for the last proton and deeply bound orbits for the other

two valence protons. The antisymmetrization among the three valence protons, however, was not treated correctly. This can lead to erroneous results.

(iv) The weak-coupling assumption stated above was fundamental for the proton-halo hypothesis in Ref. [3]. It should, however, be tested carefully whether this assumption is valid in ^8B or not. We shall estimate the coupling effects ignored in Ref. [3], which are referred to as (a) and (b) in the preceding discussion, by applying the $0\hbar\omega$ shell-model calculation with the Cohen-Kurath interaction. We obtain, for the sum of the correlation energies, $E(^7\text{Be}; \text{g.s.}) + \epsilon_p(0p_{3/2}) - E(^8\text{B}; \text{g.s.}) \sim 2 \text{ MeV}$. This value seems too large to be ignored, compared with $S_n(^8\text{Li}) - S_p(^8\text{B}) \sim 2 \text{ MeV}$.

(v) It was assumed in the theoretical interpretation of Refs. [2,3] that the effective charges should be the same among the $A = 8, 11$, and 17 nuclei. Apart from the possible arbitrariness in their selection of nuclei, the constant effective-charge assumption has to be used with extreme care to draw the proton-halo conclusion in the light mass region. Among such light nuclei, a considerable mass-number dependence of the effective charges might be possible, as we discussed in Sec. III. It can also be questioned whether or not the effective charges should be common among halo orbits and normal orbits.

(vi) The shell-model density matrix calculated with the Cohen-Kurath interaction [7] was employed in Ref. [3]. The Cohen-Kurath interaction, however, is adjusted to the levels of nonhalo nuclei. It is not consistent to apply those density matrices to the halo orbits. The influence of the halo on the density matrix should not be ignored in the case that the halo causes significant changes of physical observables.

(vii) The excited 1^+ and 3^+ states of ^8B are considered to have a certain similarity in structure to the ground 2^+ state, since these three states are regarded as the $(0p_{3/2})^4$, $T = 1$ multiplet, as a zeroth-order approximation within the jj -coupling scheme. It is desirable, therefore, that the structure of the three states should be described in a consistent manner. However, if we apply the “adjusted Woods-Saxon” method of Ref. [3] to the excited states, the following problem may occur. The 3^+ state is observed 2.2 MeV above the threshold for proton emission. Hence, Eq. (3) leads to $\epsilon_p \simeq 2.2 \text{ MeV}$ for the 3^+ state, when the state i refers to the ground state in Eq. (3) as in the 1^+ state. On the contrary, the height of a Coulomb barrier is about 1 MeV, if we calculate it by using the Woods-Saxon potential with the parameters of Ref. [18]. The higher ϵ_p than the barrier makes the state impossible to survive even as a resonance state.

(viii) Let us consider the term in Eq. (2) with the ^7Be state being the ground state. Both the proton $0p_{3/2}$ and $0p_{1/2}$ orbits can produce 2^+ states by coupling to the ^7Be ground state. If the single-particle energies of the last proton are determined from Eq. (3) as in Ref. [3], we obtain $\epsilon_p(0p_{3/2}) = \epsilon_p(0p_{1/2})$. This implies that the $L \cdot S$ splitting was ignored in Ref. [3].

(ix) The $\hbar\omega$ value adopted for the harmonic-oscillator wave functions in Ref. [3] is questionable. Although this value (and the corresponding b parameter) was important for the shell-model estimate of $Q(^8\text{B})$, the $\hbar\omega$ value was

fixed by the systematics among much heavier nuclei. We shall return to this point in Sec. IV B.

As has been mentioned at point (iii), the correct antisymmetrization among nucleons is important. It is not easy to carry out the full antisymmetrization among constituent nucleons and simultaneously reproduce the correct tail form of the ${}^7\text{Be}+p$ channel. In Secs. II and III we have been staying within the shell model, abandoning for the time being the reproduction of the tail behavior. The shell model describes the nuclear surface region pretty well, while in usual cases the tail region is beyond the scope of the model. The quadrupole moment under discussion is dominated by the surface region for normal nuclei. If we were not capable of reproducing the Q moment within the shell model, it would suggest the possibility of a halo.

B. Reconsideration of the b parameter in the $0\hbar\omega$ shell model

As mentioned at point (ix) in the previous subsection, the harmonic-oscillator wave functions in Ref. [3] would hardly reproduce the properties of the nuclei around $A = 8$. We shall reexamine this point.

The Cohen-Kurath interaction [7] has been frequently used for the $0\hbar\omega$ shell-model calculation in the p shell. The discussion in Refs. [2,3] was also based on this standard interaction. We now consider the shell-model wave function obtained from the Cohen-Kurath (8-16)TBME interaction. In order to fix the oscillator length b , we use the observed matter radius, in contrast to Ref. [3]. The rms matter radii of ${}^8\text{B}$ and ${}^8\text{Li}$ reported in Ref. [4] lead to $b \simeq 1.77$ fm, after the center-of-mass correction. This value of the b parameter differs significantly from the one adopted in Ref. [3], i.e., 1.60 fm.

The Q moments and $B(E2)$ values in $A = 6-10$ nuclei computed with $b \simeq 1.77$ fm in the $0\hbar\omega$ space with the Cohen-Kurath interaction are shown in Table I, together with the $(0+2)\hbar\omega$ results and the experimental data. This $0\hbar\omega$ calculation contains a single parameter $\delta e_{\text{IS}} = 0.5e$, which leads to $e_{\pi}^{\text{eff}} = 1.0e$. The agreement with the data seems to be rather good, indicating the validity of the present isoscalar effective charge. It is found that $Q({}^8\text{Li})$ is reproduced well, while $Q({}^8\text{B})$ is underestimated considerably.

If data of Q moments in a set of mirror nuclei are available, it is possible to determine the effective charges for those nuclei from the data, as was done in Ref. [3]. By applying this procedure to ${}^8\text{Li}$ and ${}^8\text{B}$, together with the present value $b = 1.77$ fm, we obtain $e_{\pi}^{\text{eff}} = 2.1e$ and $e_{\nu}^{\text{eff}} = 0.3e$. This set of effective charges is less surprising than those obtained from the smaller value of b in Ref. [3].

The isoscalar effective charge comes to $e_{\text{IS}}^{\text{eff}} = 1.2e$, whereas $e_{\text{IS}}^{\text{eff}} = 1.6e$ in Ref. [3] for the harmonic-oscillator wave functions. The analysis in Ref. [3] for the $A = 11$ and 17 nuclei led to $e_{\text{IS}}^{\text{eff}} = 0.9e$ and $e_{\text{IS}}^{\text{eff}} = 1.0e$, respectively. Compared with these values, $e_{\text{IS}}^{\text{eff}} = 1.2e$ for the $A = 8$ nuclei is not anomalous.

Through the above discussions we recognize that it is quite difficult to reach a conclusion when dealing with

such light nuclei, as far as we restrict ourselves to the $0\hbar\omega$ space, since both the effective charges and the single-particle wave functions give rise to ambiguities. The calculated Q moment is influenced by these ambiguities.

We further comment upon the analysis using the single-particle wave functions obtained from the ordinary Woods-Saxon potential. If the Woods-Saxon potential with the parameters in Ref. [18] is adopted, the data of $Q({}^8\text{Li})$ and $Q({}^8\text{B})$ will lead to $e_{\pi}^{\text{eff}} = 2.3e$ and $e_{\nu}^{\text{eff}} = 0.3e$. If we vary some parameters so as to fit the $\sqrt{\langle r^2 \rangle}$ data, quite similar effective charges are required to the harmonic-oscillator case giving the same $\sqrt{\langle r^2 \rangle}$.

C. Explanation of the difference between $Q({}^8\text{Li})$ and $Q({}^8\text{B})$

In this subsection we shall discuss why $Q({}^8\text{B})$ is substantially larger than $Q({}^8\text{Li})$, in an intuitive way within the $0\hbar\omega$ shell model. Suppose that Q moments are dominated by isoscalar degrees of freedom; equal values are obtained between mirror nuclei. The Q moments of ${}^{11}\text{B}$ and ${}^{11}\text{C}$ are so close, exhibiting an example of this kind. On the other hand, in experiment, $Q({}^8\text{B})$ is about twice larger than $Q({}^8\text{Li})$. Within the jj -coupling scheme without a residual interaction, the $0p_{3/2}$ orbit is partly occupied while the $0p_{1/2}$ is unoccupied, for either protons and neutrons. However, once we switch the residual interaction on, it is much easier in ${}^8\text{Li}$ for neutrons to excite to $0p_{1/2}$ than for protons, owing to the excess in number. The inspection of the density matrix and the single-particle matrix elements indicates that the excitation or deexcitation of a nucleon between $0p_{3/2}$ and $0p_{1/2}$ yields the principal contribution to the Q moment. It follows that $Q({}^8\text{Li})$ is dominated by the neutron degrees of freedom, unless e_{ν}^{eff} value is not too small. Similarly $Q({}^8\text{B})$ is governed by the proton degrees of freedom. Consequently, $Q({}^8\text{Li})$ and $Q({}^8\text{B})$ reflect the neutron and proton effective charges, respectively, giving rise to an enhanced sensitivity to the isovector charge. Note that this mechanism does not work for the $A = 11$ nuclei, primarily because of the smaller difference between proton and neutron number.

D. Sensitivity of $Q({}^8\text{B})$ to interaction

The shell-model interaction of Ref. [5] adopted for the present $(0+2)\hbar\omega$ calculation has several problems, as has been argued in Ref. [6]; the radial excitation is a primary problem in the argument. Furthermore, the present wave functions do not necessarily provide us with reasonable nuclear radii, as mentioned earlier. Although we do not discuss these points here, it should not be overlooked how sensitive the present result of $Q({}^8\text{B})$ is to the choice of interaction.

It is known that, since the quadrupole part dominates the proton-neutron correlation in low-lying states, the energy levels are sensitive to the $Q_{\pi} \cdot Q_{\nu}$ component of a two-body interaction, where

$$Q_\rho = \sum_{k \in \rho} r_k^2 Y^{(2)}(\hat{\mathbf{r}}_k) \quad (\rho = \pi, \nu), \quad (4)$$

with k being the label of each nucleon and $\hat{\mathbf{r}}_k = \mathbf{r}_k/r_k$. This $Q_\pi \cdot Q_\nu$ part plays a significant role, at the same time, in describing $E2$ properties. Hence the low-lying energy levels and the $E2$ properties are not independent of each other. Although effective charges may give rise to an ambiguity in calculating the $E2$ properties, this ambiguity will hardly influence when the effective-charge corrections (δe 's) are small. In such cases the $E2$ properties are dominated by Q_π .

The $E2$ properties of low-lying states depend predominantly on angular correlations of nucleons in the nuclear surface region. The low-lying spectrum is also dominated by the surface degrees of freedom, because the low-lying excited states are obtained usually by exciting one to a few nucleons around the surface from the ground state. The $Q_\pi \cdot Q_\nu$ component of the interaction carries an important part of this excitation. On the other hand, the radial (or monopole) excitation is much more dependent on the interior region (and excitations from this region) than the $E2$ properties and the low-lying spectra. In other words, this excitation is much of the volume character. Owing to this aspect, the low-lying spectra and $E2$ properties can be treated separately, to a certain extent, from the problem of the radial excitations.

We shall now move to the problem of the radius. This problem is more general and is not characteristic to the present scheme, as stated earlier. On the other hand, the change of the radius from a nucleus to another is treated independently of energies and $E2$ properties in most shell-model calculations. This is reasonable probably because energies and $E2$ properties are more sensitive to angular correlations in the surface region. As an example, the A dependence of the interaction and that of the single-particle wave function are not connected to each other in the successful shell-model calculation for the sd shell [19]. A more precise description of the radius seems to be a difficult task in a general term, and the goal is rather far. Note that the nuclear radius is also contributed by the interior region to a considerable extent. For a phenomenological interaction like the present one, the problems of radial modes (radius and radial excitation) may be separated from energy levels and $E2$ properties.

It has been shown in Ref. [5] that the low-lying energy levels around $A = 8$ are reproduced by the interaction of Wolters *et al.* This interaction should contain the $Q_\pi \cdot Q_\nu$ part in an effective manner. The resultant good agreement between calculated and experimental energy levels over several nuclei suggests that the effect of $Q_\pi \cdot Q_\nu$ is incorporated properly. The $E2$ properties are also reproduced in the low-lying states with only a small effective-charge correction, by the present $(0+2)\hbar\omega$ calculation. It therefore turns out that the present interaction is applicable to the investigation of $E2$ properties of the $A \sim 8$ nuclei, while there remain open issues in radii and radial excitations. This consequence is in accordance with the general discussion in the preceding paragraphs.

In the present case, the agreement in $Q(^8\text{Li})$ ensures

the amount of Q_π in ^8Li , which is exactly equal to the amount of Q_ν in ^8B . Since $Q_\pi \cdot Q_\nu$ is tested by the energy levels, it is expected that the present $(0+2)\hbar\omega$ calculation of the $Q(^8\text{B})$ value is plausible. Thus one can anticipate that the present result with respect to $Q(^8\text{B})$ will not be varied, for instance, by a future tuning of the interaction, as far as the interaction reproduces both the energy levels and other related $E2$ properties.

V. SUMMARY

As long as we try to describe the $E2$ properties of light nuclei within the $0\hbar\omega$ shell model, the ambiguities arising from single-particle wave functions and effective charges are inevitable. On the other hand, the Q moments and $B(E2)$ values of $A = 6-10$ nuclei, including $Q(^8\text{B})$, are reproduced within the present $(0+2)\hbar\omega$ shell model. It is noticed that the $(0+2)\hbar\omega$ approach is much less ambiguous in fixing the values of effective charges than the calculation restricted to the $0\hbar\omega$ space. This $(0+2)\hbar\omega$ calculation suggests that the proton halo in ^8B is very unlikely. The observed $Q(^8\text{B})$ value is not an evidence for a proton halo. This result is consistent with the data on the interaction radius [4]. Even though the proton density distribution in ^8B may be somewhat spread, its amplitude seems to be too small to be regarded as a proton halo. The expectation value of the proton number in the tail region will be much less than unity, although such a reduced tail may still produce interesting effects in astrophysical issues [20]. Though the present discussions have been based on the interaction of Wolters *et al.* [5] and this interaction has several problems [6], the conclusion on the $E2$ properties is not very sensitive to the choice of the interaction.

The effect of the $2\hbar\omega$ configuration on $E2$ is also discussed. The $2\hbar\omega$ contribution is customarily incorporated into the $0\hbar\omega$ calculation by effective charges. Indeed, the R_T values defined in Eq. (1) are rather stable for isoscalar matrix elements. However, those for isovector matrix elements are largely A dependent. This result suggests an intriguing aspect that, although the isoscalar effective charges for the $0\hbar\omega$ space could be almost constant, appreciable variation is required for the isovector effective charges, which play more important roles in nuclei far from stability with higher isospin.

Note added in proof. After completion of this manuscript, we learned that a related work has been carried out by Csoto in terms of three-cluster calculations [A. Csoto, Phys. Lett. B **315**, 24 (1993)]. A proton skin is suggested in ^8B , which is not excluded by the present work.

ACKNOWLEDGMENTS

The authors are grateful to Prof. T. Minamisono, Prof. H. Sagawa, and Dr. D. J. Millener for valuable discussions. The authors appreciate the hospitality of the Institute for Nuclear Theory at the University of Washington, and thank Prof. A. Gelberg for careful reading of the manuscript. This work was supported partly by Research Center for Nuclear Physics, Osaka University, as RCNP Computational Nuclear Physics Project (Project No. 92-B-03). This work was supported in part by Grant-in-

Aid for General Scientific Research (No. 04804012), by the Ministry of Education, Science and Culture. The code OXBASH in the program library of Computer Center, University of Tokyo is used. It is acknowledged that VAX6440, in Meson Science Laboratory, University of Tokyo, is also utilized.

APPENDIX

In this appendix we review how the tail form of a bound-state wave function is connected with separation energies. The Schrödinger equation for the nucleus A is

$$H_A \psi(A) = E(A) \psi(A). \quad (\text{A1})$$

For the sake of simplicity, we shall deal only with the ground state of A . It is straightforward to extend this discussion to excited states. If we consider the breakup channel $A \rightarrow A' + a$, where a represents a single nucleon or a cluster of nucleons, it is convenient to rewrite it as

$$\psi(A) = \mathcal{N} \sum_{i,j} \mathcal{A}[\varphi_{A',a}(\{\mathbf{R}_\alpha\}) \psi(A'; i) \psi(a; j)], \quad (\text{A2})$$

$$H_A = H_{A'} + H_a + T_R + V_{A',a}(\{\mathbf{R}_\alpha\}). \quad (\text{A3})$$

Here $\varphi_{A',a}(\{\mathbf{R}_\alpha\})$ denotes the wave function with respect to the relative coordinates between a nucleon involved in A' and another nucleon in a . The expression \mathcal{A} stands for the antisymmetrizer for all nucleons and \mathcal{N} an appropriate normalization constant. The indices i and j represent various configurations of the partition A' and a . The operator T_R in the Hamiltonian represents the kinetic energy regarding the variable \mathbf{R} , where \mathbf{R} is the relative coordinate between the center of mass of A' and that of a , and $V_{A',a}(\{\mathbf{R}_\alpha\})$ the interaction between A' and a .

For sufficiently large R in the breakup channel, we can take $\psi(A'; i)$ and $\psi(a; j)$ as eigenstates of $H_{A'}$ and H_a , respectively. Then $\varphi_{A',a}(\{\mathbf{R}_\alpha\})$ becomes a function only of \mathbf{R} and \mathcal{A} can be regarded as identity. This will be satisfied when R exceeds the range R_N outside which the nuclear force between A' and a vanishes. The potential term V consists of the nuclear part V^N and the Coulomb part V^C . Because of the finite-range nature of the nuclear force, V is described only by V^C for $R > R_N$. Hereafter we restrict ourselves to $R > R_N$, abbreviating $\varphi_{A',a}(\{\mathbf{R}_\alpha\})$ as $\varphi(\mathbf{R})$.

It is sufficient to consider only the ground states of A' and a for discussing the asymptotic behavior of $\psi(A)$. Substituting Eqs. (A2) and (A3) into Eq. (A1) and integrating out the internal variables of A' and a , we obtain

$$[T_R + V^C] \varphi(\mathbf{R}) = [E(A) - E(A') - E(a)] \varphi(\mathbf{R})$$

$$\text{for } R > R_N. \quad (\text{A4})$$

The separation energy of A for the particle a is defined by

$$S_a(A) = -[E(A) - E(A') - E(a)]. \quad (\text{A5})$$

Therefore we can rewrite Eq. (A4) as

$$\left[-\frac{\nabla_R^2}{2\mu} + V^C \right] \varphi(\mathbf{R}) = -S_a \varphi(\mathbf{R}) \quad \text{for } R > R_N, \quad (\text{A6})$$

where the reduced mass for the $A' + a$ system is denoted by μ . Remark that we are considering a bound state of A (i.e., $S_a > 0$), leading to the boundary condition of $\varphi(\mathbf{R}) \rightarrow 0$ for $R \rightarrow \infty$. The strongest damping factor in the asymptotic region is determined from the equation [by neglecting terms of $O(R^{-1})$]

$$-\frac{1}{2\mu} \frac{\partial^2}{\partial R^2} \varphi(\mathbf{R}) \sim -S_a \varphi(\mathbf{R}) \quad \text{for large } R. \quad (\text{A7})$$

Therefore, for extremely large R , the wave function $\varphi(\mathbf{R})$ necessarily has the damping form $f(R)e^{-\sqrt{2\mu S_a}R}$, where $f(R)$ represents a function without an exponential damping tail. Indeed, suppose that a is a neutron, for which $V^C = 0$; Eq. (A6) provides us with the solution

$$\varphi_{lm}(\mathbf{R}) \simeq \xi' h_l^{(1)}(i\sqrt{2\mu S_a}R) Y_m^{(l)}(\hat{\mathbf{R}}) \quad \text{for } R > R_N, \quad (\text{A8})$$

where $h_l^{(1)}(x)$ expresses the spherical Hankel function, $\hat{\mathbf{R}}$ indicates \mathbf{R}/R , and ξ' denotes an amplitude. This immediately leads to the asymptotic form of $\varphi(\mathbf{R})$ as

$$\varphi_{lm}(\mathbf{R}) \simeq \xi \frac{e^{-\sqrt{2\mu S_a}R}}{R} Y_m^{(l)}(\hat{\mathbf{R}})$$

$$\text{for } R > R_N \text{ and } R \gg 1/\sqrt{2\mu S_a}. \quad (\text{A9})$$

The asymptotic forms of other channels are obtained in a similar manner. It should be kept in mind that the value of the amplitude ξ in Eq. (A9) cannot be determined within this asymptotic treatment. It is pointed out that the channel with the smallest separation energy yields the farthest reaching tail of the total wave function of A . Thus the wave function of a bound ground state has a definite tail form given by the corresponding separation energy.

- [1] I. Tanihata, Nucl Phys. **A522**, 275c (1991).
- [2] T. Minamisono *et al.*, Phys. Rev. Lett. **69**, 2058 (1992).
- [3] H. Kitagawa and H. Sagawa, Phys. Lett. B **299**, 1 (1993).
- [4] I. Tanihata, T. Kobayashi, O. Yamakawa, S. Shimoura, K. Ekuni, K. Sugimoto, N. Takahashi, T. Shimoda, and H. Sato, Phys. Lett. B **206**, 592 (1988).
- [5] A. A. Wolters, A. G. M. van Hees, and P. W. M. Glaudemans, Phys. Rev. C **42**, 2053, 2062 (1990).
- [6] D. J. Millener, A. C. Haynes, and D. Strottman, Phys.

- Rev. C **45**, 473 (1992); A. A. Wolters, A. G. M. van Hees, and P. W. M. Glaudemans, *ibid.* **45**, 477 (1992).
- [7] S. Cohen and D. Kurath, Nucl. Phys. **73**, 1 (1965).
- [8] F. Ajzenberg-Selove, Nucl. Phys. **A413**, 1 (1984); **A433**, 1 (1985); **A449**, 1 (1986); **A460**, 1 (1986).
- [9] P. Raghavan, At. Data Nucl. Data Tables **42**, 189 (1989).
- [10] F. Ajzenberg-Selove, Nucl. Phys. **A490**, 1 (1988).
- [11] J. J. Kolata, in *Proceedings of International Symposium on Structure and Reactions of Unstable Nuclei*, Niigata,

- 1991, edited by K. Ikeda and Y. Suzuki (World Scientific, Singapore, 1991), p. 252.
- [12] P. Descouvemont and D. Baye, Phys. Lett. B **292**, 235 (1992).
- [13] H. de Vries, C. W. de Jager, and C. de Vries, At. Data Nucl. Data Tables **36**, 495 (1987).
- [14] A. G. M. van Hees and P. W. M. Glaudemans, Z. Phys. A **314**, 323 (1983); **315**, 223 (1984).
- [15] J. M. G. Gómez, J. C. Pérez Cerdán, and C. Prieto, Nucl. Phys. **A551**, 451 (1993).
- [16] J. M. G. Gómez, C. Prieto, and A. Poves, Phys. Lett. B **295**, 1 (1992).
- [17] D. J. Millener, J. W. Olness, and E. K. Warburton, Phys. Rev. C **28**, 497 (1983).
- [18] A. Bohr and B. R. Mottelson, *Nuclear Structure* (Benjamin, New York, 1969), Vol. 1, p. 239.
- [19] B. A. Brown and B. H. Wildenthal, Annu. Rev. Nucl. Part. Sci. **38**, 29 (1988).
- [20] K. Riisager and A. S. Jensen, Phys. Lett. B **301**, 6 (1993).