

Superdeformation in the bismuth nuclei

R. M. Clark,¹ S. Bouneau,² F. Azaiez,² S. Asztalos,¹ J. A. Becker,³ B. Cederwall,¹ M. A. Deleplanque,¹
 R. M. Diamond,¹ J. Duprat,² P. Fallon,¹ L. P. Farris,³ E. A. Henry,³ J. R. Hughes,³ W. H. Kelly,⁴
 I. Y. Lee,¹ A. O. Macchiavelli,¹ M. G. Porquet,⁵ J. F. Sharpey-Schafer,⁶ F. S. Stephens,¹
 M. A. Stoyer,¹ D. T. Vo,³ and A. N. Wilson⁶

¹Lawrence Berkeley Laboratory, 1 Cyclotron Road, Berkeley, California 94720

²Institut de Physique Nucléaire, Institut National de Physique Nucléaire et de Physique des Particules—CNRS, Batiment 104,
 F-91406 Orsay Cedex, France

³Lawrence Livermore National Laboratory, Livermore, California 94550

⁴Iowa State University, Ames, Iowa 50011

⁵CSNSM, Institut National de Physique Nucléaire et de Physique des Particules—CNRS, Batiment 104-108,
 F-91405 Orsay Cedex, France

⁶Oliver Lodge Laboratory, University of Liverpool, Liverpool L69 3BX, United Kingdom

(Received 17 November 1994)

High angular-momentum states in ^{196,197}Bi were populated in the reaction ¹⁸³W(¹⁹F,xn) at a beam energy of 108 MeV, and γ rays were detected with the Gammasphere array. Two weakly populated rotational bands, with energy spacings characteristic of superdeformation have been found. Both cascades can be assigned unambiguously to the Bi nuclei; however, their isotopic assignment to ¹⁹⁷Bi is tentative. The properties of the bands and their possible structures are discussed. Our results represent the first identification of superdeformed bands in a nucleus of the $A \sim 190$ mass region with $Z > 82$.

PACS number(s): 21.10.Re, 23.20.Lv, 27.80.+w

An impressive experimental and theoretical effort has been devoted to exploring the underlying physics of superdeformation in the $A \sim 190$ region. Over 40 superdeformed (SD) bands have been identified in the Au, Hg, Tl, and Pb nuclei [1–3]. The most readily measured experimental quantities for these sequences are the transition energies and intensities. In particular, transition energies, E_γ , can be used to deduce the dynamic moment of inertia, $\mathcal{A}^{(2)}$, and the behavior of this quantity can then be compared directly with theory to give information on the underlying microscopic structure of the bands. The $\mathcal{A}^{(2)}$'s for most of the bands in the $A \sim 190$ region show a steady rise with increasing rotational frequency, ω . The quadrupole moments of states in several bands have been established through lifetime measurements [4–9]. These results confirm the superdeformed nature of the structures and also show that the deformation remains constant over all the states in the measurement range. This suggests that the rise in $\mathcal{A}^{(2)}$ is not due to centrifugal stretching which would lead to a change in deformation over the band. Instead, the behavior of $\mathcal{A}^{(2)}$ has been associated with the gradual alignment of pairs of nucleons occupying specific high- N intruder orbitals (namely, $j_{15/2}$ neutrons and $i_{13/2}$ protons) in the presence of pair correlations [10]. However, mean field calculations with a monopole pair field have not been able to reproduce accurately the absolute magnitudes and slopes of the $\mathcal{A}^{(2)}$'s [11].

Until the present work, no SD bands had been found in nuclei of the $A \sim 190$ region with $Z > 82$, despite many theoretical calculations (see e.g., [12,13] and references therein) predicting well-defined secondary SD minima persisting in the bismuth nuclei. Investigating superdeformation in these

isotopes should provide the first experimental evidence concerning the nature of the proton orbitals above $Z = 82$ at large deformation ($\beta_2 \approx 0.48$). In this work we describe results from a recent experiment aimed at exploring superdeformation in ^{196,197}Bi. High-spin states in these nuclei were populated via the ¹⁸³W(¹⁹F,xn) reaction at an incident beam energy of 108 MeV. The beam, which was provided by the 88-Inch Cyclotron facility at the Lawrence Berkeley Laboratory, was incident on a target consisting of $2 \times 300 \mu\text{g cm}^{-2}$ stacked ¹⁸³W foils mounted on thin carbon backings. γ rays were detected with the Gammasphere array [14]

TABLE I. Transition energies (in keV) for the two SD bands in ¹⁹⁷Bi. Also given, in the third column, are the transition energies of band 1 in ¹⁹⁵Tl. For comparison, this latter set of energies are shown only over the observed frequency range of band 2.

Band 1	Band 2	¹⁹⁵ Tl (Band 1)
166.2(2)		
208.0(2)	186.7(2)	188.0(4)
249.7(2)	229.1(2)	228.8(3)
291.3(2)	269.9(2)	270.0(3)
332.6(2)	310.0(2)	310.5(3)
373.8(2)	351.1(2)	349.9(3)
414.3(2)	390.7(2)	389.8(3)
455.0(2)	430.8(2)	428.9(3)
495.2(2)	468.5(2)	467.2(3)
535.4(2)	507.1(4)	504.5(3)
574.3(2)	545(1)	540.8(3)
614.3(4)		
653(1)		




FIG. 1. (a) A sum of triple-gated spectra from fourfold coincidence data for band 1. The inset in the top right shows the relative intensities of transitions in the band, while the inset in the top left shows the coincident Bi x rays. (b) A sum of double-gated spectra from threefold coincidence data for band 2. The inset in the top right shows the relative intensities of transitions in the band, while the inset in the top left shows the coincident Bi x rays.

which, for this experiment, comprised 36 Compton-suppressed large-volume (~ 75 – 80% efficient) HPGe detectors. A total of 8×10^8 three- and higher-fold events were collected.

Two sequences of γ rays, with properties characteristic of SD bands in this mass region, have been observed. Spectra showing the two bands are presented in Fig. 1, while the transition energies are summarized in Table I. The insets of Fig. 1 show the relative intensities of the in-band transitions. These show behavior common to all the SD bands in the $A \sim 190$ region. Feeding occurs over the top six transitions or so, followed by a region in which the in-band states are fully fed and the intensity is constant. A rapid depopulation of the band occurs from the lowest one or two levels.

The dominant open channels in the reaction are predicted, by statistical models, to be the $5n$ (^{197}Bi) and $6n$ (^{196}Bi) channels, taking approximately 54% and 28% of the total cross section, respectively. The only other channel populated with significant intensity is predicted to be the $\alpha 4n$ (^{194}Tl)

channel taking approximately 7% of the reaction cross section. None of the known SD bands in ^{194}Tl [15] were observed in our data, implying that the two new bands are in the Bi nuclei. This is established from the coincidence of the in-band transitions with Bi x-ray lines (see Fig. 1). The decay schemes of $^{196,197}\text{Bi}$ are not well known. In addition, the presence of long-lived isomers at low spin [16,17] prevented the observation of any low-lying transitions in prompt coincidence with either of the bands. We are therefore unable to establish the mass assignments unambiguously. However, since the reaction was optimized to populate SD states in ^{197}Bi we favor this assignment for both structures. We estimate the upper limit of the total intensity of band 1 to be $\sim 2\%$ of the total $5n$ reaction channel, while band 2 has $\sim 50\%$ the intensity of band 1.

Figure 2(a) shows plots of the $\mathcal{J}^{(2)}$ moments of inertia for the two bands and also for the known SD bands in the isotones ^{196}Pb [6,18,19] and ^{195}Tl [20,21]. At rotational frequencies above $\omega \approx 0.2 \text{ MeV } \hbar^{-1}$ the $\mathcal{J}^{(2)}$ for band 1 has a




FIG. 2. (a) Plots of the dynamic moments of inertia, $\mathcal{J}^{(2)}$, as a function of rotational frequency for band 1 (open circles), band 2 (closed circles), the SD band in ^{196}Pb (closed squares), and SD bands 1 and 2 in ^{195}Tl (open and closed triangles, respectively). (b) Plots showing the difference in transition energies between: (i) the half-points of band 1 and band 2 (open circles), and (ii) band 2 and band 1 in ^{195}Tl (closed circles).

reduced slope when compared with that of any of the other bands. The $\mathcal{J}^{(2)}$ of band 2 is very similar to that of band 1 in ^{195}Tl . Indeed, on closer inspection, band 2 in ^{197}Bi and band 1 in ^{195}Tl have “identical” transition energies (to within ± 2 keV) over a sizeable spin range. This can be seen from Table I, which also summarizes the energies for band 1 in ^{195}Tl , and from Fig. 2(b) which shows the difference in transition energies between these two bands.

Estimates of the spins of states in the two bands were made using the method described in [22,23]. The 166 keV γ ray of band 1 decays to a level with a fitted spin of $6.4(1)\hbar$ ($I=13/2\hbar$, to the closest half-integer value), while the 187 keV γ ray of band 2 decays to a level with a fitted spin of $7.5(1)\hbar$ ($I=15/2\hbar$). Since the fitting procedure yields spins that are very nearly half-integer, the assignment of the bands to ^{197}Bi gains further support. Note also that the spins of the levels in the two bands differ by $1\hbar$. This may indicate that the two bands are signature partners. From the energies of the two bands (Table I) it can be seen that band 2 starts approximately at the half points of band 1. However, following the band up in spin the deviation of band 2 from the half points of band 1 steadily increases. This is illustrated in Fig. 2(b). This indicates that if the bands are signature partners they show some signature splitting. No evidence of “cross-talk” [24–26,21] between the two bands could be found.

The detailed spectroscopic behavior of a band in an odd-Bi nucleus is principally determined by which state contains the odd proton relative to the neighboring even-Pb core (^{196}Pb in the case of ^{197}Bi). Figure 3(a) presents a single-particle Woods-Saxon calculation for protons with deformation parameters $\beta_2=0.48$, $\beta_4=0.07$, and $\gamma=0^\circ$ (a representative deformation for SD bands in the $A \sim 190$ region; the values come from the calculations described in [13]). The calculation indicates that the $[514]9/2$, $[651]1/2$, and $[642]5/2$ orbitals all lie close to the Fermi surface. As discussed above, pairing plays an important role in determining the behavior of SD bands in the $A \sim 190$ region. Presented in Fig. 3(b) is a quasiproton routhian diagram calculated for the parameters $\beta_2=0.48$, $\beta_4=0.07$, $\gamma=0^\circ$, and $\Delta_p=\Delta_{BCS}(\omega=0)$. The inclusion of pairing smears the Fermi surface and the calculation indicates that one quasiproton excitations involving the $[642]5/2$, $[514]9/2$, and $[651]1/2$ levels are all energetically possible. Bands based on these states should be easy to distinguish experimentally.

The lowest calculated one quasiparticle excitations are to the $[642]5/2$ ($i_{13/2}$) signature partner levels. There is a small splitting between the two signature partners which increases with increasing rotational frequency. The inset of Fig. 3(b) shows experimental routhians for the two new bands. Over the experimentally observed range the two new bands do display a splitting in the transition energies [see also Fig. 2(b)]. The $[642]5/2$ orbital is thought to be involved in the configurations of SD bands in the Tl nuclei [15,20,27,28]. Therefore, one might reasonably expect that the $\mathcal{J}^{(2)}$'s of the ^{197}Bi bands should be similar to those of the bands in ^{195}Tl (an isotone). Indeed, as already noted band 2 in ^{197}Bi and band 1 in ^{195}Tl have “identical” transition energies over a significant range of spin. However, band 1 in ^{197}Bi has a much flatter $\mathcal{J}^{(2)}$ above $\omega \approx 0.2$ MeV \hbar^{-1} than those of the ^{195}Tl bands, which is not easy to explain. There are further problems with this interpretation. The signature partner pair of bands in ^{195}Tl [20,21] which are based on the $[642]5/2$ orbital are populated with similar intensities. In addition, dipole cross talk between these bands in ^{195}Tl has been observed [21]. Neither of these features are seen for our new bands.

The $[514]9/2$ state should give rise to a strongly coupled structure with $K=9/2$ (two signature partners with very little signature splitting). The strong coupling model predicts a large $B(M1)$ strength ($\sim 1.9\mu_N^2$) [24] which should lead to the observation of strong dipole “cross-talk” transitions between the two signature partner bands. Furthermore, the two signature partners should have $\mathcal{J}^{(2)}$ moments of inertia similar to that of the SD band in ^{196}Pb [6,18,19] since the intruder occupation is the same ($\nu 7^4 \pi 6^6$). Clearly, none of these features are displayed by the two bands we observe.

The $[651]1/2$ ($i_{11/2}$) orbital is also an energetically possible one-quasiparticle excitation. This orbital is expected to exhibit immediate signature splitting with the $\alpha = -1/2$ partner being favored. The splitting is predicted by the calculation to increase rapidly with increasing rotational frequency. The experimental routhian in Fig. 3(b) indicates that the actual splitting of the two bands is much less and starts later. It is possible that the two new bands are not signature partners. If both the bands are in ^{197}Bi and are not signature partners then it is necessary to invoke a missing band, since at least




FIG. 3. (a) Cranked Woods-Saxon single-particle diagram for protons. The deformation parameters used were: $\beta_2=0.48$, $\beta_4=0.07$, and $\gamma=0.0^\circ$. Parity and signature (π, α) of the levels are indicated in the following way: solid $=(+, +1/2)$, dotted $=(+, -1/2)$, dot-dashed $=(-, +1/2)$, and dashed $=(-, -1/2)$. (b) Cranked Woods-Saxon quasiparticle diagram for protons. The deformation parameters used were the same as those for (a), while the pairing parameter was taken as $\Delta_p = \Delta_{BCS}(\omega=0)$. Parity and signature are indicated in the same manner as 3(a). The inset shows the experimental routhians for the two new bands. A rigid rotor reference with the Harris parameter $J_0 = 94.0\hbar^2 \text{ MeV}^{-1}$ was subtracted. The absolute energy scale is arbitrary and the relative excitation energies of the bands in the diagram assumes that they are signature partners (degenerate at zero rotational frequency).

one configuration must then involve a strongly coupled orbital (either $[642]5/2$ or $[514]9/2$). Another possibility is that band 1 is in ^{197}Bi and band 2 is in ^{196}Bi .

One further comment should be made concerning the behavior of the $\mathcal{J}^{(2)}$ moments of inertia. The $\mathcal{J}^{(2)}$'s for the two new bands have much lower slopes than that for the band in ^{196}Pb [see Fig. 2(a)]. The odd proton could block the $i_{13/2}$ quasiproton alignments, flattening the $\mathcal{J}^{(2)}$'s relative to that of ^{196}Pb . However, the $j_{15/2}$ quasineutron alignments should still contribute to the rise in $\mathcal{J}^{(2)}$. If the bands were in ^{196}Bi (an odd-odd nucleus) then the $j_{15/2}$ quasineutron alignments may also be blocked giving a much flatter $\mathcal{J}^{(2)}$. It should be noted that several SD bands in $^{193,195}\text{Pb}$ [29,30] have also been found which have similar, exceptionally flat, $\mathcal{J}^{(2)}$'s. These bands are thought to have the $j_{15/2}$ quasineutron alignments blocked by an odd neutron. This behavior of the $\mathcal{J}^{(2)}$'s must be addressed, both experimentally and theoretically, in future studies.

To summarize, an experiment with the Gammasphere array using the $^{183}\text{W}(^{19}\text{F}, xn)$ reaction at a beam energy of 108

MeV has resulted in the observation of two new SD bands. The new bands can be unambiguously assigned to the Bi nuclei, but the isotopic assignments to ^{197}Bi are tentative. Our results represent the first observation of superdeformed bands in the $A \sim 190$ region with $Z > 82$. A number of important experimental questions remain. First, the isotopic and configuration assignments of the bands must be unambiguously established. Second, the superdeformed nature of these structures needs to be confirmed through lifetime measurements. Third, SD sequences in other Bi isotopes must be found in order to probe further the nature of the orbitals close to the proton Fermi surface at extreme deformation.

We would like to thank Dr. R. Wadsworth for his help in organizing the logistics of getting targets made, and to Bob Darlington of the Daresbury Laboratory for making the target foils used in the experiment. The crew and staff of the 88-Inch Cyclotron are thanked. This work has been supported in part by the U.S. Department of Energy under Contract Nos. DE-AC03-76SF00098 (LBL) and W-7405-ENG-48 (LLNL).

- [1] R. B. Firestone and B. Singh, Table of Superdeformed Nuclear Bands (unpublished).
- [2] J. F. Sharpey-Schafer, *Prog. Part. Nucl. Phys.* **28**, 187 (1992).
- [3] R. V. F. Janssens and T. L. Khoo, *Annu. Rev. Nucl. Part. Sci.* **41**, 321 (1991).
- [4] M. W. Drigert *et al.*, *Nucl. Phys.* **A530**, 452 (1991).
- [5] E. F. Moore *et al.*, *Phys. Rev. Lett.* **64**, 3127 (1990).
- [6] E. F. Moore *et al.*, *Phys. Rev. C* **48**, 2261 (1993).
- [7] M. Carpenter *et al.*, *Phys. Lett. B* **240**, 44 (1990).
- [8] P. Willsau *et al.*, *Z. Phys. A* **344**, 351 (1993).
- [9] J. R. Hughes *et al.*, *Phys. Rev. Lett.* **72**, 824 (1994).
- [10] M. A. Riley *et al.*, *Nucl. Phys.* **A512**, 178 (1990).
- [11] R. Wyss, Contribution to the Conference on Physics from Large Gamma-ray Detector Arrays, Berkeley, 1994 (unpublished).
- [12] R. R. Chasman *et al.*, *Phys. Rev. Lett. B* **219**, 308 (1989).
- [13] W. Satula *et al.*, *Nucl. Phys.* **A529**, 289 (1991).
- [14] GAMMASPHERE, A National Gamma-ray Facility, edited by M. A. Deleplanque and R. M. Diamond [Lawrence Berkeley Laboratory Report No. PUB-5202, 1988 (unpublished)].
- [15] F. Azaiez *et al.*, *Phys. Rev. Lett.* **66**, 1030 (1991).
- [16] A. H. Wapstra and K. Bos, *At. Data Nucl. Data Tables* **20**, 1 (1977).
- [17] T. Chapuran *et al.*, *Phys. Rev. C* **33**, 130 (1986).
- [18] M. J. Brinkman *et al.*, *Z. Phys. A* **336**, 115 (1990).
- [19] T. F. Wang *et al.*, *Phys. Rev. C* **43**, R2465 (1991).
- [20] F. Azaiez *et al.*, *Z. Phys. A* **338**, 471 (1991).
- [21] J. Duprat *et al.* (unpublished).
- [22] J. A. Becker *et al.*, *Phys. Rev. C* **41**, R9 (1990).
- [23] J. E. Draper *et al.*, *Phys. Rev. C* **42**, R1791 (1991).
- [24] P. B. Semmes *et al.*, *Phys. Rev. Lett.* **68**, 460 (1992).
- [25] M. J. Joyce *et al.*, *Phys. Rev. Lett.* **71**, 2176 (1993).
- [26] P. Fallon *et al.*, *Phys. Rev. Lett.* **70**, 2690 (1993).
- [27] Y. Liang *et al.*, *Phys. Rev. C* **46**, R2136 (1992).
- [28] P. B. Fernandez *et al.*, *Nucl. Phys.* **A517**, 386 (1990).
- [29] J. R. Hughes *et al.* (unpublished).
- [30] L. Farris *et al.* (unpublished).