

Model of short-range correlations in the charge response

J. E. Amaro and A. M. Lallena

Departamento de Física Moderna, Universidad de Granada, E-18071 Granada, Spain

G. Co'

Dipartimento di Fisica, Università di Lecce, and I.N.F.N. sezione di Lecce, I-73100 Lecce, Italy

A. Fabrocini

Dipartimento di Fisica, Università di Pisa, and I.N.F.N. sezione di Pisa, I-73100 Pisa, Italy

(Received 19 December 1997)

The validity of a model treating the short-range correlations up to the first order is studied by calculating the charge response of an infinite system and comparing the obtained results with those of a Fermi Hypernetted Chain calculation. [S0556-2813(98)01106-6]

PACS number(s): 21.60.-n, 21.30.Fe, 24.10.Cn, 25.30.Fj

The interest in the study of short range correlations (SRC) in nuclear systems has increased in these last few years. In this field, the experimental activity has been concentrated in the search for observables allowing a clean identification of SRC effects [1,2]. From the theoretical point of view, there has been a development of calculations which explicitly consider SRC.

The theoretical situation is quite satisfactory for the few-body systems, where Faddeev [3], correlated hyperspherical harmonics expansion [4], and Green function Monte Carlo [5] theories solve exactly the Schrödinger equation. This last technique has been recently applied to investigate light nuclei up to $A = 7$ [6]. Unfortunately, the straightforward application of these theories to the study of medium and heavy nuclei is not yet technically feasible, in spite of the rapid progress of the computer technology.

The other satisfactory situation, from the theoretical point of view, regards the opposite side of the isotope table: the infinite nuclear systems such as neutron and nuclear matter. For the study of these systems, perturbation techniques have been developed such as Brueckner Bethe Goldstone [7] or correlated basis function theories (CBF) [8]. These approaches do not provide an exact solution of the Schrödinger equation, but they can reproduce rather well the empirical properties of nuclear matter because they sum complete sets of terms of their perturbation expansions.

The application of CBF to the description of the ground state of finite nuclear systems, has been recently carried out [9] using various levels of Fermi hypernetted chain (FHNC) approximations. The results are promising and it is conceivable that CBF theories may be applied to the description of excited states in the future. For the time being, these properties have to be studied by using simpler models.

The major part of the models developed up to now to describe finite nuclear systems, treats SCR only at the lowest order in the correlation. The main field of application has been the investigation of nuclear ground state properties [10]. Recently, nuclear models dealing with SRC, have been implemented to study the electromagnetic two-nucleon emission [11,12]. Given the wide use of these models and their

relevance for predictions and comparisons with two-nucleon emission data, we believe that a test of their validity is necessary.

A model treating SRC up to the first order in the correlation, has been developed in Ref. [13] to study charge density and momentum distribution of some doubly closed shell nuclei. The model was able to reproduce rather well the finite nuclei FHNC results of Ref. [9] and it has been extended to describe the two-nucleon emission produced by the same electromagnetic operator, the charge operator, for inclusive electron scattering experiments [14]. It is, however, not obvious that the good agreement with the FHNC results obtained for the ground state can be maintained also for the excited states. Since complete FHNC calculations are available for one-particle one-hole ($1p-1h$) nuclear matter charge responses [15], we have applied our model to calculate these responses.

The basic idea of the model, already presented in [13] and [14], consists in truncating the CBF expansion in order to consider only those terms containing a single Jastrow-type correlation line, $h(r) = f^2(r) - 1$. In Fig. 1 we show the diagrams we have retained in the present calculation. It is worth pointing out that by setting in each diagram the particle line equal to the hole line, we reproduce the set of diagrams used to calculate the ground state properties of the charge operator [13]. An important property of the expansion in powers of $h(r)$ is that the normalizations of the wave function are exactly preserved at each order [16]. In Ref. [13], the nuclear charge was conserved as well as the proper normalization of the correlated many-body wave functions in the present calculation. On the contrary, the nuclear charge is not conserved in Refs. [11,12], where the expansion adopted, and truncated at the second order, is based on *the number of particles* of the cluster and not on the powers of $h(r)$.

The FHNC and the model calculations of the charge responses have been done using the same correlation, i.e., the scalar part of a complicated state dependent correlation fixed to minimize the nuclear binding energy in a FHNC calculation with the Urbana V14 nucleon-nucleon potential [8].

In Fig. 2 we compare the results of our model with those obtained with a FHNC calculations. In this figure we show




FIG. 1. Diagrams considered in our model. The dotted lines represent the correlation function. The oriented lines represent particle and hole wave functions. The black circle indicates an integration point, while the black square indicates the integration point, where the charge operator is acting.

the proton structure functions, therefore, no electromagnetic nucleon form factors have been included. The structure functions have been calculated for three values of the momentum transfer, and for Fermi momentum of 1.09 fm^{-1} . We found this value of the Fermi momentum adequate to describe the quasielastic responses of ^{12}C [17]. In the panel (a) of Fig. 2 the full lines represent the results of the model and the dashed lines those of the FHNC calculation. The difference between the two calculations are very small, and they are explicitly shown, multiplied by a factor 10^5 , in the panel (b).

In Fig. 3 we show the response functions calculated using the electromagnetic nucleon form factors of Ref. [18]. These response functions have been obtained considering, in addition to the proton structure functions shown in Fig. 2, also the neutron contribution, which in any case, turns out to be negligible in the longitudinal response. The dashed lines show the Fermi gas responses, corresponding in our model to the first diagram of Fig. 1. The dashed-dotted lines have been obtained adding the contribution of the two-point diagrams, i.e., the diagrams multiplied by the factor $1/2$ in Fig. 1. The full lines have been obtained by including all the diagrams of Fig. 1. The contribution of the two-point diagrams is partially canceled by the inclusion of the three-point diagrams. This is an effect similar to that obtained in the calculation of the ground state charge and momentum distributions [13].

The results we have presented show that a model considering only those terms with a single correlation line can reproduce extremely well the FHNC charge response functions. This conclusion is consistent with the finding of Ref. [13] for the ground state charge distribution. We should remark that our calculations have been done for the charge operator only. An extension of the calculation to evaluate responses induced by other electromagnetic operators is nec-




FIG. 2. In the panel (a) the $1p-1h$ nuclear matter proton structure functions calculated with the present model (full lines) are compared with those obtained from the FHNC calculation of Ref. [15]. The calculations have been done for $q = 300, 400,$ and $550 \text{ MeV}/c$ and $k_F = 1.09 \text{ fm}^{-1}$. In the panel (b) we show the differences, multiplied by 10^5 , between the structure functions obtained with the present model and those obtained using FHNC.

essary to test the validity and the range of applicability of the model.

We like to stress again that a good agreement with the FHNC results has been obtained, most probably, because, our model properly normalizes the many-body wave function has been conserved by evaluating both two- and three-point




FIG. 3. Nuclear matter longitudinal responses for $q = 300, 400,$ and $550 \text{ MeV}/c$ and $k_F = 1.09 \text{ fm}^{-1}$ calculated with the proton and neutron form factors of Ref. [18]. The dashed lines represent the Fermi gas responses, the dashed-dotted lines have been obtained adding the two-point diagrams, while the full lines show the results of the complete calculations, where all the diagrams of Fig. 1 have been considered.

diagrams. Calculations of $1p-1h$ responses which include the two-point diagrams only, overestimate the effect of the correlations. One may expect that the same problem could affect also the two-nucleon emission calculations, like those of Refs. [11,12], where only two-point diagrams are consid-

ered. There are, however, indications that in the $2p-2h$ responses, two- and three-point diagrams act differently than in the $1p-1h$ responses [14]. A similar analysis, as the one performed here, is needed for the $2p-2h$ responses in order to clarify the situation.

-
- [1] In *Proceedings of the Second Workshop on Two-Nucleon Emission Reactions*, Marciana Marina, Elba (Italy), 1989, edited by O. Benhar and A. Fabrocini (ETS, Pisa, 1990).
- [2] In *Proceedings of the Second Workshop on Electromagnetically Induced Two-Nucleon Emission*, Gent (Belgium), 1995, edited by J. Ryckebusch and M. Waroquier (University of Gent, 1995).
- [3] C. R. Chen, G. L. Payne, J. L. Friar, and B. F. Gibson, *Phys. Rev. C* **33**, 1740 (1986); A. Stadler, W. Glöckle, and P. U. Sauer, *ibid.* **44**, 2319 (1991).
- [4] A. Kievsky, M. Viviani, and S. Rosati, *Nucl. Phys.* **A551**, 241 (1993).
- [5] J. Carlson, *Phys. Rev. C* **38**, 1879 (1988).
- [6] B. Pudliner, V. R. Pandharipande, J. Carlson, S. C. Pieper, and R. B. Wiringa, *Phys. Rev. C* **56**, 1720 (1997).
- [7] M. Baldo, I. Bombaci, S. Ferreira, G. Giansiracusa, and U. Lombardo, *Phys. Rev. C* **43**, 2605 (1991).
- [8] R. B. Wiringa, V. Fiks, and A. Fabrocini, *Phys. Rev. C* **38**, 1010 (1988); A. Fabrocini and S. Fantoni, *Phys. Lett. B* **298**, 263 (1993).
- [9] F. Arias de Saavedra, G. Co', A. Fabrocini, and S. Fantoni, *Nucl. Phys.* **A605**, 359 (1996); A. Fabrocini, F. Arias de Saavedra, G. Co', and P. Folgarait, *Phys. Rev. C* **57**, 1668 (1998).
- [10] F. C. Khanna, *Phys. Rev. Lett.* **20**, 871 (1968); M. Gaudin, J. Gillespie, and G. Ripka, *Nucl. Phys.* **A176**, 237 (1971); O. Bohigas and S. Stringari, *Phys. Lett.* **95B**, 9 (1980); F. Dellagiacoma, G. Orlandini, and M. Traini, *Nucl. Phys.* **A393**, 95 (1983); O. Benhar, C. Ciofi degli Atti, S. Liuti, and G. Salmè, *Phys. Lett. B* **177**, 135 (1986); M. V. Stoitsov, A. N. Antonov, and S. S. Dimitrova, *Phys. Rev. C* **47**, R455 (1993); *ibid.* **48**, 74 (1993); *Z. Phys. A* **345**, 359 (1993).
- [11] C. Giusti and F. D. Pacati, *Nucl. Phys.* **A535**, 573 (1991); C. Giusti, F. D. Pacati, and M. Radici, *ibid.* **A546**, 607 (1992); S. Boffi, C. Giusti, F. D. Pacati, and M. Radici, *ibid.* **A564**, 473 (1995); C. Giusti and F. D. Pacati *ibid.* **A571**, 694 (1994); C. Giusti and F. D. Pacati, *ibid.* **A585**, 618 (1995).
- [12] J. Ryckebusch, M. Vanderhaeghen, K. Heyde, and M. Waroquier, *Phys. Lett. B* **350**, 1 (1995); M. Vanderhaeghen, L. Machenil, J. Ryckebusch, V. Van der Sluys, and M. Waroquier, in *Proceedings of the Second Workshop on Electromagnetically Induced Two-Nucleon Emission*, edited by J. Ryckebusch and M. Waroquier (University of Gent, Gent, 1995).
- [13] G. Co', *Nuovo Cimento A* **108**, 623 (1995); F. Arias de Saavedra, G. Co', and M. M. Renis, *Phys. Rev. C* **55**, 673 (1997).
- [14] G. Co' and A. M. Lallena, *Phys. Rev. C* **57**, 145 (1998).
- [15] S. Fantoni and V. R. Pandharipande, *Nucl. Phys.* **A473**, 234 (1987); A. Fabrocini and S. Fantoni, *ibid.* **A503**, 375 (1989); A. Fabrocini, *Phys. Rev. C* **55**, 338 (1997).
- [16] S. Fantoni and A. Fabrocini, in *Lecture Notes in Physics*, Proceedings of the European Summer School on Microscopic Quantum Many Body Theories and their Applications, Valencia, 1997, edited by J. Navarro and A. Polls (Springer-Verlag, Berlin, in press).
- [17] J. E. Amaro, G. Co', and A. M. Lallena, *Int. J. Mod. Phys. E* **3**, 735 (1994).
- [18] G. Höhler, E. Pietarinen, I. Sabba-Stefanescu, F. Borkowski, G. G. Simon, V. H. Walter, and R. D. Wendling, *Nucl. Phys.* **B114**, 505 (1976).